King's Research Portal

DOI:

[10.1016/j.nbd.2018.12.011](https://doi.org/10.1016/j.nbd.2018.12.011)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Paternoster, V., Svanborg, M., Edhager, A. V., Rajkumar, A. P., Eickhardt, E. A., Pallesen, J., Grove, J., Qvist, P., Fryland, T., Wegener, G., Nyengaard, J. R., Mors, O., Palmfeldt, J., Børglum, A. D., & Christensen, J. H. (2018). Brain proteome changes in female Brd1^{+/+} mice unmask dendritic spine pathology and show enrichment for schizophrenia risk. *Neurobiology of Disease*. <https://doi.org/10.1016/j.nbd.2018.12.011>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Brain proteome changes in female Brd1+/- mice unmask dendritic spine pathology and show enrichment for schizophrenia risk

Veerle Paternoster, Maria Svanborg, Anders Valdemar Edhager, Anto P. Rajkumar, Esben Ahlburg Eickhardt, Jonatan Pallesen, Jakob Grove, Per Qvist, Tue Fryland, Gregers Wegener, Jens Randel Nyengaard, Ole Mors, Johan Palmfeldt, Anders Dupont Børglum, Jane Hvarregaard Christensen

PII: S0969-9961(18)30767-8
DOI: <https://doi.org/10.1016/j.nbd.2018.12.011>
Reference: YNBDI 4351

To appear in: *Neurobiology of Disease*

Received date: 25 April 2018
Revised date: 23 November 2018
Accepted date: 18 December 2018

Please cite this article as: Veerle Paternoster, Maria Svanborg, Anders Valdemar Edhager, Anto P. Rajkumar, Esben Ahlburg Eickhardt, Jonatan Pallesen, Jakob Grove, Per Qvist, Tue Fryland, Gregers Wegener, Jens Randel Nyengaard, Ole Mors, Johan Palmfeldt, Anders Dupont Børglum, Jane Hvarregaard Christensen, Brain proteome changes in female Brd1+/- mice unmask dendritic spine pathology and show enrichment for schizophrenia risk. *Ynbdi* (2018), <https://doi.org/10.1016/j.nbd.2018.12.011>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Brain proteome changes in female *Brdl*^{+/-} mice unmask dendritic spine pathology and show enrichment for schizophrenia risk

Veerle Paternoster, PhD^{a-d,*} veerle@biomed.au.dk, Maria Svanborg, MSc^{a-c}, Anders Valdemar Edhager, PhD^e, Anto P. Rajkumar, MD, MRCPsych, PhD^{a-c,f,g}, Esben Ahlburg Eickhardt, PhD^{a-c}, Jonatan Pallesen, PhD^{a-c}, Jakob Grove, PhD^{a-c,h}, Per Qvist, PhD^{a-c}, Tue Fryland, PhD^{a-c}, Gregers Wegener, MD, PhD^d, Jens Randel Nyengaard, PhDⁱ, Ole Mors, MD, PhD^{a,b,d}, Johan Palmfeldt, PhD^c, Anders Dupont Børghlum, MD, PhD^{a-c}, Jane Hvarregaard Christensen, PhD^{a-c}

^aThe Lundbeck Foundation Initiative for Integrative Psychiatric Research, iPSYCH, Aarhus, Denmark

^bCentre for Integrative Sequencing, iSEQ, Aarhus University, Aarhus, Denmark

^cDepartment of Biomedicine, Aarhus University, Aarhus, Denmark

^dTranslational Neuropsychiatry Unit, Department of Clinical Medicine, Aarhus University, Aarhus, Denmark

^eResearch Unit for Molecular Medicine, Aarhus University Hospital, Aarhus, Denmark

^fMental Health of Older Adults and Dementia Clinical Academic Group, South London and Maudsley NHS foundation trust, London, UK

^gDepartment of Old Age Psychiatry, Institute of Psychiatry, Psychology, & Neuroscience, King's College London, London, UK

^hBioinformatics Research Centre, BiRC, Aarhus University, Aarhus, Denmark

ⁱStereology and Electron Microscopy Laboratory, Centre for Stochastic Geometry and Advanced Bioimaging, Aarhus University Hospital, Aarhus, Denmark

*Corresponding author at: Department of Biomedicine, Wilhelm Meyers Allé 4, building 1242, room 235, 8000 Aarhus C, Denmark.

ACCEPTED MANUSCRIPT

Abstract

Genetic and molecular studies have implicated the Bromodomain containing 1 (*BRD1*) gene in the pathogenesis of schizophrenia and bipolar disorder. Accordingly, mice heterozygous for a targeted deletion of *Brd1* (*Brd1*^{+/-} mice) show behavioral phenotypes with broad translational relevance to psychiatric disorders. *BRD1* encodes a scaffold protein that affects the expression of many genes through modulation of histone acetylation. BRD1 target genes have been identified in cell lines; however the impact of reduced *Brd1* levels on the brain proteome is largely unknown. In this study, we applied label-based quantitative mass spectrometry to profile the frontal cortex, hippocampus and striatum proteome and synaptosomal proteome of female *Brd1*^{+/-} mice. We successfully quantified between 1537 and 2196 proteins and show widespread changes in protein abundancies and compartmentalization. By integrative analysis of human genetic data, we find that the differentially abundant proteins in frontal cortex and hippocampus are enriched for schizophrenia risk further linking the actions of BRD1 to psychiatric disorders. Affected proteins were further enriched for proteins involved in processes known to influence neuronal and dendritic spine morphology e.g. regulation of cytoskeleton dynamics and mitochondrial function. Directly prompted in these findings, we investigated dendritic spine morphology of pyramidal neurons in anterior cingulate cortex and found them significantly altered, including reduced size of small dendritic spines and decreased number of the mature mushroom type. Collectively, our study describes known as well as new mechanisms related to BRD1 dysfunction and its role in psychiatric disorders, and provides evidence for the molecular and cellular dysfunctions underlying altered neurosignalling and cognition in *Brd1*^{+/-} mice.

Keywords: Dendritic spine, cytoskeleton, proteomics, TMT10plex, bromodomain containing 1, animal model, schizophrenia, mitochondria.

Abbreviations:

BD: bipolar disorder; BH: Benjamini-Hochberg; DAP: differentially abundant protein; DCP: differentially compartmentalized protein; FDR: false-discovery rate; LD: linkage disequilibrium; LFC: log₂ fold change; GWAS: genome-wide association study; IAA: 2-Iodoacetamide; MS: mass spectrometry; SZ: schizophrenia; TCEP: tris(2-carboxyethyl)phosphine; FrC: frontal cortex; HC: hippocampus; ST: striatum; WT: wild type; Cell: cell lysate; Syn: synaptosome; LTP: long term potentiation; LTD: long term depression

Introduction

Repeated evidence from large genetic studies have implied that the bromodomain containing 1 (*BRD1*) gene has a role in susceptibility to schizophrenia (SZ) and possibly also to bipolar disorder (BD)¹⁻⁸. The SZ risk allele of the *BRD1* promoter SNP rs138880 and SNPs in high linkage disequilibrium (LD) have been shown to be correlated with reduced *BRD1* mRNA expression⁹ as well as increased DNA methylation of the *BRD1* promoter¹⁰, suggesting that reduced *BRD1* expression might contribute to increase SZ risk. This is further indicated, by the identification of a SZ case with a disruptive nonsense mutation in *BRD1*⁴, even though *BRD1* is otherwise highly intolerant to loss of function mutations¹¹. *BRD1*, which is a scaffold protein that interacts with histone modifiers and chromatin remodelling proteins^{12,13}, seems to be functionally interconnected with a large number of schizophrenia risk genes by means of direct protein-protein interactions¹² as well as by its genomic binding sites¹² and gene regulatory effects^{9,12,14}. The expression of *BRD1* itself is highly regulated during brain development² and stem cell differentiation¹⁵ as well as by environmental stimuli like chronic restraint stress¹⁶, electroconvulsive seizures¹⁷ and commonly used mood stabilizers¹⁰. Underlining the impact of *BRD1* on mental health, mice with a monoallelic inactivation of *Brd1* (*Brd1*^{+/-} mice) are characterized by behavioral-, neuromorphological and neurochemical changes with broad translational relevance to psychiatric disorders^{9,14,18,19}. In female *Brd1*^{+/-} mice, this includes reduced cortical serotonin, despair-like behaviors, impaired pre-attentive processing and neurocognitive deficits accompanied by aberrant morphology of pyramidal neurons and reduced dendritic spine density in anterior cingulate cortex (aCC)^{14,18}. Furthermore, genomic and transcriptomic studies have revealed that *BRD1* might act as a modulator of nuclear receptor mediated signaling, which may explain sex-biased symptom profiles^{12,14}. Although, expression profiles have been generated by RNAseq for selected brain regions to characterize molecular signatures in *Brd1*^{+/-} mice^{9,14}, the effects of *BRD1* deficiency on protein abundancies in the brain remains largely unknown. Here, we thus investigated the brain proteome in *Brd1*^{+/-} mice using quantitative mass spectrometry (MS) and labelling with isobaric tags (Tandem Mass Tags, TMT). We set out to map the proteome composition of three different brain regions: frontal cortex (FrC), hippocampus (HC) and

striatum (ST), and additionally applied synaptosomal proteomics of the same tissues to further decipher differences in protein abundance and cellular compartmentalization caused by BRD1 deficiency in the brain. Our findings are followed up by integrative analyses using human genetic data and detailed analysis of dendritic spine morphology in mice.

Materials and Methods

Animals

Brd1^{+/-} mice are heterozygous for an inactivated allele of *Brd1*⁹. Female *Brd1*^{+/-} and wild type (WT) littermates were sacrificed at age 6-8 weeks by cervical dislocation and brain tissues (frontal cortex, hippocampus and striatum) were collected by free-hand dissection, snap frozen and stored at -80°C upon further use (**Suppl. Information 1**). All studies were carried out in accordance with Danish legislation, granted by the animal welfare committee, appointed by the Danish Ministry of Food, Agriculture and Fisheries – Danish Veterinary and Food Administration.

Protein Isolation

Brain regions were homogenized using pestles from the Grinding Kit (GE Healthcare, Chicago, Illinois, USA) in core buffer as previously described²⁰ supplemented with 1 Complete Mini Protease Inhibitor Cocktail tablet (Roche, Penzberg, Germany) per 10 ml buffer. The homogenate was split in two fractions for total cell lysate and for enrichment of synaptosomes, respectively. The total cell lysate was prepared by two repeated freeze-thaw cycles, bath sonication and centrifugation at 12000 rpm for 5 min. The supernatant was isolated and stored at -20°C upon further use. The synaptosomal fraction was prepared as described before with minor adaptations²⁰. In short, the homogenate was filtered through a 5 µm membrane (Millipore, Billerica, Massachusetts, USA). Thereafter the flow-through was centrifuged to pellet the synaptosomal fraction. The frontal cortex samples for MS analysis were redissolved in core buffer. The frontal cortex (for validation), hippocampal and striatal samples were redissolved in 0.1% SDS in core buffer and sonicated at maximum intensity for 1 min. Total protein concentration of all

fractions was measured with the Fluoroprobe Protein Quantification Kit (Sigma Aldrich, St. Louis, Missouri, USA) according to manufacturer's recommendations.

Western Blotting Analysis

Protein samples were diluted to 2.5 mg/ml in reducing loading buffer (Life Technologies, Waltham, Massachusetts, USA), boiled at 95°C for 10 min and loaded on a Criterion™ TGX Stain-Free™ SDS PAGE gel (Biorad, Hercules, California, USA). After size separation, the gel was activated for 4 min under UV light. The proteins were then transferred to a low fluorescence PVDF membrane (Biorad). The membrane was blocked in 10% skimmed milk buffer (Millipore) and incubated with one of the following primary and appropriate secondary antibodies: anti-beta actin (Abcam, Cambridge, United Kingdom, #ab6276, 1:25000), anti-PSD95 (Millipore #MAB1596, 1:700), anti-VDAC1 (Abcam #ab15895, 1:25000), anti-GAP43 (Santa Cruz, Dallas, Texas, USA #SC10786, 1:1000), anti-HSP60 (Sigma Aldrich, #H3524, 1:25000), anti-NDUFA9 (Mitoscience, Eugene, Oregon, USA, #MS111, 1:25000), anti-NF200 (Sigma Aldrich, #N4142, 1:1000), anti-GFAP (Sigma Aldrich, #G3893, 1:10000), goat anti-rabbit antibody (Dako, Glostrup, Denmark, #P0448), and goat anti-mouse antibody (Dako, #P0447). The membrane was developed with Pierce ECL Plus Western Blotting Substrate (Thermo Scientific, Waltham, Massachusetts, USA) and visualized with Image Quant LAS-4000 Imager (GE Healthcare). Antibody sensitivity was tested for by checking for linear response to increased protein amounts (standard curve from a pool of all samples, four points between 12.5 and 50 µg total protein loaded). Protein abundances of samples within an extrapolated range of the standard curve (± 7 µg of the most extreme points from the standard curve) were calculated based on the standard curve and normalized to total protein intensity of transferred proteins (minimum 3 samples per group).

Sample Preparation for Mass Spectrometry

Protein samples were prepared for labelling by TMT10plex Mass Tag Labelling (Thermo Scientific) according to the manufacturer's protocol. In short, 100 µg of total protein was reduced and alkylated with tris(2-carboxyethyl)phosphine (TCEP) and 2-Iodoacetamide (IAA), followed by overnight acetone

precipitation. After redissolving the pellet, proteins were digested overnight with trypsin (Promega, Fitchburg, Wisconsin, USA). Resulting peptides were labelled with the TMT10plex Mass tags and the 10 samples were pooled. Labelled peptide pools were dissolved in 0.1% formic acid pH 2.3, before being fractionated into 8 fractions using Oasis MCX μ Elution (Waters Corporation, Milford, Connecticut, USA). Peptides were stepwise eluted in seven 50 mM ammonium formate buffers, containing 15% (first eluate) or 35% (all other eluates) methanol and with acidity ranging between pH 2.6 and pH 9. The final fraction was eluted with a buffer containing 35% methanol and 5% NH_4OH .

NanoLC-MS/MS

Liquid chromatography tandem mass spectrometry (LC-MS/MS) was performed on an EASY nanoLC-1000 coupled to Q Exactive™ Plus Hybrid Quadrupole-Orbitrap™ Mass Spectrometer (Thermo Scientific). TMT tagged peptide samples were trapped on a pre-column (PepMap 100, C18, 2 cm, 75 μm i.d., 3 μm , 100 Å, Thermo Scientific) followed by separation on a C18 reverse phase column with integrated emitter (EASY-Spray column, Plug and spray, PepMap 25 cm, 75 μm i.d., 2 μm , 100 Å, Thermo Scientific). Separation was conducted using Buffer B containing 90/10 acetonitrile/water and 0.1% formic acid at a flowrate of 300 nL/min in a 90 minute linear gradient (4 to 40% Buffer B). The MS was operated in a positive data dependent mode, automatically switching between MS1 (full scan) and MS2 (fragmentation scan) acquisition. Resolution of MS1 was set to 70000 and MS2 to at least 35000. In MS1, AGC target was set to 1×10^6 ions and scan range between 380-1800 m/z. In MS2, AGC target was set at 2×10^5 ions, with fixed first mass set to 120 m/z. Dynamic exclusion was set to 20 sec, isolation window at 2.0 m/z, and up to ten of the most intense ions were fragmented after every MS1 scan, by higher-energy C-trap dissociation (HCD). Ions with single charge or unassigned charge states were excluded from fragmentation.

LC-MS/MS analyses were performed three times for each peptide sample, with the third analysis applying an exclusion list of the 100 most intense peptide peaks. All three LC-MS analyses of the same sample set were merged and submitted to database searches for protein identification and quantification.

Data Analysis

Peptide identification was performed in Proteome Discoverer 1.4 (Thermo Scientific) using the Mascot (Matrix Science, Boston, Massachusetts, USA) database algorithm and the Percolator algorithm²¹, with UniProt Knowledgebase release 2014_10 (Number of sequences after taxonomy (*Mus musculus*): 16 693) as reference proteome. The following settings were applied: total intensity threshold of 20000; precursor mass tolerance of 7 ppm; fragment mass tolerance of 10 mmu; peak integration tolerance of 10 ppm; minimum MS/MS peak count of 20; maximum 2 missed trypsin cleavages allowed. Only protein-unique peptides with FDR <0.01 and co-isolation <40% were included in the analysis. Proteins with at least 2 unique peptides and minimum 3 scans were considered for quantification. Raw peptide intensities were normalized using LOESS regression (0.4 window width) in the InfernoRDN software²². Protein abundances were then obtained by the RRollup algorithm²². Differentially abundant proteins (DAPs) were identified by PLGEM^{23,24}. Log2 fold change (LFC) cut-off was set as two times the global standard error of the mean of the dataset. Proteins were considered differential abundant between *Brd1*^{+/-} and WT mice if both $p < 0.05$ and $LFC > \text{cut-off value}$. Proteins were considered as 'top DAPs' when $p < 0.001$ and $LFC > \text{cut-off}$. To identify differentially compartmentalized proteins (DCPs), the synaptosome-to-cell lysate abundancy ratio was calculated for all proteins detected in both datasets from the same brain region. Ratios were compared between genotypes using PLGEM. Proteins are considered DCPs when $p < 0.05$.

Mitochondrial proteins were selected according to Pagliarini *et al.* with a cut-off threshold corrected false discovery rate (cFDR) set at 7% according to previously reported research²⁵ and to MitoMiner with positive score in at least 4 of the evidence lines²⁶. Mouse protein IDs were converted to the human homologue Ensembl Gene ID for bioinformatic analysis using the db2db online tool²⁷. DAVID bioinformatics tools were used for Functional Annotation Clustering²⁸ with standard settings and using all identified proteins of the specific brain region as background. Protein-protein interaction network analysis was performed using the STRING database employing standard settings²⁹. Enrichment analysis

for transcription factor binding was performed using Enrich³⁰ (ENCODE and ChEA Consensus TFs from ChIP-X) using default settings. Risk gene enrichment analysis was performed using MAGMA³¹ with default settings and using all identified proteins of the specific brain region as background. The analysis was based in summary statistics from single-marker genome-wide association studies (GWASs)^{8,32–38} (excluding the broad MHC region (chr6: 25 M-35 M)) and filtered for info score ≥ 0.8 if available. Genes were annotated using Ensemble (GRCh38.p12). Information about linkage disequilibrium was obtained using the 1000 Genomes European panel³⁹.

Availability of data

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD009688; PXD009689; PXD009690; PXD009691; PXD009692; PXD009961.

Dendritic Spine Morphology

Dendritic spine morphology analysis was performed using the Golgi-cox stained brain slices derived from an independent set of mice and 3D reconstructed pyramidal neurons described previously (**Suppl. Information 1**)¹⁴. Identification and analysis of dendritic spines was done with IMARIS 7.7.1 software. Classification of spine types was done with following settings: Stubby (length < 1 μm), Mushroom (length < 3 μm and max diameter (head) > 2x mean width (neck)), Long Thin (mean width (head) $\geq$ mean width (neck)), and Filopodia / Dendrite (rest).

Continuous parameters were analyzed using a linear mixed-effects model (lmer function in the R package lme4⁴⁰), binary parameters were analyzed using a generalized linear mixed-effects model (glmer function in the R package lme4⁴⁰) as previously described⁴¹. All models included spine level and depth (as defined by IMARIS) and genotype as explanatory variables, and neuron and mouse as level identifiers. Significance testing of contribution of parameters was conducted using the lmerTest package in R⁴⁰ and BH correction was applied to correct for multiple testing.

Results

BRD1 deficiency is associated with widespread changes in protein abundancies in brain

To discover novel BRD1 dependent molecular changes in the brain, we profiled differentially abundant proteins (DAPs) in three brain regions which show differential neurochemistry and mRNA expression levels and which are involved in observed behavioral changes in female *Brd1*^{+/-} mice^{14,18}: frontal cortex (FrC), hippocampus (HC) and striatum (ST). By subjecting total cell lysates (Cell) and extracts enriched for synaptosomes (Syn) to MS-based quantitative proteomics using TMT10plex (**Figure 1**) we obtained six protein datasets (FrC-cell, FrC-syn, HC-cell, HC-syn, ST-cell, and ST-syn) (**Suppl. Information 2, Suppl. Figure 2**). All datasets were checked for gross differences in cell composition using cell type specific markers and one sample from FrC-Syn was removed from further analysis due to an apparent diverging proportion of oligodendrocytes (**Suppl. Information 2, Suppl. Figure 2**). We observed a shift in the mitochondrial subproteome (69/234 (29%) detected mitochondrial proteins are identified as DAPs, of which 66 are downregulated in *Brd1*^{+/-} mice compared to WT) in the FrC-syn dataset, which was subsequently removed to avoid bias in normalization (**Suppl. Information 3, Suppl. Figure 4**). Overall, we detected 1880 unique proteins in FrC (both datasets combined), 2370 unique proteins in HC (both datasets combined) and 1978 unique proteins in ST (both datasets combined) (**Suppl. Table 2**). Because no protein remained significant after BH-correction for multiple testing in any of the datasets, we classified 16 proteins as ‘top DAPs’ (**Figure 2** and **Table 1**). These top DAPs could be categorized in six groups based on function (**Figure 2** and **Table 1**): Transcription & chromatin (6 top DAPs), neuronal morphology and growth (4 top DAPs), cellular transport (2 top DAPs), neuroendocrine (2 top DAPs), and other (2 top DAPs). We selected 4 proteins from frontal cortex for validation by Western blotting analysis in an independent set of samples: 3 mitochondrial proteins (VDAC1, HSP60 and NDUFA9) and GAP43. After exclusion of HSP60 for technical reasons, we were able to confirm the differential abundancies of 2 out of the 3 remaining proteins (NDUFA9 and GAP43) (**Suppl. Information 4**). Additionally, we found significantly less PSD95 in *Brd1*^{+/-} mice in FrC-syn (LFC=-0.10, p=0.0480). This is in line with

previously observed reduced dendritic spine density in anterior cingulate cortex (aCC),¹⁴ thereby further validating our findings.

Figure 1: Overview of the sample preparations and MS pipeline. Abbreviations: wild type (wt); female heterozygous *Brd1* knockout mice (*Brd1*^{+/-}); frontal cortex (FrC); hippocampus (HC); striatum (ST); cell lysate (cell); synaptosome (syn); mass spectrometry (MS); mixed-mode cationic exchange (MCX); nano liquid chromatography – tandem mass spectrometry (nanoLC-MS/MS); quality control (QC); not significant (ns); differentially abundant proteins (DAPs); top differentially abundant proteins (top DAPs).

Figure 2: A) Total number of unique proteins detected in female *Brd1*^{+/-} mouse brain regions. Differentially abundant proteins (DAPs, $p < 0.05$ and $LFC > \text{cut-off}$); top differentially abundant proteins (Top DAPs $p < 0.001$ and $LFC > \text{cut-off}$). LFC cut-off: ± 0.231 (FrC-Cell), ± 0.104 (FrC-Syn), ± 0.099 (FrC-Syn (excl mito)), ± 0.191 (HC-Cell), ± 0.072 (HC-Syn), ± 0.189 (ST-Cell), ± 0.102 (ST-Syn). B) Total number of unique proteins shared between cell lysate and synaptosomal preparations of the same brain

region and differentially compartmentalized proteins (DCPs, $p < 0.05$). **C)** Distribution of the proteins among fractions and statistical significant groups. Abbreviations: frontal cortex (FrC); hippocampus (HC); striatum (ST); cell lysate (cell); synaptosome (syn); not significant (ns).

Table 1: Overview of the top differential abundant proteins (top DAPs) detected in female *Brd1*^{+/-} mouse brain.

UniProt Name ¹	Entry	Gene Symbol	UniProt Protein Name	P- value	LF C	Datas et
Transcription & chromatin						
TB182_MOUSE		Tnks1bp1	182 kDa tankyrase-1-binding protein	0.0003	0.70	Frc- cell
TMA7_MOUSE		Tma7	Translation machinery-associated protein 7	0.0003	0.93	Frc- cell
TCAL5_MOUSE		Tceal5	Transcription elongation factor A protein- like 5	0.0004	1.05	Frc- cell
HMGN1_MOUSE		Hmgn1	Non-histone chromosomal protein HMG- 14	0.0005	0.82	Frc- cell
T22D1_MOUSE		Tsc22d1	TSC22 domain family protein 1	0.0005	0.92	Frc- cell
CNBP_MOUSE		Cnbp	Cellular nucleic acid-binding protein	0.0010	0.75	Frc- cell
Cellular transport						
SYNRG_MOUSE		Synrg	Synergina gamma	0.0008	0.79	Frc- cell
MON2_MOUSE		Mon2	Protein MON2 homolog	0.0006	0.53	HC- syn
Neuroendocrine						
CMGA_MOUSE		Chga	Chromogranin-A	0.0004	1.05	Frc- cell

PENK_MOUSE	Penk	Proenkephalin-A	0.0004	0.85	Frc-cell
Neuronal morphology and growth					
OGFR_MOUSE	Ogfr	Opioid growth factor receptor	0.0005	0.93	Frc-cell
S10A5_MOUSE	S100a5	Protein S100-A5	0.0007	- 0.67	Frc-syn
MT3_MOUSE	Mt3	Metallothionein-3	0.0008	1.07	Frc-cell
NEUM_MOUSE	Gap43	Neuromodulin	0.0002	0.47	Frc-syn
Other					
PTMA_MOUSE	Ptma	Prothymosin alpha	0.0005	1.12	Frc-cell
PRC2C_MOUSE	Prrc2c	Protein PRRC2C	0.0007	0.56	Frc-cell

1. The listed proteins passed the criteria of both $p < 0.001$ and $\log_2$ fold change ($LFC > \text{cut-off}$).
LFC: $\log_2$ fold change compared to WT mice. Abbreviations: frontal cortex (FrC); hippocampus (HC); striatum (ST); cell lysate (cell); synaptosome (syn).

DAPs are interconnected and cluster in functions related to mitochondria and the cytoskeleton

To inform about a potential biological role of the DAPs, we first performed a protein-protein interaction (PPI) enrichment score analysis using the STRING algorithm (**Suppl. Information 5**). DAPs in all brain regions, except HC-cell were significantly enriched for PPIs, with enrichment ratios ranging from 3.31 (FrC-cell) to 26.14 (ST-cell), meaning that DAPs within these datasets are at least partly biologically interconnected. The lack of significant results among DAPs from HC-cell is likely due to the lack of power as a result of the low number of DAPs ($n = 18$) in this dataset.

Next, in order to gain more insight into associated functions, we performed DAVID Functional Annotation Clustering analysis. We found eight clusters significantly enriched in ST-cell, five in FrC-cell and three in ST-syn (**Suppl. Table 5**). Five of the ST-cell clusters are related to mitochondria (Enrichment Scores between 10.42 and 2.1). Two of the FrC-cell cluster are related to the proteasome (Enrichment Scores 4.5 and 1.7). Among the other enriched cluster are calmodulin-binding, microtubule, and synapse.

Subsequently, in order to investigate the underlying mechanism and cascades that may explain the observed DAPs, we tested whether the genes encoding DAPs were enriched for transcription factor binding. We found enrichment for 33 unique transcription factors (**Suppl. Table 6**). Six of the transcription factors are enriched more than once: TAF1 (FrC-cell and HC-syn), ATF2 (FrC-cell and HC-syn), CEBPD (FrC-cell and HC-syn), NELFE (FrC-cell and HC-syn), PBX3 (FrC-cell and HC-syn) and REST (FrC-cell, ST-cell and ST-syn).

DAPs are enriched for schizophrenia risk

We hypothesized that DAPs, like BRD1 target genes and differentially expressed genes in mouse brains^{9,12,14}, are enriched for psychiatric disorder risk genes. Because the strongest evidence for genetic association is found between SZ and *BRD1*, we first tested whether DAPs are enriched for SZ risk using MAGMA³¹. We found that DAPs from FrC-cell and HC-cell are significantly enriched for SZ risk ($p_{\text{FrC-cell}}=0.049$ and $p_{\text{HC-cell}}=2.6\text{E-}5$) (**Figure 3**). Next, we investigated the enrichment of for mental health risk genes among DAPs, testing 7 other GWAS datasets, comprising 5 psychiatric disorders and 2 unrelated phenotypes (**Suppl. Information 6**).

Figure 3: MAGMA disease risk enrichment analysis for schizophrenia. Black dashed line: significance threshold ($p=0.05$). Abbreviations: frontal cortex (FrC); hippocampus (HC); striatum (ST); cell lysate (cell); synaptosome (syn).

BRD1 deficiency is associated with differential protein compartmentalization

Next we wondered whether we could identify differences in protein abundance between the cell lysate and synaptosomal preparations, indicating that certain proteins are affected by processes such as trafficking or local protein degradation resulting in differential compartmentalization. To do so, we calculated the synaptosome-to-cell lysate abundance ratio (syn/cell) for all proteins detected in both datasets from the same region ($n_{\text{FrC}}=1054$, $n_{\text{HC}}=1586$, $n_{\text{ST}}=1308$) and compared these ratios between genotypes (**Figure 2B and 2C**). We identified 48, 5 and 104 differentially compartmentalized proteins (DCPs) in FrC, HC and ST, respectively, when comparing genotypes (**Suppl. Table 7**). The vast majority of these (40/48 in FrC, 3/5 in HC and 99/104 in ST) of these have a negative LFC, indicating that these proteins are relatively less abundant in synaptosomal preparations compared to cell lysate in $\text{Brd1}^{+/-}$ mice. In line with this finding, several of the DCPs had been identified as DAPs (**Figure 2C**). In FrC, 39 DCPs were also DAPs (37 upregulated) in cell lysate, and 7 were DAPs (5 downregulated) in synaptosome. Among the 5 DCPs in HC, we identified 1 DAP (upregulated) from cell lysate and 3 DAPs (1 downregulated) from synaptosome. In ST, 56 DAPs (54 upregulated) from cell lysate were among the DCPs, and 11 (6 downregulated) from synaptosome. We then used DAVID Functional Annotation

Clustering analysis to identify the function of DCPs. This analysis revealed 1 cluster (enrichment score = 2.0) related to calcium signaling among DCPs in FrC, none in HC and 10 enriched clusters among DCPs in ST (enrichment scores between 1.4 and 13.7). Eight out of ten DCP clusters in ST were related to mitochondria.

BRD1 deficiency is associated with changes in dendritic spine morphology

We note that recurrent results point in the direction of aberrant neuromorphology in *Brd1*^{+/-} mice: 1) Four top DAPs (OGFR, S100A5, MT3, GAP43) in this study have a function related to neuronal morphology and growth 2) DAVID cluster analysis of DAPs showed “microtubule” among the enriched clusters for the FrC-cell dataset, including DAPs like Tau, Microtubule-associated Protein 4 (MAP4) and Microtubule-associated protein 2 (MAP2); microtubule dynamics have been shown to influence dendritic spine morphology^{42,43} and 3) We have previously reported altered morphology of pyramidal neurons in the aCC of female *Brd1*^{+/-} mice¹⁴. We hypothesized here that dendritic spine morphology in addition is altered in *Brd1*^{+/-} mouse brain, and performed a morphology analysis of dendritic spines derived from aCC pyramidal neurons.

We first classified the spines into 4 classes (stubby, mushroom, long thin and filopodia) and observed an overall difference in distribution between genotypes (Chi square test, $p < 0.001$) (**Figure 4A**). This overall difference seems to be driven by a loss of mushroom types (-5.2%) and a gain of stubby type spines (+4.3%) in *Brd1*^{+/-} mice. To determine the underlying differences between genotypes that cause this class shift, we measured spine length, area, volume, minimum and maximum diameter for the whole dendritic spine as well as for the subparts spine head, neck and ground (the subparts not for spine area) from the same dendritic spines. After correcting for multiple testing, we found that dendritic spines from *Brd1*^{+/-} mice were smaller with respect to minimum (total spine, $p_{\text{adjusted}} < 0.0001$, $\beta_{\log\text{odds}} = -28.8 \pm 2.1$) and maximum (total spine, $p_{\text{adjusted}} = 0.01216$, $\beta_{\log\text{odds}} = -4.8 \pm 1.5$) diameter for the total spine, spine head, neck and ground (**Figure 4B, 4C**, and **Suppl. table 8**). The observed differences in maximum diameter and maximum ground diameter was found in the smaller subpopulation (spines with diameter ≤ 234.0

nm), but not in the larger subpopulation (**Suppl. table 8**). Additionally, spines with length $< 1.2 \mu\text{m}$ were significantly smaller in *Brd1*^{+/-} mice ($p_{\text{adjusted}} = 0.01391$, $\beta = -48.3 \pm 16.6 \text{ nm}$) (**Figure 4D**). **Figure 4E** illustrates and summarizes the effects of reduced *Brd1* levels on pyramidal neuron spine morphology in aCC.

A

B

C

D

E

Figure 4: Analysis of dendritic spine morphology in aCC pyramidal neurons. A) Relative distribution of 4 dendritic spine types. B) Comparison of relative frequency per genotype of maximum diameter of dendritic spines per category (category 1: maximum spine diameter < 117.0 nm, category 2: $117.0 \text{ nm} \leq \text{maximum spine diameter}$) between wild type (n=27182) and *Brd1*^{+/-} (n=11820) spines. P-value as determined using a generalized linear mixed-effect model. C) Comparison of relative frequency of minimum diameter of dendritic spines per category (category 1: minimum spine diameter < 117.0 nm, category 2: $117.0 \text{ nm} \leq \text{minimum spine diameter} \leq 234.0 \text{ nm}$) between wild type and *Brd1*^{+/-} dendritic spines. P-value as determined using a generalized linear mixed-effect model. D) Comparison of the cumulative frequency of spine length between wild type and *Brd1*^{+/-} spines. P-value as determined using a linear mixed-effect model. Small spines are defined as spines with spine length < 1.2 μm . E) Overview of measured parameters and comparison between dendritic spines derived from wild type and *Brd1*^{+/-} mice. Dotted line represents parameters with a significant difference between genotypes. Ns: not significant, * $p < 0.05$, **** $p < 0.001$.

Discussion

In this study, we use proteomic profiling to demonstrate widespread changes in protein abundancies and compartmentalization in brain regions of female *Brd1*^{+/-} mice. We found that affected proteins are highly interconnected and they cluster in functions related to transcription and chromatin, in line with the putative role of BRD1 as a scaffold protein in HAT complexes^{12,13}, as well as mitochondria and the cytoskeleton. Directly based on these findings, we studied dendritic spines in aCC and found evidence of aberrant morphology. Upstream of DAPs, we found enrichment for several transcription factors. In line with the current and previous findings that proteins important for mitochondrial function are regulated upon reduced BRD1 levels in mouse brain regions¹⁴, the PPARG transcription factor is involved in the nuclear control of proteins important for mitochondrial function. Of specific interest with respect to previous findings are CREB1 and SMAD4, the latter found in complexes with among others c-FOS. In male *Brd1*^{+/-} mice, CREB has been identified as an likely upstream regulator of differentially expressed genes in striatum and aCC and has been found hyperphosphorylated in whole brain extracts⁹. Its transcriptional target, c-FOS, was furthermore significantly more abundant in the same samples⁹. Of special note, we found enrichment for schizophrenia risk among identified DAPs in frontal cortex and hippocampus further supporting the functional link between BRD1 and schizophrenia risk genes^{9,12,14}. In close analogy to the experimental setup in the present study, review-analyses of proteomic studies of *post-mortem* brain tissue from patients with SZ and BD have further revealed that, indifferent for the used platform or studied brain region, mitochondrial and cytoskeletal components are consistently dysregulated⁴⁴⁻⁴⁶.

Previously we have shown that the reduced expression of *Brd1* leads to phenotypes with translational relevance to SZ in mice, including many aspects from positive, negative and cognitive symptoms^{9,14,18}. In the present study, we profiled BRD1 dependent changes in protein abundancies and compartmentalization in mouse brain to further decipher the potential impact of its deficiency. Proteomic profiles represent the true state of the cell better than transcriptomic profiles, as they assess the final sum of protein synthesis,

degradation and localization. On the other hand, researchers in proteomics have to compromise between sensitivity and risk for false positives, therefore tending to be rather conservative, and face limited coverage of low abundant proteins⁴⁷ and limited sample size in the case of quantification based on isobaric labelling. To deal with low coverage, we employed the combination of distinct sample preparation, each enriching for proteins from different cell compartments, as well as repeated measurement of the same samples with different LC-MS/MS settings. To avoid false positives, we employed a rather conservative approach during QC, only including MS scans, peptides and proteins with sufficient evidence in the final datasets.

Both cytoskeleton dynamics and mitochondria are known to influence neuronal and dendritic spine morphology. Firstly, spine morphology is regulated tightly by both the F-actin and microtubule network. F-actin is mostly located within the spine, whereas microtubule provide a dynamic support for transport both long-distance and activity-dependent transport of cargoes in and out the spine⁴³. Inadequate functioning of the cytoskeleton dynamics could lead to aberrant spine morphology and to disease. For example, drug-induced destabilization of microtubule modulated spine morphology from the mature mushroom-type into the immature filopodia-type⁴⁸, knockout of PAR1B in mice reduced the number of mature spines and induced spatial memory formation⁴⁹. Furthermore, decreased immunoreactivity of MAP2 has been repeatedly shown in *post-mortem* brains from SZ patients and is associated with reduced spine density⁵⁰.

Secondly, besides their well-known function in energy metabolism, calcium homeostasis, production of reactive oxygen species (ROS) and apoptosis, mitochondria play an essential role in regulating neurogenesis and neuronal plasticity⁵¹. Disruption of mitochondrial protein translation in *Drosophila* neurons, reduced dendritic arborization, but axon morphology remained relatively unchanged⁵², which is similar to the aberrant pyramidal neuronal morphology observed in female *Brdl*^{+/-} mice¹⁴. The differences in fold change ratios of mitochondrial proteins between cell lysate and synaptosome purification from the same brain region that we find in the present study could be indicative for altered

mitochondrial morphology in female *Brdl*^{+/-} mice, as the filtration steps during the synaptosome preparation favor the extraction of smaller mitochondria. Interestingly, presynaptic mitochondrial morphology has been shown to correlate with impaired working memory in monkey brain⁵³. Moreover, the mitochondrial apoptotic pathway can be moderately and locally activated to induce LTD, resulting in shrinkage and loss of dendritic spines in the absence of cell death^{54,55}. This system as well as mitochondria morphological changes are sensitive to sublethal mitochondrial stress^{55,56}, which can be induced by excess of Ca²⁺ through disinhibition of glutamate release as a result of GABAergic dysfunction or by loss of Ca²⁺ buffering capacity through loss of parvalbumin⁵⁷, both observed in *Brdl*^{+/-} mice⁹. Finally, mitochondrial function has been shown to be linked to synaptic plasticity, cell resilience and psychiatric phenotypes^{58,59}. Mitochondrial genes are in fact significantly enriched among the 108 schizophrenia genome-wide associated loci⁶⁰ and psychiatric disorders are comorbid to mitochondrial dysfunction and can be diagnosed years before any symptoms of mitochondrial disease^{61,62}.

When we investigated dendritic spine morphology in aCC from female *Brdl*^{+/-} mice, we found that reduced *Brdl* levels coincide with a loss of the mature mushroom-type and an increase of the immature stubby spines⁶³. Furthermore, dendritic spines in *Brdl*^{+/-} mice are generally smaller, and this is especially pronounced in the small spines subgroup. Whereas the spine head size is directly correlated to the post synaptic density size, presynaptic vesicle amount⁶⁴ and postsynaptic receptor abundance^{65,66}, reduced neck and ground diameter have been shown to greatly limit the signal transduction and diffusion rates of molecules into the dendritic shaft^{65,67}. Under normal circumstances, small spines with thin necks are more likely to limit efflux from the spine head and, hereby making them more optimized for selective induction of long term potentiation (LTP)⁶⁵, and large spines are more resistant to LTP hereby securing memory formation⁶⁸. On the other hand, Ca²⁺ spread from the dendritic spine into the dendritic shaft and neighboring spines contributes to the potentiation and stabilization of co-active spines during development of the brain⁶⁹. Similarly, in the adult brain Rho GTPases can diffuse to neighboring spines and change the conditions for LTP or long term depression (LTD) locally, greatly influencing the

establishment and maintenance of neuronal networks^{70,71}. The smaller spines could therefore suggest memory formation impairments as observed in both male and female *Brdl*^{+/-} mice^{9,14,18}.

Conclusion

In summary, we show that reduced BRD1 levels have widespread effects on protein abundancies and compartmentalization in mouse brain regions. Further linking the actions of BRD1 to psychiatric disorders, we find that affected proteins in frontal cortex and hippocampus are enriched for schizophrenia risk. Our study also principally corroborate previous indications from RNAseq studies in *Brdl*^{+/-} mice that reduced *Brdl* expression in mouse brain might disturb mitochondrial function and cytoskeletal organization, both of which have been shown to play a role in neuromorphology, as well as density and morphology of spines. Lastly, we show that female *Brdl*^{+/-} mice display reduced dendritic spine size, which is likely to affect memory formation and inhibit clustered spine plasticity, affecting brain network formation and information processing. Future studies using for example two photon microscopy investigating live spine response to stimuli and the effect of malformed spines on spine crosstalk could supplement our data and help us identify the role of BRD1 in these brain processes.

Legends Suppl. Tables (online)

Suppl. Table 1: GO cellular component annotation for unique proteins from synaptosomal and cell lysate preparations. Green: $p < 0.05$.

Suppl. Table 2: Overview of all proteins per brain region and purification method. Statistical significance according to PLGEM^{23,24}. Marked in red: $p < 0.05$. Marked in green: $LFC > \text{cut-off}$. Marked in orange: $LFC < -\text{cut-off}$.

Suppl. Table 3-4: see suppl. Information

Suppl. Table 5: Overview of DAVID Annotation Cluster analysis. Disease, function and keyword and other annotations are included according to standard settings of DAVID Annotation Clustering. A Cluster is defined as a group of terms having similar biological meaning due to sharing similar gene or protein members. Statistical enrichment is calculated by the EASE Score, a modified Fisher Exact P-value. The Group Enrichment Score is calculated as the geometric mean (in $-\log_{10}$ scale) of member's P-values in a corresponding annotation cluster. List Total = number of genes in the gene list. Population (Pop) Hits = number of genes in the total group of genes assayed that belong to the specific Gene Category. Population (Pop) Total = number of genes in the total group of genes assayed that belong to any Gene Category within the System.

Suppl. Table 6: Overview of enrichment for transcription factor binding. P-values are calculated by Fisher's exact test and corrected for multiple testing by the BH method.

Suppl. Table 7: Overview of DCPs per brain region. Synaptosomal-to-cell lysate abundant ratios (syn/cell) are calculated for each sample and compared between genotypes using PLGEM. DCPs are identified when $p < 0.05$ and $LFC < 0$. Marked in red: $p < 0.05$. Marked in green: $LFC > \text{cut-off}$. Marked in orange: $LFC < -\text{cut-off}$.

Suppl. Table 8: Overview of dendritic spine morphology parameter calculations.

Acknowledgements

This study was funded by the Lundbeck Foundation, Denmark (grant number R155-2014-1724) (ADB), The Novo Nordisk Foundation (ADB & JHC), The Danish Council for Independent Research–Medical Sciences (ADB and JHC), Aarhus University (ADB) and John and Birthe Meyer Foundation (JhP). Centre for Stochastic Geometry and Advanced Bioimaging is supported by Villum Foundation. The funders had no involvement in any aspect of the study.

We thank Anne Hedemand (Aarhus University) and Tanja Østergaard (Aarhus University) for genotyping of mice, Margrethe Kjeldsen (Aarhus University Hospital) and Tanja Østergaard (Aarhus University) for assistance with Western Blotting and Abdel-Rahman Al-Absi (Aarhus University Hospital) for assistance with dendritic spine morphology analysis.

Contributions

VP, MS, APR, GW, JRN, JhP, ADB, JHC designed and directed experiments. VP, MS, APR collected samples and data. VP, MS, AVE, EAE, JhP analysed mass spectrometry results. VP, MS, APR, JG, JhP, ADB, JHC performed statistical analysis. JhP performed MAGMA analysis. VP, APR, JG, GW, JRN, JHC designed, performed and analysed spine morphology experiments. VP, MS, PQ, TF, JhP, ADB, JHC interpreted results. VP wrote the first draft of the manuscript. All authors contributed to the finalization of the manuscript.

Conflict of interest

OM and ADB are co-inventors on a patent application submitted by Aarhus University entitled “Method for diagnosis and treatment of a mental disease” (EP2287340) that includes claims relating to BRD1 among other genes. OM, ADB, JHC, APR, and PQ are coinventors on a patent application submitted by Capnova A/S entitled “Genetically modified non-human mammal and uses thereof” (PCT/EP2013/069524) that includes the *Brd1*^{+/-} mouse.

The other authors declare no conflict of interest.

References

- 1 Nyegaard M, Severinsen JE, Als TD, Hedemand A, Straarup S, Nordentoft M *et al.* Support of association between BRD1 and both schizophrenia and bipolar affective disorder. *Am J Med Genet Part B Neuropsychiatr Genet* 2010; **153**: 582–591.
- 2 Severinsen JE, Bjarkam CR, Kiaer-Larsen S, Olsen IM, Nielsen MM, Blechingsberg J *et al.* Evidence implicating BRD1 with brain development and susceptibility to both schizophrenia and bipolar affective disorder. *Mol Psychiatry* 2006; **11**: 1126–1138.
- 3 Aberg K a, Liu Y, Bukszar J, McClay JL, Khachane AN, Andreassen O a *et al.* A comprehensive family-based replication study of schizophrenia genes. *JAMA psychiatry* 2013; **70**: 573–81.
- 4 Purcell SM, Moran JL, Fromer M, Ruderfer D, Solovieff N, Roussos P *et al.* A polygenic burden of rare disruptive mutations in schizophrenia. *Nature* 2014; **506**: 185–90.
- 5 Jorgensen TH, Børghlum a. D, Mors O, Wang a. G, Pinaud M, Flint TJ *et al.* Search for common haplotypes on chromosome 22q in patients with schizophrenia or bipolar disorder from the Faroe Islands. *Am J Med Genet - Neuropsychiatr Genet* 2002; **114**: 245–52.
- 6 Andreassen OA, Thompson WK, Dale AM. Boosting the power of schizophrenia genetics by leveraging new statistical tools. *Schizophr Bull* 2014; **40**: 13–7.
- 7 Pardiñas AF, Holmans P, Pocklington AJ, Escott-Price V, Ripke S, Carrera N *et al.* Common schizophrenia alleles are enriched in mutation-intolerant genes and maintained by background selection. 2016. doi:10.1101/068593.
- 8 Ripke S, Neale BM, Corvin A, Walters JTR, Farh K-H, Holmans PA *et al.* Biological insights from 108 schizophrenia-associated genetic loci. *Nature* 2014; **511**: 421–427.
- 9 Qvist P, Christensen JH, Vardya I, Rajkumar AP, Mørk A, Patemoster V *et al.* The Schizophrenia-Associated BRD1 Gene Regulates Behavior, Neurotransmission, and Expression of Schizophrenia Risk Enriched Gene Sets in Mice. *Biol Psychiatry* 2016. doi:10.1016/j.biopsych.2016.08.037.
- 10 Dyrvig M, Qvist P, Lichota J, Larsen K, Nyegaard M, Børghlum AD *et al.* DNA Methylation Analysis of BRD1 Promoter Regions and the Schizophrenia rs138880 Risk Allele. *PLoS One* 2017; **12**: e0170121.
- 11 Lek M, Karczewski KJ, Minikel E V., Samocha KE, Banks E, Fennell T *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 2016; **536**: 285–291.
- 12 Fryland T, Christensen JH, Pallesen J, Mattheisen M, Palmfeldt J, Bak M *et al.* Identification of the BRD1 interaction network and its impact on mental disorder risk. *Genome Med* 2016; **8**: 53.
- 13 Mishima Y, Miyagi S, Saraya A, Negishi M, Endoh M, Endo T a. *et al.* The Hbo1-Brd1/Brpf2 complex is responsible for global acetylation of H3K14 and required for fetal liver erythropoiesis. *Blood* 2011; **118**: 2443–2453.
- 14 Rajkumar AP, Qvist P, Larsen SH, Lazarus R, Pallesen J, Nava N *et al.* The behavioral and neurobiological effects of reduced Brd1 expression in mice are sex-biased and implicate gender dependent dysregulation of nuclear receptor mediated signaling in mental disorders. *bioRxiv* 2018; : 257170.
- 15 Cho HI, Kim MS, Jang YK. The BRPF2/BRD1-MOZ complex is involved in retinoic acid-induced differentiation of embryonic stem cells. *Exp Cell Res* 2016; **346**. doi:10.1016/j.yexcr.2016.05.022.
- 16 Christensen JH, Elfving B, Müller HK, Fryland T, Nyegaard M, Corydon TJ *et al.* The Schizophrenia and Bipolar Disorder associated BRD1 gene is regulated upon chronic restraint stress. *Eur Neuropsychopharmacol* 2012; **22**: 651–656.
- 17 Fryland T, Elfving B, Christensen JH, Mors O, Wegener G, Børghlum AD. Electroconvulsive seizures regulates the Brd1 gene in the frontal cortex and hippocampus of the adult rat. *Neurosci Lett* 2012; **516**: 110–113.
- 18 Qvist P, Rajkumar AP, Redrobe JP, Nyegaard M, Christensen JH, Mors O *et al.* Mice heterozygous for an inactivated allele of the schizophrenia associated Brd1 gene display selective cognitive deficits with translational relevance to schizophrenia. *Neurobiol Learn Mem* 2017; **141**: 44–52.
- 19 Qvist P, Eskildsen SF, Hansen B, Baragji M, Ringgaard S, Roovers J *et al.* Brain volumetric alterations accompanied with loss of striatal medium-sized spiny neurons and cortical parvalbumin expressing interneurons in Brd1 +/- mice. *Sci Rep* 2018; **8**: 16486.
- 20 Chang JW, Arnold MM, Rozenbaum A, Caputo A, Schweizer FE, Huynh M *et al.* Synaptoneurosome micromethod for fractionation of mouse and human brain, and primary neuronal cultures. *J Neurosci Methods* 2012; **211**: 289–295.
- 21 Brosch M, Yu L, Hubbard T, Choudhary J. Accurate and sensitive peptide identification with Mascot Percolator. *J Proteome Res* 2009; **8**: 3176–81.
- 22 Polpitiya AD, Qian W-JJ, Jaitly N, Petyuk V a., Adkins JN, Camp DG *et al.* DAnTE: A statistical tool for quantitative analysis of -omics data. *Bioinformatics* 2008; **24**: 1556–1558.
- 23 Pavelka N, Fournier ML, Swanson SK, Pelizzola M, Ricciardi-Castagnoli P, Florens L *et al.* Statistical similarities between transcriptomics and quantitative shotgun proteomics data. *Mol Cell Proteomics* 2008; **7**: 631–644.
- 24 Pavelka N, Pelizzola M, Vizzardelli C, Capozzoli M, Splendiani A, Granucci F *et al.* A power law global error model for the identification of differentially expressed genes in microarray data. *BMC Bioinformatics* 2004; **5**: 203.
- 25 Pagliarini DJ, Calvo SE, Chang B, Sheth SA, Vafai SB, Ong S-E *et al.* A mitochondrial protein compendium elucidates complex I disease biology. *Cell* 2008; **134**: 112–23.

- 26 Smith AC, Blackshaw JA, Robinson AJ. MitoMiner: a data warehouse for mitochondrial proteomics data. *Nucleic Acids Res* 2012; **40**: D1160-7.
- 27 Mudunuri U, Che A, Yi M, Stephens RM. bioDBnet: the biological database network. *Bioinformatics* 2009; **25**: 555-6.
- 28 Huang DW, Sherman BT, Lempicki RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 2009; **4**: 44-57.
- 29 Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J *et al.* STRING v10: protein-protein interaction networks, integrated over the tree of life. *Nucleic Acids Res* 2015; **43**: D447-52.
- 30 Chen EY, Tan CM, Kou Y, Duan Q, Wang Z, Meirelles GV *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics* 2013; **14**: 128.
- 31 de Leeuw CA, Mooij JM, Heskes T, Posthuma D. MAGMA: generalized gene-set analysis of GWAS data. *PLoS Comput Biol* 2015; **11**: e1004219.
- 32 Lambert JC, Ibrahim-Verbaas CA, Harold D, Naj AC, Sims R, Bellenguez C *et al.* Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nat Genet* 2013; **45**: 1452-8.
- 33 Major Depressive Disorder Working Group of the Psychiatric GWAS Consortium, Ripke S, Wray NR, Lewis CM, Hamilton SP, Weissman MM *et al.* A mega-analysis of genome-wide association studies for major depressive disorder. *Mol Psychiatry* 2013; **18**: 497-511.
- 34 Cross-Disorder Group of the Psychiatric Genomics Consortium. Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet* 2013; **381**: 1371-9.
- 35 Psychiatric GWAS Consortium Bipolar Disorder Working Group. Large-scale genome-wide association analysis of bipolar disorder identifies a new susceptibility locus near ODZ4. *Nat Genet* 2011; **43**: 977-83.
- 36 Wood AR, Esko T, Yang J, Vedantam S, Pers TH, Gustafsson S *et al.* Defining the role of common variation in the genomic and biological architecture of adult human height. *Nat Genet* 2014; **46**: 1173-86.
- 37 Demontis D, Walters RK, Martin J, Mattheisen M, Als TD, Agerbo E *et al.* Discovery Of The First Genome-Wide Significant Risk Loci For ADHD. *bioRxiv* 2017; : 145581.
- 38 Grove J, Ripke S, Als TD, Mattheisen M, Walters R, Won H *et al.* Common risk variants identified in autism spectrum disorder. *bioRxiv* 2017; : 224774.
- 39 1000 Genomes Project Consortium, Abecasis GR, Auton A, Brooks LD, DePristo MA, Durbin RM *et al.* An integrated map of genetic variation from 1,092 human genomes. *Nature* 2012; **491**: 56-65.
- 40 Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest: Tests in Linear Mixed Effects models. 2016.
- 41 Paternoster V, Rajkumar AP, Nyengaard JR, Børglum AD, Grove J, Christensen JH. The importance of data structure in statistical analysis of dendritic spine morphology. *J Neurosci Methods* 2017; **296**: 93-98.
- 42 Penazzi L, Bakota L, Brandt R. Microtubule Dynamics in Neuronal Development, Plasticity, and Neurodegeneration. *Int Rev Cell Mol Biol* 2016; **321**: 89-169.
- 43 Dent EW. Of microtubules and memory: implications for microtubule dynamics in dendrites and spines. *Mol Biol Cell* 2017; **28**: 1-8.
- 44 English JA, Pennington K, Dunn MJ, Cotter DR. The neuroproteomics of schizophrenia. *Biol Psychiatry* 2011; **69**: 163-72.
- 45 Saia-Cereda VM, Cassoli JS, Martins-de-souza D, Nascimento JM. Psychiatric disorders biochemical pathways unraveled by human brain proteomics. *Eur Arch Psychiatry Clin Neurosci* 2017; **267**: 3-17.
- 46 Davaliev K, Maleva Kostovska I, Dwork AJ. Proteomics Research in Schizophrenia. *Front Cell Neurosci* 2016; **10**: 18.
- 47 Sidoli S, Kulej K, Garcia BA. Why proteomics is not the new genomics and the future of mass spectrometry in cell biology. *J Cell Biol* 2016. doi:10.1083/jcb.201612010.
- 48 Jaworski J, Kapitein LC, Gouveia SM, Dortland BR, Wulf PS, Grigoriev I *et al.* Dynamic Microtubules Regulate Dendritic Spine Morphology and Synaptic Plasticity. *Neuron* 2009; **61**: 85-100.
- 49 Segu L, Pascaud A, Costet P, Darmon M, Buhot M-C. Impairment of spatial learning and memory in ELKL Motif Kinase1 (EMK1/MARK2) knockout mice. *Neurobiol Aging* 2008; **29**: 231-40.
- 50 Shelton MA, Newman JT, Gu H, Sampson AR, Fish KN, MacDonald ML *et al.* Loss of Microtubule-Associated Protein 2 Immunoreactivity Linked to Dendritic Spine Loss in Schizophrenia. *Biol Psychiatry* 2015; **78**. doi:10.1016/j.biopsych.2014.12.029.
- 51 Morava E, Kozicz T. Mitochondria and the economy of stress (mal)adaptation. *Neurosci Biobehav Rev* 2013; **37**: 668-80.
- 52 Chihara T, Luginbuhl D, Luo L. Cytoplasmic and mitochondrial protein translation in axonal and dendritic terminal arborization. *Nat Neurosci* 2007; **10**: 828-37.
- 53 Hara Y, Yuk F, Puri R, Janssen WGM, Rapp PR, Morrison JH. Presynaptic mitochondrial morphology in monkey prefrontal cortex correlates with working memory and is improved with estrogen treatment. *Proc Natl Acad Sci U S A* 2014; **111**: 486-91.
- 54 Sheng M, Ertürk A. Long-term depression: a cell biological view. *Philos Trans R Soc Lond B Biol Sci* 2014; **369**: 20130138.
- 55 D'Amelio M, Cavallucci V, Middei S, Marchetti C, Pacioni S, Ferri A *et al.* Caspase-3 triggers early synaptic dysfunction in a mouse model of Alzheimer's disease. *Nat Neurosci* 2011; **14**: 69-76.

- 56 Barbour JA, Turner N. Mitochondrial stress signaling promotes cellular adaptations. *Int J Cell Biol* 2014; **2014**: 156020.
- 57 Davis J, Moylan S, Harvey BH, Maes M, Berk M. Neuroprogression in schizophrenia: Pathways underpinning clinical staging and therapeutic corollaries. *Aust New Zeal J Psychiatry* 2014; **48**: 512–529.
- 58 Manji H, Kato T, Di Prospero N a., Ness S, Beal MF, Krams M *et al.* Impaired mitochondrial function in psychiatric disorders. *Nat Rev Neurosci* 2012; **13**: 293–307.
- 59 Palmfeldt J, Henningsen K, Eriksen SA, Müller HK, Wiborg O. Protein biomarkers of susceptibility and resilience to stress in a rat model of depression. *Mol Cell Neurosci* 2016; **74**: 87–95.
- 60 Hjelm BE, Rollins B, Mamdani F, Lauterborn JC, Kirov G, Lynch G *et al.* Evidence of Mitochondrial Dysfunction within the Complex Genetic Etiology of Schizophrenia. *Mol neuropsychiatry* 2015; **1**: 201–19.
- 61 Anglin RE, Tarnopolsky MA, Mazurek MF, Rosebush PI. The psychiatric presentation of mitochondrial disorders in adults. *J Neuropsychiatry Clin Neurosci* 2012; **24**: 394–409.
- 62 Fattal O, Link J, Quinn K, Cohen BH, Franco K. Psychiatric comorbidity in 36 adults with mitochondrial cytopathies. *CNS Spectr* 2007; **12**: 429–38.
- 63 Qiao H, Li M-X, Xu C, Chen H-B, An S-C, Ma X-M *et al.* Dendritic Spines in Depression: What We Learned from Animal Models. *Neural Plast* 2016; **2016**: 1–26.
- 64 Harris ' KM, Stevens JK, Harris KM. Dendritic Spines of CA1 Pyramidal Cells in the Rat Hippocampus: Serial Electron Microscopy with Reference to Their Biophysical Characteristics. *J Neurosci* 1989; : 2982–2997.
- 65 Noguchi J, Matsuzaki M, Ellis-Davies GCR, Kasai H. Spine-neck geometry determines NMDA receptor-dependent Ca²⁺ signaling in dendrites. *Neuron* 2005; **46**: 609–22.
- 66 Matsuzaki M, Ellis-Davies GCR, Nemoto T, Miyashita Y, Iino M, Kasai H. Dendritic spine geometry is critical for AMPA receptor expression in hippocampal CA1 pyramidal neurons. *Nat Neurosci* 2001; **4**: 1086–1092.
- 67 Ramirez SA, Raghavachari S, Lew DJ. Dendritic spine geometry can localize GTPase signaling in neurons. *Mol Biol Cell* 2015; **26**: 4171–81.
- 68 Matsuzaki M, Honkura N, Ellis-Davies GCR, Kasai H. Structural basis of long-term potentiation in single dendritic spines. *Nature* 2004; : 761–766.
- 69 Lee KFH, Soares C, Thivierge J-P, Béique J-C. Correlated Synaptic Inputs Drive Dendritic Calcium Amplification and Cooperative Plasticity during Clustered Synapse Development. *Neuron* 2016; **89**: 784–799.
- 70 Govindarajan A, Israely I, Huang S-Y, Tonegawa S. The Dendritic Branch Is the Preferred Integrative Unit for Protein Synthesis-Dependent LTP. *Neuron* 2011; **69**: 132–146.
- 71 van Bommel B, Mikhaylova M. Talking to the neighbours: The molecular and physiological mechanisms of clustered synaptic plasticity. *Neurosci Biobehav Rev* 2016; **71**: 352–361.

Highlights

- Differentially abundant proteins could be identified in frontal cortex, hippocampus and striatum of *Brdl*^{+/-} mice.
- Differentially abundant proteins are enriched for proteins involved in cytoskeleton dynamics and mitochondria.
- Reduced *Brdl* expression affects proteins encoded by schizophrenia risk genes.
- *Brdl*^{+/-} mice have less mature mushroom-type and more immature stubby-type dendritic spines.
- Small dendritic spines are even smaller in *Brdl*^{+/-} mice.