King's Research Portal

DOI:

[10.1016/j.niox.2019.01.011](https://doi.org/10.1016/j.niox.2019.01.011)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Rokkedal-Lausch, T., Franch, J., Poulsen, M. K., Thomsen, L. P., Weitzberg, E., Kamavuako, E. N., Karbing, D. S., & Larsen, R. G. (2019). Chronic high-dose beetroot juice supplementation improves time trial performance of well-trained cyclists in normoxia and hypoxia. *NITRIC OXIDE*, 85, 44-52.
<https://doi.org/10.1016/j.niox.2019.01.011>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

1 **Title**

2 **Chronic high-dose beetroot juice supplementation improves time trial**
3 **performance of well-trained cyclists in normoxia and hypoxia**

4 **Torben, Rokkedal-Lausch¹, Jesper Franch¹, Mathias K. Poulsen², Lars P. Thomsen², Eddie**
5 **Weitzberg³, Ernest N. Kamavuako^{4,5}, Dan S. Karbing², Ryan, G. Larsen¹**

6 ¹Sport Sciences, Department of Health Science and Technology, Aalborg University, DK-9220,
7 Aalborg, Denmark

8 ²Respiratory and Critical Care Group, Center for Model-based Medical Decision Support,
9 Department of Health Science and Technology, Aalborg University, DK-9220, Aalborg, Denmark

10 ³Department of Physiology and Pharmacology, Karolinska Institutet, 171 77 Stockholm, Sweden.

11 ⁴Center for Robotics Research, Department of Informatics, King's College London, London,
12 United Kingdom

13 ⁵SMI, Department of Health Science and Technology, Aalborg University, Aalborg, Denmark

14

15

16

17 Corresponding author: Torben@hst.aau.dk (Torben Rokkedal-Lausch)

18 Declarations of interest: EW is a co-applicant on patents related to the therapeutic
19 use of nitrate and nitrite. Other authors, none.

20

21

22

23

24

25

26

27

28 **Abstract**

29 Dietary nitrate (NO_3^-) supplementation via beetroot juice (BR) is known to
30 improve endurance performance in untrained and moderately trained individuals.
31 However, conflicting results exist in well-trained individuals. Evidence suggests
32 that the effects of NO_3^- are augmented during conditions of reduced oxygen
33 availability (e.g., hypoxia), thereby increasing the probability of performance
34 improvements for well-trained athletes in hypoxia vs. normoxia. This randomized,
35 double-blinded, counterbalanced-crossover study examined the effects of 7 days
36 of BR supplementation with 12.4 mmol NO_3^- per day on 10-km cycling time trial
37 (TT) performance in 12 well-trained cyclists in normoxia (N) and normobaric
38 hypoxia (H). Linear mixed models for repeated measures revealed increases in
39 plasma NO_3^- and NO_2^- after supplementation with BR (both $p < 0.001$). Further, TT
40 performance increased with BR supplementation ($\sim 1.6\%$, $p < 0.05$), with no
41 difference between normoxia and hypoxia ($p = 0.92$). For respiratory variables
42 there were significant effects of supplementation on VO_2 ($p < 0.05$) and VE
43 ($p < 0.05$) such that average VO_2 and VE during the TT increased with BR, with no
44 difference between normoxia and hypoxia ($p \geq 0.86$). We found no effect of
45 supplementation on heart rate, oxygen saturation or muscle oxygenation during
46 the TT. Our results provide new evidence that chronic high-dose NO_3^-
47 supplementation improves cycling performance of well-trained cyclists in both
48 normoxia and hypoxia.

49
50 Keywords:

51 Nitrate,

52 Nitrite,

53 Endurance exercise,

54 Cycling performance,

55 Hypoxia,

56

57 **1.1 Introduction**

58 There is general consensus regarding the physiological factors that limit
59 endurance performance [1,2]. These factors include maximal oxygen consumption
60 ($\text{VO}_{2\text{max}}$), the fractional utilization of $\text{VO}_{2\text{max}}$, and exercise efficiency. Even
61 minor improvements in these factors can enhance performance of endurance
62 athletes. One strategy proposed to improve performance is inorganic nitrate (NO_3^-
63) supplementation, most often in the form of concentrated beetroot juice (BR) [3].
64 When ingested, nitrate is reduced to nitrite and nitric oxide (NO). This pathway
65 differs from the classical pathway for NO generation which involves specific
66 enzymes, NO-synthases (NOS) that use L-arginine and molecular oxygen to
67 generate NO. Nitric oxide has been demonstrated to alter several physiological
68 processes such as blood flow, mitochondrial function and contractile properties
69 [3-8]. Recently, several studies have provided evidence that dietary intake of NO_3^-
70 can improve exercise efficiency (reduction in VO_2 at same work rate) [9-12] and
71 endurance performance [9,10,13-17]. Notably, the majority of studies reporting
72 beneficial effects of NO_3^- has been conducted in untrained and moderately trained
73 individuals ($\text{VO}_{2\text{max}} < 60 \text{ ml/min/kg}$) [10,15,16,18], whereas studies in highly
74 trained individuals ($\text{VO}_{2\text{max}} > 60 \text{ ml/min/kg}$) have shown minor [16,19-21] or no
75 improvements [22-27], indicating that NO_3^- may be less effective in this
76 population [28,29]. In addition to this, recent studies in hypoxia have also
77 provided evidence that NO_3^- improves exercise efficiency [17,21,30,31], muscle
78 oxygenation [31] and elevates oxygen saturation (SpO_2) [21,30,31]. The lower O_2
79 availability in hypoxia impairs the L-Arginine-NOS pathway, and potentiates the
80 nitrate-nitrite-NO pathway, suggesting that BR may be more effective in hypoxia

81 than in normoxia [3,32-34]. Supporting the notion that BR is more effective in
82 hypoxia, Kelly et al. [30] showed that, in healthy individuals, BR improved time
83 to exhaustion during severe intensity exercise in hypoxia but not in normoxia. In
84 addition, BR has been shown to attenuate the decrease in muscle oxygenation and
85 muscle metabolic perturbation in hypoxia in untrained and moderately trained
86 subjects [31,35]. Hence, highly trained athletes may also experience greater
87 performance improvements with BR in hypoxia compared with normoxia.
88 Recently, few studies have examined this idea with conflicting results. In well-
89 trained athletes NO_3^- supplementation had no effect on 10-km or 15-km cycling
90 performance, 10-km running performance or roller-skiing treadmill performance
91 in hypoxia [36-39]. Contrary to this, two studies have reported positive effects of
92 BR in hypoxia on 16.1-km cycling performance and 1500m running performance
93 in trained athletes [17,21]. The discrepancy could be due to different
94 supplementation strategies for NO_3^- . Specifically, the effects of NO_3^-
95 supplementation seems to be potentiated with BR as source of NO_3^- [40,41], with
96 chronic loading over several days [42,43], and by using a dose of >8mmol per day
97 [13,20,44]. Optimizing the supplementation strategy of NO_3^- may be even more
98 important in trained athletes, as this population already exhibit adaptations elicited
99 by endurance training and diet, including higher NO_3^- plasma levels [45,46], NO
100 release [47], NOS activity[48] and a higher percentage of type I fibers [8,49], that
101 altogether may attenuate the response to NO_3^- supplementation.
102 The purpose of the present study was to examine the effects of several days
103 supplementation with a high-dose BR on cycling time trial performance in well-
104 trained cyclists, with continuous measurements of SpO_2 , muscle oxygenation and

oxygen uptake in normoxia and normobaric hypoxia. We hypothesized that BR would improve TT cycling performance in hypoxia but not in normoxia.

2.1 Material and Methods

2.1.1 Participants

Twelve healthy male cyclists at the age of 29.1 ± 7.7 yrs (range 22 to 44 yrs) were enrolled in the study. Participants had a $\text{VO}_{2\text{max}}$ of $5.09 \pm 0.47 \text{ L}\cdot\text{min}^{-1}$ corresponding to $66.4 \pm 5.3 \text{ ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$ and a wattmax of $430 \pm 35 \text{ watt}$ corresponding to $5.6 \pm 0.3 \text{ watt}\cdot\text{kg}^{-1}$ (mean $\pm$ SD). Participants were best classified as well-trained in performance level 4 as defined by Jeukendrup et al. [50] and De Pauw et al. [51], respectively. The protocol and test procedures used in the current study were conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of Northern Jutland (N-20150049). All participants signed informed consent prior to enrollment.

2.1.2 Study design

Participants reported to the laboratory on five separate occasions. Experimental trials followed a randomized counterbalanced-crossover design and were double-blinded for supplementation and single-blinded for inspiratory conditions. The first visit consisted of a maximal exercise performance test to ensure participants were familiar with testing procedures and to ensure participants met the inclusion criteria (i.e., $\text{VO}_{2\text{max}} > 60 \text{ ml}\cdot\text{kg}^{-1}\cdot\text{min}^{-1}$ or wattmax $\geq 5 \text{ w/kg}$). Visits 2-5 involved four experimental trials (Fig 1). Each trial consisted of a 10-km time trial performed in conditions of normoxia or hypoxia, with supplementation of BR or nitrate-depleted BR as placebo (PLA). Specifically, supplementations were ingested in periods of seven days, separated by a wash out period of at least seven

129 days. During each supplementation period, 10-km time trials were performed on
 130 day four and day seven, in different conditions. The order of condition was
 131 maintained for each individual for the first and second supplementation period
 132 such that visits 1 and 3 (and visit 2 and 4) were performed in the same condition.
 133 The design was counterbalanced for condition and supplementation such that half
 134 of the participants started with normoxia and half of the participants started with
 135 BR. All exercise trials were performed on the Cyclus2 ergometer (RBM Cyclus 2,
 136 Germany) using the participants' own bike.

137 **Figure 1: Experimental design**

138 2.1.3 Maximal exercise performance

139 Participants completed a 10-minute warm up at 100 watts and hereafter an
 140 incremental exercise test to exhaustion to determine gas exchange threshold
 141 (GET[30]), VO_{2max} and wattmax (Fig 1). The incremental exercise test
 142 commenced at 100 watts and increased by 30 watts each minute until voluntary
 143 exhaustion. Following a 10-minute rest, participants completed a familiarization
 144 trial for the 10-km TT. While a VO_{2max} validation bout is recommended [52], this

was not performed in this present study as these well-trained cyclists routinely achieve maximal effort during exercise. Respiratory breath-by-breath data were measured throughout the test using a metabolic cart (Jaeger, Vyntus CPX, Carefusion). The metabolic cart was calibrated before each test according to the manufacturer's recommendations. Maximal oxygen uptake ($\text{VO}_{2\text{max}}$) was determined as the highest 30-second average, Wattmax as peak power output from the last minute of the test ($((\text{watt}) + \text{time in last stage (s)})/60 \times 30 \text{ (W)})$ and heart rate (HR) as the peak value attained during the test. GET was determined from a number of measurements, including 1) the first disproportionate increase in VCO_2 from visual inspection of plotting VCO_2 and VO_2 and 2) an increase in expired ventilation ($\text{V}_\text{E}/\text{VO}_2$) with no increase in $\text{V}_\text{E}/\text{VCO}_2$ [30]. HR was recorded continuously using a heart rate sensor (Polar Electro, Oy, Finland).

2.1.4 Experimental trials

Participants ingested BR or PLA for seven consecutive days (Fig 1). Specifically, participants consumed 140ml of concentrated BR (~12.4 mmol nitrate) or 140ml of nitrate-depleted BR (PLA; ~0 mmol nitrate) (Beet It Sport, James White Drinks Ltd., Ipswich, UK) per day; one dose (70 ml) in the morning and one dose (70 ml) in the evening. On the days of the experimental trials (i.e., days four and seven), participants were instructed to consume the total dose (i.e., 140 ml) 2-h prior to arriving at the laboratory (approx. 2.75-h. before commencing the time trial). During the 24-h preceding the first experimental trial, each participant recorded their diet and was told to replicate this diet for the remaining three trials. Participants were also instructed to avoid the intake of specific nitrate-rich foods.

169 The use of antibacterial mouthwash products was not permitted and caffeine
170 intake was prohibited for 12-h preceding each test. For each individual, all
171 experimental trials were performed at the same time of day.

172 Upon arrival at the laboratory, participants rested for 5-minutes before a resting
173 blood sample was drawn into two 4 ml lithium heparin vacutainers
174 (Becton Dickinson, Plymouth, UK). Blood samples were immediately centrifuged
175 for 10 min at 4°C, 3000g after which plasma was extracted and stored at -80 °C
176 for later determination of plasma nitrate and nitrite according to the method
177 described by Hezel et al. [53]. A near infrared spectroscopy (NIRS) probe
178 (Oxymon MK III, Artinis Medical Systems, Netherlands) was placed on the belly
179 of the Vastus Lateralis of the right leg in order to measure changes in muscle
180 oxygenation. Probe position was marked with a permanent pen to ensure identical
181 probe placement for subsequent trials, and the NIRS probe was placed with
182 double-sided adhesive tape. Further, elastic bandages were used to ensure a fixed
183 placement of the probe. An earlobe pulse oximeter (Nonin XPod 8000Q2, Nonin
184 Medical, Inc, Plymouth, MN) was used to measure SpO₂ throughout the tests.

185 Participants then rested 5-minutes on the bike while breathing the gas mixture
186 corresponding to the condition for that specific trial. Throughout each trial,
187 participants breathed through a facemask (Hans Rudolph, V-982185) connected to
188 a low resistance y-valve (Hans Rudolph, two way Y-shape non-rebreathing valve,
189 2730L), with the inspiration valve connected to a closed reservoir. The inspired
190 gas was modified via the closed reservoir using a custom built setup consisting of
191 a mechanical ventilator (SV-300, Maquet, Solna, Sweden) modified such that
192 mixing of gas (pressurized room air and nitrogen) was controlled by manipulating

the inspired oxygen setting on the ventilator. The participants breathed through the same circuit for all experimental trials. The fraction of inspired oxygen was adjusted to $15 \pm 0.1\%$ in hypoxia (~2500m of altitude) and $20.9 \pm 0.1\%$ in normoxia (sea level). Warm-up consisted of three six-minute exercise bouts at the power output corresponding to 70% of GET measured in normoxia. A six-minute rest separated each bout. After the third bout, participants rested for 10 minutes without the facemask. Prior to the TT, participants sat on the bike for five minutes while breathing the gas mixture corresponding to the conditions for that specific trial. Then participants completed a 10-km TT with the instruction of finishing with the highest average power output and as fast as possible. Participants were blinded to all information except cadence and remaining distance of the TT, and were verbally encouraged at each km completed. VO_2 and HR were measured continuously during the TT. For all physiological variables, average values from the 10km-TT were calculated and used for further analyses. Further, peak values for VO_2 , RER (both highest 30-s average) and HR (highest 1-s value) during the TT were calculated and used for further analyses. The ratio of average power to average oxygen uptake (PO/VO_2) during the time trial was used as an index of exercise efficiency [15]. NIRS variables of oxygenated (HbO_2), deoxygenated (HHb) and total (THb) hemoglobin were recorded continuously at 2 Hz and expressed as relative changes (Δ) from the baseline value measured during the final 90-seconds pre-exercise rest period.

2.1.5 Statistical analysis

Differences in performance and physiological parameters were analyzed using linear mixed models for repeated measures. This method of data analysis was

217 used as it has the advantage of preventing listwise deletion due to missing data
 218 (md). For clarification, md for each variable has been noted in table 1. As the
 219 dependent variable, the variable of interest was entered (watt, VO_2 , VE, VCO_2 ,
 220 SpO_2 , etc.) into the model. To investigate the effects of supplementation (BR vs.
 221 PLA), condition (hypoxia vs. normoxia) and supplementation-by-condition, these
 222 were entered as fixed effects. Subject id was included in the model as a random
 223 effect to control for the within-subject nature of the 4 trials. Further, paired t-tests
 224 were used to compare differences between the $\text{VO}_{2\text{peak}}$ obtained during the
 225 normoxic time trials and the $\text{VO}_{2\text{max}}$ from the ramp incremental test. Within group
 226 effect sizes were calculated as the difference in means (BR vs. PLA) divided by
 227 the pooled SD of the change score, using the following definitions: trivial effect d
 228 < 0.2 , small effect > 0.2 , moderate effect > 0.5 , large effect > 0.8 [54].
 229 Associations between changes in TT performance and changes in NO_3^- , NO_2^- ,
 230 VO_2 , and SpO_2 from PLA to BR were assessed using Pearson correlation
 231 coefficient.
 232 All data are presented as means $\pm$ SE, unless stated otherwise, with statistical
 233 significance being accepted when $P \leq 0.05$. All statistical tests were performed
 234 using SPSS 25 (IBM Corp., Armonk, USA) or STATA (Texas, USA) version SE
 235 12.1.

236 **3.1 Results**

237 *3.1.1 Plasma nitrate and nitrite*

238 There were significant main effects of supplementation on NO_3^- and NO_2^- (both
 239 $p < 0.001$) such that BR elevated NO_3^- and NO_2^- (Fig 2). There were no effects of

condition (NO_3^- $p=0.858$; NO_2^- $p=0.542$) or supplementation-by-condition
interaction (NO_3^- $p<0.907$; NO_2^- $p=0.687$).
Further, there were no differences in levels of NO_3^- ($p=0.234$) or NO_2^- ($p=0.231$)
between 4 and 7 days of supplementation (Fig 3).

Figure 2: Individual and mean plasma levels of NO_3^- (A) and NO_2^- (B) (mean $\pm$ SE) prior to time trial tests in normoxia (open bars) and hypoxia (filled bars), after supplementation with beetroot juice (BR) or placebo (PLA). (#, $p < 0.001$, PLA vs. BR, N=11 in hypoxic conditions).

251

252 **Figure 3: Individual and mean plasma levels of NO₃⁻ (A) and NO₂⁻ (B) (mean±SE) prior to**
 253 **time trial tests at day 4 and day 7 after supplementation with beetroot juice (filled bars) or**
 254 **placebo (open bars). (#, p < 0.001, PLA vs. BR, N=11 in hypoxic conditions).**

255

256 3.1.2 Time trial performance

257 All participants completed all four TT's. However, two tests were discarded due
 258 to measurement error (n=1 in N-BR and n=1 in H-PLA). Time trial performance
 259 data are presented in Table 1. There was a main effect of condition (p<0.001) on
 260 time trial performance such that hypoxia lowered power output by ~15% and ~6
 261 %, respectively. Further, there was a main effect of supplementation on time trial
 262 power output (p=0.019) and completion time (p=0.024) showing an overall 1.6%

263 increase in power output and 0.6% reduction in completion time with BR (Fig 4),
 264 with no condition-by-supplementation interaction (both $p=0.923$). Notably, 10 out
 265 of 11 participants increased power output in H-BR compared to H-PLA, whereas
 266 6 out of 11 increased power output in N-BR compared to N-PLA (Fig 4). Effect
 267 size calculations for within group differences between BR and PLA show
 268 moderate (0.703) and small (0.398) effects for hypoxia and normoxia,
 269 respectively.

270
 271 **Figure 4. Individual and mean differences in power output (watt) during 10 km TT**
 272 **performance between placebo and beetroot supplementations in normoxic and hypoxic**
 273 **conditions. Bold horizontal lines indicate mean values for each condition. Single dotted line**
 274 **indicates no difference between beetroot and placebo supplementation**

	Md	N-PLA	N-BR	H-PLA	H-BR	Linear mixed model effects		
						Supplement	Condition	Interaction
	<u>Time Trial</u>							
Performance variable								
Power output, Watt	2	311.3 ± 13.2	315.8 ± 13.2	264.4 ± 13.2	269.3 ± 13.2	p=0.019	p<0.001	p=0.923
Completion time, sec	2	890.1 ± 16	884.5 ± 16	945.6 ± 16	939.5 ± 16	p=0.024	p=0.001	p=0.923
Average values								
PO/VO ₂ , W/L ⁻¹ ·min ⁻¹	10	71.1 ± 1.8	70.8 ± 1.8	68.0 ± 1.8	68.0 ± 1.8	p=0.777	p=0.001	p=0.757
VO ₂ , ml· min ⁻¹	10	4364 ± 140	4443 ± 139	3855 ± 142	3948 ± 142	p=0.030	p<0.001	p=0.862
% VO _{2max}	10	85.9 ± 1.6	87.4 ± 1.6	75.8 ± 1.7	77.7 ± 1.7	p=0.038	p<0.001	P=0.798
VCO ₂ , ml· min ⁻¹	10	4300 ± 151	4498 ± 150	4012 ± 153	4067 ± 153	p=0.005	p<0.001	P=0.120
VE, L· min ⁻¹	10	129.9 ± 7.0	135.8 ± 7.0	136.4 ± 7.1	142.4 ± 7.1	p=0.019	p=0.010	P=0.998
RER	10	0.99 ± 0.01	1.01 ± 0.01	1.04 ± 0.01	1.03 ± 0.01	p=0.462	p=0.003	P=0.082
HR· min ⁻¹ ,	3	168.5 ± 3.1	171.2 ± 3.1	169.4 ± 3.1	169.5 ± 3.1	p=0.118	p=0.486	P=0.072
SpO ₂ , %	9	97.1 ± 0.9	97.1 ± 0.9	84.5 ± 0.9	84.3 ± 0.9	p=0.787	p=0.000	P=0.779
Peak values								
VO _{2peak} , ml· min ⁻¹	10	4925 ± 151	4895 ± 150	4225 ± 152	4304 ± 152	p=0.443	p<0.001	p=0.111
HR _{peak} ,· min ⁻¹	3	183.9 ± 2.9	185.5 ± 2.9	181.1 ± 2.9	181.5 ± 2.9	p=0.153	p<0.001	p=0.308
RER _{peak}	10	1.07 ± 0.02	1.1 ± 0.02	1.14 ± 0.02	1.14 ± 0.02	p=0.334	p=0.003	p=0.246
NIRS								
ΔHbO ₂ , AU	3	-28.5 ± 2.6	-27.6 ± 2.6	-30.7 ± 2.6	-29.4 ± 2.6	p=0.543	p=0.061	p=0.849
ΔHHb, AU	3	24.5 ± 2.6	23.9 ± 2.6	26.3 ± 2.6	26.6 ± 2.6	p=0.885	p=0.042	p=0.633
ΔTHb, AU	3	-4.3 ± 2.0	-3.4 ± 2.0	-3.9 ± 2.0	-2.7 ± 1.9	p=0.527	p=0.766	p=0.934
ΔHHb/VO ₂ , AU· L·min ⁻¹	12	5.68 ± 0.73	5.75 ± 0.71	7.01 ± 0.78	6.78 ± 0.74	p=0.851	p=0.017	p=0.728

279 **Table 1- Average and peak performance, ventilatory and cardiopulmonary data during the**
280 **TT. md denotes the number of missing data points from each variable (complete number of**
281 **data points = 48).**

282 *3.1.3 TT physiological data*

283 Physiological data obtained during the TT are presented in Table 1. There were
284 significant effects of condition on SpO₂ (p<0.001), VE (p=0.010), RER

285 (p=0.003), VCO₂ (p=0.001), VO₂ (p<0.001), PO/VO₂(p=0.001) and % VO_{2max}
286 (p<0.001) such that hypoxia decreased SpO₂, VCO₂, VO₂, PO/VO₂, VO_{2peak},
287 HR_{peak} and % VO_{2max} while VE, RER and RER_{peak} increased. There were
288 significant effects of supplementation on VO₂ (p=0.030) (Fig 5), VE (p=0.019),
289 VCO₂ (p=0.005) and % VO_{2max} (p=0.038) such that VO₂, VE, VCO₂ and % VO_{2max}
290 increased with BR. The VO_{2peak} attained during the time trials in normoxia were
291 significantly lower than the VO_{2max} measured from the incremental test (N-PLA
292 ~3.3%, p=0.03; N-BR ~3.7%, p=0.02).

293

294

295

296

297 **Figure 5- Oxygen uptake profiles from an exemplar subject (A) and mean data (B) from all**
 298 **conditions.**

299

300 3.1.4 Near infrared spectroscopy measures of muscle oxygenation

301 Data reflecting changes in muscle oxygenation during the TT are presented in

302 Table 1. There was a main effect of condition on ΔHHb ($p=0.042$) and

303 $\Delta\text{HHb}/\text{VO}_2$ ($p=0.017$) such that the increase in ΔHHb and $\Delta\text{HHb}/\text{VO}_2$ during the

304 TT was greater in hypoxia (Table 1). We also found a near-significant main effect

of condition on ΔHbO_2 ($p=0.061$) indicating a greater reduction of ΔHbO_2 during TT in hypoxia.

3.1.5 Correlations

There were no significant correlations between changes in performance and changes in plasma NO_3^- or NO_2^- after BR supplementation in normoxia or hypoxia. Further, there were no significant correlations between changes in performance (BR vs. PLA) and changes in VO_2 or SpO_2 nor between changes in performance (BR vs. PLA) and $\text{VO}_{2\max}$.

4.1 Discussion

This is the first study to examine the effects of chronic supplementation with high-dose NO_3^- , in the form of BR, on time trial performance in well-trained athletes in both hypoxia and normoxia.

We show a significant main effect of BR on 10-km TT performance, indicating that well-trained cyclists improve power output and completion time with BR in both normoxia and hypoxia. Supplementation with BR also increased VO_2 during the TT in hypoxia and normoxia, showing that the participants were able to utilize a higher fraction of $\text{VO}_{2\max}$ with BR.

4.1.1 Effects of BR supplementation on TT performance

We found a main effect of BR supplementation on TT performance with no condition-by-supplementation interaction, indicating that BR increased TT performance with no difference between hypoxia and normoxia. However, from a practical perspective, it is worth highlighting that 10 out of 11 participants had higher power output in H-BR vs. H-PLA, while only 6 out of 11 had higher power output in N-BR vs. N-PLA (Figure 3). In support of a small effect of BR, a recent

meta-analysis, including studies performed in hypoxia and normoxia, reported a non-significant 0.8% improvement in time trial endurance performance following BR supplementation [55]. The improvement in 10-km TT completion time and power output of 0.6% and 1.6%, respectively, in the present study, is of practical relevance for elite and well-trained athletes. Specifically, only 0.9% separated first and fourth position during the 13.8-km TT of stage 1 at the 2015 Tour De France cycling race [56], and only 0.3% separated the first and third position during the 9.7-km TT of stage 1 at the 2018 Giro d'Italia cycling race [57]. Further, 0.6% is the smallest worthwhile change in completion time for road TT cyclists proposed by Paton and Hopkins [58].

Few other studies have examined the effects of NO_3^- on TT performance in well-trained athletes in both normoxia and hypoxia within the same study. None of these studies have reported significant improvements in TT performance after BR supplementation [36,38,39]. Nonetheless, the study by Bourdillion et al. [39] reported statistically non-significant improvements in 15-km TT performance of 16s (~1%) and 151s (~7%) in normoxia and hypoxia, respectively.

In general, studies on TT performance performed in well-trained athletes in hypoxia or in normoxia have reported mixed results. In hypoxia, two studies found statistically significant improvements of 2.2-3.2% (~2.2%) [17,21], while one study reported no effect [37]. In normoxia, numerous studies show no effect [22-27,59-61], while a few studies report a significant effect [15,16,20]. The discrepancy in the literature may partly be due to the use of different NO_3^- supplementation strategies that vary in terms of source, dose, and duration (e.g., chronic vs. acute). Many of the previous TT studies have not used an optimized

353 supplementation strategy. Specifically, some studies have used sodium nitrate as
354 the source of NO_3^- [23,39], while there is evidence suggesting that
355 supplementation with NO_3^- in concentrated BR is more effective [40,62]. Several
356 studies have used an acute dose of BR [17,25,26,36-38,59-61], however, a chronic
357 loading protocol consisting of BR supplementation over several days, as used in
358 the present study, has been suggested to be more effective in raising plasma levels
359 of NO_3^- and NO_2^- , and improving performance [11,43]. Finally, several studies
360 have used a low-to-moderate dose of NO_3^- [36,37,59-61], while a higher dose (8-
361 16 mmol), as used in the present study, may be more effective in raising plasma
362 levels and improving performance [13,20,44]. The high dose of NO_3^- used in the
363 present study was tolerated without any adverse events or complaints,
364 demonstrating the efficacy of this supplementation strategy for 7 days. However,
365 there is currently no evidence demonstrating additional benefits with doses higher
366 than 8 mmol. In support of the notion that supplementation strategy is important,
367 studies utilizing an optimized supplementation strategy with chronic
368 supplementation of high dose NO_3^- in the source of BR have reported a significant
369 2.1% [16] and a non-significant 1.7% [24] improvement in TT power output in
370 trained cyclists.

371 4.1.2 Plasma levels of NO_3^- and NO_2^-

372 In the present study, plasma levels of NO_3^- and NO_2^- after placebo (i.e., nitrate-
373 depleted BR) supplementation, were similar to results from other studies using
374 nitrate-depleted BR [17,21,22,37,38,63].

375 Four and seven days of BR supplementation increased NO_3^- and NO_2^- to levels
376 reported in studies using a similar supplementation strategy [13,22], with no

377 differences between 4 and 7 days. Notably, NO_3^- and NO_2^- levels, in the present
 378 study, were higher than those reported in studies using acute supplementation
 379 [17,21,37,38,63]) or lower dosage of NO_3^- [17,37,59,60]. Taken together,
 380 markedly elevated levels of NO_3^- and NO_2^- , in the present study, indicate that BR
 381 supplementation was effective in providing an abundant source of NO via the
 382 nitrate-nitrite-NO pathway. Plasma levels of nitrite displayed a higher variability
 383 compared to plasma nitrate (Fig 2 and Fig 3). This is a common finding and is
 384 most likely due to the shorter half-life of nitrite (less than 1h)[64] compared to
 385 nitrate (5-8h)[65]. This may be explained by a much higher reactivity of nitrite
 386 being subjected to both enzymatic reduction to NO and oxidation to nitrate [33].
 387 Moreover, due to the markedly lower concentration of nitrite in plasma,
 388 measuring techniques display more variable results compared to nitrate.

389 *4.1.3 Physiological effects of beetroot juice supplementation*

390 We found a main effect of supplementation on VO_2 , VE, VCO_2 and % $\text{VO}_{2\text{max}}$
 391 such that BR supplementation resulted in higher VO_2 , VE, VCO_2 and % $\text{VO}_{2\text{max}}$
 392 during the TT in both hypoxia and normoxia. As studies generally show
 393 unchanged [10,12,13,30] or reduced [66,67] $\text{VO}_{2\text{max}}$ following BR
 394 supplementation, these results indicate that the participants were able to utilize a
 395 higher proportion of their maximal aerobic capacity during the TT with BR.
 396 Further, in the present study, a proxy of exercise efficiency (PO/VO_2) during the
 397 TT was unaffected by BR supplementation, suggesting that changes in exercise
 398 efficiency did not contribute to improved TT performance. In agreement with this,
 399 several studies, in well-trained athletes ($>60 \text{ ml}\cdot\text{min}^{-1}\cdot\text{kg}^{-1}$), have shown
 400 unchanged exercise efficiency during submaximal exercise following BR

401 supplementation [24,37,38,63], while only a single study has reported improved
402 efficiency (lower VO_2 during submaximal exercise) in well-trained athletes [21].
403 In club-level cyclists ($56.0 \text{ ml} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$) [15], BR supplementation improved
404 power output with unchanged VO_2 (greater PO/VO_2), indicating improved
405 exercise efficiency. The discrepancy between these results could be due to the
406 training level of the subjects, as our study included well-trained athletes (66.4
407 $\text{ml} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$). Thus, the increase in $\% \text{VO}_{2\text{max}}$ with BR was likely the main factor
408 contributing to increased TT performance. In accordance with these results,
409 Bourdillion et al. [39] reported greater VO_2 and VE with nitrate supplementation
410 in trained cyclists during a 15-km TT in normoxia and hypoxia, which was
411 accompanied by a non-significant increase (1-7%) in performance (discussed
412 above). Contributing to the increased VO_2 with BR, the increase in VE ($\sim 6 \text{ L/min}$)
413 is estimated to account for 10-15 $\text{ml}/\text{O}_2/\text{min}$ ($\sim 10\text{-}20\%$) of the increase in VO_2 ,
414 due to greater oxygen demands of the respiratory muscles [68-70].
415 The active skeletal muscles are the primary site for O_2 usage during the TT, and
416 oxygenation in the vastus lateralis was monitored continuously using NIRS.
417 During the TT, ΔHHb increased in hypoxia compared with normoxia, indicating
418 increased O_2 extraction. However, in agreement with Kelly et al. [30] and
419 Bourdillion et al. [39], ΔHHb was unaffected by BR supplementation, indicating
420 that fractional O_2 extraction in vastus lateralis was not different between BR and
421 PLA. Hence, according to the Fick principle, the increased oxygen uptake in the
422 present study may be a result of increased total O_2 extraction due to increased
423 blood flow. This interpretation is consistent with results demonstrating that NO_3^-

424 supplementation enhances vascular control and muscle blood flow redistribution
425 during exercise [8,49,72].

426 **5.1 Conclusion**

427 In summary, our results provide novel evidence that chronic high-dose BR
428 supplementation improves 10 km time trial performance of well-trained cyclists in
429 both normoxia and hypoxia. Further, BR supplementation resulted in higher VO_2
430 and VE during the TT, suggesting that utilization of a greater proportion of the
431 aerobic capacity contributed to the improved performance. While our results do
432 not identify the underlying mechanisms, enhanced vascular control and muscle
433 blood flow redistribution may contribute to higher VO_2 and improved time trial
434 performance with BR supplementation.

435 **6.1 Acknowledgements**

436 We would like to thank all the participants for the contribution to the present
437 study. Further, we thank Merete Fredsgaard, Brita Holst Serup, Hanne Krone
438 Nielsen and Ditte Beck Christensen for the support in blood sample collections.
439 We thank Carina Nihlen for support in analyzing blood samples. All authors
440 approved the final version of the paper, and have no conflicts of interest.

441 **7.1 Conflict of interest statement**

442 The authors declare: no support from any organization for the submitted work; no
443 financial relationships with any organizations that might have an interest in the
444 submitted work in the previous 3 years; no other relationships or activities that
445 could appear to have influenced the submitted work. EW is a co-applicant on
446 patents related to the therapeutic use of nitrate and nitrite.

447

- 449 [1] Joyner MJ, Coyle EF. Endurance exercise performance: the physiology of
450 champions. *J Physiol* 2008;586:35-44.
- 451 [2] McLaughlin JE, Howley ET, Bassett DR, Jr, Thompson DL, Fitzhugh EC. Test
452 of the classic model for predicting endurance running performance. *Med Sci*
453 *Sports Exerc* 2010;42:991-7.
- 454 [3] Jones AM. Dietary nitrate supplementation and exercise performance. *Sports*
455 *Med* 2014;44 Suppl 1:S35-45.
- 456 [4] James PE, Willis GR, Allen JD, Winyard PG, Jones AM. Nitrate
457 pharmacokinetics: Taking note of the difference. *Nitric Oxide* 2015;48:44-50.
- 458 [5] Haider G, Folland JP. Nitrate supplementation enhances the contractile
459 properties of human skeletal muscle. *Med Sci Sports Exerc* 2014;46:2234-43.
- 460 [6] Hoon MW, Johnson NA, Chapman PG, Burke LM. The effect of nitrate
461 supplementation on exercise performance in healthy individuals: a systematic
462 review and meta-analysis. *Int J Sport Nutr Exerc Metab* 2013;23:522-32.
- 463 [7] Larsen FJ, Schiffer TA, Borniquel S, Sahlin K, Ekblom B, Lundberg JO,
464 Weitzberg E. Dietary inorganic nitrate improves mitochondrial efficiency in
465 humans. *Cell Metab* 2011;13:149-59.
- 466 [8] Ferguson SK, Hirai DM, Copp SW, Holdsworth CT, Allen JD, Jones AM,
467 Musch TI, Poole DC. Impact of dietary nitrate supplementation via beetroot juice
468 on exercising muscle vascular control in rats. *J Physiol* 2013;591 Pt 2:547-57.
- 469 [9] Lansley KE, Winyard PG, Fulford J, Vanhatalo A, Bailey SJ, Blackwell JR,
470 DiMenna FJ, Gilchrist M, Benjamin N, Jones AM. Dietary nitrate
471 supplementation reduces the O₂ cost of walking and running: a placebo-controlled
472 study. *J Appl Physiol* (1985) 2011;110:591-600.
- 473 [10] Bailey SJ, Winyard P, Vanhatalo A, Blackwell JR, Dimenna FJ, Wilkerson
474 DP, Tarr J, Benjamin N, Jones AM. Dietary nitrate supplementation reduces the
475 O₂ cost of low-intensity exercise and enhances tolerance to high-intensity exercise
476 in humans. *J Appl Physiol* (1985) 2009;107:1144-55.
- 477 [11] Wylie LJ, Ortiz de Zavallos J, Isidore T, Nyman L, Vanhatalo A, Bailey SJ,
478 Jones AM. Dose-dependent effects of dietary nitrate on the oxygen cost of
479 moderate-intensity exercise: Acute vs. chronic supplementation. *Nitric Oxide*
480 2016;57:30-9.
- 481 [12] Larsen FJ, Weitzberg E, Lundberg JO, Ekblom B. Effects of dietary nitrate
482 on oxygen cost during exercise. *Acta Physiol (Oxf)* 2007;191:59-66.

- 483 [13] Wylie LJ, Kelly J, Bailey SJ, Blackwell JR, Skiba PF, Winyard PG,
484 Jeukendrup AE, Vanhatalo A, Jones AM. Beetroot juice and exercise:
485 pharmacodynamic and dose-response relationships. *J Appl Physiol* (1985)
486 2013;1153:325-36.
- 487 [14] Dominguez R, Cuenca E, Mate-Munoz JL, Garcia-Fernandez P, Serra-Paya
488 N, Estevan MC, Herreros PV, Garnacho-Castano MV. Effects of Beetroot Juice
489 Supplementation on Cardiorespiratory Endurance in Athletes. A Systematic
490 Review. *Nutrients* 2017;91:10.3390/nu9010043.
- 491 [15] Lansley KE, Winyard PG, Bailey SJ, Vanhatalo A, Wilkerson DP, Blackwell
492 JR, Gilchrist M, Benjamin N, Jones AM. Acute dietary nitrate supplementation
493 improves cycling time trial performance. *Med Sci Sports Exerc* 2011;436:1125-
494 31.
- 495 [16] Cermak NM, Gibala MJ, van Loon LJ. Nitrate supplementation's
496 improvement of 10-km time-trial performance in trained cyclists. *Int J Sport Nutr*
497 *Exerc Metab* 2012;221:64-71.
- 498 [17] Muggeridge DJ, Howe CC, Spendiff O, Pedlar C, James PE, Easton C. A
499 single dose of beetroot juice enhances cycling performance in simulated altitude.
500 *Med Sci Sports Exerc* 2014;461:143-50.
- 501 [18] Murphy M, Eliot K, Heuertz RM, Weiss E. Whole beetroot consumption
502 acutely improves running performance. *J Acad Nutr Diet* 2012;1124:548-52.
- 503 [19] Bond H, Morton L, Braakhuis AJ. Dietary nitrate supplementation improves
504 rowing performance in well-trained rowers. *Int J Sport Nutr Exerc Metab*
505 2012;224:251-6.
- 506 [20] Peeling P, Cox GR, Bullock N, Burke LM. Beetroot Juice Improves on-
507 Water 500 m Time-Trial Performance, and Laboratory-Based Paddling Economy
508 in National and International-Level Kayak Athletes. *Int J Sport Nutr Exerc Metab*
509 2014.
- 510 [21] Shannon OM, Duckworth L, Barlow MJ, Woods D, Lara J, Siervo M, O'Hara
511 JP. Dietary nitrate supplementation enhances high-intensity running performance
512 in moderate normobaric hypoxia, independent of aerobic fitness. *Nitric Oxide*
513 2016;59:63-70.
- 514 [22] Boorsma RK, Whitfield J, Spriet LL. Beetroot juice supplementation does
515 not improve performance of elite 1500-m runners. *Med Sci Sports Exerc*
516 2014;4612:2326-34.
- 517 [23] Bescos R, Ferrer-Roca V, Galilea PA, Roig A, Drobic F, Sureda A,
518 Martorell M, Cordova A, Tur JA, Pons A. Sodium nitrate supplementation does
519 not enhance performance of endurance athletes. *Med Sci Sports Exerc*
520 2012;4412:2400-9.

- 521 [24] Christensen PM, Nyberg M, Bangsbo J. Influence of nitrate supplementation
522 on VO₂ kinetics and endurance of elite cyclists. *Scand J Med Sci Sports*
523 2013;231:e21-31.
- 524 [25] Peacock O, Tjonna AE, James P, Wisloff U, Welde B, Bohlke N, Smith A,
525 Stokes K, Cook C, Sandbakk O. Dietary nitrate does not enhance running
526 performance in elite cross-country skiers. *Med Sci Sports Exerc* 2012;4411:2213-
527 9.
- 528 [26] Cermak NM, Res P, Stinkens R, Lundberg JO, Gibala MJ, van Loon LJC. No
529 improvement in endurance performance after a single dose of beetroot juice. *Int J*
530 *Sport Nutr Exerc Metab* 2012;226:470-8.
- 531 [27] Lane SC, Hawley JA, Desbrow B, Jones AM, Blackwell JR, Ross ML,
532 Zemski AJ, Burke LM. Single and combined effects of beetroot juice and caffeine
533 supplementation on cycling time trial performance. *Appl Physiol Nutr Metab*
534 2014;399:1050-7.
- 535 [28] Porcelli S, Ramaglia M, Bellistri G, Pavei G, Pugliese L, Montorsi M, Rasica
536 L, Marzorati M. Aerobic Fitness Affects the Exercise Performance Responses to
537 Nitrate Supplementation. *Med Sci Sports Exerc* 2014.
- 538 [29] Jones AM, Ferguson SK, Bailey SJ, Vanhatalo A, Poole DC. Fiber Type-
539 Specific Effects of Dietary Nitrate. *Exerc Sport Sci Rev* 2016;442:53-60.
- 540 [30] Kelly J, Vanhatalo A, Bailey SJ, Wylie LJ, Tucker C, List S, Winyard PG,
541 Jones AM. Dietary nitrate supplementation: effects on plasma nitrite and
542 pulmonary O₂ uptake dynamics during exercise in hypoxia and normoxia. *Am J*
543 *Physiol Regul Integr Comp Physiol* 2014;3077:R920-30.
- 544 [31] Masschelein E, Van Thienen R, Wang X, Van Schepdael A, Thomis M,
545 Hespel P. Dietary nitrate improves muscle but not cerebral oxygenation status
546 during exercise in hypoxia. *J Appl Physiol (1985)* 2012;1135:736-45.
- 547 [32] Dauncey S. Can dietary nitrate supplements improve tolerance to hypoxia?
548 *The Intensive Care Society* 2012;133:198-204.
- 549 [33] Lundberg JO, Weitzberg E, Gladwin MT. The nitrate-nitrite-nitric oxide
550 pathway in physiology and therapeutics. *Nat Rev Drug Discov* 2008;72:156-67.
- 551 [34] Shannon OM, McGawley K, Nyback L, Duckworth L, Barlow MJ, Woods D,
552 Siervo M, O'Hara JP. "Beet-ing" the Mountain: A Review of the Physiological
553 and Performance Effects of Dietary Nitrate Supplementation at Simulated and
554 Terrestrial Altitude. *Sports Med* 2017;4711:2155-69.
- 555 [35] Vanhatalo A, Fulford J, Bailey SJ, Blackwell JR, Winyard PG, Jones AM.
556 Dietary nitrate reduces muscle metabolic perturbation and improves exercise
557 tolerance in hypoxia. *J Physiol* 2011;589Pt 22:5517-28.

558 [36] MacLeod KE, Nugent SF, Barr SI, Koehle MS, Sporer BC, MacInnis MJ.
559 Acute Beetroot Juice Supplementation Does Not Improve Cycling Performance in
560 Normoxia or Moderate Hypoxia. *Int J Sport Nutr Exerc Metab* 2015;254:359-66.

561 [37] Arnold JT, Oliver SJ, Lewis-Jones TM, Wylie LJ, Macdonald JH. Beetroot
562 juice does not enhance altitude running performance in well-trained athletes. *Appl*
563 *Physiol Nutr Metab* 2015;406:590-5.

564 [38] Nyback L, Glannerud C, Larsson G, Weitzberg E, Shannon OM, McGawley
565 K. Physiological and performance effects of nitrate supplementation during roller-
566 skiing in normoxia and normobaric hypoxia. *Nitric Oxide* 2017;70:1-8.

567 [39] Bourdillon N, Fan JL, Uva B, Muller H, Meyer P, Kayser B. Effect of oral
568 nitrate supplementation on pulmonary hemodynamics during exercise and time
569 trial performance in normoxia and hypoxia: a randomized controlled trial. *Front*
570 *Physiol* 2015;6:288.

571 [40] Flueck JL, Bogdanova A, Mettler S, Perret C. Is beetroot juice more effective
572 than sodium nitrate? The effects of equimolar nitrate dosages of nitrate-rich
573 beetroot juice and sodium nitrate on oxygen consumption during exercise. *Appl*
574 *Physiol Nutr Metab* 2016;414:421-9.

575 [41] Thompson C, Vanhatalo A, Kadach S, Wylie LJ, Fulford J, Ferguson SK,
576 Blackwell JR, Bailey SJ, Jones AM. Discrete physiological effects of beetroot
577 juice and potassium nitrate supplementation following 4 weeks sprint interval
578 training. *J Appl Physiol* (1985) 2018.

579 [42] Jo E, Fischer M, Auslander AT, Beigarten A, Daggy B, Hansen K, Kessler L,
580 Osmond A, Wang H, Wes R. The effects of multi-day vs. Single pre-exercise
581 nitrate supplement dosing on simulated cycling time trial performance and
582 skeletal muscle oxygenation. *J Strength Cond Res* 2017.

583 [43] Vanhatalo A, Bailey SJ, Blackwell JR, DiMenna FJ, Pavey TG, Wilkerson
584 DP, Benjamin N, Winyard PG, Jones AM. Acute and chronic effects of dietary
585 nitrate supplementation on blood pressure and the physiological responses to
586 moderate-intensity and incremental exercise. *Am J Physiol Regul Integr Comp*
587 *Physiol* 2010;2994:R1121-31.

588 [44] Hoon MW, Jones AM, Johnson NA, Blackwell JR, Broad EM, Lundy B,
589 Rice AJ, Burke LM. The effect of variable doses of inorganic nitrate-rich beetroot
590 juice on simulated 2,000-m rowing performance in trained athletes. *Int J Sports*
591 *Physiol Perform* 2014;94:615-20.

592 [45] Jungersten L, Ambring A, Wall B, Wennmalm A. Both physical fitness and
593 acute exercise regulate nitric oxide formation in healthy humans. *J Appl Physiol*
594 (1985) 1997;823:760-4.

- 595 [46] Poveda JJ, Riestra A, Salas E, Cagigas ML, Lopez-Somoza C, Amado JA,
596 Berrazueta JR. Contribution of nitric oxide to exercise-induced changes in healthy
597 volunteers: effects of acute exercise and long-term physical training. *Eur J Clin*
598 *Invest* 1997;2711:967-71.
- 599 [47] Vassalle C, Lubrano V, Domenici C, L'Abbate A. Influence of chronic
600 aerobic exercise on microcirculatory flow and nitric oxide in humans. *Int J Sports*
601 *Med* 2003;241:30-5.
- 602 [48] McConell GK, Bradley SJ, Stephens TJ, Canny BJ, Kingwell BA, Lee-
603 Young RS. Skeletal muscle nNOS mu protein content is increased by exercise
604 training in humans. *Am J Physiol Regul Integr Comp Physiol* 2007;2932:R821-8.
- 605 [49] Ferguson SK, Holdsworth CT, Wright JL, Fees AJ, Allen JD, Jones AM,
606 Musch TI, Poole DC. Microvascular oxygen pressures in muscles comprised of
607 different fiber types: Impact of dietary nitrate supplementation. *Nitric Oxide*
608 2015;48:38-43.
- 609 [50] Jeukendrup AE, Craig NP, Hawley JA. The bioenergetics of World Class
610 Cycling. *J Sci Med Sport* 2000;34:414-33.
- 611 [51] De Pauw K, Roelands B, Cheung SS, de Geus B, Rietjens G, Meeusen R.
612 Guidelines to classify subject groups in sport-science research. *Int J Sports*
613 *Physiol Perform* 2013;82:111-22.
- 614 [52] Poole DC, Jones AM. Measurement of the maximum oxygen uptake $\text{VO}_{2\text{max}}$:
615 $\text{VO}_{2\text{peak}}$ is no longer acceptable. *J Appl Physiol* (1985) 2017;1224:997-1002.
- 616 [53] Hezel M, Peleli M, Liu M, Zollbrecht C, Jensen BL, Checa A, Giulietti A,
617 Wheelock CE, Lundberg JO, Weitzberg E, Carlstrom M. Dietary nitrate improves
618 age-related hypertension and metabolic abnormalities in rats via modulation of
619 angiotensin II receptor signaling and inhibition of superoxide generation. *Free*
620 *Radic Biol Med* 2016;99:87-98.
- 621 [54] Cohen J. *Statistical Power Analysis for the Behavioral Sciences*, Routledge,
622 Hillsdale, N.J., . 1988;2 edition.
- 623 [55] McMahon NF, Leveritt MD, Pavey TG. The Effect of Dietary Nitrate
624 Supplementation on Endurance Exercise Performance in Healthy Adults: A
625 Systematic Review and Meta-Analysis. *Sports Med* 2017;474:735-56.
- 626 [56] Cyclingnews. Tour De France 2015, 13.8 time trial stage 1.
627 Available from:
628 <http://www.cyclingnews.com/races/tour-de-france-2015/stage-1/results/>.
- 629 [57] Cyclingnews. Giro d'Italia 2018, 9.7km time trial stage 1.
630 Available from:
631 <http://www.cyclingnews.com/races/giro-ditalia-2018/stage-1/results/>.

- 632 [58] Paton C, G. Hopkins W. Variation in performance of elite cyclists from race
633 to race. *European Journal of Sport Science* 2006;6:25-31.
- 634 [59] Glaister M, Pattison JR, Muniz-Pumares D, Patterson SD, Foley P. Effects of
635 dietary nitrate, caffeine, and their combination on 20-km cycling time trial
636 performance. *J Strength Cond Res* 2015;291:165-74.
- 637 [60] Wilkerson DP, Hayward GM, Bailey SJ, Vanhatalo A, Blackwell JR, Jones
638 AM. Influence of acute dietary nitrate supplementation on 50 mile time trial
639 performance in well-trained cyclists. *Eur J Appl Physiol* 2012;11212:4127-34.
- 640 [61] Hoon MW, Hopkins WG, Jones AM, Martin DT, Halson SL, West NP,
641 Johnson NA, Burke LM. Nitrate supplementation and high-intensity performance
642 in competitive cyclists. *Appl Physiol Nutr Metab* 2014;399:1043-9.
- 643 [62] Clifford T, Howatson G, West DJ, Stevenson EJ. Beetroot juice is more
644 beneficial than sodium nitrate for attenuating muscle pain after strenuous
645 eccentric-bias exercise. *Appl Physiol Nutr Metab* 2017;4211:1185-91.
- 646 [63] Christensen PM, Petersen NK, Friis SN, Weitzberg E, Nybo L. Effects of
647 nitrate supplementation in trained and untrained muscle are modest with initial
648 high plasma nitrite levels. *Scand J Med Sci Sports* 2017;2712:1616-26.
- 649 [64] Oldfield EH, Loomba JJ, Monteith SJ, Crowley RW, Medel R, Gress DR,
650 Kassell NF, Dumont AS, Sherman C. Safety and pharmacokinetics of sodium
651 nitrite in patients with subarachnoid hemorrhage: a phase IIa study. *J Neurosurg*
652 2013;1193:634-41.
- 653 [65] McKnight GM, Smith LM, Drummond RS, Duncan CW, Golden M,
654 Benjamin N. Chemical synthesis of nitric oxide in the stomach from dietary
655 nitrate in humans. *Gut* 1997;402:211-4.
- 656 [66] Bescos R, Rodriguez FA, Iglesias X, Ferrer MD, Iborra E, Pons A. Acute
657 administration of inorganic nitrate reduces $\text{VO}_{2\text{peak}}$ in endurance athletes. *Med Sci*
658 *Sports Exerc* 2011;4310:1979-86.
- 659 [67] Larsen FJ, Weitzberg E, Lundberg JO, Eklom B. Dietary nitrate reduces
660 maximal oxygen consumption while maintaining work performance in maximal
661 exercise. *Free Radic Biol Med* 2010;482:342-7.
- 662 [68] Aaron EA, Seow KC, Johnson BD, Dempsey JA. Oxygen cost of exercise
663 hyperpnea: implications for performance. *J Appl Physiol* (1985) 1992;725:1818-
664 25.
- 665 [69] Dominelli PB, Render JN, Molgat-Seon Y, Foster GE, Sheel AW. Precise
666 mimicking of exercise hyperpnea to investigate the oxygen cost of breathing.
667 *Respir Physiol Neurobiol* 2014;201:15-23.

- 668 [70] Vella CA, Marks D, Robergs RA. Oxygen cost of ventilation during
669 incremental exercise to $\text{VO}_{2\text{max}}$. *Respirology* 2006;112:175-81.
- 670 [71] Calbet JA, Lundby C. Air to muscle O_2 delivery during exercise at altitude.
671 *High Alt Med Biol* 2009;102:123-34.
- 672 [72] Richards JC, Racine ML, Hearon CM, Jr, Kunkel M, Luckasen GJ, Larson
673 DG, Allen JD, Dinunno FA. Acute ingestion of dietary nitrate increases muscle
674 blood flow via local vasodilation during handgrip exercise in young adults.
675 *Physiol Rep* 2018;62:10.14814/phy2.13572.
- 676 [73] Van Beekvelt MC, Colier WN, Wevers RA, Van Engelen BG. Performance
677 of near-infrared spectroscopy in measuring local O_2 consumption and blood flow
678 in skeletal muscle. *J Appl Physiol* (1985) 2001;90:511-9.
- 679