King's Research Portal

DOI:

[10.1093/sleep/zsz048](https://doi.org/10.1093/sleep/zsz048)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Bubu, O. M., Pirraglia, E., Andrade, A. G., Sharma, R. A., Gimenez-Badia, S., Umasabor-Bubu, O. Q., Hogan, M. M., Shim, A. M., Mukhtar, F., Sharma, N., Mbah, A. K., Seixas, A. A., Kam, K., Zizi, F., Borenstein, A. R., Mortimer, J. A., Kip, K. E., Morgan, D., Rosenzweig, I., ... Osorio, R. S. (2019). Obstructive Sleep Apnea and Longitudinal Alzheimer's disease biomarker changes. *Sleep*, 42(6), [zsz048].
<https://doi.org/10.1093/sleep/zsz048>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Title: Obstructive Sleep Apnea and Longitudinal Alzheimer's disease biomarker changes

Author Block: Omonigho M. Bubu^{1,2,5,7}, Elizabeth Pirraglia², Andreia G. Andrade², Ram A. Sharma², Sandra Gimenez-Badia³, Ogie Q. Umasabor-Bubu⁴, Megan M. Hogan⁵, Amanda M. Shim⁵, Fahad Mukhtar¹, Nidhi Sharma⁶, Alfred K. Mbah¹, Azizi A. Seixas⁷, Korey Kam⁸, Ferdinand Zizi⁷, Amy R. Borenstein⁹, James A. Mortimer¹, Kevin E. Kip¹, David Morgan^{10,11}, Ivana Rosenzweig¹², Indu Ayappa⁸, David M. Rapoport⁸, Girardin Jean-Louis⁷, Andrew W. Varga⁸, Ricardo S. Osorio^{2,13} for the Alzheimer's Disease Neuroimaging Initiative*

Author Affiliations

1. Department of Epidemiology and Biostatistics, College of Public Health, University of South Florida, Tampa, FL
2. Center for Brain Health, Department of Psychiatry, NYU Langone Medical Center, New York, NY
3. Department of Clinical Psychobiology, University of Barcelona, Spain
4. State University New York (SUNY) Downstate Medical Center, Brooklyn, NY
5. Department of Applied Health Sciences, Wheaton College, Wheaton, IL
6. Dipartimento Di Scienze Biochimica A. Rossi Fanelli, Rome, University of Rome
7. Department of Population Health, New York University School of Medicine, Center for Healthful Behavior Change, New York, USA.
8. Division of Pulmonary, Critical Care and Sleep Medicine at the Icahn School of Medicine at Mount Sinai, New York, NY
9. Department of Family Medicine and Public Health, University of California-San Diego, San Diego, CA.
10. Department of Molecular Science and Translational Medicine, Michigan State University, Grand Rapids, MI
11. Byrd Alzheimer's Disease Institute, College of Medicine, Psychiatry and Behavioral Neurosciences, University of South Florida Health
12. Sleep Disorders Centre, Guy's Hospital, Great Maze Pond, London, SE1 9RT, United Kingdom; Sleep and Brain Plasticity Centre, Department of Neuroimaging, IoPPN, King's College London, United Kingdom.
13. Nathan S. Kline Institute for Psychiatric Research, Orangeburg, New York, USA.

*Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). The investigators within the ADNI contributed to the design and implementation of ADNI projects and provided data, but did not participate in the design, analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

Corresponding Authors: Omonigho M Bubu MD, MPH, PhD. 180 Madison Ave, 7th Floor, New York, NY 10016. E-mail: omonigho.bubu@nyulangone.org, & Ricardo S. Osorio, M.D., 145 East 32nd Street, 5th Floor, New York, NY 10016-6055. E-mail: ricardo.osorio@nyumc.org.

ABSTRACT

Study Objectives: To determine the effect of self-reported clinical diagnosis of Obstructive Sleep Apnea (OSA) on longitudinal changes in brain amyloid-PET and CSF-biomarkers (A β 42, T-tau and P-tau) in cognitively normal (NL), mild cognitive impairment (MCI) and Alzheimer's Disease (AD) elderly.

Methods: Longitudinal study with mean follow-up time of 2.52 $\pm$ 0.51 years. Data was obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database. Participants included 516 NL, 798 MCI and 325 AD elderly. Main Outcomes were annual rate-of-change in brain amyloid-burden (*i.e.* longitudinal increases in florbetapir-PET uptake or decreases in CSF-A β 42 levels); and tau-protein aggregation (*i.e.* longitudinal increases in CSF total-tau (T-tau) and phosphorylated-tau (P-tau)). Adjusted multi-level mixed effects linear regression models with randomly varying intercepts and slopes was used to test whether the rate-of-biomarker-change differed between participants with and without OSA.

Results: In NL and MCI groups, OSA+ subjects experienced faster annual increase in florbetapir uptake (B=.06, 95% CI .02, .11 and B=.08, 95% CI .05, .12 respectively) and decrease in CSF-A β 42 levels (B=-2.71, 95% CI -3.11, -2.35 and B=-2.62, 95% CI -3.23, -2.03, respectively); as well as increases in CSF T-tau (B=3.68, 95% CI 3.31, 4.07 and B=2.21, 95% CI 1.58, 2.86, respectively) and P-tau (B=1.221, 95% CI 1.02, 1.42 and, B=1.74, 95% CI 1.22, 2.27, respectively); compared to OSA- participants. No significant variations in the biomarker changes over time were seen in the AD group.

Conclusions: In both NL and MCI, elderly, clinical interventions aimed to treat OSA are needed to test if OSA treatment may affect the progression of cognitive impairment due to AD.

Key Words: Obstructive sleep apnea; Alzheimer's disease; Longitudinal study, CSF-biomarkers; Brain amyloid-PET; A β 42; T-tau; and P-tau

STATEMENT OF SIGNIFICANCE

Recent studies show that obstructive sleep apnea (OSA) is associated with increased Alzheimer's disease (AD) risk. This study adds to the literature by providing evidence that OSA is related to longitudinal increases in amyloid and tau burden in cognitively normal and mild cognitive impairment (MCI) OSA patients when compared to healthy controls. These novel findings are directly relevant to the emerging literature examining for evidence of a causal relationship between OSA and AD, and are of interest to further understand the various possible mechanistic links that explain this relationship. More importantly, it suggests that clinical interventions aimed to treat OSA in the elderly may slow the progression of cognitive impairment due to AD.

INTRODUCTION

Intermittent hypoxia and sleep fragmentation have been implicated as possible mechanistic links in a causal pathway between obstructive sleep apnea (OSA) and Alzheimer's disease (AD).¹⁻⁴ Recent cross-sectional studies have demonstrated associations between OSA and AD-biomarkers (*i.e.* brain amyloid-PET and CSF-biomarkers (A β 42, T-tau and P-tau)) in both cognitively normal (NL) and mild cognitive impairment (MCI) elderly.⁵⁻⁷ Clearly establishing whether OSA individuals are at heightened risk to develop AD is critical for preventing AD. Positron emission tomography (PET) amyloid-tracer uptake and decreases in cerebrospinal fluid (CSF) amyloid-beta42 (A β 42) levels are robust predictors of amyloid burden as well as of future development of AD.^{8,9} Significant increases in CSF P-tau and CSF T-tau are also documented in AD patients compared with controls.¹⁰ An important question is whether OSA-AD association is related to changes in AD neuropathology over time. Recently, our group demonstrated that objectively measured OSA was associated with markers of increased amyloid burden over a 2-year follow-up in the NYU cohort which consists exclusively of community-dwelling healthy cognitively normal elderly. In this study, we examined self-reported clinical diagnosis of OSA's association with longitudinal changes of brain amyloid-burden and tau-protein aggregation in a sample of cognitively normal (NL), mild cognitive impairment (MCI) and Alzheimer's Disease (AD) elderly in the larger Alzheimer's Disease Neuroimaging Initiative (ADNI) cohort, to provide additional evidence for a possible causal relationship between OSA and AD.

METHODS

Data were obtained from the ADNI database (adni.loni.usc.edu). ADNI was launched in 2003 as a public-private partnership, led by Michael W. Weiner, MD. The primary goal of ADNI has been to test whether serial MRI, PET, other biological markers, as well as clinical and neuropsychological assessments can be combined to measure the progression of MCI and early AD. Thus far, ADNI has recruited over 2,000 adults aged 55-90, consisting of NL, MCI, and early AD. Follow-up for CSF sampling and PET typically occurs every 1-2 years.

Standard Protocol Approvals, Registrations and Patient Consent

The Institutional Review Board at each of the ADNI participating centers approved the ADNI study. Written informed consent was obtained from all patients participating in ADNI. ADNI inclusion and exclusion criteria are detailed elsewhere (http://adni.loni.ucla.edu/wp-content/uploads/2010/09/ADNI_GeneralProceduresManual.pdf).

Study Participants

Participant data used for this study were based on medical history obtained from the ADNI database on December 2016. This study included 1,639 participants: 516 NL, 798 MCI and 325 AD. Subjects missing biomarker data were excluded. Subjects with pre-existing comorbid sleep disorders other than OSA, and those with body mass index (BMI) change greater than 5 between visits, were further excluded, since both could modify group allocation (*i.e.* OSA+ vs. OSA-).¹¹ Other exclusions included previous OSA surgery and having a reversible diagnosis (*i.e.* had a MCI or AD diagnosis at any time point but a NL diagnosis or MCI diagnosis, respectively, at their last visit), thereby allowing us to exclude reversible conditions and unspecified diagnoses.

OSA diagnosis

Presence or absence of OSA was based on self-reported clinical diagnosis of OSA (variable name: *MHDESC*) during a clinical interview. Patients with reported “sleep apnea”, “sleep disordered breathing”, “OSA” or “SDB” were labeled OSA+ and the remaining participants were considered OSA-. To ensure that patients were allocated into the correct groups, three physicians (R.O., S.A. and O.B.) reviewed medical history clinical notes from the ADNI download, for group allocation.

NL, MCI and AD Diagnosis

ADNI criteria for subject classification are described elsewhere.¹² NL and MCI subjects scored between 24-30 on the Mini-Mental State Examination (MMSE) while AD subjects scored between 20-26. MCI and AD participants had global Clinical Dementia Rating (CDR) scores of 0.5 and 1, respectively. The diagnosis of AD was made using established clinical criteria.¹³

Florbetapir-PET Imaging Acquisition and Interpretation

ADNI florbetapir summary data was uploaded to the Laboratory of Neuroimaging (LONI) by the University of California at Berkeley.^{14,15} For more information on the methods see: https://adni.loni.usc.edu/wp-content/uploads/2010/05/ADNI2_PET_Tech_Manual_0142011.pdf, <http://adni.loni.usc.edu/updated-florbetapir-av-45-pet-analysis-results/>. Briefly, native-space segmented and parcellated magnetic resonance imaging (MRI) scan with Freesurfer (version 5.3.0) was used for each subject to define cortical grey matter regions of interest (frontal, anterior/posterior cingulate, lateral parietal, lateral temporal) that make up a summary cortical region of interests (ROI). Five reference regions (cerebellar grey matter, whole cerebellum,

brainstem/pons, eroded subcortical white matter, and a composite reference region) were defined. Each florbetapir scan was then matched to the corresponding MRI and the mean florbetapir uptake within the cortical and reference regions is calculated. The baseline MRI was used to define regions for all subsequent florbetapir scans. Image data are available at 3 levels of preprocessing (raw, unsmoothed, and smoothed) as described online at <http://adni.loni.usc.edu/data-samples/mri/>. Calculation of Florbetapir standardized uptake value ratios (SUVRs) were done by obtaining means across the 4 cortical regions and dividing this cortical summary ROI by one of the five reference regions. Cortical ROI selection and reference region is predicated on the goals of the analysis. Two summary SUVRs were provided. The first is the cortical summary ROI divided by the whole cerebellum reference region (SUMMARYSUVR_WHOLECEREBNORM), which as recommended we used for cross-sectional florbetapir analyses. The second is the cortical summary ROI divided by a composite reference region (SUMMARYSUVR_COMPOSITE_REFNORM), which as recommended we used for our longitudinal florbetapir analyses.

Cerebrospinal Fluid Methods (CSF)

CSF bio-specimen data collection details can be found at <http://adni.loni.usc.edu/data-samples/biospecimen-data/>. Briefly, A β 42, t-tau and p-tau181 were measured from CSF that was collected at each ADNI site, and transferred into polypropylene transfer tubes followed by freezing on dry ice within 1 h after collection and shipped overnight to the ADNI Biomarker Core laboratory at the University of Pennsylvania Medical Centre on dry ice. A standardized protocol was implemented to quantify biomarker concentrations in each of the CSF baseline aliquots using a multiplex xMAP Luminex platform (Luminex Corp, Austin,TX) with Innogenetics (INNO-BIA AlzBio3, Ghent, Belgium; for research use only reagents)

immunoassay kit-based reagents, validated in Vanderstichele et al.¹⁶ and Shaw et al.¹⁷ Further details can be found at (<http://www.adni-info.org/index.php>).

Covariates/potential confounders

As our study tried to answer an etiological question, we employed a broad selection of covariates based on published literature.^{18,19} Covariates were selected *a priori* and included factors that might be more common in cases with OSA and that might contribute to either A β -pathology (CSF A β 42 and Florbetapir PET) or neurodegeneration (CSF T-tau), including age, sex, BMI, education, CPAP-use, ApoE4 status, alcohol intake, baseline biomarker data, history of respiratory disease, hypertension, diabetes and history of cardiovascular disease (e.g. including ischemic heart disease, heart failure, and stroke/TIA), as well as history of traumatic brain injury.

Data analyses

All analyses were conducted separately for each clinical group (*i.e.* NL, MCI, and AD). Cross-sectional analyses of baseline AD-biomarker levels by OSA status were conducted using generalized linear methods.

Multivariate analysis of variance (MANOVA) was conducted to test differences in time-trend, groups (OSA+ *vs.* OSA-) and time points. To test differences in time-trend, baseline biomarker data was subtracted from data at each time point.

ADNI data is unbalanced with unequal numbers of measurement for each participant. We therefore used multi-level mixed effects linear regression models with normal errors,²⁰ examining the relationship between OSA and the rate-of-change in AD biomarkers. This allowed

the incorporation of all available information and possibly reduced bias resulting from using only the complete cases.²¹

To determine the specific type of modeling, we examined profile plots for the biomarker levels over time by OSA status, using both the original dataset and within-subject residuals. We also examined trajectory plots obtained by subtracting the baseline measurement from the original measurements (*i.e.* $d_{ij} = Y_{ij} - Y_{i1}$) and examined plots of mean and variance of the biomarkers at each time point by OSA status. Further, using the slopes as summary statistics, we conducted different formal tests to compare groups. Our assumption was that the rate-of-change of biomarker data is approximately linear over time. Thus, we used a parametric model with time as a continuous variable. We fitted the models with randomly varying intercepts and slopes and allowed them to depend on exposure group (*i.e.*, OSA status). Based on the findings, we then fitted a random coefficients model to analyze the relationship between OSA status and time using unstructured covariance. To find a parsimonious covariance structure, we fitted many different covariance models and conducted a likelihood ratio test of each nested model using information criteria for non-nested models. Finally, using the unstructured, independence covariance model (*i.e.*, the preferred covariance structure), we tested whether the rate-of-change in biomarkers differed between OSA groups. Final models were adjusted for age, sex, BMI, education, CPAP-use, ApoE4 status, alcohol intake, baseline biomarker data, history of respiratory disease, hypertension, diabetes and history of cardiovascular disease (e.g. including ischemic heart disease, heart failure, and stroke/TIA), and history of traumatic brain injury.

Adjustment for the extensive list of covariates had little impact on the estimates. The age and sex only-adjusted model and the extensively-adjusted model showed similar results, indicating little to no risk of bias due to over-adjustment. Sensitivity analysis removing CPAP-

users (NL: n=10; MCI: n=16; AD: n=9) from OSA+ participants had also a negligible impact on the estimates (*e.g.* NL florbetapir estimate of .06 changed to .08). Statistical analyses were performed using SAS (*version 9.4; SAS Institute Inc., Cary, NC*).

These analyses allowed us to examine whether significant variation between OSA subjects in mean AD biomarker level at baseline (intercept) existed, as well as whether significant variation in the change in AD biomarker level over time (slope) occurred. Furthermore, the covariance between the baseline AD biomarker level (intercept) and AD biomarker change over time indicated whether OSA+ or OSA- subjects had experienced a faster increase or decrease in AD biomarker level over time (significant slope). It also allowed for assessment of significant differences in the rate of change in AD biomarker level between OSA+ and OSA- patient groups over time.

Data Availability Statement

Authors state that anonymized data will be shared by request from any qualified investigator.

RESULTS

Demographic and Clinical Characteristics

Figure 1 shows the stepwise exclusion process for NL, MCI and AD participants used in the study. Table 1 shows the demographic and clinical characteristics of study participants at baseline. Overall, the mean ages of OSA+ and OSA- were 72.3 ± 7.1 and 73.9 ± 7.3 years, respectively. Female participants represented 49%, 40% and 37% of the NL, MCI and AD groups, respectively. Six percent were OSA+ in the NL group, while 13% and 7% were OSA+ in

the MCI and AD groups. Participants differed markedly in ApoE4 status with 28%, 51% and 66% being ApoE4 positive in the NL, MCI and AD groups. NL, MCI and AD participants were similar in age, education, BMI, and the three medical conditions across groups. Mean follow up time was 2.52 ± 0.51 years.

Baseline AD Biomarker levels by clinical group and OSA status

Table 1 and Supplementary Figures 1a-c show raw baseline CSF A β 42, T-tau and P-tau levels, and florbetapir PET uptake in NL, MCI and AD groups, by OSA status. There were significant differences in CSF A β 42 levels for both MCI ($F=4.37$, $p=.04$), and AD ($F=6.89$, $p<.01$) groups respectively (Figure 1), with OSA+ participants having significantly higher levels at baseline for the MCI group. Significant difference in T-tau levels was seen for the MCI patients ($F=5.08$, $p=.02$) (Figure 1) with OSA+ individuals having significantly lower levels. These differences remained after controlling for covariates, but the magnitude of the differences was small. For florbetapir values there were significant differences between OSA groups for both NL and MCI participants ($F=5.53$, $F=5.15$, $p \leq .02$ for all respectively) (Figure 1) with OSA+ participants having significantly lower florbetapir values. However, after controlling for age, sex, BMI, education, CPAP use, ApoE4 status, and other medical conditions, these differences in florbetapir values were no longer significant. For P-tau levels, no significant difference was seen across all groups in the uncontrolled and controlled analyses.

Differences in Time Trend, Groups (OSA+ versus OSA-) And Time-points

Table 2, Figures 2 - 4, and Supplementary Figures 2a-c show MANOVA results testing the differences in the mean change in AD-biomarker values over time, based on OSA status. The Pillai's trace test values presented examines time-point*OSA effect and provides the exact F

statistics for the time trend of mean-change in AD biomarker across OSA groups. The time-point effect provides the exact F statistics for the mean change (increase or decrease) in Alzheimer's disease biomarker over time. Other stats interpretations can be seen in the Table 2 footnote.

In NL and MCI participants, the time trend of mean change in florbetapir PET value indicated the change was not parallel across groups (Pillai's trace test, $p=.01$ for all). Across all subjects, mean change in florbetapir uptake increased significantly over time (Pillai's trace test, $p<.001$ for all). There were significant differences in mean change in florbetapir values across the OSA groups when the previous time-point was compared to the next ($p\leq.01$ for all).

In AD participants, the only significant finding was that the mean change in florbetapir uptake increased over time across all subjects (Pillai's trace test, $p<.001$).

In NL and MCI participants, the time trend of mean change in CSF A β 42, T-tau and P-tau values indicated the change was not parallel across groups (Pillai's trace test, $p\leq.05$ for all). Across all subjects, mean change in CSF A β 42 values decreased while T-tau and P-tau values increased over time (Pillai's trace test, $p<.001$ for all). There were significant differences in mean change in CSF A β 42, and T-tau values across the OSA groups when the previous time-point was compared to the next ($p\leq.01$ for all). For P-Tau, significant differences were only seen when timepoint_1 (year 2) was compared to timepoint_0 (year 1) ($p<.001$).

In AD participants, the time trend of mean change in CSF T-tau indicated the change was not parallel across groups (Pillai's trace test, $p\leq.01$). Across all subjects, mean change in CSF A β 42 values decreased while P-tau values increased significantly over time (Pillai's trace test, $p<.001$ for all). There were significant differences in mean change in T-tau values across the OSA

groups only when timepoint_1 (year 2) was compared to timepoint_0 (year 1) ($p < .001$). No significant differences were seen for CSF A β 42.

Rate-of-change in AD Biomarker by OSA status

Table 3 reports the between-subjects variation and covariance parameter estimates in AD biomarker levels over time by OSA group.

In both NL and MCI participants at baseline, OSA+ participants had lower levels of CSF T-tau, CSF P-tau and brain Florbetapir values and higher CSF- A β 42 levels, compared to OSA- subjects (Table 1). Interpretation of the covariance parameters is such that, if the signs of these estimates are negative, they show that persons with higher baseline values tend to have less rates of change over time. For example, in NL participants, for the florbetapir values, the covariance parameter estimate of -.06 (-.09, -.04) indicates that OSA- participants who had a higher baseline tended to have slower rates of amyloid deposition over time (Figure 2). The covariance parameter estimate for CSF A β 42 of 3.93 (3.56, 4.31) indicates that OSA+ who had higher baseline values tended to have higher rates of changes over time (in this case decline because of the negative slope)

In NL participants, there was significant variation by OSA status in the annual change in brain florbetapir uptake (mean SUVR; $B=0.06$, $p<.0001$). In MCI participants, there was also significant variation by OSA status in the annual change in florbetapir values (mean SUVR; $B=0.08$, $p<.0001$). The covariance between the baseline florbetapir uptake and the annual florbetapir value change was -.06, indicating that OSA+ subjects experienced faster increase in amyloid deposition over time ($p<.0001$) when compared to OSA- (Figure 3 and supplementary

2b). No significant variation in the change in brain florbetapir-PET volumes by OSA status was seen in the AD group (Table 3 and Figure 4).

For the NL and MCI groups, there was significant variation in the annual change in CSF A β 42 by OSA status (mean CSF levels; B=-2.708, mean CSF levels; B =-2.264, $p<.0001$ for all respectively). Significant differences in the annual change in T-tau and P-tau levels over time were also observed in both groups. Covariance parameters between the baseline CSF A β 42, T-tau and P-tau level change over time indicated that OSA+ subjects experienced a faster longitudinal decrease in CSF A β 42 and faster longitudinal increases in T-tau and P-tau levels ($p<.0001$ for all) compared to OSA- participants, in both the NL and MCI groups (Table 3 and Figures 2 - 4). No significant variations in the change in CSF A β 42, T-tau and P-tau levels over time were seen for the AD group (Table 3).

DISCUSSION

The major objective of this study was to examine the association of OSA and longitudinal changes in brain amyloid-PET deposition and CSF biomarkers. There were significant differences in the annual rate-of-change in florbetapir uptake as well as CSF A β 42, T-tau and P-tau levels over the follow-up period for the NL and MCI groups, with OSA+ subjects experiencing significantly faster increase in brain amyloid load (as measured by increases in florbetapir uptake and decreases in CSF A β 42 levels) and tau aggregates (as measured by increases in CSF T-tau and P-tau levels). The direction of these longitudinal changes is in accordance with growing evidence described below showing a link between disturbed sleep, OSA and AD pathogenesis. In a cross-sectional study, Spira et al.²² found that MCI individuals with higher apnea-hypopnea and oxygen desaturation indices demonstrated higher amyloid

deposition on Pittsburgh compound B (PiB)-PET uptake globally and regionally in the precuneus. Recently, Yun et al.⁷ found that OSA patients had a higher amyloid burden in the right posterior cingulate gyrus and right temporal cortex relative to controls, suggesting possible contribution of OSA. In a population of NL individuals, Liguori et al.⁵ compared CSF biomarkers in OSA+ vs. OSA+ treated with CPAP as well as OSA- controls. Findings demonstrated lower CSF A β 42 and higher T-tau/A β 42 ratio in OSA+ compared to CPAP treated and OSA- control subjects. We also have recently documented that OSA severity is associated with increased amyloid burden (measured as longitudinal decreases in CSF A β 42 and increases in PiB uptake) over a 2-year follow-up in community-dwelling NL-elderly.²³

Possible mechanisms responsible for these findings include intermittent hypoxia, sleep fragmentation and intrathoracic pressure swings (see Figure 5). Studies of cerebral ischemia suggest that both acute and intermittent hypoxia may promote A β accumulation,^{2,4,24-26} possibly mediated by hypertension,²⁷ cardiovascular disease,²⁸ diabetes,²⁹ chronic inflammation,³⁰ and oxidative stress.^{31,32} Sleep fragmentation, arousals and circadian rhythm disruption have also been shown to increase the risk of developing MCI/AD,^{1,33} possibly through disruption of slow wave sleep (SWS).^{34,35} Findings from a recent prospective study on subjects from the Framingham Heart Study cohort; suggest that OSA-related REM disruption may also contribute to this association.³⁶ Other possible mechanisms include elevated intrathoracic, intracranial pressures and/or increased venous pressure, which have been hypothesized to acutely and repetitively impede the circulation of brain metabolites from the interstitial fluid (ISF) into the CSF through the glymphatic system, leading to increases in A β accumulation.^{24,31,32,37} Lastly, depression has been reported as both a consequence of OSA and an AD risk factor and may also contribute to this association.³⁸

Intriguingly, at baseline, our data showed that OSA+ patients had significantly higher CSF A β 42 in MCIs, lower florbetapir uptake in both NL and MCIs, and lower T-tau levels in MCIs, compared to OSA- subjects. A possible concern here is a regression toward the mean, since the OSA+ group starts with significantly lower values of amyloid load in some of the biomarkers used. However, there are two factors worth addressing in this regard. First, the absolute difference in these levels between groups was numerically quite small (*e.g.*, in NL subjects, the difference in the median CSF A β 42 levels was 17 pg/mL). Second, the longitudinal analyses examined the “rate-of-change” as an outcome and not the mean levels of the AD-biomarkers. The longitudinal mean beta statistics presented refer to the mean change incorporating within and between group data. In addition, what we examined was the slope statistics incorporating the effects of time, which looked at the rate-of-change for each individual between OSA groups. Furthermore, one could expect faster progression of amyloid deposition in the NL group that has already higher amyloid burden at baseline,^{39,40} which would have driven our findings toward the null. Possible explanations for the reduced baseline AD biomarker presentation in the OSA+ group include; first, if OSA accelerates AD pathology build-up this may simply be a sign of selection (survival) bias as most OSA+ will have transitioned to MCI or AD and only those with very low AD pathology burden at disease onset would remain as cognitively normal. Second, particularly in those with MCI, is the presence of increased vascular risk in this group (*i.e.* higher rates of diabetes and obesity) (Table 1), as onset of cognitive impairment is likely to occur at a lower severity of Alzheimer lesions in individuals with more vascular pathology.⁴¹ Notably, after age-stratification (55 – 73.8, and 74 - 92) based on age distribution (we categorized based on the 50th percentile), compared to OSA- individuals, OSA+ individuals in the 55 – 73.8 years old category were more likely to have higher vascular burden

(i.e. composite score for BMI, hypertension, diabetes and cardiovascular disease), higher CSF A β 42 in MCIs, lower florbetapir uptake in both NL and MCIs, and lower T-tau levels in MCIs (see supplementary Table 1). In addition, among ages 74 years and above, OSA+ participants had a worse AD biomarker presentation (i.e. lower CSF A β 42 in MCIs, higher florbetapir uptake in both NL and MCIs, and higher T-tau levels in MCIs) at baseline. These findings suggest that the younger population especially in the OSA+ with higher vascular burden were driving the noted differences at baseline between OSA+ vs. OSA- groups. Third, specifically for CSF A β 42; slow wave sleep (SWS) occurring during non-rapid eye movement (NREM) sleep, a known A β modulator,^{35,42,43} may be responsible. OSA is associated with reduced SWS.⁴⁴ Sleep is involved in the clearance of A β ,³⁵ and fragmented SWS limits brain A β clearance, thereby leaving higher levels that reflect in the CSF.^{45,46} At cross-section, disrupted SWS has been shown to be associated with higher CSF A β 42 in middle-aged^{24,47} and older adults.³⁴ Fourth, in both the NL and MCI groups, OSA- had more APOE4 positive participants, which is known to confer higher risk and worse AD biomarker burden.⁴⁸ It is important to note that as stated above (see Results section: Baseline AD Biomarker levels by clinical group and OSA status), after controlling for age, sex, BMI, education, CPAP use, ApoE4 status, and other medical conditions, the differences in florbetapir values were no longer significant. For P-tau levels, no significant difference was seen across all groups in the uncontrolled and controlled analyses. In other words, these differences are numerically but not statistically different from each other, and where they were “statistically” significantly different, it is not clear that they are clinically meaningfully different.

Our findings in a cohort of self-reported OSA patients suggest that untreated OSA can predispose to neurodegenerative processes in the long-term by gradually weakening brain structure and altering its functioning.^{49,50} This hypothesis would be consistent with an

overarching model of late-onset AD with brain amyloid deposition and tau aggregates proceeding at different rates, influenced by a combination of protective/risk factors of which OSA is part.⁵¹ This model of AD implies a contributory role of OSA severity in affecting and/or accelerating AD biomarker change and possibly increasing neurodegeneration. In agreement with these findings, our previous study performed in the ADNI cohort showed that the presence of OSA was associated with accelerated cognitive decline and an earlier age of MCI or AD-dementia onset.⁵² As expected, there was no difference in the rate-of-biomarker change in the AD group by OSA status. This may be because in AD patients, brain amyloid is exhibiting a ceiling effect at higher levels of brain atrophy.⁵³ OSA's effect as it relates to AD biomarker changes may therefore be attenuated after it has reached its maximum neuronal damage.

Strengths and Limitations

Our study possesses several strengths including a well-defined cohort, and objective assessment of amyloid burden and CSF P-tau/T-tau, which allowed for a high degree of certainty regarding measurement of clinical groups and outcomes.⁵⁴ Our statistical analytic methods were also robust with respect to unbalanced number of observations per subject over time.

We acknowledge we were limited by the measurement of OSA by self-report. Self-reported sleep measures can be impacted by diminished cognition⁵⁵ and in certain situations might not be correlated with objective methods.⁵⁶ The prevalence of reported OSA was also significantly lower than expected in this elderly cohort, which can relate to an under-diagnosis effect in this population, as epidemiological and sleep laboratory studies document much higher OSA prevalence in elderly populations. Therefore, some OSA+ subjects were likely misclassified into the OSA- group; however, this would have driven our findings towards the

null. However, one implication of OSA classification by self-report is that those with self-reported OSA very likely had associated symptoms (*i.e.* excessive daytime sleepiness) that prompted these subjects to seek diagnosis. The overall prevalence of self-reported OSA in this group (6%) is similar to the U.S. prevalence of the OSA Syndrome (OSAS) (4%) defined by the presence of $AHI4\% \geq 5/\text{hour}$ and daytime symptoms across all ages.⁵⁷ The prevalence of OSA (with or without symptoms) in the elderly is much higher, estimated at 30-50% in older subjects,⁵⁸ thus additional work may be required to differentiate the risk of OSA for AD with and without associated daytime symptoms. Notably, all-cause excessive daytime sleepiness in elderly subjects defined by Epworth sleepiness scores ≥ 10 was associated with longitudinal brain beta amyloid accumulation in a recent study.⁵⁹

CONCLUSIONS

OSA appears to accelerate increases in amyloid deposition, CSF T-tau and P-tau levels over time, both in NL and MCI individuals. Sleep fragmentation, intermittent hypoxia and intrathoracic pressure swings from OSA are likely candidate mechanisms. Thus, clinical interventions aimed at OSA, such as treatment with CPAP or dental appliances, in cognitive normal and MCI patients, could possibly mitigate or slow the progression of cognitive impairment to AD. Further studies examining the mechanisms underlying these observed effects are needed.

ACKNOWLEDGMENTS

The authors would like to gratefully acknowledge the dedication, time, and commitment of the participants and staff of the Alzheimer's Disease Neuroimaging Initiative (ADNI). Collection and sharing of data for this project was funded by ADNI (National Institutes of Health Grant

U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: AbbVie, Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica, Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California.

DISCLOSURE STATEMENT

Financial Disclosure: RSO gave a paid talk to Eisai Global at the SLEEP 2017 meeting.

Non-financial Disclosure: none

Study Funding: This work was supported by grants from the NIH/NIA/NHLBI (T32HL129953, R01HL118624, R21AG049348, R21AG055002, R01AG056031, R01AG022374,

R21AG059179, R01AG056682, P30AG008051 and 1R01AG056531-01A1); Foundation for Research in Sleep Disorders; American Sleep Medicine Foundation, the American Thoracic Society Foundation, the Friedman Brain Institute. The Wellcome Trust (103952/Z/14/Z) supports I.R.

AUTHORS CONTRIBUTION

OMB, RSO, EP, AA, and RAS were involved in the conception and design of the study, acquisition and analysis of the data, as well as drafting a significant portion of the manuscript or figures. SG, OQU, MMH, AMS, FM, NS, and AKM were involved in the literature review, conception and design of the study, and revision of the manuscript for intellectual content. AAS, KK, FZ, ARB, JAM, KEK, DM, IR, IA, DMR, GJ, and AWV, interpreted the data; revised the manuscript for intellectual content and drafted significant portions of the manuscript.

References

1. Lim AS, Kowgier M, Yu L, Buchman AS, Bennett DA. Sleep Fragmentation and the Risk of Incident Alzheimer's Disease and Cognitive Decline in Older Persons. *Sleep*. 2013;36(7):1027-1032.
2. Nalivaeva NN, Fisk L, Kochkina EG, et al. Effect of hypoxia/ischemia and hypoxic preconditioning/reperfusion on expression of some amyloid-degrading enzymes. *Annals of the New York Academy of Sciences*. 2004;1035:21-33.

3. Polsek D, Gildeh N, Cash D, et al. Obstructive sleep apnoea and Alzheimer's disease: In search of shared pathomechanisms. *Neuroscience and biobehavioral reviews*. 2018;86:142-149.
4. Sun X, He G, Qing H, et al. Hypoxia facilitates Alzheimer's disease pathogenesis by up-regulating BACE1 gene expression. *Proceedings of the National Academy of Sciences of the United States of America*. 2006;103(49):18727-18732.
5. Liguori C, Mercuri NB, Izzi F, et al. Obstructive Sleep Apnea is Associated With Early but Possibly Modifiable Alzheimer's Disease Biomarkers Changes. *Sleep*. 2017;40(5).
6. Spira AP, Gamaldo AA, An Y, et al. Self-reported Sleep and beta-Amyloid Deposition in Community-Dwelling Older Adults. *JAMA neurology*. 2013.
7. Yun CH, Lee HY, Lee SK, et al. Amyloid Burden in Obstructive Sleep Apnea. *Journal of Alzheimer's disease : JAD*. 2017;59(1):21-29.
8. Brys M, Pirraglia E, Rich K, et al. Prediction and longitudinal study of CSF biomarkers in mild cognitive impairment. *Neurobiol Aging*. 2009;30(5):682-690.
9. Waragai M, Okamura N, Furukawa K, et al. Comparison study of amyloid PET and voxel-based morphometry analysis in mild cognitive impairment and Alzheimer's disease. *Journal of the neurological sciences*. 2009;285(1-2):100-108.
10. Buerger K HH. Evaluation of phosphorylated tau protein as a core biomarker of Alzheimer's disease. . In: *Iqbal I, Winblad B, editors Proceedings of the 9th International Conference on Alzheimer's Disease and Related Disorders Philadelphia, Pennsylvania: Alzheimer's Association; 2004.p. 46–58.*
11. Fitzpatrick AL, Kuller LH, Lopez OL, et al. Midlife and late-life obesity and the risk of dementia: cardiovascular health study. *Archives of neurology*. 2009;66(3):336-342.

12. Petersen RC, Aisen PS, Beckett LA, et al. Alzheimer's Disease Neuroimaging Initiative (ADNI): clinical characterization. *Neurology*. 2010;74(3):201-209.
13. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology*. 1984;34(7):939-944.
14. Landau SM, Breault C, Joshi AD, et al. Amyloid-beta imaging with Pittsburgh compound B and florbetapir: comparing radiotracers and quantification methods. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine*. 2013;54(1):70-77.
15. Landau SM, Lu M, Joshi AD, et al. Comparing positron emission tomography imaging and cerebrospinal fluid measurements of beta-amyloid. *Annals of neurology*. 2013;74(6):826-836.
16. Vanderstichele H DMG SF, et al. . Biomarkers for early diagnosis of Alzheimer's disease. Hauppauge, NY. *Nova Science Publishers*. 2008.
17. Shaw LM, Vanderstichele H, Knapik-Czajka M, et al. Cerebrospinal fluid biomarker signature in Alzheimer's disease neuroimaging initiative subjects. *Annals of neurology*. 2009;65(4):403-413.
18. Hernan MA, Hernandez-Diaz S, Werler MM, Mitchell AA. Causal knowledge as a prerequisite for confounding evaluation: an application to birth defects epidemiology. *American journal of epidemiology*. 2002;155(2):176-184.
19. Lee PH. Should we adjust for a confounder if empirical and theoretical criteria yield contradictory results? A simulation study. *Scientific reports*. 2014;4:6085.

20. Jennrich RI, Schluchter MD. Unbalanced repeated-measures models with structured covariance matrices. *Biometrics*. 1986;42(4):805-820.
21. Ten Have TR, Kunselman AR, Pulkstenis EP, Landis JR. Mixed effects logistic regression models for longitudinal binary response data with informative drop-out. *Biometrics*. 1998;54(1):367-383.
22. Spira AP, Yager C, Brandt J, et al. Objectively Measured Sleep and beta-amyloid Burden in Older Adults: A Pilot Study. *SAGE open medicine*. 2014;2.
23. Sharma RA, Varga AW, Bubu OM, et al. Obstructive Sleep Apnea Severity Affects Amyloid Burden in Cognitively Normal Elderly: A Longitudinal Study. *American journal of respiratory and critical care medicine*. 2017.
24. Ju YE, Finn MB, Sutphen CL, et al. Obstructive sleep apnea decreases central nervous system-derived proteins in the cerebrospinal fluid. *Annals of neurology*. 2016;80(1):154-159.
25. Kalaria RN, Bhatti SU, Palatinsky EA, et al. Accumulation of the beta amyloid precursor protein at sites of ischemic injury in rat brain. *Neuroreport*. 1993;4(2):211-214.
26. Zhang X, Zhou K, Wang R, et al. Hypoxia-inducible factor 1alpha (HIF-1alpha)-mediated hypoxia increases BACE1 expression and beta-amyloid generation. *The Journal of biological chemistry*. 2007;282(15):10873-10880.
27. Hou H, Zhao Y, Yu W, et al. Association of obstructive sleep apnea with hypertension: A systematic review and meta-analysis. *Journal of global health*. 2018;8(1):010405.
28. Lurie A. Endothelial dysfunction in adults with obstructive sleep apnea. *Advances in cardiology*. 2011;46:139-170.

29. Nagayoshi M, Punjabi NM, Selvin E, et al. Obstructive sleep apnea and incident type 2 diabetes. *Sleep medicine*. 2016;25:156-161.
30. Mosconi L, Herholz K, Prohovnik I, et al. Metabolic interaction between ApoE genotype and onset age in Alzheimer's disease: implications for brain reserve. *Journal of neurology, neurosurgery, and psychiatry*. 2005;76(1):15-23.
31. Bero AW, Yan P, Roh JH, et al. Neuronal activity regulates the regional vulnerability to amyloid-beta deposition. *Nature neuroscience*. 2011;14(6):750-756.
32. Kiviniemi V, Wang X, Korhonen V, et al. Ultra-fast magnetic resonance encephalography of physiological brain activity - Glymphatic pulsation mechanisms? *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism*. 2016;36(6):1033-1045.
33. Schlosser Covell GE, Dhawan PS, Lee Iannotti JK, et al. Disrupted daytime activity and altered sleep-wake patterns may predict transition to mild cognitive impairment or dementia: a critically appraised topic. *The neurologist*. 2012;18(6):426-429.
34. Varga AW, Wohlleber ME, Gimenez S, et al. Reduced Slow-Wave Sleep Is Associated with High Cerebrospinal Fluid Abeta42 Levels in Cognitively Normal Elderly. *Sleep*. 2016;39(11):2041-2048.
35. Xie L, Kang H, Xu Q, et al. Sleep drives metabolite clearance from the adult brain. *Science (New York, NY)*. 2013;342(6156):373-377.
36. Pase MP, Himali JJ, Grima NA, et al. Sleep architecture and the risk of incident dementia in the community. *Neurology*. 2017;89(12):1244-1250.
37. Dreha-Kulaczewski S, Joseph AA, Merboldt KD, Ludwig HC, Gartner J, Frahm J. Identification of the Upward Movement of Human CSF In Vivo and its Relation to the

- Brain Venous System. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2017;37(9):2395-2402.
38. Green RC, Cupples LA, Kurz A, et al. Depression as a risk factor for Alzheimer disease: the MIRAGE Study. *Archives of neurology*. 2003;60(5):753-759.
39. Donohue MC, Sperling RA, Salmon DP, et al. The preclinical Alzheimer cognitive composite: measuring amyloid-related decline. *JAMA neurology*. 2014;71(8):961-970.
40. Edmonds EC, Bangen KJ, Delano-Wood L, et al. Patterns of Cortical and Subcortical Amyloid Burden across Stages of Preclinical Alzheimer's Disease. *Journal of the International Neuropsychological Society : JINS*. 2016;22(10):978-990.
41. Snowdon DA, Greiner LH, Mortimer JA, Riley KP, Greiner PA, Markesbery WR. Brain infarction and the clinical expression of Alzheimer disease. The Nun Study. *JAMA : the journal of the American Medical Association*. 1997;277(10):813-817.
42. Cirrito JR, Yamada KA, Finn MB, et al. Synaptic activity regulates interstitial fluid amyloid-beta levels in vivo. *Neuron*. 2005;48(6):913-922.
43. Nir Y, Staba RJ, Andrillon T, et al. Regional slow waves and spindles in human sleep. *Neuron*. 2011;70(1):153-169.
44. Ju YS, Zangrilli MA, Finn MB, Fagan AM, Holtzman DM. Obstructive sleep apnea treatment, slow wave activity, and amyloid-beta. *Annals of neurology*. 2018.
45. Hosselet J, Ayappa I, Norman RG, Krieger AC, Rapoport DM. Classification of sleep-disordered breathing. *American journal of respiratory and critical care medicine*. 2001;163(2):398-405.

46. Hosselet JJ, Norman RG, Ayappa I, Rapoport DM. Detection of flow limitation with a nasal cannula/pressure transducer system. *American journal of respiratory and critical care medicine*. 1998;157(5 Pt 1):1461-1467.
47. Ju YS, Ooms SJ, Sutphen C, et al. Slow wave sleep disruption increases cerebrospinal fluid amyloid-beta levels. *Brain : a journal of neurology*. 2017;140(8):2104-2111.
48. Osorio RS, Ayappa I, Mantua J, et al. The interaction between sleep-disordered breathing and apolipoprotein E genotype on cerebrospinal fluid biomarkers for Alzheimer's disease in cognitively normal elderly individuals. *Neurobiol Aging*. 2014;35(6):1318-1324.
49. Heneka MT, Carson MJ, El Khoury J, et al. Neuroinflammation in Alzheimer's disease. *Lancet neurology*. 2015;14(4):388-405.
50. Rosenzweig I, Glasser M, Polsek D, Leschziner GD, Williams SC, Morrell MJ. Sleep apnoea and the brain: a complex relationship. *The Lancet Respiratory medicine*. 2015;3(5):404-414.
51. Jack CR, Jr., Lowe VJ, Weigand SD, et al. Serial PIB and MRI in normal, mild cognitive impairment and Alzheimer's disease: implications for sequence of pathological events in Alzheimer's disease. *Brain : a journal of neurology*. 2009;132(Pt 5):1355-1365.
52. Osorio RS, Gumb T, Pirraglia E, et al. Sleep-disordered breathing advances cognitive decline in the elderly. *Neurology*. 2015;84(19):1964-1971.
53. Jack CR, Jr., Wiste HJ, Vemuri P, et al. Brain beta-amyloid measures and magnetic resonance imaging atrophy both predict time-to-progression from mild cognitive impairment to Alzheimer's disease. *Brain : a journal of neurology*. 2010;133(11):3336-3348.

54. Johnson KA, Fox NC, Sperling RA, Klunk WE. Brain imaging in Alzheimer disease. *Cold Spring Harbor perspectives in medicine*. 2012;2(4):a006213.
55. Van Den Berg JF, Van Rooij FJ, Vos H, et al. Disagreement between subjective and actigraphic measures of sleep duration in a population-based study of elderly persons. *Journal of sleep research*. 2008;17(3):295-302.
56. Unruh ML, Redline S, An MW, et al. Subjective and objective sleep quality and aging in the sleep heart health study. *Journal of the American Geriatrics Society*. 2008;56(7):1218-1227.
57. Young T, Palta M, Dempsey J, Skatrud J, Weber S, Badr S. The occurrence of sleep-disordered breathing among middle-aged adults. *The New England journal of medicine*. 1993;328(17):1230-1235.
58. Heinzer R, Marti-Soler H, Haba-Rubio J. Prevalence of sleep apnoea syndrome in the middle to old age general population. *The Lancet Respiratory medicine*. 2016;4(2):e5-6.
59. Carvalho DZ, St Louis EK, Knopman DS, et al. Association of Excessive Daytime Sleepiness With Longitudinal beta-Amyloid Accumulation in Elderly Persons Without Dementia. *JAMA neurology*. 2018.

Figure Captions

Figure 1: Stepwise exclusion process for NL, MCI and AD participants used in the study

Figure 2: Mean change variations & Trajectory Plots in AD-biomarker values over time in NL, MCI, and AD-subjects by OSA status (Cognitive Normal – NL)

Figure 3: Mean change variations & Trajectory Plots in AD-biomarker values over time in NL, MCI, and AD-subjects by OSA status (Mild Cognitive Impairment – MCI)

Figure 4: Mean change variations & Trajectory Plots in AD-biomarker values over time in NL, MCI, and AD-subjects by OSA status (Alzheimer's Disease – AD)

Figure 5: Proposed Mechanisms Linking OSA with Amyloid Deposition in Late-life

Supplemental Figures 1a-c: Baseline AD-Biomarker burden in NL, MCI and AD subjects by OSA status

Supplemental Figure 2a-c: MANOVA results showing mean change differences in AD-biomarker values over time by OSA status

Table 1: Descriptive Characteristics of Participants by Self-reported Obstructive Sleep Apnea Status at Baseline

Cognitive Normal			
Characteristics	All	OSA-	OSA+
Number of participants (%)	516 (100)	487 (94)	29 (6)
Female sex, number (%)	253 (49)	244 (50)	9 (32)
Age, years, median (interquartile range)	74 (71, 78)	71 (70, 78)	71 (70, 76)
ApoE4 positive, number (%)	145 (28)	140 (29)	5 (17)
Education, years, median (interquartile range)	16 (14, 18)	16 (14, 18)	16 (15, 18)
BMI (kg/m ²)	27.2 ± 4.8	27.1 ± 4.7	29.8 ± 5.9
Hypertension, number (%)	251 (49)	235 (48)	16 (55)
Diabetes, number (%)	45 (9)	40 (8)	5 (17)
Thyroid Disease, number (%)	112 (22)	109 (22)	3 (10)
Respiratory Disease, number (%)	123 (24)	100 (21)	23 (79)
Cardiovascular Disease, number (%)	358 (69)	338 (69)	20 (69)
TBI, number (%)	13 (3)	12 (3)	1 (3)
Alcohol, number (%)	23 (5)	22 (5)	1(3)
CPAP use	10 (19)		10 (35)
CSF- Amyloid Beta 42 pg/ml median (interquartile range)	210 (155, 241)	209 (155, 241)	226 (199, 259)
T-Tau pg/ml median (interquartile range)	59 (45, 84)	59 (45 83)	56 (48, 84)
P-Tau pg/ml median (interquartile range)	27 (20, 40)	27 (20, 42)	27 (20, 32)
Florbetapir SUVR, median (interquartile range) ^s	1.1 (1.0, 1.2)	1.1 (1.0, 1.2)	1.0 (1.0, 1.0)
MMSE median (interquartile range)	29 (28, 29)	29 (28, 30)	29 (27, 30)
CDR median (interquartile range)	0 (0, 0)	0 (0, 0)	0 (0, 0)
Mild Cognitive Impairment			
Characteristics	All	OSA-	OSA+
Number of participants (%)	798 (100)	695 (87)	103 (13)
Female sex, number (%)	319 (40)	294 (42)	25 (25)
Age, years, median (interquartile range)	74 (68, 79)	71 (70, 78)	71 (70, 76)
ApoE4 positive, number (%)	410 (51)	368 (53)	42 (41)

Education, years, median (interquartile range)	16 (14, 18)	16 (14, 18)	16 (14, 18)
BMI (kg/m2)	26.9 ± 4.6	26.5 ± 4.4	29.4 ± 5.3
Hypertension, number (%)	395 (49)	337 (48)	58 (56)
Diabetes, number (%)	75 (9)	57 (8)	18 (17)
Thyroid Disease, number (%)	156 (20)	133 (19)	23 (22)
Respiratory Disease, number (%)	195 (24)	133 (19)	62 (60)
Cardiovascular Disease, number (%)	577 (72)	490 (71)	87 (85)
TBI, number (%)	18 (2)	14 (2)	4 (4)
Alcohol, number (%)	35 (4)	31 (5)	4 (4)
CPAP use	16 (20)		16 (16)
CSF- Amyloid Beta 42 pg/ml median (interquartile range) ^s	153 (130, 209)	150 (128, 206)	169 (139, 214)
T-Tau pg/ml median (interquartile range) ^s	80 (54, 116)	81 (54, 122)	67 (51, 97)
P-Tau pg/ml median (interquartile range)	36 (23, 51)	37 (23, 52)	32 (21, 44)
Florbetapir SUVR, median (interquartile range) ^s	1.2 (1.0, 1.4)	1.2 (1.0, 1.4)	1.1 (1.0, 1.3)
MMSE median (interquartile range)	27 (24, 28)	28 (25, 29)	26 (25, 28)
CDR median (interquartile range)	0.5 (0.5, 0.5)	0.5 (0.5, 0.5)	0.5 (0.5, 0.5)

Alzheimer's Disease

Characteristics	All	OSA-	OSA+
Number of participants (%)	325 (100)	303 (93)	22 (7)
Female sex, number (%)	119 (37)	113 (37)	6 (27)
Age, years, median (interquartile range)	76 (71, 80)	76 (71, 80)	71 (64, 76)
ApoE4 positive, number (%)	216 (66)	198 (65)	18 (82)
Education, years, median (interquartile range)	16 (14, 18)	16 (14, 18)	16 (14, 18)
BMI (kg/m2)	25.9 ± 4.6	25.6 ± 4.3	29.0 ± 6.8
Hypertension, number (%)	165 (51)	149 (49)	16 (73)
Diabetes, number (%)	32 (10)	30 (10)	2 (9)
Thyroid Disease, number (%)	65 (20)	61 (20)	4 (18)
Respiratory Disease, number (%)	67 (21)	52 (17)	15 (68)
Cardiovascular Disease, number (%)	222 (68)	203 (67)	19 (86)
TBI, number (%)	18 (2)	14 (2)	4 (4)
Alcohol, number (%)	20 (6)	17 (6)	3 (14)
CPAP use	9 (3)		9 (41)
CSF-Amyloid Beta 42 pg/ml median (interquartile range) ^s	132 (116, 151)	115 (96, 182)	113 (79, 151)
T-Tau pg/ml median (interquartile range)	80 (54, 116)	67 (51, 97)	81 (54, 122)
P-Tau pg/ml median (interquartile range)	42 (33, 61)	41 (33, 61)	54 (35, 66)
Florbetapir SUVR, median (interquartile range)	1.4 (1.3, 1.5)	1.4 (1.3, 1.5)	1.4 (1.3, 1.6)

MMSE median (interquartile range)	24 (20, 26)	24 (22, 26)	24 (20, 25)
CDR median (interquartile range)	2.0 (1.0, 2.5)	1.5 (1.0, 2.5)	2.0 (1.0, 2.5)

Abbreviation: A β : amyloid beta, ApoE4: Apolipoprotein epsilon4, BMI: body mass index, CDR: Clinical Dementia Rating; CPAP: continuous pulmonary airway pressure; CSF: cerebrospinal fluid, MMSE: Mini Mental State Exam; TAU: tau protein, PTAU: phosphorylated tau, TBI: traumatic brain injury. Superscript s: indicates significant differences between groups (CSF A β 42 levels for both MCI (F=4.37, p =.04), and AD (F=6.89, p <.01 and T-tau levels for the MCI patients (F=5.08, p =.02)

Accepted Manuscript

Table 2: Multivariate ANOVA Results Testing Biomarker Mean Change Differences in Time Trend, Groups and Time Points

MANOVA Test Criteria	Statistic	Value	F-value	P-value	MANOVA Test Criteria	Statistic	Value	F-value	P-value
Florbetapir SUVR					CSF Aβ42				
Normal Cognition (NL)					Normal Cognition (NL)				
Timepoint*OSA Effect	Pillai's Trace	0.035	4.46	0.0125*	Timepoint*OSA Effect	Pillai's Trace	0.122	5.23	0.0075**
Timepoint Effect	Pillai's Trace	0.162	23.66	<.0001***	Timepoint Effect	Pillai's Trace	0.579	51.77	<.0001***
Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-					Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-				
Timepoint_1 (Year 2)			0.14	0.7125	Timepoint_1 (Year 2)			9.99	0.0023**
timepoint_2 (Year 3)			8.93	0.0031**	timepoint_2 (Year 3)			4.12	0.046*
timepoint_1 vs. timepoint_0			34.22	<.0001***	timepoint_1 vs. timepoint_0			69.4	<.0001***
timepoint_2 vs. timepoint_1			6.51	0.0113*	timepoint_2 vs. timepoint_1			8.08	0.0006***
Mild Cognitive Impairment (MCI)					Mild Cognitive Impairment (MCI)				
Timepoint*OSA Effect	Pillai's Trace	0.047	4.24	0.0143*	Timepoint*OSA Effect	Pillai's Trace	0.219	5.06	0.0116*
Timepoint Effect	Pillai's Trace	0.129	18.06	<.0001***	Timepoint Effect	Pillai's Trace	0.791	68.23	<.0001***
Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-					Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-				
Timepoint_1 (Year 2)			0.08	0.7786	Timepoint_1 (Year 2)			4.95	0.0323*
timepoint_2 (Year 3)			4.63	0.0323*	timepoint_2 (Year 3)			7.38	0.01*

timepoint_1 vs. timepoint_0		25.63	<.0001***	timepoint_1 vs. timepoint_0		132.92	<.0001***		
timepoint_2 vs. timepoint_1		6.16	0.0137*	timepoint_2 vs. timepoint_1		13.12	0.0009***		
Alzheimer's Disease (AD)				Alzheimer's Disease (AD)					
Timepoint*OSA Effect	Pillai's Trace	0.016	1.94	0.1457	Timepoint*OSA Effect	Pillai's Trace	0.092	2.48	0.0944
Timepoint Effect	Pillai's Trace	0.207	31.86	<.0001***	Timepoint Effect	Pillai's Trace	0.933	341.26	<.0001***
Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-				Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-					
Timepoint_1 (Year 2)		0.64	0.4228	Timepoint_1 (Year 2)		3.78	0.0575		
timepoint_2 (Year 3)		3.39	0.067	timepoint_2 (Year 3)		1.16	0.2873		
timepoint_1 vs. timepoint_0		3.39	0.067	timepoint_1 vs. timepoint_0		1.16	0.2873		
timepoint_2 vs. timepoint_1		0.55	0.4593	timepoint_2 vs. timepoint_1		0.12	0.7253		
CSF TAU				CSF PTAU					
Normal Cognition (NL)				Normal Cognition (NL)					
Timepoint*OSA Effect	Pillai's Trace	0.096	3.99	0.0226*	Timepoint*OSA Effect	Pillai's Trace	0.142	4.06	0.0235*
Timepoint Effect	Pillai's Trace	0.302	16.24	<.0001***	Timepoint Effect	Pillai's Trace	0.545	29.29	<.0001***
Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-				Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-					
Timepoint_1 (Year 2)		3.1	0.083	Timepoint_1 (Year 2)		5.74	0.0204*		
timepoint_2 (Year 3)		8.04	0.0006***	timepoint_2 (Year 3)		5.74	0.0203*		
timepoint_1 vs.		32.9	<.0001***	timepoint_1 vs.		53.39	<.0001***		

timepoint_0					timepoint_0				
timepoint_2 vs. timepoint_1		17.34	<.0001***		timepoint_2 vs. timepoint_1		3.64	0.0621	
Mild Cognitive Impairment (MCI)					Mild Cognitive Impairment (MCI)				
Timepoint*OSA Effect	Pillai's Trace	0.198	4.44	0.0189*	Timepoint*OSA Effect	Pillai's Trace	0.272	6.73	0.0033**
Timepoint Effect	Pillai's Trace	0.321	8.46	0.001**	Timepoint Effect	Pillai's Trace	0.762	57.72	<.0001***
Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-					Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-				
Timepoint_1 (Year 2)		1.13	0.2957		Timepoint_1 (Year 2)		12.72	0.0001***	
timepoint_2 (Year 3)		9.1	0.0046**		timepoint_2 (Year 3)		4.2	0.0474*	
timepoint_1 vs. timepoint_0		17.39	0.0002***		timepoint_1 vs. timepoint_0		62.45	<.0001***	
timepoint_2 vs. timepoint_1		12.9	0.001**		timepoint_2 vs. timepoint_1		3.89	0.056	
Alzheimer's Disease (AD)					Alzheimer's Disease (AD)				
Timepoint*OSA Effect	Pillai's Trace	0.114	4.83	0.0106*	Timepoint*OSA Effect	Pillai's Trace	0.074	1.63	0.2086
Timepoint Effect	Pillai's Trace	0.751	112.37	<.0001***	Timepoint Effect	Pillai's Trace	0.939	312.8	<.0001***
Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-					Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA-				
Timepoint_1 (Year 2)		3.42	0.0684		Timepoint_1 (Year 2)		3.23	0.0797	
timepoint_2 (Year 3)		8.68	0.0043**		timepoint_2 (Year 3)		0.09	0.7622	
timepoint_1 vs. timepoint_0		8.68	0.0043**		timepoint_1 vs. timepoint_0		0.09	0.7622	
timepoint_2 vs.		0.52	0.4712		timepoint_2 vs.		2.24	0.1423	

timepoint_1			timepoint_1		
-------------	--	--	-------------	--	--

SUVr: Standard Uptake Value Ratio of Brain Florbetapir; CSF: Cerebrospinal fluid; PTAU: phosphorylated TAU. MANOVA: Multivariate Analysis of Variance. MANOVA test criteria interpretations: Timepoint*OSA Effect provides the exact F statistics for the time trend of mean change in Alzheimer's disease biomarker across OSA groups; Timepoint Effect provides the exact F statistics for the mean change (increase or decrease) in Alzheimer's disease biomarker over time. Repeated Measures Analysis of Variance of Contrast Variables OSA+ vs. OSA- interpretations: Timepoint_1 (Year 2) provides the F statistic for the difference in mean change in Alzheimer's disease biomarker across OSA groups from baseline (Year 1) to Year 2; Timepoint_2 (Year 3) provides the F statistic for the difference in mean change in Alzheimer's disease biomarker across OSA groups from baseline (Year 1) to Year 3.; timepoint_1 vs. timepoint_0 provides the F statistic for differences in mean change in Alzheimer's disease biomarker across OSA groups when timepoint 1 (Year 2) is compared to baseline (Year 1); timepoint_2 vs. timepoint_1 provides the F statistic for differences in mean change in Alzheimer's disease biomarker across OSA groups when timepoint 2 (Year 3) is compared to timepoint 1 (Year 2). **P-value* <=.05; ***P-value* <=.01; ****P-value* <=.001

Accepted Manuscript

Table 3 Between Subject Variation in AD Biomarkers and Covariance Parameter Estimates

Parameters	Estimate	95% CI	P-value
OSA+ vs. OSA- (Cognitive Normal Patients)			
Florbetapir SUVR over time	0.06	.02, .11	<.0001
Florbetapir SUVR over time (covariance)	-0.06	-.09, -.04	<.0001
CSF A β -42 over time	-2.71	-3.11, -2.35	<.0001
CSF A β -42 over time (covariance)	3.93	3.56, 4.31	<.0001
CSF TTAU over time	3.68	3.31, 4.07	<.0001
CSF TTAU over time (covariance)	-2.89	-3.51, -2.29	<.0001
CSF PTAU over time	1.22	1.02, 1.42	<.0001
CSF PTAU over time (covariance)	-1.21	-1.71, -.74	<.0001
OSA+ vs. OSA- (Mild Cognitive Impairment Patients)			
Florbetapir SUVR over time	0.08	.05, .12	<.0001
Florbetapir SUVR over time (covariance)	-0.06	-.09, -.04	<.0001
CSF A β -42 over time	-2.62	-3.23, -2.03	<.0001
CSF A β -42 over time (covariance)	2.69	2.02, 3.36	<.0001
CSF TAU volume over time	2.21	1.58, 2.86	<.0001
CSF TAU over time (covariance)	-1.89	-2.91, -.87	<.0001
CSF PTAU over time	1.74	1.22, 2.27	<.0001
CSF PTAU over time (covariance)	-1.48	-2.05, -.94	<.0001
OSA+ vs. OSA- (Alzheimer's disease Patients)			
Florbetapir SUVR over time	0.07	-1.19, 1.33	0.33
Florbetapir SUVR over time (covariance)	-0.29	-2.07, 1.49	0.31
CSF A β -42 over time	-1.11	-3.31, 1.09	0.53
CSF A β -42 over time (covariance)	-1.14	-3.38, 1.63	0.56
CSF TTAU over time	0.26	-1.02, 1.28	0.47
CSF TTAU over time (covariance)	-0.15	-1.94, 1.64	0.47
CSF PTAU over time	0.94	0.23, 1.65	0.11
CSF PTAU over time (covariance)	-0.16	-1.66, 1.34	0.11

These analyses allowed us to examine whether there was significant variation between OSA+ and OSA- subjects in mean AD biomarker level at baseline, as well as whether significant variation in the change in AD biomarker level over time occurred. Furthermore, the covariance between the baseline AD biomarker level and AD biomarker change over time indicated whether OSA+ or OSA- subjects had experienced a faster increase or decrease in AD biomarker level over time. It also allowed for assessment of significant differences in the rate-of-change in AD biomarker level between OSA groups over time.

Abbreviations: A β : amyloid beta, ApoE4: Apolipoprotein epsilon4, BMI: body mass index, CSF: cerebrospinal fluid, TAU: tau protein, PTAU: phosphorylated tau. Models were adjusted for age, sex, BMI, education, CPAP-use, ApoE4 status, alcohol intake, baseline biomarker data, history of respiratory disease, hypertension, diabetes and history of cardiovascular disease (e.g. including ischemic heart disease, heart failure, and stroke/TIA), and history of traumatic brain injury

Figure 1: ADNI: Alzheimer's Disease Neuroimaging Initiative; NL: Cognitive Normal; MCI: Mild Cognitive Impairment; AD: Alzheimer's disease

Figure 2: The MANOVA (Multivariate Analysis of Variance) analysis allowed assessment of whether mean changes in AD-biomarker levels were parallel across OSA-groups, whether the mean change in biomarker-data increased/decreased over time, and whether there are significant differences in mean biomarker-data across OSA-groups for all time-points. Blue line: OSA-; Red line: OSA+; NL: Cognitive Normal; SUVR: Standard Uptake Value Ratio of Brain Florbetapir; SUVR_COM, SUVR_COM2 and SUVR_COM3: Mean change Brain Florbetapir Uptake Value at time 1, 2 & 3 respectively; CSF: Cerebrospinal fluid; ABETA: amyloid beta; ABETA1_new, ABETA2_new, ABETA3_new: Mean change CSF Aβ42 at time 1, 2 & 3 respectively. TAU1_new, TAU2_new, TAU3_new: Mean change TAU at time 1, 2 & 3 respectively; PTAU1_new, PTAU2_new, PTAU3_new: Mean change phosphorylated TAU at time 1, 2 & 3 respectively. Time 1 is baseline; Time 1, 2 & 3 values obtained by subtracting baseline data at each time point

Figure 3: The MANOVA (Multivariate Analysis of Variance) analysis allowed assessment of whether mean changes in AD-biomarker levels were parallel across OSA-groups, whether the mean change in biomarker-data increased/decreased over time, and whether there are significant differences in mean biomarker-data across OSA-groups for all time-points. Blue line: OSA-; Red line: OSA+; MCI: Mild Cognitive Impairment; SUVR: Standard Uptake Value Ratio of Brain Florbetapir; SUVR_COM, SUVR_COM2 and SUVR_COM3: Mean change Brain Florbetapir Uptake Value at time 1, 2 & 3 respectively; CSF: Cerebrospinal fluid, ABETA: amyloid beta; ABETA1_new, ABETA2_new, ABETA3_new: Mean change CSF Aβ42 at time 1, 2 & 3 respectively. TAU1_new, TAU2_new, TAU3_new: Mean change TAU at time 1, 2 & 3 respectively; PTAU1_new, PTAU2_new, PTAU3_new: Mean change phosphorylated TAU at time 1, 2 & 3 respectively. Time 1 is baseline; Time 1, 2 & 3 values obtained by subtracting baseline data at each time point

Figure 4: The MANOVA (Multivariate Analysis of Variance) analysis allowed assessment of whether mean changes in AD-biomarker levels were parallel across OSA-groups, whether the mean change in biomarker-data increased/decreased over time, and whether there are significant differences in mean biomarker-data across OSA-groups for all time-points. Blue line: OSA-; Red line: OSA+; AD: Alzheimer's disease; SUVR: Standard Uptake Value Ratio of Brain Florbetapir; SUVR_COM, SUVR_COM2 and SUVR_COM3: Mean change Brain Florbetapir Uptake Value at time 1, 2 & 3 respectively; CSF: Cerebrospinal fluid, ABETA: amyloid beta; ABETA1_new, ABETA2_new, ABETA3_new: Mean change CSF A β 42 at time 1, 2 & 3 respectively. TAU1_new, TAU2_new, TAU3_new: Mean change TAU at time 1, 2 & 3 respectively; PTAU1_new, PTAU2_new, PTAU3_new: Mean change phosphorylated TAU at time 1, 2 & 3 respectively. Time 1 is baseline; Time 1, 2 & 3 values obtained by subtracting baseline data at each time point

Fig. 5. Possible intermediate mechanisms in the relationship between OSA and AD. Chronic exposure to intermittent hypoxia may lead to increased inflammation and oxidative stress, diabetes, hypertension and CVD, all potentially contributing to AD pathology development. Sleep fragmentation, both by itself and by leading to decreased REM and SWS stages, can additionally promote AD pathogenesis. Intrathoracic pressure swings associated with OSA may disrupt CSF-ISF exchange integrity and lead to AD neuropathology accumulation. Lastly, depression may also contribute to this association. OSA, obstructive sleep apnea; CVD, cardiovascular disease; REM, rapid eye movement; SWS, slow wave sleep; CSF-ISF, cerebrospinal fluid-interstitial fluid; AD, Alzheimer's disease.