



King's Research Portal

DOI:

[10.1016/j.earlhumdev.2018.11.004](https://doi.org/10.1016/j.earlhumdev.2018.11.004)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Dassios, T., Hickey, A., Krokidis, M., & Greenough, A. (2019). Congenital diaphragmatic hernia in newborn infants: Variable endotracheal tube and umbilical venous catheter positions. *Early Human Development*, 128, 12-14. <https://doi.org/10.1016/j.earlhumdev.2018.11.004>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Congenital diaphragmatic hernia in newborn infants: Variable endotracheal tube and umbilical venous catheter positions

Theodore Dassios PhD^{1,2}, Ann Hickey MBBs¹, Miltiadis Krokidis PhD³, Anne Greenough MD^{2,4,5}

¹ Neonatal Intensive Care Centre, King's College Hospital NHS Foundation Trust, London, UK; ² Women and Children's Health, School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London, UK; ³ Department of Radiology, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK; ⁴ MRC-Asthma UK Centre in Allergic Mechanisms of Asthma, King's College London, UK; ⁵ National Institute for Health Research (NIHR) Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London; UK

Address for Correspondence: Dr Theodore Dassios, NICU, 4th Floor Golden Jubilee Wing, King's College Hospital, Denmark Hill, London, SE5 9RS, Tel: 0203 299 4644; fax: 0203 299 8284; email: theodore.dassios@kcl.ac.uk

Short title: Endotracheal tube and umbilical venous catheters in CDH

Key words: chest and abdomen radiograph, neonatal radiography

Funding source: The research was supported by the National Institute for Health Research (NIHR) Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

Conflict of interest: None to declare.

Role of the study sponsor: No involvement.

Dr Theodore Dassios wrote the first draft of the manuscript. No honorarium, grant, or other form of payment was given to anyone to produce the manuscript.

Data sharing statement: We agree to sharing data on request.

LIST OF ABBREVIATIONS

CDH Congenital diaphragmatic hernia

ETT Endotracheal tube

UVC Umbilical venous catheter

In fetuses with congenital diaphragmatic hernia (CDH), the herniation of abdominal contents in the chest alters the relative position of the thoracic and abdominal organs (1) and could give rise to an unusual radiographic appearance of the indwelling support apparatus [endotracheal tube (ETT), umbilical venous catheters (UVC)] (2). In this study we reviewed the radiographs of infants with CDH and reported the position of the ETT and the UVC. The preoperative chest and abdominal radiographs of infants born with CDH at King's College Hospital (KCH), London, UK were evaluated by a Consultant Radiologist (MK) and a Consultant Neonatologist (TD). The deviation of the ETT and UVC from the midline was measured with the Sectra PACS software (Sectra AB, Linköping, Sweden) at the level of the second-third thoracic vertebra for the ETT and at the level of the tenth thoracic vertebra for the UVC (figure 1). Rotated radiographs were excluded from the study. The insertion of the ETT in the trachea was confirmed with end-tidal carbon dioxide (ETCO₂) monitoring. The position of the umbilical artery catheter was not measured as it has been reported to be universally normal (2). The study was registered as a service evaluation with the Clinical Governance Department of KCH. From 1 January 2006 to 31 July 2018, 143 infants with CDH were treated at KCH. Twelve infants were excluded because of rotated radiographs or lack of images. One hundred and thirty-one (77 male) infants with a median (IQR) age of 36 (34-39) weeks (range 25-41 weeks) were included in the study. They had a median (IQR) birth weight of 2.68 (2.20-3.14) kg (range 0.75-4.01 kg). **Infants with left-sided CDH (n=116, 89%):** the ETT deviated to the right in 104 (90%) with a median (IQR) distance from the midline of 7 (6-10) mm (range 1-11mm), deviated to the left in six (5%) infants with a median (range) distance from the midline of 2 (1-14) mm and in another six (5%) infants the ETT did not deviate from the midline. The UVC deviated to the left in 86 (74%) with a median (IQR) distance from the midline of 16 (10-24) mm (range 1-45 mm), deviated to the right in 25 (22%) with a median (IQR) distance from the midline of 10 (3-14) mm (range 1-

30 mm) and in 5 (4%) infants the UVC did not deviate from the midline. **Infants with right-sided CDH (n=15, 11%):** the ETT deviated to the left in 12 (80%) with a median (IQR) distance from the midline of 3 (1-6) mm (range 1-8mm), deviated to the right in one (7%) infant with a distance from the midline of 1 mm and in another two (13%) infants the ETT did not deviate from the midline. The UVC deviated to the right in 12 (80%) with a median (IQR) distance from the midline of 14 (8-18) mm (range 1-37 mm), deviated to the left in two (13%) (17 and 19 mm) and in one (1%) infant the UVC did not deviate from the midline.


In conclusion, we report that in infants with CDH there is a wide variation in the radiographical position of the ETT and the UVC and the ETT or UVC can be found on either side of the midline. Predominantly, in left-sided CDH the umbilical vein is deviated to the left and the trachea to the right. Conversely, in right-sided CDH the umbilical vein is usually deviated to the right and the trachea to the left. As our results show, however, this is not a universal finding and there is significant variation in the magnitude of deviation. This knowledge is clinically important so that catheters are not removed unnecessarily as affected infants are commonly severely unwell and require central venous access, (3) but equally we hope these data may reduce reintubation or at least promote the use of ETCO₂ monitoring to identify correct ETT placement (4).

REFERENCES

1. Levin TL, Liebling MS, Ruzal-Shapiro C, Berdon WE, Stolar CJ. Midgut malfixation in patients with congenital diaphragmatic hernia: what is the risk of midgut volvulus? *Pediatric radiology*. 1995;25(4):259-61. PubMed PMID: 7567230.
2. Sakurai M, Donnelly LF, Klosterman LA, Strife JL. Congenital diaphragmatic hernia in neonates: variations in umbilical catheter and enteric tube position. *Radiology*. 2000 Jul;216(1):112-6. PubMed PMID: 10887235.
3. Snoek KG, Reiss IK, Greenough A, Capolupo I, Urlesberger B, Wessel L, et al. Standardized Postnatal Management of Infants with Congenital Diaphragmatic Hernia in Europe: The CDH EURO Consortium Consensus - 2015 Update. *Neonatology*. 2016;110(1):66-74. PubMed PMID: 27077664.
4. Hunt KA, Yamada Y, Murthy V, Srihari Bhat P, Campbell M, Fox GF, et al. Detection of exhaled carbon dioxide following intubation during resuscitation at delivery. *Archives of disease in childhood Fetal and neonatal edition*. 2018 Mar 17. PubMed PMID: 29550769.

FIGURE LEGEND

Figure 1: A chest and abdominal radiograph of an infant with a left-sided CDH taken on the first day of life. The measurements of the distance of the ETT and the UVC from the midline are schematically presented along with the median values of the study cohort with left-sided CDH.



7mm

16mm