King's Research Portal

DOI:

[10.1080/09637486.2019.1600664](https://doi.org/10.1080/09637486.2019.1600664)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Gill, S. K., Lever, E., Emery, P. W., & Whelan, K. (2019). Nutrient, fibre, sorbitol and chlorogenic acid content of prunes (*Prunus domestica*): an updated analysis and comparison of different countries of origin and database values. *International Journal of Food Sciences and Nutrition*, 70(8), 924-931.
<https://doi.org/10.1080/09637486.2019.1600664>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

1 **Nutrient, fibre, sorbitol and chlorogenic acid content of prunes**
2 **(*Prunus domestica*): an updated analysis and comparison of**
3 **different countries of origin and database values**
4

5 **Samantha K Gill*, Ellen Lever*, Peter W Emery, Kevin Whelan**

6 King's College London, Department of Nutritional Sciences, London, United Kingdom
7

8 Corresponding author: Professor Kevin Whelan, King's College London, Department of
9 Nutritional Sciences, 150 Stamford Street, London, SE1 9NH, United Kingdom, + 44 20 78 48
10 38 58, kevin.whelan@kcl.ac.uk
11

12 Short running head: Prune composition

13 Key words: dried fruit, dried plums, prunes, gastrointestinal health, fibre, polyphenols
14

15 *Joint first authors
16

17 **Disclosure of interest:** This study was funded by the California Dried Plum Board. The funders
18 played no role in study conception, design, analysis, interpretation or writing of the
19 manuscript. KW has served as a consultant for Danone, has received speaker fees from Alpro
20 and the National Dried Fruit Trade Association UK Ltd and has received research funding from
21 Almond Board of California, California Dried Plum Board, Clasado Biosciences, Nestec Ltd,
22 International Nut and Dried Fruit Council, and receives royalties from FoodMaestro. The
23 remaining authors report no conflicts of interest.

Abstract

Current prune composition data is outdated and requires a comprehensive and comparative re-analysis. This novel study aimed to: (i) analyse and compare prune composition from major countries of origin; and (ii) provide a comprehensive compositional analysis of prunes of USA origin and compare this with UK and USA database data. Prune samples were analysed for major nutrients and bioactive compounds and compared between countries of origin. Total fibre was higher in prunes from the USA (12.0 g/100g) and Chile (11.5 g/100g) compared with France (8.4 g/100g) and Argentina (8.9 g/100g), while prunes from all countries contained high levels of sorbitol (11.2-15.5 g/100g). Differences of energy and starch values compared with national databases reflected different approaches to sampling and analysis. In conclusion, prunes contain high levels of fibre and other bioactive compounds. Variations between country of origin and database values highlight the importance of transparency in documenting sampling and analysis methods.

Introduction

Studies have highlighted the potential benefits of dried fruits on a variety of health outcomes (Chang et al. 2016). In particular their high fibre content has led to investigation of the role of dried fruit in the maintenance and promotion of gastrointestinal health (Lever et al. 2015) which is considered of major public health importance (DuBois 2004; Wald et al. 2007). The impact of dietary fibre on health is affected by variations in its chemical composition (e.g. distribution of different fibre fractions) and physical structure (e.g. degree of polymerisation, molecular weight and linkages) that alter its solubility, viscosity and fermentability. Given that dried fruits are nutritionally comparable to whole fresh fruits, only provided in a smaller and more concentrated form, they may be a convenient and versatile option for increasing fruit consumption across population groups (Sadler et al. 2019).

Plums are taxonomically diverse stone fruits of *Prunus domestica* L. and are commonly consumed in their dried form, termed prunes. Data from various sources, including the United Kingdom (Finglas 2015) and the United States of America (USA) (US Department of Agriculture 2018), indicate that prunes are naturally high in a variety of poorly-fermented and readily-fermented dietary fibres (>6 g/100g including hemicellulose, pectin, cellulose). In addition, prunes contain other bioactive compounds such as polyphenols, which may stimulate colonic proliferation of microorganisms such as Bifidobacteria and Lactobacilli. Furthermore, prunes contain high amounts of sorbitol (~12 g/100g) which is known to have laxative effects (Yao et al. 2014). Indeed, a systematic review concluded that prunes may play a role in gastrointestinal health by increasing stool frequency and improving stool consistency (Lever et al. 2014).

Data on prune composition require updating for several reasons. Firstly, existing databases, such as McCance and Widdowson (UK) and USDA (USA) databases, do not report a wide range of components relevant to gut health (e.g. different fibre fractions and sorbitol content), and secondly the current data were compiled between 1980 and 2001 and therefore may no longer accurately reflect present-day prune composition. Thirdly, the USDA data calculates total carbohydrate 'by difference', which does not account for the lower energy contributions from unavailable carbohydrates. Fourthly, the composition of prunes may vary depending on

a variety of factors including growing and harvesting conditions and post-harvest processes (e.g. drying, dehydration and rehydration, storage conditions). Given that the vast majority of global supply of prunes originates from four countries: the USA (largely California, 43%), Chile (24%), France (16%) and Argentina (15%) (Buncher 2012), currently-available prune composition data may be confounded by variations in origin. For example, standard yellow plums have been shown to contain higher vitamin and phenolic compound content than organically grown plums (Lombardi-Boccia et al. 2004), while prunes from Australia have been shown to contain higher iron and folate contents than prunes from USA and Chile (Bennett et al. 2011). Finally, variations in the nutrient composition of prunes of different countries of origin will impact the database values in each country. For example, databases in the USA (USDA) and France (CIQUAL), both of whom are large producers of prunes, will reflect the composition of prunes from those countries, whereas the database in the UK (McCance and Widdowson), which does not grow large supplies of prunes, will reflect the composite of prunes from different countries of origin.

With this in mind, we aimed to investigate the energy, macronutrient, micronutrient, fibre, sorbitol and polyphenol composition of prunes by: (i) analysing and comparing the composition of prunes from major countries of origin (USA, Chile, France and Argentina); and (ii) undertaking an in-depth analysis of prunes of USA (Californian) origin and comparing this with data from food composition databases.

Materials and Methods

Sample collection

Prune samples grown by the four largest producers of prunes were collected in order that composition could be both globally representative and compared between country of origin (USA, Chile, France and Argentina). Prune samples were purchased from major population centres in five countries across Europe (France, Germany, Italy, Spain, United Kingdom) as these are five major European markets for prunes and thus data would reflect the composition of prunes available across Europe, as well as meeting the Food Information to Consumers Legislation (European Commission 2011). Prune samples were purchased as sold to the customer from major retail outlets including supermarkets, department stores and health food stores and including a range of brands (where available) to ensure purchase of

prunes representative of those most frequently consumed, i.e. with the highest volume of sales. Prune samples were purchased at the same time of year and within use-by-dates. Samples were required to be in unopened packets of ≥ 100 g with a remaining shelf life of ≥ 6 months. Prunes were stored unopened until analysis to minimise drying, water absorption and contamination. Prunes were purchased pitted (stone removed) or whole. If purchased whole, stones were removed prior to compositional analysis.

In total, the goal was to purchase 10 to 12 samples from each of the four countries of origin, with at least 3 samples from each sampling country. This number is recommended for nutrient composition database data and based upon guidance from Greenfield and Southgate (2003), though this depends on the variability of the nutrients being measured.

Prune samples from each country of origin were pooled prior to analysis. Funding restrictions meant that the study could either: (i) individually analyse a number of prune samples from a single country, thus allowing measurement of within-country variation but not between-country variation; or (ii) analyse a pooled sample from a number of sampling countries, thus allowing measurement of between-country variation albeit without statistical comparison. Given the wide geographic difference in countries of origin (USA, France, Chile, Argentina), it was felt that between-country variations, rather than within-country variations, were likely to be larger and therefore of greater nutritional relevance.

Sample preparation and analysis

Samples were pooled according to country of origin (**Table 1**). This pooled sample comprised an equal weight of 500 g (i.e. 100 g adjusted weight from each sampling country), of prunes from each of the five sampling countries. Pooled samples were homogenised using a hand mincer, divided into aliquots, stored frozen at -80°C and defrosted prior to analysis.

Prune samples were analysed at Leatherhead Food Research, Surrey, UK. The pooled sample from each of the four countries of origin was analysed using standard methods for energy (calculated from macronutrient data), protein (total nitrogen), fat (Soxhlet), carbohydrate (calculated by difference), sugars and sorbitol (ion-exchange chromatography), a range of fibre classifications (AOAC methods, Englyst), and chlorogenic and neochlorogenic acid (ultra-

performance liquid chromatography tandem mass spectrometry, UPLC with MS-MS) (**Table 2**). In addition, further in-depth analyses were performed on the pooled sample of prunes from the largest global producer (California, USA) including fatty acids (gas chromatography with flame ionisation detection), sugars (ion-exchange chromatography) and major micronutrients (inductively coupled plasma optical emission spectrometry ICP-OES, high performance liquid chromatography HPLC) (**Table 2**).

In terms of the chromatographic methods, for sugar and sorbitol, extraction from the prune samples was performed by sonication in hot water and treatment with Carrez reagents. The filtered solution was then analysed using high-performance anion-exchange chromatography coupled with pulsed electrochemical detection (HPAEC-PED) using a Dionex PA20 column (Corradini et al, 2012). For chlorogenic and neochlorogenic acids, extraction from the prune samples was performed in hot water and methanol and the solution analysed using UPLC with MS-MS equipped with an ethylene bridged hybrid column (C18 2.1 x 50 mm, 1.7 µm). For fatty acids, transmethylation was undertaken to form methyl esters which were analysed using gas-liquid chromatography with a flame ionisation detection (Seppänen-Laakso, et al, 2002). For vitamin B analysis, extraction was performed using HCl and the solution analysed using HPLC with fluorescence detection using a C18 conventional column (250 x 4.6 mm, 5 µm).

Duplicate analyses were carried out for analytes that were not routinely measured at the research centre. However, routine analyses were not performed in duplicate as these had criteria defining the limits of repeatability.

Results

Sample purchases

Sample purchases were made in France (Normandy), Germany (Bonn), Italy (Milan, Novara), Spain (Madrid), and the UK (London) between March and June 2013. The pack sizes of the purchased prune samples varied between 120 g and 1000 g. **Table 1** shows the number of prune samples purchased and analysed from each sampling country and by country of origin. One sample was excluded as it exceeded the use-by-date by the time of analysis and four

samples were excluded because the country of origin was unclear. Eighteen different samples were pooled and analysed for USA and French prunes, fifteen for Chilean prunes, but only five for Argentinian prunes (all purchased from Spain) due to their lack of availability in Europe at that time. The amount analysed from each individual sample was weighted so that an equal amount from each sampling country was included and pooled to make up a total of 500g from each country of origin (**Table 1**).

Composition Data

The composition of prunes from the pooled samples originating from USA, Chile, France and Argentina are shown in **Table 2**. In general there were few major differences in nutrients and fibre fractions between prunes of different countries of origin.

Differences in starch content were observed between countries, being lower in prunes of French origin (1.9 g/100 g) compared with others (5.7-6.6 g/100g). Total fibre (measured using AOAC 2011.25) was higher in prunes from the USA (12.0g/100g) and Chile (11.5g/100g) compared with those from France (8.4g/100g) and Argentina (8.9g/100g).

Prunes contained high levels of sorbitol (11.2-15.5g/100g) with broadly similar values across the countries of origin. High levels of the phenolic compounds chlorogenic acid and neochlorogenic acid were also found in prunes, however, in general these were higher in prunes from the USA (3.6 and 89.3 g/100g) and France (3.9 and 92.0 g/100g) compared with prunes from Chile (1.3 and 39.8 g/100g) and Argentina (1.3 and 40.3 g/100g) (**Table 2**).

The composition of prunes from the USA pooled sample compared with data from the USDA nutrient database and McCance and Widdowson's the composition of foods is shown in **Table 3**. Concentrations per wet weight are presented in order to be consistent with these databases. Energy and starch values (230 kcals/100g and 6.6g/100g, respectively) were closer to values published by the USDA database (240 kcals/100g and 5.1 g/100g, respectively), than McCance and Widdowson (141 kcals/100g and 0.0 g/100g, respectively).

Discussion

The current study aimed to investigate the energy, macronutrient, micronutrient, fibre, sorbitol and polyphenol composition of prunes by: (i) analysing and comparing the composition of prunes from major countries of origin (USA, Chile, France and Argentina); and (ii) undertaking an in-depth analysis of prunes of USA (Californian) origin and comparing this with data from food composition databases.

In regards to the measured composition of prunes from different countries of origin, while the pooling of samples precluded statistical comparisons, at face value there were few differences in energy and macronutrient content between them, apart from starch which was lower in prunes of French origin (**Table 2**). Given that the same methods of analysis were used for prunes of all countries of origin and analysis occurred at the same time and in the same run, these differences likely reflect true compositional differences in prune samples between countries.

There were no major differences in dietary fibre content when measured using the AOAC 991.43 method (which excludes low molecular weight fibres and most types of resistant starch) nor using the Englyst method (non-starch polysaccharides). However, when measured using the AOAC 2011.25 method, fibre was higher in prunes from USA and Chile compared with France and Argentina. The AOAC 2011.25 method includes all categories of dietary fibre, high and low molecular weight fibres and all types of resistant starch. Taken together, this suggests that US and Chilean prunes likely contain greater low molecular weight fibres and resistant starch than French and Argentinian prunes. Low molecular weight fibres are soluble, explaining the higher soluble fibre content in USA (7.6 g/100 g) and Chilean (6.3 g/100 g) prunes compared with others (4.4-4.6 g/100 g). In addition, French prunes contained less sorbitol, while Chilean and Argentinian prunes had lower chlorogenic and neochlorogenic acid content than the prunes of other origins. As previously mentioned, these differences may be due to variations in soil management, plum ripeness or storage conditions (Donen 1939; Piga et al. 2003), although the reasons for any differences was not investigated here.

In regards to prunes of USA (Californian) origin (**Table 3**), there were minor differences between the current analytical data and those published by USDA, though these were small and likely negligible from a nutritional perspective. Given that the USDA database is updated regularly through submission of independent analyses from food manufacturers, the minor differences observed may simply reflect seasonal variation in composition. However, there were larger differences in both the current analytical data and the USDA data compared with the UK data provided by McCance and Widdowson, the latter reporting lower energy and starch values. Notably, little information is provided on the sourcing of prune samples reported in McCance and Widdowson and so it is unknown whether prune origin could be responsible for differences in starch content. Water content was comparable between the data (30.9-31.1%), therefore any differences were not due to variation in water content. Rather, the difference in energy content is explained by differences in the components included in the energy calculation and different conversion factors used.

In the current study, energy content is calculated based upon the contribution of 'available carbohydrate', fat, protein, fibre and polyols, as per European Union labelling regulations (EC, 2011). The USDA data includes 'total carbohydrate' in the energy calculation (rather than 'available carbohydrate'), and therefore does not take into account the lower energy contribution from fibre and polyols, and this is reflected in the slightly higher energy value published by USDA (240 kcal/100 g) compared with the current analytical data (230 kcal/100 g). In stark contrast, the UK data from McCance and Widdowson (141 kcal/100 g) excludes fibre and polyols from the energy calculation.

Prunes contained high levels of sorbitol (11.2-15.5 g/100g), these values being similar to USDA values (12.0 g/100g) and other studies in the scientific literature (10.8 g/100g) (Yao et al. 2014). The sorbitol content of prunes is therefore higher than that of its non-dried counterpart plums (2.4 g/100g), as well as other non-dried stone fruits such as cherries (0.7 g/100g) and dried fruits such as dried apricots (6g/100g), dried pear (8.1 g/100g) and dried apple (1.9g/100g) (Yao et al. 2014). Some polyols have been shown to induce increases in small intestinal water, although this has not been confirmed for sorbitol. For example, a fourfold increase in small intestinal water was observed in healthy individuals 60 minutes following ingestion of 17.5 g of mannitol (Marciani et al. 2010).

255

256 Prunes also contained high levels of chlorogenic acid (1.3-3.9 g/100g) and neochlorogenic acid
257 (39.8-92.0 g/100g), particularly those from the USA and France. This reflects data from
258 previous studies reporting high levels of phenolic compounds in prunes (Donovan et al. 1998;
259 Stacewicz-Sapuntzakis 2013). These phenolic compounds are partially absorbed in the small
260 intestine and the remainder enter the colon where they undergo biotransformation by the
261 microbiota into caffeic acid and quinic acid (Olthof et al. 2001). A recent systematic review
262 suggests that polyphenols and their degradation products can modulate the gut microbiota
263 and have prebiotic effects (Nash et al. 2018). Chlorogenic acid has been shown to inhibit the
264 growth and adhesion of selected gut pathogens to a human gut cell line and to enhance the
265 proliferation and adhesion of the probiotic *L. rhamnosus* (Parkar et al. 2008). Taken together,
266 it is plausible that the combination of different dietary fibres, sorbitol and polyphenols
267 naturally abundant in prunes create a synergistic effect, which in part, may be the reason why
268 prunes are considered beneficial for gastrointestinal health (European Food Safety Authority
269 2014).

270

271 The current analytical data and the USDA data calculate carbohydrate values by difference
272 (subtracting amounts of the other proximates from the total weight), while McCance and
273 Widdowson calculate available carbohydrate using monosaccharide equivalents of each
274 measured component. In the McCance and Widdowson UK data, available carbohydrate is
275 equal to total sugars since no polyols, oligosaccharides or starch are reported. Though the
276 reason for the lack of starch in prunes (0.0 g/100 g) reported in the McCance and Widdowson
277 UK data is unclear, it is possible that when analysis was undertaken, prunes of French origin
278 (which in the current analysis contained less starch) were more readily available and
279 accessible. Notably, the current analytical data and the data from the USDA database may
280 considerably overestimate the available carbohydrate content by including unmeasured
281 components which are not absorbed or not metabolised in the body to produce energy (e.g.
282 sorbitol). In the present analytical data the sum of starch and sugars is 12 g/100g less than
283 the value for available carbohydrate by difference, while in the USDA data the sum of starch
284 and sugars is around 20 g/100g less than the value for total carbohydrate by difference that
285 has been used to calculate energy content. True energy values (kcal) for prunes appear to be
286 between 230-240 kcal/100g in accordance with the present analytical data (230 kcal/100g)

and the USDA database (240 kcals/100g). This is in contrast to McCance and Widdowson that presents noticeably lower energy values (141 kcals/100g).

Limitations and strengths

The major limitation of this study was that, due to financial constraints, we did not analyse multiple prune samples from each country of origin that would have enabled both within-country variation and between-country statistical comparisons to be performed. In contrast, our approach enabled only between-country variation to be analysed, albeit not statistically compared. However, this approach allowed for a wide range of important nutrients and other compounds relevant to health to be included, which we felt outweighed the limitations of pooling samples. Despite the limitation of pooling prune samples from each country of origin, a robust sampling methodology was adopted based upon standards used for food composition databases to ensure high levels of representativeness in each pooled sample, including sourcing from a range of major retail centres in numerous sampling countries.

Further limitations include the small number of samples from Argentina, which may therefore not be fully representative of Argentinian prunes available across Europe. Any differences attributable to country of origin can only be ascertained by controlling other factors that can influence variation in nutrient composition. The sampling protocol attempted to account for seasonal variation and storage conditions by purchasing samples at the same time of year and within use by dates, and minimised changes in composition between purchase and analysis. However, given that all prunes were sampled at point of sale, pre-purchase confounding variables such as exposure to heat, light and humidity could not be controlled for. This might be relevant if these factors influence nutrient composition as some previous data suggests, however, although such analyses are of important academic and commercial interest, from a practical perspective the consumer cannot currently impact post-harvest/pre-purchasing processing.

Conclusion

The current study provides evidence that small differences in dietary fibre, sorbitol and phenolic content may exist between prunes of different countries of origin. To our knowledge, this is the first study to provide a comprehensive and comparable compositional analysis of

prunes of USA (Californian) origin, updating the currently available data reported in the USDA and McCance and Widdowson UK databases. This allows for a more accurate measurement of nutrient intake for future dietary intervention studies. The current study has highlighted the need for thorough and transparent documentation of sampling methods used to produce data for national databases. Furthermore, to eliminate artificial differences in energy content between different databases, carbohydrate values should be expressed using the same method and energy should be calculated using the same conversion factors.

References

- Bennett LE, Singh DP, Clingeffer PR. 2011. Micronutrient mineral and folate content of Australian and imported dried fruit products. *Crit Rev Food Sci Nutr*. 51(1): 38-49.
- Buncher RP. 2012. Prune Production Manual. UCANR Publications.
- Chang SK, Alasalvar C, Shahidi F. 2016. Review of dried fruits: Phytochemicals, antioxidant efficacies, and health benefits. *J Funct Foods*. 21: 113-132.
- Corradini C, Cavazza A, Bignardi C. 2012. High-performance anion-exchange chromatography coupled with pulsed electrochemical detection as a powerful tool to evaluate carbohydrates of food interest: principles and applications. *Inter J Carbohydrate Chem*. 2012.
- Donen I. 1939. The role of sorbitol in the carbon-metabolism of the Kelsey plum: Changes in chemical composition during growth and storage. *Biochem J*. 33(10): 1611-20.
- Donovan JL, Meyer AS, Waterhouse AL. 1998. Phenolic Composition and Antioxidant Activity of Prunes and Prune Juice (*Prunus domestica*). *Journal of Agricultural and Food Chemistry*. 46(4): 1247-1252.
- Dubois RN. 2004. Disease prevention: The future is here. *Gastroenterology*. 127: 1289.
- European Commission 2011. The provision of food information to consumers, Regulation (EU) No 1169/2011 of 25 October 2011 of the European Parliament and Council. *Official Journal of the European Union* L304/18.
- European Food Safety Authority. 2014. Scientific Opinion on the substantiation of a health claim related to prunes and contribution to normal bowel function pursuant to Article 14 of Regulation (EC) No 1924/2006.
- Finglas PM, Roe MA, Pinchen HM, Berry R, Church SM, Dodhia SK, Farron-Wilson M, Swan G. 2015. McCance and Widdowson's The Composition of Foods, Seventh summary edition. Cambridge: Royal Society of Chemistry.
- Lever E, Cole J, Scott SM, Emery PW, Whelan K. 2014. Systematic review: the effect of prunes on gastrointestinal function. *Aliment Pharmacol Ther*. 40(7): 750-8.
- Lever E, Scott SM, Emery PW, Whelan K. 2015. The effect of prunes on stool output, whole gut transit time and gastrointestinal symptoms: a randomised controlled trial. *Proceedings of the Nutrition Society*. 74, E2.
- Lombardi-Boccia G, Lucarini M, Lanzi S, Aguzzi A, Cappelloni M. 2004. Nutrients and antioxidant molecules in yellow plums (*Prunus domestica* L.) from conventional and organic productions: a comparative study. *J Agric Food Chem*. 52(1): 90-4.

- Marciani L, Cox EF, Hoad CL, Pritchard S, Totman JJ, Foley S, Mistry A, Evans S, Gowland PA, Spiller R C. 2010. Postprandial changes in small bowel water content in healthy subjects and patients with irritable bowel syndrome. *Gastroenterology*. 138(2): 469-77, 477.e1.
- Nash V, Ranadheera CS, Georgousopoulou EN, Mellor DD, Panagiotakos DB, Mckune AJ, Kellett J. Naumovski N. 2018. The effects of grape and red wine polyphenols on gut microbiota – A systematic review. *Food Research International*. 113: 277-287.
- Olthof MR, Hollman PC, Katan MB. 2001. Chlorogenic acid and caffeic acid are absorbed in humans. *J Nutr*. 131(1): 66-71.
- Parkar SG, Stevenson DE, Skinner MA. 2008. The potential influence of fruit polyphenols on colonic microflora and human gut health. *Int J Food Microbiol*. 124(3): 295-8.
- Piga A, Del Caro A, Corda G. 2003. From plums to prunes: influence of drying parameters on polyphenols and antioxidant activity. *J Agric Food Chem*. 51(12): 3675-81.
- Sadler MJ, Gibson S, Whelan K, Ha MA, Lovegrove J, Higgs J. 2019. Dried fruit and public health – what does the evidence tell us? *Inter J Food Sci Nutr* (in press). DOI 10.1080/09637486.2019.1568398
- Seppänen-Laakso T, Laakso I, Hiltunen R. 2002. Analysis of fatty acids by gas chromatography, and its relevance to research on health and nutrition. *Analytica Chimica Acta*. 465 (1-2): 39-62.
- Stacewicz-Sapuntzakis, M. 2013. Dried plums and their products: composition and health effects--an updated review. *Crit Rev Food Sci Nutr*. 53(12): 1277-302.
- US Department Of Agriculture, A. R. S., Nutrient Data Laboratory 2018. USDA National Nutrient Database for Standard Reference, Legacy.
- Wald A, Scarpignato C, Kamm MA, Mueller-Lissner S, Helfrich I, Schuijt C, Bubeck J, Limoni C, Petrini O. 2007. The burden of constipation on quality of life: results of a multinational survey. 26(2): 227-236.
- Yao CK, Tan HL, Van Langenberg DR, Barrett JS, Rose R, Liels K, Gibson PR, Muir JG. 2014. Dietary sorbitol and mannitol: food content and distinct absorption patterns between healthy individuals and patients with irritable bowel syndrome. *J Hum Nutr Diet*. 27 Suppl 2, 263-75.

Table 1: Number of prune samples purchased, pooled and analysed, by sampling country and country of origin.

Sampling country	USA (California)			France			Chile			Argentina		
	Number of samples		Weight contributing to analysis	Number of samples		Weight contributing to analysis	Number of samples		Weight contributing to analysis	Number of samples		Weight contributing to analysis
	Purchased	Analysed		Purchased	Analysed		Purchased	Analysed		Purchased	Analysed	
Germany	4	4	125 g (31.25 g per sample)	3	3	125 g (41.6 g per sample)	4	4	125 g (31.25 g per sample)	0	0	-
Italy	5	5	125 g (25 g per sample)	2	2	125 g (62.5 g per sample)	2	2	125 g (62.5 g per sample)	0	0	-
UK	5	5	125 g (25 g per sample)	5	5	125 g (25 g per sample)	5	5	125 g (25 g per sample)	0	0	-
France	0	0	-	16	8**	125 g (15.625 g per sample)	0	0	-	0	0	-
Spain	5	4*	125 g (31.25 g per sample)	4	0***	-	4	4	125 g (31.25 g per sample)	5	5	500 g (100 g per sample)
Total	19	18	500 g	30	18	500 g	15	15	500 g	5	5	500 g

* One sample exceeded the best before date and was excluded

** Only 8 samples required

*** Samples all labelled Spanish/French origin and thus excluded

Table 2: Composition of prunes from the pooled samples from USA, Chile, France and Argentina, purchased in Europe

	Method of analysis	Country of origin (per 100 g wet weight)				Country of origin (per 100 g dry weight)			
		USA	Chile	France	Argentina	USA	Chile	France	Argentina
Moisture (g)	Oven drying	30.9	30.5	33.2	28.7	-	-	-	-
Ash (g)	Incineration (muffle furnace)	1.58	1.53	1.36	1.38	-	-	-	-
Energy (kcal)	Multiplying macronutrients by Atwater factors	230	235	228	241	333	337	340	338
Protein (g)	Total N (Dumas, TruSpec analyser) x 6.25	2.5	2.1	1.6	2.0	3.6	2.9	2.4	2.8
Fat (g)	Soxhlet method	<0.2	<0.2	<0.2	<0.2	<0.2	<0.2	<0.2	<0.2
Total carbohydrate (g)	Calculated 'by difference'	65.1	65.9	63.9	68.0	94.1	94.9	95.4	95.3
Available carbohydrate (g)	Calculated 'by difference'	56.9	58.2	55.6	59.9	82.2	83.8	83.0	84.0
Starch (g)	Enzymatic hydrolysis (Megazyme)	6.6	5.7	1.9	6.1	9.5	8.1	2.8	8.5
Total sugars (g)	Ion-exchange chromatography	38.2	41.3	40.7	42.2	55.2	59.4	60.7	59.1
Fructose	Ion-exchange chromatography	14.0	16.2	16.0	16.9	20.2	23.3	23.9	23.6
Glucose	Ion-exchange chromatography	24.2	25.1	24.6	25.3	35.0	36.1	36.8	35.5
Galactose	Ion-exchange chromatography	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Lactose	Ion-exchange chromatography	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Maltose	Ion-exchange chromatography	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Sucrose	Ion-exchange chromatography	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
Sugar alcohols (g)									
Sorbitol	Ion-exchange chromatography	14.8	13.8	11.2	15.5	21.4	19.9	16.7	21.7
Dietary Fibre (g)									
Total fibre	Enzymatic-gravimetric (AOAC 991.43)	8.2	7.7	8.3	8.1	11.9	11.1	12.4	11.4
Total fibre	Enzymatic-gravimetric (AOAC 2011.25)	12.0	11.5	8.4	8.9	17.4	16.5	12.5	12.5
Insoluble fibre	Enzymatic-gravimetric (AOAC 2011.25)	4.5	5.7	4.0	4.3	6.5	8.1	6.0	6.0
Soluble fibre	Enzymatic-gravimetric (AOAC 2011.25)	7.6	6.3	4.4	4.6	11.0	9.1	6.5	6.4
Total NSP	Englyst et al (1994)	6.2	5.8	5.8	5.9	9.0	8.3	8.7	8.3
Insoluble NSP	Englyst et al (1994)	2.0	1.8	2.0	1.6	2.9	2.6	3.0	2.2
Soluble NSP	Englyst et al (1994)	4.3	4.1	3.9	4.3	6.1	5.8	5.7	6.0
Cellulose	Englyst et al (1994)	0.2	0.3	0.3	0.3	0.3	0.4	0.4	0.4

Method of analysis		Country of origin (per 100 g wet weight)				Country of origin (per 100 g dry weight)			
		USA	Chile	France	Argentina	USA	Chile	France	Argentina
Lignin	Enzymatic-gravimetric (AOAC 994.13)	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Fructans	Enzymatic spectrophotometric (AOAC 999.03)	0.3	0.2	0.3	0.3	0.4	0.3	0.5	0.4
Phenolic compounds (mg)									
Chlorogenic acid	UPLC with MS-MS	3.6	1.3	3.9	1.3	5.2	1.9	5.8	1.8
Neochlorogenic acid	UPLC with MS-MS	89.3	39.8	92.0	40.3	129.1	57.2	137.3	56.4

UPLC with MS-MS: Ultra-performance liquid-chromatography tandem mass spectrometry

Table 3: Composition of prunes originating from the USA (California), as analysed in the current study, compared with data from the USDA nutrient database and McCance and Widdowson's The Composition of Foods. Values are units per 100 g wet weight.

Analysis in the current study			Database values	
	Method of analysis in the current study	USA (Californian) (wet weight per 100 g)	USDA (wet weight per 100 g)	McCance & Widdowson (wet weight per 100 g)
Water (g)	Oven drying	30.9	30.9	31.1
Ash (g)	Incineration (muffle furnace)	1.58	NR	2.64
Energy (kcal)	Multiplying macronutrients by Atwater factors	230	240	141
Protein (g)	Total N content (Dumas, TruSpec analyser) x 6.25	2.5	2.2	2.5
Fat (g)	Soxhlet method	<0.2	0.4	0.4
Fatty acids (g)				
Saturated Fatty Acids	Gas chromatography with flame ionisation detection	<0.1	<0.1	NR
Mono-unsaturated Fatty Acids	Gas chromatography with flame ionisation detection	<0.1	<0.1	NR
Polyunsaturated Fatty Acids	Gas chromatography with flame ionisation detection	<0.1	<0.1	NR
Trans-unsaturated Fatty Acids	Gas chromatography with flame ionisation detection	<0.1	NR	NR
Total carbohydrate (g)	Calculated 'by difference'	65.1	63.9	NR
Available carbohydrate (g)	Calculated 'by difference'	56.9	NR	34.0
Starch (g)	Enzymatic hydrolysis (Megazyme)	6.6	5.1	0.0
Total sugars (g)	Ion-exchange chromatography, HPAEC-PED	38.2	38.1	34.0
Fructose	Ion-exchange chromatography, HPAEC-PED	14.0	12.5	12.1
Glucose	Ion-exchange chromatography, HPAEC-PED	24.2	25.5	17.9
Galactose	Ion-exchange chromatography, HPAEC-PED	<0.01	0.0	NR
Lactose	Ion-exchange chromatography, HPAEC-PED	<0.01	0.0	0.0
Maltose	Ion-exchange chromatography, HPAEC-PED	<0.01	0.1	0.0
Sucrose	Ion-exchange chromatography, HPAEC-PED	<0.01	0.2	4.1

Analysis in the current study		Database values		
	Method of analysis in the current study	USA (Californian) (wet weight per 100 g)	USDA (wet weight per 100 g)	McCance & Widdowson (wet weight per 100 g)
Sugar alcohols (g)				
Sorbitol	Ion-exchange chromatography, HPAEC-PED	14.8	NR	NR
Dietary Fibre (g)				
Total dietary Fibre	AOAC 991.43	8.2	7.1	NR
Total dietary Fibre	AOAC 2011.26	12.0	NR	NR
Insoluble dietary fibre	AOAC 2011.26	4.5	NR	NR
Soluble dietary fibre	AOAC 2011.26	7.6	NR	NR
Non-starch polysaccharides	Englyst et al (1994)	6.2	NR	5.7
Insoluble NSP	Englyst et al (1994)	2.0	NR	NR
Soluble NSP	Englyst et al (1994)	4.3	NR	NR
Cellulose	Englyst et al (1994)	0.2	NR	NR
Lignin	Enzymatic-gravimetric method (AOAC 994.13)	0.009	NR	NR
Fructans	Enzymatic spectrophotometric (AOAC 999.03)	0.3	NR	NR
Phenolic compounds (mg)				
Chlorogenic acid	UPLC with MS-MS	3.6	NR	NR
Neochlorogenic acid	UPLC with MS-MS	89.3	NR	NR
Minerals				
Calcium (mg)	ICP-OES	45.0	43.0	34.0
Iron (mg)	ICP-OES	0.7	0.9	2.6
Potassium (mg)	ICP-OES	622	732	760
Magnesium (mg)	ICP-OES	47.0	41.0	24.0
Sodium (mg)	ICP-OES	9.8	2.0	11.0

Analysis in the current study			Database values	
	Method of analysis in the current study	USA (Californian) (wet weight per 100 g)	USDA (wet weight per 100 g)	McCance & Widdowson (wet weight per 100 g)
Phosphorus (mg)	ICP-OES	68.1	69.0	73.0
Zinc (mg)	ICP-OES	0.4	0.4	0.4
Iodine (mg)	ICP-OES	3.0	NR	NR
Selenium (µg)	ICP-OES	30.0	0.3	3.0
Vitamins				
Riboflavin (mg)	HPLC	0.0	0.2	0.2
Niacin (mg)	HPLC	1.1	1.9	1.3
Vitamin B6 (mg)	HPLC	0.3	0.2	0.2
Biotin (µg)	Plasmon resonance technology	20.0	NR	Tr

NR = Not reported

Tr = trace

ICP-OES Inductively Coupled Plasma Optical Emission Spectrometry

HPAEC-PED High-Performance Anion-Exchange Chromatography Coupled with Pulsed Electrochemical Detection

HPLC High Performance Liquid Chromatography