King's Research Portal

Document Version
Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Doostdar, N., Kim, E., Grayson, B., Harte, M. K., Neill, J. C., & Vernon, A. C. (2019). Global brain volume reductions in a subchronic phencyclidine animal model for schizophrenia and their relationship to recognition memory. *Journal of Psychopharmacology*.

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Neuroscience-based Nomenclature (NbN) for *Journal of Psychopharmacology*

David J Nutt¹ and Pierre Blier²

Journal of Psychopharmacology

2016, Vol. 30(5) 413–415

© The Author(s) 2016

Reprints and permissions:

sagepub.co.uk/journalsPermissions.nav

DOI: 10.1177/0269881116642903

jop.sagepub.com

As of May 2016, the *Journal of Psychopharmacology* will fully adopt Neuroscience-based Nomenclature (NbN) for all publications and correspondence. In this, we join many other leading journals in our field, including *European Neuropsychopharmacology*, *Biological Psychiatry*, *CNS Spectrums*, *European Psychiatry*, *International Journal of Neuropsychopharmacology*, *Journal of Clinical Psychopharmacology*, *Neuropsychopharmacology*, *Pharmacopsychiatry*, *World Journal of Biological Psychiatry* and others that will also recommend the use of NbN. This decision has been ratified by the British Association for Psychopharmacology (BAP) council.

For the *Journal of Psychopharmacology* and the BAP, this step marks the output of a process that we have been involved in developing with the European College of Neuropsychopharmacology (ECNP). Part of the impetus to this initiative came from an editorial in *Journal of Psychopharmacology* back in 2009 (Nutt, 2009). Most of the vital background data collecting and organisation of the knowledge base has been done by a BAP member, Dr Sue Wilson.

Under the leadership of the ECNP in 2008, a taskforce for psychotropic nomenclature composed of representatives from five international organisations: the ECNP, American College of Neuropsychopharmacology, Asian College of Neuropsychopharmacology, International College of Neuropsychopharmacology and International Union of Basic and Clinical Pharmacology. The group tasked itself 'to examine ways of improving the current nomenclature in psychopharmacology'. Specifically, the new nomenclature was to (a) be based on contemporary scientific knowledge, (b) help clinicians to make informed choices when working out the next 'pharmacological step', (c) provide a system that does not conflict with the use of medications and (d) be future proof to accommodate new types of compounds. An initial proposal (Zohar et al., 2014) was discussed in the scientific community and accordingly revised (Zohar et al., 2015). It is this revised system that *Journal of Psychopharmacology* will use.

NbN is a pharmacologically driven system. In this, it is scientifically precise and extensible: new drug targets or modes of action can be easily added to the system once accepted by the scientific community. It is fairly comprehensive: at the present time, it includes 108 compounds which span the great majority of what is currently used in the practice of psychopharmacology. However, this very precision requires of authors and readers to adjust some well-worn habits: for example, use of the terms 'second-generation' or 'atypical antipsychotic', or even more so a group reference such as 'anxiolytics' referring to a group of

substances with quite heterogeneous receptor targets and mode of action. From an editor's point of view, we realize that this will pose some additional challenges for our authors, but we believe that readers and ultimately the field will benefit greatly – first, by the self-imposed terminology precision itself, and second, by sometimes making it clear where gaps in our knowledge still exist, especially when the modes of action of drugs with relationship to our evolving understanding of the neurobiology of mental illness are concerned (Millan et al., 2015).

Using NbN will require authors to clarify their meaning when they use a term for a drug. To ease the transition, *Journal of Psychopharmacology* will adopt NbN in stages: the first, effective as of May 2016, requires authors to define their usage, in the paper, of a term such as 'antipsychotics' using NbN at the point that it first appears in the main text of the paper. Furthermore, to make all new papers searchable by NbN, the NbN nomenclature of the substances that the paper covers has to be added to the keywords of the paper. To this end, we are dropping our previous limitation (of five) for keywords and have added a specific sub-category NbN to the keyword finder.

The new keywords include 11 pharmacological domains and 10 modes of action which are the building stones of NbN. It will make this process easier for authors and readers searching for one of our publications.

How does NbN work in practice? To 'translate' between old and new nomenclature, the easiest and recommended way is to use the approved app, which is available on the project's website (<http://nbomenclature.org/>), as well as in the software repositories of the various platforms for which it is available. On this website, there is a special tag – For Authors. A link to these resources is also now included in our Instructions For Authors.

The NbN effort is still in its infancy. Some necessary steps, such as the inclusion of paediatric psychopharmacology and neuropsychopharmacology in domains such as epilepsy are already being

¹Centre for Neuropsychopharmacology, Division of Brain Sciences, Imperial College London, London, UK

²Department of Psychiatry and Cellular and Molecular Medicine, University of Ottawa, Ottawa, ON, Canada

Corresponding author:

David Nutt, Centre for Neuropsychopharmacology, Division of Brain Sciences, Imperial College London, Burlington Danes Building, London W12 0NN, UK.

Email: d.nutt@imperial.ac.uk

worked on and will be included in the nomenclature in the near future. Most importantly, we believe that a clearer understanding of pharmacology will greatly benefit translational neuroscience and the discovery of new treatments for brain disorders.

References

- Millan MJ, Goodwin GM, Meyer-Lindenberg A, et al. (2015) 60 years of advances in neuropsychopharmacology for improving brain health, renewed hope for progress. *Eur Neuropsychopharmacol* 25: 591–598.
- Nutt DJ 2009 Beyond psychoanaleptics – can we improve antidepressant drug nomenclature? *J Psychopharmacol* 23: 343–345. Erratum in: *J Psychopharmacol* 2009; 23: 861.
- Zohar J, Nutt DJ, Kupfer DJ, et al. (2014) A proposal for an updated neuropsychopharmacological nomenclature. *Eur Neuropsychopharmacol* 24: 1005–1014.
- Zohar J, Stahl S, Moller HJ, et al. (2015) A review of the current nomenclature for psychotropic agents and an introduction to the Neuroscience-based Nomenclature. *Eur Neuropsychopharmacol* 25: 2318–2325.

Table 1. Neuroscience-based Nomenclature – NbN Glossary^a.

Former terminology	NbN – Pharmacological based		Drugs
Indication based	Pharmacology	Mode of action ^b	
Antidepressant (TCA)	Drugs for depression norepinephrine norepinephrine, serotonin	reuptake inhibitor (NET) reuptake inhibitor (NET and SERT)	desipramine protriptyline, lofepramine, amoxapine, nortriptyline
	serotonin, norepinephrine serotonin	reuptake inhibitor (SERT and NET) reuptake inhibitor (SERT)	imipramine, dosulepin clomipramine
	serotonin, norepinephrine	MM; reuptake inhibitor (SERT and NET), receptor antagonist (5-HT ₂)	amitriptyline
	norepinephrine, serotonin	MM; reuptake inhibitor (NET and SERT), receptor antagonist (5-HT ₂)	Doxepin
(MAOI)	serotonin, dopamine serotonin, norepinephrine, dopamine	receptor antagonist (5-HT ₂ and D ₂) enzyme inhibitor (MAO-A and -B) reversible enzyme inhibitor (MAO-A) MM; enzyme inhibitor (MAO-A and -B), releaser (DAT, NET)	trimipramine isocarboxazid, phenelzine moclobemide tranylcypromine
	dopamine, norepinephrine, serotonin serotonin	enzyme inhibitor (MAO-B and -A) reuptake inhibitor (SERT)	selegiline citalopram, escitalopram, fluoxetine, fluvoxamine, paroxetine, sertraline
(SSRI)			venlafaxine, duloxetine
(SNRI)	serotonin, norepinephrine norepinephrine, serotonin	reuptake inhibitor (SERT and NET) reuptake inhibitor (NET and SERT)	milnacipran
Stimulants	Dopamine and norepinephrine	Reuptake inhibitors and release	amphetamine (D) and (D,L), lisdexamfetamine, methylphenidate (D) and (D, L)
Antipsychotic (Neuroleptics, Major tranquilisers)	Drugs for psychosis		
Typical (1st generation)	dopamine	receptor antagonist (D ₂)	flupenthixol, fluphenazine, haloperidol, perphenazine, pimozide, pipotiazine, sulpiride, trifluoperazine, zuclopenthixol
Atypical (2nd generation)	dopamine, serotonin dopamine dopamine, serotonin	receptor antagonist (D ₂ , 5-HT ₂) receptor antagonist (D ₂) receptor antagonist (D ₂ , 5-HT ₂)	chlorpromazine, thioridazine amisulpiride iloperidone, loxapine, lurasidone, olanzapine, perospirone, sertindole, ziprasidone, zotepine
	dopamine, serotonin	receptor partial agonist (D ₂ , 5-HT _{1A})	aripiprazole
	dopamine, serotonin, noradrenaline	receptor antagonist (D ₂ , 5-HT ₂ , NE alpha-2) MM; receptor antagonist (D ₂ , 5-HT ₂) and reuptake inhibitor (NET) (metabolite)	asenapine, clozapine, risperidone, paliperidone quetiapine

Table 1. (Continued)

Former terminology	NbN – Pharmacological based		Drugs
Indication based	Pharmacology	Mode of action ^b	
Anxiolytic (benzodiazepine)	Drugs for anxiety		
	GABA	Positive Allosteric Modulator (GABA-A receptor, benzodiazepine site)	alprazolam, chlordiazepoxide, clonazepam, clorazepate, diazepam, flunitrazepam, lorazepam, oxazepam
	serotonin glutamate	receptor partial agonist (5-HT1A) voltage-gated calcium channel blocker	buspirone gabapentin, pregabalin
Hypnotic (benzodiazepine)	histamine	receptor antagonist (H1)	hydroxyzine
	Drugs for insomnia		
	GABA	Positive Allosteric Modulator (GABA-A receptor, benzodiazepine site)	estazolam, eszopiclone, flunitrazepam, lormetazepam, midazolam, quazepam, temazepam, triazolam, zaleplon, zolpidem, zopiclone
Mood stabilizers	melatonin	receptor agonist (M1, M2)	melatonin, ramelteon
	Drugs for relapse prevention		
	glutamate	voltage-gated sodium and calcium channel blocker	carbamazepine, oxcarbazepine
	glutamate	voltage-gated sodium channel blocker	lamotrigine
	glutamate	yet to be determined enzyme interactions	valproate lithium

^aThe glossary includes only the psychotropics relevant to former terminology. Newer psychotropics not included here could be found in NbN by their name.

^bMM (Multi-Modal)=more than one mode of action.