King's Research Portal

DOI:

[10.1111/dom.13771](https://doi.org/10.1111/dom.13771)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Hakim, O., Bello, O., Bonadonna, R. C., Mohandas, C., Shojee-moradie, F., Jackson, N., Boselli, L., Whitcher, B., Shuaib, M. H., Alberti, G., Peacock, J. L., Umpleby, A. M., Charles-Edwards, G., Amiel, S. A., & Goff, L. (2019). Ethnic differences in intrahepatic lipid and its association with hepatic insulin sensitivity and insulin clearance between men of Black and White ethnicity with early type 2 diabetes. *DIABETES OBESITY AND METABOLISM*, 21(9), 2163-2168. <https://doi.org/10.1111/dom.13771>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Ethnic differences in intrahepatic lipid and its association with hepatic insulin sensitivity and insulin clearance between men of Black and White ethnicity with early type 2 diabetes

Short title: Ethnicity, hepatic fat and type 2 diabetes

Olah Hakim¹, Oluwatoyosi Bello¹, Riccardo C. Bonadonna², Cynthia Mohandas¹, Fariba Shojee-Moradie³, Nicola Jackson³, Linda Boselli⁴, Brandon Whitcher⁵, Haris Shuaib^{6,7}, K. George M.M. Alberti¹, Janet L. Peacock⁸, A. Margot Umpleby³, Geoff Charles-Edwards^{6,9}, Stephanie A. Amiel¹, Louise M. Goff¹

¹Department of Diabetes, School of Life Course Sciences, Faculty of Life Sciences & Medicine, King's College London, London, UK; ²Department of Medicine & Surgery, University of Parma and Azienda Ospedaliera Universitaria di Parma, Parma, Italy; ³Faculty of Health and Medical Sciences, University of Surrey, Guildford, UK; ⁴Division of Endocrinology and Metabolic Disease, University of Verona School of Medicine, Verona, Italy; ⁵Research Centre for Optimal Health, School of Life Sciences, University of Westminster, London, UK; ⁶Medical Physics, Guy's & St Thomas' NHS Foundation Trust, London, UK; ⁷Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK; ⁸School of Population Health and Environmental Sciences, King's College London, London, UK; ⁹School of Biomedical Engineering & Imaging Sciences, King's College London, London, UK

Corresponding author: Dr. Louise M. Goff, Diabetes Research Group, Departments of Diabetes & Nutritional Sciences, School of Life Course Sciences, Faculty of Life Sciences &

26 Medicine, King's College London, Waterloo Campus, Franklin-Wilkins Building, Room 3.87,
27 London, SE1 9NH, United Kingdom. T: +44 (0)20 7848 6111; E: louise.goff@kcl.ac.uk
28 **Word count: main body 1957, abstract: 187**
29 **Tables & Figures:** 2 (1 table, 1 figure)
30 **References: 18**

ABSTRACT

Intrahepatic lipid (IHL) is linked with reduced hepatic insulin sensitivity and insulin clearance. Despite their high risk for type 2 diabetes (T2D), there have been limited investigations of these relationships in Black populations. We investigated these relationships in 18 White European (WE) and 18 Black West African (BWA) men with T2D <5 years. They underwent magnetic resonance imaging to quantify IHL, a hyperinsulinemic-euglycemic clamp with [6,6 $^2\text{H}_2$] glucose infusion to assess hepatic insulin sensitivity and a hyperglycemic clamp to assess insulin clearance. BWA men had lower IHL than WE men (3.7 (5.3) vs 6.6 (10.6) %, $p=0.03$). IHL was inversely associated with basal hepatic insulin sensitivity in WE but not BWA men (BWA: $r=-0.01$, $P=0.96$; WE: $r=-0.72$, $P=0.006$) with a significant interaction by ethnicity ($P_{\text{interaction}}=0.05$), however, IHL was not associated with % suppression of endogenous glucose production by insulin in either ethnicity. IHL showed a trend to an association with insulin clearance in BWA only (BWA: $r=-0.42$, $P=0.09$; WE: $r=-0.14$, $P=0.58$). The lack of association between IHL and hepatic insulin sensitivity in BWA men indicates IHL may play a lesser detrimental role in T2D in BWA men.

KEY WORDS: Ethnicity, hepatic fat, insulin sensitivity, insulin clearance, African, lipotoxicity

49 **ABBREVIATIONS**

50 ALT: Alanine aminotransferase

51 BSA: Body surface area

52 BWA: Black West African

53 HbA1C: Glycated hemoglobin

54 IHL: Intrahepatic lipids

55 MRI: Magnetic resonance imaging

56 SAT: Subcutaneous adipose tissue

57 VAT: Visceral adipose tissue

58 WE: White European

INTRODUCTION

Black populations are disproportionately affected by type 2 diabetes (T2D) with 2-3 times greater prevalence compared to white populations¹ despite typically having lower intrahepatic lipids (IHL) and visceral adipose tissue (VAT)^{1,2}. IHL is usually elevated in individuals with T2D and is inversely associated with both hepatic insulin sensitivity and insulin clearance³. Consistently, insulin clearance has been shown to be lower in Black populations compared to White populations⁴. However, investigations of ethnic differences in hepatic insulin sensitivity have shown inconsistent findings⁵⁻⁷. We have previously reported similar hepatic insulin sensitivity but lower insulin clearance in Black West African (BWA) compared to White European (WE) men^{8,9}. Despite the literature reporting ethnic differences in IHL, hepatic insulin sensitivity, and insulin clearance between Black and White populations, these have not previously been investigated in a single study to understand their relationships and how ethnicity impacts on these in the development of T2D in Black populations. Therefore, our aim was to investigate ethnic differences in IHL and its relationship with hepatic insulin sensitivity and insulin clearance in BWA and WE men with early T2D.

METHODS

This investigation was conducted as part of the South London Diabetes and Ethnicity Phenotyping study (Soul-Deep)¹⁰. Data on metabolic parameters for 92% of the present cohort have been previously reported^{8,9}, the present analyses relate to the whole cohort in whom relevant data were available. Participant recruitment and data collection took place April 2013 to January 2015. The study was approved by the London Bridge National Research Ethics Committee (12/LO/1859); all participants provided written informed consent.

Participants

Participants were recruited from primary care practices in London and deemed eligible to participate if they were 1) 18-65 years old, 2) BMI of 25-40 kg/m², 3) self-reported WE or BWA ethnicity, 4) diagnosis of T2D (less than 5 years), and 5) treated with lifestyle and/or metformin only. Further details of eligibility criteria are published in the protocol¹⁰. Participants attended all assessments after an overnight fast. If on metformin, participants were instructed to cease taking it for 7-days prior to each visit. Physical activity was measured as hours per day of moderate intensity activity using accelerometry watches worn for 4 consecutive days (MotionWatch 8.0, CamTech).

Magnetic Resonance Imaging

A Dixon-based MRI sequence was used on a 1.5 Tesla Siemens scanner to obtain images for the quantification of IHL and VAT. Participants were scanned lying supine on a spine RF coil with body phased array RF coils placed over the chest, abdomen and pelvis. While abdominal images were acquired participants were instructed to complete three 17-second breath-holds. From each participant, contiguous, axial T1-weighted gradient-echo images (repetition time: 6.77ms; echo times: 4.77ms (in-phase), 2.39ms (out-of-phase), flip angle: 10°) each with a slice thickness of 3mm were acquired, from which water and fat images were produced. Images were analyzed using HOROS v1.1.7 (www.horosproject.org; accessed 21/10/2017). IHL was

measured by selecting two abdominal MRI images representing the superior and inferior parts of the liver. Four circular regions of interest (ROIs) in identical positions were placed within the liver tissue of each pair of water and fat images (supplementary material, Figure 1). ROIs were positioned to include the posterior, anterior, medial and lateral sections of the liver. ROI areas ranged from 20 to 30cm², intending to cover as large an area of liver as possible while avoiding blood vessels, bile ducts and artefacts. Using the formula: %IHL = (F/(F+W))*100, where F is the pixel signal intensity of the fat image and W is the pixel signal intensity of the water image, the hepatic fat fraction was calculated in each ROI and IHL was calculated as the mean of all 8 ROIs. Total abdominal VAT and body SAT (neck to knee, excluding arms) was determined using an automated MRI analysis technique (Klarismo Ltd., London, UK) as previously described¹¹.

Clamp assessments

Whole-body insulin sensitivity (M-value) during the high dose insulin infusion (40 mU m⁻² BSA min⁻¹), hepatic insulin sensitivity (% suppression of endogenous glucose production (EGP) during the low dose insulin infusion (10 mU m⁻² BSA min⁻¹) and basal hepatic insulin sensitivity index) were measured using a two-step hyperinsulinemic-euglycemic clamp with the infusion of [6,6 ²H₂] glucose, according to previously described methodology⁹. The basal hepatic insulin sensitivity index was calculated as the reciprocal of the product of basal EGP rate (mmol/BSA min⁻¹) and fasting insulin concentration (pmol/l). To assess and model insulin clearance, each participant underwent a hyperglycemic clamp, described in detail elsewhere⁸.

Statistical analysis

Ethnic differences were determined using independent samples t-test for normally distributed variables or a Mann-Whitney test for variables that could not be log transformed to normal. ANCOVA was used, with VAT, BMI and age as separate covariates, to investigate ethnic differences in IHL and VAT. Correlations were assessed using Pearson's correlation; partial

124 correlation was used to investigate associations while adjusting for VAT, BMI and age.
125 Significance of an interaction by ethnicity was assessed using multiple regression with
126 ethnicity*logIHL used as an interaction term. Analyses were conducted with SPSS version
127 25.0; $P \leq 0.05$ were considered statistically significant.

RESULTS

Participant characteristics

The 18 BWA and 18 WE men were well-matched for age and BMI, Table 1. The BWA men had significantly lower IHL and total VAT mass, Table 1 (Figure 2, supplementary data). After adjustment for BMI, the ethnic differences in VAT remained significant ($P=0.008$) but not for IHL ($P=0.18$). After adjustment for VAT, there were no ethnic differences in IHL (WE: 6.07 (SE 1.16) vs BWA: 5.56 (SE 1.16) %, $P=0.70$). Non-alcoholic fatty liver disease, defined as liver fat above 5% determined by Dixon-MRI¹², was present in 33% of BWA men compared to 67% of WE men ($P=0.047$).

Metabolic characteristics

There were no ethnic differences in whole-body insulin sensitivity (M-value) or hepatic insulin sensitivity, expressed as % suppression of EGP during the low dose insulin infusion, Table 1; consistent with earlier findings we reported from a smaller sample from this cohort⁹. However, there was a trend towards higher basal hepatic insulin sensitivity index in the BWA men, Table 1. Insulin clearance was not different between the BWA and WE men (Table 1), again consistent with our earlier report⁸.

Relationships between IHL and insulin sensitivity

Relationships between IHL and the measures of insulin sensitivity are presented in Figure 1 (A, B and C). The inverse associations between IHL and both whole-body insulin sensitivity (M-value) and basal hepatic insulin sensitivity reached statistical significance in only the WE men. In multiple regression analysis a significant ethnicity interaction was found in the relationship between IHL and basal hepatic insulin sensitivity ($P_{\text{interaction}}=0.05$); no other significant ethnicity interactions were found. There were no changes in the associations after adjustment for VAT, BMI or age except for the relationship between IHL and M-value which reduced in significance in WE men after adjustment for BMI ($P=0.13$) (supplementary data).

153 **Relationships between IHL and insulin clearance**

154 Relationships between IHL and insulin clearance, are presented in Figure 1 (D). IHL was
155 inversely associated with insulin clearance, which neared significance, in BWA but not WE
156 men; partial correlation adjusting for VAT reduced the significance of this relationship (BWA:
157 $r=-0.41$, $P=0.11$; WE: $r=-0.29$, $P=0.27$); no significant ethnicity interaction was found
158 ($P_{\text{interaction}}=0.40$).

CONCLUSIONS

In this study of White European and Black West African men with early T2D, we investigated ethnic differences in hepatic fat and its relationship with hepatic insulin sensitivity and insulin clearance. Consistent with published data¹³, BWA men had lower IHL and VAT. We found additional ethnic differences in relationships between IHL and hepatic insulin sensitivity and insulin clearance whereby in WE men, IHL was inversely related to basal hepatic insulin sensitivity and whole-body insulin sensitivity, which was not the case in BWA men. In BWA men we found a trend towards an inverse relationship between IHL and insulin clearance which was not found in the WE men. Our findings suggest that IHL is implicated in the metabolic derangements of the liver in T2D differently according to ethnicity. To our knowledge, this is the first study to investigate relationships between IHL and insulin clearance in a Black population; the trend towards an inverse relationship in BWA but not WE men suggests that the reduction of insulin clearance may be modulated differently depending on ethnicity. Despite relationships between IHL and whole-body insulin resistance being commonly reported, the mechanisms that link the two are less understood. Current investigations show an excess of liver fat leads to accumulation of lipid intermediates causing hepatic mitochondrial dysfunction, inflammation and increased VLDL-TAG production which may result in hepatic and systemic insulin resistance¹⁴. Our finding of IHL being inversely associated with basal hepatic insulin sensitivity and whole-body insulin sensitivity which reached significance in WE but not BWA men may indicate the above detrimental effects of lipid intermediates occurring to a greater extent in WE men. There was no relationship between IHL and suppression of EGP in either ethnic group. This could indicate a decreased effect of IHL on hepatic insulin sensitivity in the insulin stimulated state compared to the basal state in WE men. To our knowledge only one other study has investigated the relationship between IHL and hepatic insulin sensitivity using the

hyperinsulinemic-euglycemic clamp with infusion of isotopically labelled glucose⁶; they found that IHL was associated with hepatic insulin sensitivity in obese Black South African women but not in obese White South African women, which contradicts our findings. There are several potential explanations for this, such as glycemic state; our study included participants with T2D whereas the South African women were normal glucose tolerant. The disparities may also be due to gender differences as there is consistent evidence demonstrating that the phenotype of T2D differs by gender within populations of African descent¹⁵.

The presence of NAFLD was comparable to that reported in other large multi-ethnic cohorts¹⁶ and was significantly lower in the BWA men. One of the main theories that explains how IHL accumulates is the “portal theory”, which states that excess VAT releases free fatty acids directly into the portal vein, subsequently depositing as IHL¹⁷. Our study may support the portal theory as after adjustment for VAT, IHL no longer differed by ethnicity, suggesting that the lower IHL in BWA men may be driven by lower VAT. Indeed, ethnic differences in the mechanisms of SAT expansion may explain the differences we found in VAT as others have suggested¹⁸; however, we did not directly measure adipogenesis in our study which may be an implication for further research.

The strengths of this study include the use of the rigorous hyperinsulinemic-euglycemic clamp method combined with the infusion of [6,6 ²H₂] glucose to determine both whole-body and hepatic insulin sensitivity. However, our study is not without its limitations. Our sample size is relatively small; in these secondary analyses we may not have sufficient power to reliably detect ethnic differences. Our measurement of insulin clearance does not differentiate hepatic from extrahepatic insulin clearance, rather it is a measure of whole-body insulin clearance. However, it has been shown that approximately 80% of endogenous insulin is degraded in the liver³. Our WE men had greater statin use which may have resulted in lower hepatic fat accumulation and reduced the ethnic discrepancies due to the lipid lowering effects of statins.

209 Another limitation is studying only men with T2D, however previous studies have mostly
210 focused on women, due to the greater prevalence of T2D in Black women compared to men.
211 Our study redresses this.

212 In conclusion, our study demonstrates ethnic differences in the relationships between IHL and
213 metabolic parameters of the liver. The lack of inverse association between IHL and basal
214 hepatic insulin sensitivity in the BWA men, found in the WE men, suggests that fasting hepatic
215 insulin resistance occurs independently of IHL in BWA men. However, the reduction of insulin
216 clearance may be influenced by IHL more so in Black men with T2D compared to White men.

217 **Author contributions**

218 L.M.G. formulated the research question and designed the study, supervised data collection
219 and interpretation, and performed the minimal modelling analysis. S.A.A. formulated the
220 research question and designed the study, supervised data collection and interpretation. J.L.P.
221 formulated the research question, designed the study, and provided statistical advice. A.M.U.
222 formulated the research question and designed the study. K.G.M.M.A. supervised data
223 collection and interpretation. C.M. coordinated the study and data acquisition, and performed
224 the metabolic assessments. T.B. undertook data acquisition and analysis. G.C.E. coordinated
225 MRI data acquisition. B.W. and H.S. undertook MRI data analysis. F.S. and N.J. undertook
226 data acquisition. R.B. and L.B. performed the modelling analysis. O.H. undertook data
227 analysis, statistical analysis and drafted the manuscript. All authors contributed to the
228 intellectual content and reviewed the final version of the submitted manuscript.

229 **Acknowledgements**

230 The authors thank Andrew Pernet, Bula Wilson and Ines De Abreu (research nurses, Diabetes
231 Research Group, King's College Hospital, UK) for assisting with the metabolic assessments;
232 Anne-Catherine Perz (King's College London, UK), Daniel Curtis (University of Surrey, UK)
233 and Tracy Dew (ViaPath, UK) for assistance with sample processing and laboratory analysis;
234 Elka Giemsa (CRF manager, King's College Hospital, UK) for accommodating the participant
235 visits; Maddalena Trombetta (University of Verona, Italy) for assisting with the minimal
236 modelling analysis. The staff of the Clinical Research Facility at King's College Hospital for
237 help in performing the studies; and the study participants for their time and commitment.

238 JLP is supported by the NIHR Biomedical Research Centre based at Guy's and St Thomas'
239 NHS Foundation Trust and King's College London. JLP is a NIHR Senior Investigator. OH
240 was supported by the National Institute for Health Research (NIHR) Collaboration for
241 Leadership in Applied Health Research and Care South London at King's College Hospital

242 NHS Foundation Trust. The views expressed are those of the authors and not necessarily those
243 of the NHS, the NIHR or the Department of Health.

244 LB is supported in part by funds of the Italian Ministry of Education, University and Research
245 (MIUR) PRIN 2015 2015373Z39_004 and with University of Parma research funds, both to
246 RCB.

247 Louise Goff is the guarantor of this work, had full access to all the data, and takes full
248 responsibility for the integrity of the data and the accuracy of data analysis.

249 **Funding source:** this work was funded by a Diabetes UK project grant: #12/0004473, and in
250 part by funds of the Italian Ministry of Education, University and Research (MIUR) PRIN 2015
251 2015373Z39_004 and with University of Parma research funds, both to RCB.

252 **Duality of interests:** The authors declare that there is no duality associated with this
253 manuscript.

254 **Table 1: Clinical and metabolic characteristics of Black West African and White**
255 **European men**

	BWA (n=18)	WE (n=18)	P
Age (years)†	54.9 (9.3)	58.5 (6.3)	0.67
Weight (kg)	92.3 ± 12.3	99.8 ± 16.7	0.14
BMI (kg/m ²)	29.8 ± 3.5	31.5 ± 4.1	0.18
Waist circumference (cm)	104.9 ± 10.2	111.9 ± 13.0	0.08
SAT (neck to knee) (kg)‡	12.6 (10.5-15.2)	14.6 (12.2-17.6)	0.24
VAT, total (kg) ^a	3.99 ± 1.54	6.09 ± 2.46	0.006
IHL (%)†	3.7 (5.3)	6.6 (10.6)	0.03
Diabetes duration (years)†	3.0 (2.2)	3.0 (1.3)	0.42
Statin use§	10/18	16/18	0.026
Fasting glucose (mmol/l)	6.63 ± 0.67	6.88 ± 1.38	0.50
HbA1c (%)	6.67 ± 0.68	6.64 ± 0.70	0.90
ALT‡ (IU/l)	26.7 (21.8-32.5)	31.2 (25.7-37.7)	0.24
Systolic BP (mm Hg)	136.7 ± 13.8	130.9 ± 14.2	0.22
Diastolic BP (mm Hg)†	89.0 (8.7)	83.0 (12.5)	0.06
Total cholesterol (mmol/l)	4.11 ± 0.73	4.27 ± 0.70	0.50
LDL-cholesterol (mmol/l)	2.32 ± 0.56	2.28 ± 0.66	0.85
HDL-cholesterol (mmol/l)	1.18 ± 0.38	1.19 ± 0.25	0.92
Triglyceride (mmol/l)†	1.05 (0.70)	1.60 (1.25)	0.03
Moderate activity time (hours/day) ^c	2.1 ± 0.66	1.9 ± 0.90	0.74
Metabolic characteristics			
M value (mg/m ² BSA min ⁻¹) ^d	162.0 ± 75.0	128.5 ± 63.7	0.17
Hepatic basal insulin sensitivity index ((mmol/m ² BSA min pmol l ⁻¹) ⁻¹) ^e	68.0 ± 24.6	49.1 ± 29.4	0.09
Suppression of endogenous glucose production (%) ^e	37.9 ± 19.5	34.7 ± 20.7	0.70
Average insulin clearance (mL/m ² BSA min ⁻¹)†	732.8 (505.7)	814.6 (450.2)	0.61

256 Data presented as mean ± SD or geometric mean (95% CI) for log transformed data (§) or median (interquartile
257 range) for non-parametric data (†) or number of participants for ordinal data (§). *P* values determined using
258 independent samples t-tests for normally distributed data, Mann-Whitney test for non-parametric data or chi-

259 squared test for ordinal data. N for ^aWE=17, BWA=17; ^bWE=17, BWA=16; ^cWE=10, BWA=7; ^dWE=18,
260 BWA=16; ^eWE=12, BWA=14.

261 Abbreviations: ALT, alanine aminotransferase; BP, blood pressure; BWA, Black West African; HbA1c, glycated
262 hemoglobin; HDL, high density lipoprotein; IHL, intrahepatic lipid; LDL, low density lipoprotein; SAT,
263 subcutaneous adipose tissue; VAT, visceral adipose tissue; WE, White European.

Figure 1: Relationships between hepatic fat fraction and (A) hepatic insulin sensitivity index (basal) ($((\text{mmol}/\text{m}^2 \text{ BSA min pmol l})^{-1})$), (B) suppression of hepatic glucose production (%), (C) whole-body insulin sensitivity (M-value) ($(\text{mg}/\text{m}^2 \text{ BSA min}^{-1})$), and (D) insulin clearance ($\text{mL}/\text{m}^2 \text{ BSA min}^{-1}$) in WE and BWA men. Relationships between hepatic insulin clearance and (E) suppression of endogenous glucose production, and (F) hepatic insulin sensitivity index (basal) in WE and BWA men. Black circles with dotted line = BWA men, white circles with solid line = WE men.

REFERENCES

1. *Health Survey for England 2004: The Health of Minority Ethnic Groups– headline tables*. NHS Health and Social Care Information Centre, Public Health Statistics 2005.
2. Liska D, Dufour S, Zern TL, et al. Interethnic Differences in Muscle, Liver and Abdominal Fat Partitioning in Obese Adolescents. *PLoS ONE*. 2007;2(6).
3. Kotronen A, Juurinen L, Tiikkainen M, Vehkavaara S, Yki-Jarvinen H. Increased liver fat, impaired insulin clearance, and hepatic and adipose tissue insulin resistance in type 2 diabetes. *Gastroenterology*. 2008;135(1):122-130.
4. Piccinini F, Polidori DC, Gower BA, Bergman RN. Hepatic but Not Extrahepatic Insulin Clearance Is Lower in African American Than in European American Women. *Diabetes*. 2017;66(10):2564-2570.
5. Ellis AC, Alvarez JA, Granger WM, Ovalle F, Gower BA. Ethnic differences in glucose disposal, hepatic insulin sensitivity, and endogenous glucose production among African American and European American women. *Metabolism: clinical and experimental*. 2012;61(5):634-640.
6. Goedecke JH, Keswell D, Weinreich C, et al. Ethnic differences in hepatic and systemic insulin sensitivity and their associated determinants in obese black and white South African women. *Diabetologia*. 2015;58(11):2647-2652.
7. DeLany JP, Dube JJ, Standley RA, et al. Racial differences in peripheral insulin sensitivity and mitochondrial capacity in the absence of obesity. *The Journal of clinical endocrinology and metabolism*. 2014;99(11):4307-4314.
8. Mohandas C, Bonadonna R, Shojee-Moradie F, et al. Ethnic differences in insulin secretory function between Black African and White European men with early type 2 diabetes. *Diabetes Obes Metab*. 2018.
9. Bello O, Mohandas C, Shojee-Moradie F, et al. Black African men with early type 2 diabetes have similar muscle, liver and adipose tissue insulin sensitivity to white European men despite lower visceral fat. *Diabetologia*. 2019.
10. Goff L. Soul-Deep: the South London Diabetes and Ethnicity Phenotyping Study Protocol. In:2013.
11. Hakim O, Bonadonna RC, Mohandas C, et al. Associations between pancreatic lipids and β -cell function in Black African and White European men with type 2 diabetes. *The Journal of Clinical Endocrinology & Metabolism*. 2018;jc.2018-01809-jc.02018-01809.
12. Nouredin M, Lam J, Peterson MR, et al. Utility of Magnetic Resonance Imaging Versus Histology for Quantifying Changes in Liver Fat in Nonalcoholic Fatty Liver Disease Trials. *Hepatology (Baltimore, Md)*. 2013;58(6):1930-1940.
13. Alderete T, Toledo-Corral C, Goran M. Metabolic Basis of Ethnic Differences in Diabetes Risk in Overweight and Obese Youth. *Current diabetes reports*. 2014;14(2):455.
14. Trouwborst I, Bowser SM, Goossens GH, Blaak EE. Ectopic Fat Accumulation in Distinct Insulin Resistant Phenotypes; Targets for Personalized Nutritional Interventions. *Frontiers in nutrition*. 2018;5:77.
15. Carnethon MR, Palaniappan LP, Burchfiel CM, Brancati FL, Fortmann SP. Serum insulin, obesity, and the incidence of type 2 diabetes in black and white adults: the

- atherosclerosis risk in communities study: 1987-1998. *Diabetes care*. 2002;25(8):1358-1364.
16. Browning JD, Szczepaniak LS, Dobbins R, et al. Prevalence of hepatic steatosis in an urban population in the United States: impact of ethnicity. *Hepatology (Baltimore, Md)*. 2004;40(6):1387-1395.
 17. Item F, Konrad D. Visceral fat and metabolic inflammation: the portal theory revisited. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2012;13 Suppl 2:30-39.
 18. Goedecke JH, Levitt NS, Evans J, et al. The role of adipose tissue in insulin resistance in women of African ancestry. *Journal of obesity*. 2013;2013:952916.

Supplementary data:

Table 1: Pearson's correlation and partial correlation coefficients showing relationships between hepatic fat fraction and measures of insulin sensitivity and insulin clearance in white European and black West African men

	WE		BWA	
	r	p	r	p
M-Value	-0.64	0.004	-0.35	0.18
Adjusted for age	-0.07	0.003	-0.39	0.15
Adjusted for BMI	-0.38	0.13	-0.05	0.85
Adjusted for VAT	-0.56	0.02	-0.06	0.85
Hepatic insulin sensitivity index	-0.72	0.006	-0.01	0.96
Adjusted for age	-0.70	0.01	-0.12	0.96
Adjusted for BMI	-0.60	0.04	-0.16	0.60
Adjusted for VAT	-0.73	0.01	0.35	0.26
% Suppression of endogenous glucose production	-0.11	0.73	-0.34	0.23
Adjusted for age	-0.12	0.72	-0.51	0.07
Adjusted for BMI	-0.04	0.91	-0.21	0.50
Adjusted for VAT	-0.03	0.93	-0.02	0.95
Insulin clearance	-0.14	0.58	-0.42	0.09
Adjusted for age	-0.15	0.57	-0.44	0.07
Adjusted for BMI	0.14	0.59	-0.27	0.29
Adjusted for VAT	-0.29	0.27	-0.41	0.11

Abbreviations: BMI, body mass index; BWA, black West African; VAT, visceral adipose tissue; WE, white European

28 **Figure 1:** Quantification of intrahepatic lipids by selection of two axial MRI images with
 29 regions of interest positioned on the right and left lobes of the liver as well as the posterior,
 30 anterior, medial and lateral sections.

31 Panel A shows a coronal MRI image with the horizontal line depicting the position of axial
 32 images C and D. Panel B shows a coronal MRI image with the horizontal line depicting the
 33 position of axial images E and F. Panel C shows 4 circular regions of interest on an axial
 34 abdominal MRI water image on the superior section of the liver. Panel D shows the axial
 35 abdominal MRI fat image that corresponds to image C with 4 identical regions of interest.
 36 Panel E shows 4 circular regions of interest on an axial abdominal MRI water image on the
 37 inferior section of the liver. Panel F shows the axial abdominal MRI fat image that corresponds
 38 to image E with 4 identical regions of interest.

40

41 **Figure 2:** Boxplot of hepatic fat fraction in White European (WE) and Black West African
42 men (BWA) with early type 2 diabetes matched for both age and BMI

43