King's Research Portal

DOI:

[10.1097/RLU.0000000000002811](https://doi.org/10.1097/RLU.0000000000002811)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Flaus, A., Riaño Barros, D. A., Hinz, R., Myers, J. F., Lingford-Hughes, A., Koepp, M. J., Hammers, A., & McGinnity, C. J. (2019). Decreased GABA-A Receptor Binding in Association With -Lactam Antibiotic Use. *Clinical Nuclear Medicine*, 44(12), 981-982. <https://doi.org/10.1097/RLU.0000000000002811>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Decreased GABA-A receptor binding in association with β -lactam antibiotic use

Flaus A^{a, b}, Riano Barros DA^{c, d}, Hinz R^e, Myers JF^c, Lingford-Hughes A^f, Koepp MJ^{g, h}, Hammers A^{b, c, d, i}, McGinnity CJ^{b, c, d}.

^a Department of Nuclear Medicine, University Hospital, Saint-Etienne, France

^b School of Biomedical Engineering & Imaging Sciences, King's College London, London, UK and Guy's and St Thomas' PET Centre, St Thomas' Hospital, London, UK

^c Centre for Neuroscience, Department of Medicine, Imperial College London, London, UK

^d MRC Clinical Sciences Centre Hammersmith Hospital, London, UK

^e Wolfson Molecular Imaging Centre, University of Manchester, Manchester, UK

^f Neuropsychopharmacology Unit, Department of Medicine, Imperial College London, London, UK

^g Department of Clinical and Experimental Epilepsy, Institute of Neurology, University College London, UK

^h Epilepsy Society, Chalfont St Peter, UK

ⁱ Neurodis Foundation, CERMEP, Imagerie du Vivant, Lyon, France

Corresponding author:

Flaus Anthime

Service de Médecine Nucléaire, Hopital Nord, CHU de St Etienne
42055 Saint-Etienne Cedex 2 FRANCE.

Email address : anthime.flaus@gmail.com – Telephone number: +33477828318

16-digit ORCID identifier is 0000-0001-5877-2159

Compliance with Ethical Standards

- The authors declare that they have no conflict of interest.
- Informed consent was obtained from all individual participants included in the study.
- All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Decreased GABA-A receptor binding in association with β -lactam antibiotic use

Abstract : β -lactam antibiotics are proconvulsive. In laboratory animals, this effect appears to be predominantly mediated through inhibition of GABA-A receptors, but it has not been demonstrated in humans in vivo. We report images of a [^{11}C]Ro15-4513 PET from a 40 years old male who had completed a one week course of flucloxacillin before it. Relative to healthy controls, the participant had significantly lower mean grey matter binding. These novel data suggest that in humans, the proconvulsive effect of β -lactam antibiotics is mediated via either competition for the same benzodiazepine binding site as [^{11}C]Ro15-4513, or downregulation of GABA-A receptor expression.

Keywords : PET; Ro15-4513; β -lactam; GABA-A

FIGURE 1. β -lactam antibiotics such as flucloxacillin are proconvulsive, and seizures are a well-recognized manifestation of penicillin neurotoxicity[1,2]. In laboratory animals, this effect appears to be predominantly mediated through inhibition of GABA-A receptors[3–5]. However, to our knowledge, the effect of β -lactam antibiotics on GABA-A receptor binding has not been demonstrated in humans *in vivo*.

[^{11}C]Ro15-4513 was used to acquire positron emission (PET) data from a 40-year-old male who had completed a one-week course of flucloxacillin 250mg q.i.d. one day before the scan. Data were acquired as described previously[6]. [^{11}C]Ro15-4513 volume-of-distribution (V_T ; i.e. binding) reflects the availability of GABA-A receptor subunit: total, $\alpha 1$ (mostly fast), and $\alpha 5$ (mostly slow) components [7]. We corrected for partial volume effects using SFS-RR[8].

Relative to 23 unmedicated healthy controls, the participant had significantly lower mean grey matter total V_T (-30% [3.39 vs. mean $\pm$ standard deviation 4.86 ± 0.51], $p=0.002$). The reduction was of similar magnitude for both fast and slow mean grey matter V_T s (fast: -29% [2.10 vs. 2.94 ± 0.29], $p=0.002$; slow: -31% [1.24 vs. 1.81 ± 0.60], $p=0.174$).

The participant's injectate had contained 17.05 μg of ligand, whereas $3.44 \pm 1.34 \mu\text{g}$ was typical for the healthy controls. This cannot explain our findings, as $\alpha 5$ subunit occupancy was $\sim 11.9\%$, and $\alpha 1$ occupancy only $\sim 1.0\%$, with a near-identical reduction for both components.

These novel data suggest that in humans, the proconvulsive effect of β -lactam antibiotics is mediated via either competition for the same benzodiazepine binding site as [^{11}C]Ro15-4513, or downregulation of GABA-A receptor expression.

1. Grill MF, Maganti RK. Neurotoxic effects associated with antibiotic use: management considerations. *Br J Clin Pharmacol.* 2011;72:381–393.
2. Chow KM, Hui AC, Szeto CC. Neurotoxicity induced by beta-lactam antibiotics: from bench to bedside. *Eur J Clin Microbiol Infect Dis.* 2005;24:649–653.
3. Sugimoto M, Uchida I, Mashimo T, et al. Evidence for the involvement of GABA(A) receptor blockade in convulsions induced by cephalosporins. *Neuropharmacology.* 2003;45:304–314.
4. Fujimoto M, Munakata M, Akaike N. Dual mechanisms of GABAA response inhibition by beta-lactam antibiotics in the pyramidal neurones of the rat cerebral cortex. *Br J Pharmacol.* 1995;116:3014–3020.
5. Rossokhin AV, Sharonova IN, Bukanova JV, et al. Block of GABAA receptor ion channel by penicillin: Electrophysiological and modeling insights toward the mechanism. *Molecular and Cellular Neuroscience.* 2014;63:72–82.
6. McGinnity CJ, Riaño Barros DA, Rosso L, et al. Test-retest reproducibility of quantitative binding measures of [^{11}C]Ro15-4513, a PET ligand for GABA A receptors containing $\alpha 5$ subunits. *NeuroImage.* 2017;152:270–282.
7. Myers JF, Rosso L, Watson BJ, et al. Characterisation of the contribution of the GABA-benzodiazepine $\alpha 1$ receptor subtype to [^{11}C]Ro15-4513 PET images. *J Cereb Blood Flow Metab.* 2012;32:731–744.
8. McGinnity CJ, Shidahara M, Feldmann M, et al. Quantification of opioid receptor availability following spontaneous epileptic seizures: Correction of [^{11}C]diprenorphine PET data for the partial-volume effect. *NeuroImage.* 2013;79:72–80.