King's Research Portal

DOI:

[10.1016/j.wneu.2019.08.262](https://doi.org/10.1016/j.wneu.2019.08.262)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Patel, E., Aydin, A., Cearn, M., Dasgupta, P., & Ahmed, K. (2020). A Systematic Review of Simulation-based Training in Neurosurgery, Part 1: Cranial Neurosurgery. *World Neurosurgery*, 133, e850-e873. <https://doi.org/10.1016/j.wneu.2019.08.262>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Manuscript Number:

Title: Current Status of Simulation-based Training in Neurosurgery: A Systematic Review

Article Type: Original Article

Keywords: Education; Neurosurgery; Simulation; Training

Corresponding Author: Dr. Abdullatif Aydın, B.Sc. (Hons), M.B.B.S.

Corresponding Author's Institution: King's College London

First Author: Ebrahim A Patel, BSc

Order of Authors: Ebrahim A Patel, BSc; Abdullatif Aydın, B.Sc. (Hons), M.B.B.S.; Michael Cearns, BSc (Hons), MBBS, MRCS, AFHEA; Kamran Ahmed, FRCS, PhD

Abstract: Objective: The recent emphasis on simulation-based training in neurosurgery has led to the development of many simulation models and training courses. We aim to identify the currently available simulators and training courses for neurosurgery, assess their validity and determine their effectiveness.

Design: Both Medline and EMBASE were searched for English language articles which validate simulation models for neurosurgery. Each study was screened according to Messick's validity framework, and rated in each domain. McGaghie's model of translational outcomes was then used to determine a level of effectiveness (LoE) for each simulator or training course.

Results: Upon screening of 6006 articles, 102 were identified either validating or determining a LoE for 97 simulation-based training models or courses. Achieving the highest rating for each validity domain were: six models and training courses for content validity; 19 for response processes; 5 for internal structure; 12 for relations to other variables and only 2 for consequences. For translational outcomes, 45 simulators or training courses were given a LoE of 1, 34 a LoE of 2, 1 a LoE of 3 and 1 a LoE of 4. Three models and one training course achieved the highest LoE of 5.

Conclusions: With the advent of increasing neurosurgery simulators and training tools, there is a need for more validity studies. Further attempts to investigate translational outcomes to the operating theatre when using these simulators is particularly warranted. Finally, more training tools incorporating full immersion simulation and non-technical skills training are recommended.

Dear Editor,

Please find submitted our systematic review on simulation-based training in neurosurgery entitled “Current Status of Simulation-based Training in Neurosurgery: A Systematic Review”. We hope it fulfils the criteria to be considered for publication in your journal. I, Abdullatif Aydin, certify that this manuscript is a unique submission and is not being considered for publication, in part or in full, with any other source in any medium.

Yours truly,

Dr. Abdullatif Aydin

Current Status of Simulation-based Training in Neurosurgery: A Systematic

Review

Authors:

Ebrahim Patel (1) BSc (Hons)

Abdullatif Aydin (2) BSc (Hons), MBBS

Michael Cearnns (3) BSc (Hons), MBBS, MRCS, AFHEA

Kamran Ahmed (2) FRCS, PhD

Affiliations:

- (1) Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, United Kingdom
- (2) MRC Centre for Transplantation, Guy's Hospital, King's College London, London, United Kingdom
- (3) Department of Neurosurgery, Institute of Neurological Sciences, Glasgow, United Kingdom

Correspondence:

Dr Abdullatif Aydin BSc, MBBS

MRC Centre for Transplantation,

5th Floor Southwark Wing, Guy's Hospital,

King's College London, London SE1 9RT

Ph: +44 (0)20 7188 8580

Fax: +44 (0)20 3312 6787

Email: abdullatif.aydin@kcl.ac.uk

Keywords: Education, Neurosurgery, Simulation, Training

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Abbreviations

ACDF: Anterior cervical discectomy and fusion

AR: Augmented Reality

CNS: Congress of neurological surgeons

CSF: Cerebrospinal fluid

ETV: Endoscopic third ventriculostomy

EVD: External Ventricular Drain

FNS: Fundamentals of neurosurgery

ICA: Internal Carotid Artery

LoE: Level of effectiveness

NET: Neuro-Endo Trainer

OSATS: Objective Structured Assessment of Technical Skills

PCF: Posterior cervical fusion

PPDIS: Physician Performance Diagnostic Inventory scale

PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-Analyses

SNS: Society of neurological surgeons

V-PAT: Ventriculostomy Procedural Assessment Tool

VR: Virtual Reality

Objective: The recent emphasis on simulation-based training in neurosurgery has led to the development of many simulation models and training courses. We aim to identify the currently available simulators and training courses for neurosurgery, assess their validity and determine their effectiveness.

Results: Upon screening of 6006 articles, 102 were identified either validating or determining a LoE for 97 simulation-based training models or courses. Achieving the highest rating for each validity domain were: six models and training courses for content validity; 19 for response processes; 5 for internal structure; 12 for relations to other variables and only 2 for consequences. For translational outcomes, 45 simulators or training courses were given a LoE of 1, 34 a LoE of 2, 1 a LoE of 3 and 1 a LoE of 4. Three models and one training course achieved the highest LoE of 5.

Conclusions: With the advent of increasing neurosurgery simulators and training tools, there is a need for more validity studies. Further attempts to investigate translational outcomes to the operating theatre when using these simulators is particularly warranted. Finally, more training tools incorporating full immersion simulation and non-technical skills training are recommended.

Introduction

Neurosurgical training has traditionally assumed the Halstedian model of 'see one, do one, teach one' ¹. While this model of teaching has proven invaluable over the years, it is facing increasing challenges in the modern era ². Issues such as cost, reduced working hours, increased expectations of patient safety, surgeon performance and transparency have led to an emphasis on protecting patients from being used as training tools themselves. The use of simulation-based training within neurosurgery has thus been advocated to enhance the current model of teaching ³. Indeed, this mode of training has demonstrated merit for surgical skill acquisition in other specialties ⁴. While many reviews exist for simulation-based training in neurosurgery, none rigorously determine the validity and effectiveness of these training models within the specialty.

The aim of this review is to identify the current simulation-based training models described in the literature, quantitatively assess their validity and determine their effectiveness for training in neurosurgery.

Methods

Information Sources and Search

A broad search of the PubMed and EMBASE databases was performed to identify articles that described neurosurgery simulators and training models. The search terms included “neurosurgery” and “simulation”. Additionally, procedure specific searches were conducted including a combination of the following terms; “ventriculostomy”, “external ventricular drain”, “burr hole”, “craniotomy”, “craniectomy”, “ventriculo-peritoneal shunt”, “lumbo-peritoneal shunt”, “tumour resection”, “dural repair”, “laminectomy”, “discectomy”, “pedicle screw” and “neurosurgery simulation”. Titles and abstracts were screened according to the preferred reporting items for systematic reviews and meta-analysis (PRISMA) guidelines⁵.

Study Eligibility Criteria

Original articles describing the validation and use of a neurosurgery training simulator were included. Models and simulators were then classified into the following categories: Virtual Reality (VR); Augmented Reality (AR); bench; cadaver; human tissue; and animal models. Abstracts, duplicates and non-English articles were excluded, as well as those describing the development of simulators only.

Data Extraction

Articles were first screened based on title and abstract. The remaining results were then examined, and articles were included only if they described a validation process or determined translational outcomes of the simulator.

Data Analysis

Upon selection, training models were identified and outcomes for validation studies were noted. A rating scale devised by Beckman et al.⁶ was used to evaluate the strength of each source of validity evidence based on Messick's framework⁷. A level of effectiveness (LoE) was then assigned to each training model using McGaghie's proposed levels of simulation-based translational outcomes⁸ (Supplementary Tables 1 and 2).

Results

Description of studies

From the 6006 articles retrieved from the search, 102 articles were identified upon screening (Figure 1). Ninety-seven simulators or simulation based training courses were either validated or had a translational outcome described (Table 1). Results were categorised into general neurosurgery; spine surgery; paediatric neurosurgery; endonasal/transsphenoidal surgery; training courses; and non-technical skills.

General neurosurgery

Overall, 47 articles validated 43 simulators for general neurosurgery (Table 2). Of these, 3 simulators were used for multiple procedures, 7 for general neurosurgery procedures, 19 for ventriculostomy, 10 for tumour resection and 19 for vascular neurosurgery.

Mixed simulators & general procedures

Three simulators for multiple procedures were validated or demonstrated a LoE in 4 articles. The remaining 7 articles described a corresponding simulator for a general neurosurgical procedure.

The S.I.M.O.N.T. was developed for training neuroendoscopy, rhinological and endonasal skull base surgery ⁹. In an initial validation study, among 37 surgeons with 9 experienced, the model gained favourable responses and thus achieved a rating 1 for content, with a LoE of 1 also. The S.I.M.O.N.T. was then validated for ventriculostomy and tumour resection among 22 experienced and novice neurosurgeons ¹⁰. A rating 2 for both internal structure and content was achieved, as retest reliability and interrater reliability were determined, and expert opinions were favourable. However, a rating 0 for relations to other variables was awarded as this was discussed, but no data was presented. Finally, a LoE of 2 was awarded as the probability of error reduced in subsequent attempts.

A perfusion-based cadaveric model was used to simulate a number of procedures within a mock operating theatre such as: extracranial-intracranial bypass; endonasal surgery with ICA injury; external ventricular drain (EVD) insertion; spinal dural repair; and endonasal surgery with CSF leak repair ¹¹. Trainees reported improved confidence scores for each procedure, thus demonstrating a LoE of 1. However, no validity was investigated for the model.

The MARTYN bench model was designed to train craniotomy and EVD insertion via burr holes ¹². Both realism and confidence were assessed, with an improvement demonstrated in both parameters. Thus, the model achieved a LoE of 1, and achieved a rating of 1 for content as it was assessed by 5 experts.

The ImmersiveTouch trigeminal rhizotomy simulator demonstrated a rating 2 relations to other variables among 71 residents of differing experience ¹³. However, translational outcomes were not investigated in this study. A rapid-prototyped 3D model for anterior clinoidectomy was evaluated by 10 neurosurgeons, and demonstrated level 1 content ¹⁴. However, no LoE could be determined as trainees were not surveyed. A foramen ovale puncture model achieved a rating 1 content, as five experienced neurosurgeons concluded that it mimicked the human face and was sufficient for training neurosurgeons ¹⁵. Again, no LoE was determined in this study. Another study used the Medtronic platform with a custom 3D model, to simulate navigation, burr hole placement and frameless biopsy ¹⁶. From 6 participants, the model demonstrated a LoE of 2. While there was difference in time and attempts between surgeon and trainees, the lack of quantitative analysis resulted in only a rating 1 for relations to other variables.

The PeriopSim™ by Conquer Mobile has two modules for burr hole surgery and instrument recognition ¹⁷. The simulator was used to improve instrument recognition during burr hole surgery, and demonstrated a LoE of 2 as fewer errors and improved total score occurred on subsequent use. Further, performance was measured using an algorithm to calculate a total score. This earned the model a rating of 2 for response processes. USim is an app-based training approach for intra-operative ultrasound ¹⁸. Among 14 residents, understanding and comprehension levels improved when asked to repeat the initial task, thereby demonstrating a LoE of 2 with contained effects. However, no validity was determined in this preliminary study.

A 3D-printed brain model based on MR images and designed for low cost was rated as useful for training and realistic by 10 neurosurgeons and residents ¹⁹. The model thus earned level 1 content validity as well as a LoE of 1.

Ventriculostomy

Twelve validation studies were identified for 11 ventriculostomy simulators. Nine additional simulators for ventriculostomy are described in their respective sections. These include, two simulators for paediatric endoscopic third ventriculostomy (ETV) ^{20 21}, an adolescent hydrocephalus model for ETV training ²², the perfusion-based cadaveric model, MARTYN bench model and the S.I.M.O.N.T. as described earlier ^{11 12 10}. Finally, in the training course section, the Society of Neurological Surgeons (SNS) Bolivian boot camp, the Fundamentals of Neurosurgery (FNS) training module for ventriculostomy and the Congress of Neurological Surgeons (CNS) trauma module with ETV training have all been validated and are discussed accordingly ^{23 24 25}.

The ImmersiveTouch, a VR simulator, was validated for EVD placement in a multicentre study involving 92 participants ²⁶. The simulator achieved a rating of 2 for both content and relations to other variables, however only achieved a LoE of 1.

Another group described the development of an ImmersiveTouch ventriculostomy placement task and compared it to clinical scenarios to determine how accurate the VR simulation is ²⁷. In the same study, the authors investigated the relationship between catheter placement and year of training, however no relationship was found. Thus a rating 0 was given for relations to other variables and no LoE was investigated in this study. The

same group describe the use of the ImmersiveTouch to simulate a ventriculostomy in a shifted ventricle ²⁸. While no validity was determined, a LoE of 2 was demonstrated among 48 residents as a significant improvement was found on subsequent attempts. Additionally, the ImmersiveTouch was used in another study, in which the same authors describe a virtual brain library for ventriculostomy simulation ²⁹. Here, a rating 2 relations to other variables was achieved, and a LoE of 3 was achieved as participants were followed-up and demonstrated improved rates of first attempt catheter placement.

The introduction of simulation-based workshops using the Rowena model for EVD placement was performed as part of a quality improvement project, and led to a reduction in infection rates, increased satisfactory EVD placement rates, and reduction in displacement ³⁰. While validity was not investigated here, the workshops and simulator notably achieved a LoE of 4.

A hollow 3D-printed head model, designed with fast assembly and low cost, was used to simulate EVD placement for junior residents ³¹. Consequences were discussed by the authors, however no data was presented to support this. A LoE of 2 was however awarded.

A 3D hydrocephalus model for teaching ETV demonstrated content validity with a rating of 1, as well as a LoE of 1 as it was assessed by 15 participants ³². A mixed reality ventriculostomy simulator demonstrated both response processes and relations to other variables with a rating of 2 as an algorithm for timing and grading was used, and a difference between interns and residents was demonstrated, as well as a LoE of 1 ³³. A cost

effective 3D model for cerebral lateral ventriculostomy training was evaluated by 10 medical students and residents, demonstrating a LoE of 1, with no validity investigated ³⁴.

The Medtronic StealthStation, a neuronavigational tool, was used with a resin head model for ventriculostomy among 31 participants ³⁵. In this study, stress, performance and workload were measured earning the simulator a rating 2 for response processes, and while content was discussed, no data was presented. A LoE of 2 was also awarded as results improved with subsequent attempts.

In an initial validation study, Tai et al. demonstrated content validity (rating 1) among 4 faculty neurosurgeons as well as a LoE of 1 for a low-cost ventriculostomy simulator ³⁶. This model was later used with the Ventriculostomy Procedural Assessment Tool (V-PAT) and Objective Structured Assessment of Technical Skills (OSATS) and achieved rating 2 validity for all domains besides content (rating 1) among 14 participants ³⁷. Translational outcomes however, were not investigated.

Tumour resection

Seven simulators were validated in 10 articles for tumour resection. Three simulators are discussed elsewhere, and these include the earlier mentioned S.I.M.O.N.T. ¹⁰. The Fundamentals of Neurosurgery also involved a tumour debulking module ²⁴. Finally, the skull base injectable tumour model is discussed in the endonasal surgery section ³⁸.

Several agar-gelatin models for tumour resection of varying hardness were assessed by Mashiko et al ³⁹. The models demonstrated level 1 content validity among 4 neurosurgeons,

and level 1 relations to other variables as they were able to discriminate between neurosurgeons and residents for tumour resection and operative time, however could not for tumour eloquent area resection. Finally, the model achieved a LoE of 2.

Formalin-fixated and latex-injected cadavers were used for a navigation-guided endoscopic intraventricular injectable tumour model ⁴⁰. The model was used to simulate endoscopic resection of an intraventricular tumour, achieving a LoE of 1, with no validity investigated.

Placenta used for simulating tumour resection demonstrated a rating 1 for content and relations to other variables due to limited expert participant number and also using only time as a measuring tool ⁴¹. The model also achieved a LoE of 1.

An initial study on a phantom-based training system for planning tumour resection, as well as the corresponding craniotomy was assessed by 5 residents ⁴². The model demonstrated a LoE of 1, however validity was not investigated. In another study among 9 participants, the same model demonstrated a rating 2 for response processes, as a standardised protocol was used to assess participants, and a rating 1 for relations to other variables as the model could distinguish between level of training for time to suture and craniotomy ⁴³. Additionally, improvement in time and score was demonstrated after multiple attempts and thus the model achieved a LoE of 2.

The NeuroTouch was again assessed in a study involving a meningioma-like tumour scenario ⁴⁴. Here, the simulator was given favourable responses by survey, demonstrating a LoE of 1, and when analysing performance metrics such as volume resected and duration of excessive

force, the simulator achieved a rating 2 response processes, but only rating 1 for relations to other variables as it could only discriminate between students and residents, but not junior and senior residents.

The NeuroTouch glioblastoma simulation was able to differentiate between 71 medical students and 12 residents by measuring surgical effectiveness and efficiency, as well as demonstrate skill improvement among students ⁴⁵. In doing so, the simulator demonstrated relations to other variables (rating 2) and a LoE of 2.

Another group used other glioma scenarios and identified simulation metrics which assess psychomotor variables such as force applied and volume resected. These metrics were then studied using the NeuroTouch to identify proficiency benchmarks ⁴⁶. Thus the simulator was well validated achieving both relations to other variables (rating= 2) and response processes (rating=2). Translational outcomes were not investigated in this study. More complex scenarios were assessed by the same authors using the same metrics, and the NeuroTouch again demonstrated relations to other variables (rating 2), content (rating 1), response processes (rating 2) and a LoE of 1 as neurosurgeons and residents alike confirmed its usefulness for training and realism ⁴⁷. The NeuroTouch was used with the same metrics to investigate performance among 17 medical students ⁴⁸. Again, the simulator achieved rating 2 for response processes and also for relations to other variables as a difference in metric results was found between medical student deciles, as well as with residents. No LoE was investigated in the study.

Vascular neurosurgery

Twelve models were validated in 13 articles for aneurysm repair, and 3 models for general vascular neurosurgery (Table 3). Four additional models are discussed elsewhere. The earlier mentioned Perfusion-based Cadaveric model was used for extracranial-intracranial bypass and ICA injury in endonasal surgery, and is discussed in the mixed simulators section¹¹. Three models for ICA injury in endonasal surgery were validated, including the adult cadaver head perfusion model⁴⁹, laser-sintered model⁵⁰, and another perfusion-based human cadaver model^{51 52}.

Mashiko et al. developed a series of 3D hollow elastic models, which demonstrated level 1 content, and a LoE of 1 was achieved as the models were rated favourably by junior neurosurgeons⁵³.

Wang et al. compared both a whole and regional 3D-printed model of cerebral aneurysm⁵⁴. No significant difference was found between either models, however they demonstrated a rating 1 for content amongst an unspecified number of neurosurgeons, as well as achieved a LoE of 1. These models were then modified to simulate MCA aneurysm and were evaluated by 6 residents, for the tasks of planning, craniotomy and aneurysm clipping⁵⁵. As the study involved a qualitative survey by non-experts, no validity was demonstrated, however the model did achieve a LoE of 1.

The ImmersiveTouch aneurysm clipping simulator was evaluated by 17 residents as a useful training tool, thus earning a LoE of 1⁵⁶. However, no validity was investigated in the study.

A virtual cerebral aneurysm-clipping model was evaluated by 18 neurosurgeons of differing experience⁵⁷. While a survey demonstrated adequate craniotomy simulation, the clipping procedure was deemed adequate by only 22%. Overall, the model achieved content validity (rating 1) and a LoE of 1.

Human placenta was used for aneurysm clipping and was well validated among 30 participants of differing experience⁵⁸. Indeed, the model achieved a rating 2 for each of response processes, internal structure, content and relations to other variables. Consequences was not discussed in the study, and while the participants were surveyed as to their perceived improvement in skills on using the model, this was not objectively assessed. Thus the model achieved a LoE of 1 only.

Another study investigated the use of placenta to simulate intracranial-intracranial bypass surgery⁵⁹. The model was well validated, achieving a rating 2 for internal structure as test-retest validity was assessed among 50 healthcare professionals, rating 1 for content among 5 experts, and rating 2 for relations to other variables. No LoE was determined in the study.

Cadaver heads were compared to human placenta by another group in a later study, with human placenta demonstrating superiority for teaching sylvian fissure splitting and aneurysm dissection⁶⁰. The study involved blinding examiners, demonstrating level 2 response processes for both models. Further, level 2 content was achieved by the placenta model as well as a LoE of 5 as use of the placenta model demonstrated satisfactory task completion during live surgery when compared to video training alone or cadaver simulation. The cadaver only achieved level 1 content as some parts of the task were not

rated by the neurosurgeons and also achieved a LoE of 5, however, it should be noted that for fissure and aneurysm dissection no skill benefit was found in live surgery.

A live cadaver model with artificial aneurysms and an immersive simulation environment were used in several training courses⁶¹. Upon evaluation by 91 participants, the live model achieved level 2 content and a LoE of 1.

Liu et al. describe the validation of a 3D cerebral aneurysm simulator, and demonstrated level 1 content validity among 6 neurosurgeons, as well as a LoE of 1 from 4 other medical students⁶².

A training program for aneurysm clipping involved practice on 3D models followed by feedback from a senior neurosurgery and a subsequent trial⁶³. Trainees who had performed clipping previously demonstrated superior skill, and skill improved on subsequent trials. Overall the program achieved a LoE of 2, and demonstrated limited validity for relations to other variables due to low participant number (n=6).

Venous sinus injury with a complication of air embolus was simulated on a bench model and assessed by 12 participants⁶⁴. Heart rate was monitored as a surrogate measure of anxiety and compared with a faculty neurosurgeon, and thus a rating 1 for both response processes and relations to other variables was awarded. A LoE of 1 was achieved as the simulator was rated favourably by trainees. The ImmersiveTouch hemostasis simulation was surveyed by 54 participants, and achieved a LoE of 1 with no validity investigated⁶⁵.

Spine surgery

Eighteen simulators for spine surgery were validated, with 5 training modules incorporated into training courses as well (Table 4).

Three simulators and training courses are discussed elsewhere. The perfusion-based cadaveric model for spinal dural repair is discussed in the mixed simulator section ¹¹. The CURE fellowship for spina bifida training is discussed in the paediatric neurosurgery section. Finally, the SNS fundamentals curriculum included lumbar torso drain insertion kits, and is discussed in the training program section ⁶⁶.

The CNS developed 5 training modules that were incorporated into simulation courses or fellowship training. This includes a durotomy repair module used to train CSF leak repair which demonstrated a LoE of 2 among 4 participants, as improvements in mean time for closure as well as leak rate occurred ⁶⁷. However, validity was not investigated in the study. Both cervical foraminotomy and dural repair modules from a simulation course by the CNS were incorporated into a residency program at Thomas Jefferson University Hospital ⁶⁸. In doing so, the modules demonstrated a LoE of 2, as users improved in both knowledge and skills, as well as response processes (rating 2), however the modules failed relations to other variables, as no significant different in OSATS was found with advanced years of training. Additionally, the impact of microanastomosis, anterior cervical discectomy and fusion (ACDF), posterior cervical fusion (PCF), and durotomy repair modules on cognitive and technical skills was assessed with significant improvements in OSATS and knowledge ⁶⁹. This course was tested on 20 residents at the 2013 Neurological Society of India Meeting. Overall, it achieved a LoE of 2 as well as rating 2 for response processes. The anterior

cervical discectomy and fusion simulator training module developed by the CNS, involved didactic teaching and use of a simulator and was assessed amongst 6 participants of varying experience ⁷⁰. An improvement in knowledge was demonstrated and thus the simulator was awarded a LoE of 2. No validity was investigated in this pilot study. The CNS also developed a cervical spine simulator for posterior foraminotomy and laminectomy, and the simulator was assessed amongst 11 participants ⁷¹. The OSATS was used to assess technical ability, and an improvement was demonstrated after both didactic and technical components. Thus the simulation module achieved a LoE of 2, with a rating 2 for response processes.

A combined bench and virtual reality model for pedicle screw placement and lumbar stenosis decompression was evaluated by experienced spinal surgeons, using a validated questionnaire and demonstrated content validity (rating 2), with no LoE determined in the study ⁷².

A low cost dural repair model was used among 13 trainees, who demonstrated improved time and OSATS rating by two blinded consultant surgeons, and thus the model achieved a LoE of 2 and a rating 2 for response processes ⁷³.

A randomized trial to assess the efficacy of a 3D software-based pedicle screw simulator was performed with subjects practising on the simulator prior to screw insertion into a cadaver ⁷⁴. From 17 respondents, the majority gave favourable responses and the simulator thus achieved a LoE of 1. It should also be noted that the subject group failed to show any improvement, and no validity was demonstrated.

A minimally invasive spine surgery model was evaluated by 8 residents and involved both animal and bench components for bilateral laminectomy training⁷⁵. The model was given a LoE of 1 as it received favourable ratings by survey, with no validity investigated.

The Simulated Lumbar Minimally Invasive Surgery Education Model involved didactic teaching as well as use of an image-guided pedicle screw placement simulator⁷⁶. Scores in knowledge and technical placement improved with repeated attempts, and thus the model was awarded a LoE of 2. No validity was investigated.

In a randomised trial involving 10 participants, the study group was exposed to a laboratory based spinal fixation program involving a cadaveric specimen with the Stealth S7 and Sawbones Models⁷⁷. Both groups then performed screw placement on cadavers. Examiners were blinded, and so a rating of 2 for response processes was allocated, and a LoE of 2 was allocated as the study group produced fewer errors.

The ImmersiveTouch was used to simulate thoracic pedicle screw placement, and learning retention was then evaluated⁷⁸. Amongst 51 participants, performance accuracy significantly improved, and the simulator was thus awarded a LoE of 2, although no validity was investigated.

The same authors then assessed the ImmersiveTouch for percutaneous spinal placement amongst 63 participants⁷⁹. Again, an improvement in performance accuracy was demonstrated and the simulator achieved a LoE of 2, with no validity investigated.

A Saw Bones model for pedicle screw placement training in scoliosis demonstrated a rating 2 for response processes as an adapted version of the OSATS was used, however, a difference between junior and senior trainees was not found overall, but only in the domain of time and motion, thus achieving a rating 1 for relations to other variables was achieved⁸⁰. No LoE was determined in the study.

The ImmersiveTouch was assessed for pedicle screw placement in another randomised trial involving 26 medical students⁸¹. Here, the simulator group demonstrated a reduced average error rate when compared to traditional teaching. Thus, the simulator demonstrated an LoE of 2, and no validity was determined in the study.

A Desktop-based computer assisted orthopaedic training system for pedicle screw insertion was assessed among 12 participants⁸². While the authors claimed their scoring system was validated among 6 experts, no data was presented, however examiners were blinded and thus a rating 2 for response processes, and rating 0 for internal structure was allocated. Mean score improved with training and thus a LoE of 2 was achieved.

A Life-Sized 3D Spine Model for pedicle screw training demonstrated a learning effect and thus achieved a LoE of 2 as placement accuracy and time both improved amongst 2 novice surgeons⁸³. No validity was determined in the study.

Bioskills training modules involving sawbone models were used to train both lumbar pedicle screw placement⁸⁴ and lumbar laminectomy⁸⁵ by the same group and assessed in a randomised trial. Both studies described the use of the OSATS, and the PPDIS and thus a

rating 2 for response processes was achieved. It should be noted that while both modules achieved a LoE of 2 as knowledge improved, only the lumbar laminectomy module demonstrated superior OSATS and thus skills when compared to controls.

An immersive artificial wetlab training system for lumbar discectomy was described by Adermann et al⁸⁶ and achieved a rating 1 for content among 17 spinal surgeons, as only the realism and haptic element was validated. An adapted form of the OSATS was referred to but not detailed, and thus a rating 1 for response processes was allocated. A LoE of 1 was also allocated to the model.

One study compared 3 types of cadavers, embalmed with either the Thiel method, the Crosado method, or with formaldehyde⁸⁷. Here, the Thiel cadaver rated the highest, and the formaldehyde the lowest, and as the surgeons provided a favourable rating for cadaver use in spinal surgery simulation, a rating 1 for content was allocated. The cadavers also achieved a LoE of 1.

The virtual surgical training system for cervical pedicle screw placement training was assessed, and when compared with controls, use of the virtual system led to more accurate insertion of the pedicle screw⁸⁸. Thus a LoE of 2 was achieved, however, no validity was determined.

A 3D patient specific rendering for pedicle screw insertion was used by 2 junior surgeons and 2 experts⁸⁹. Here the simulator achieved a LoE of 2 as well as a rating 1 for relations to

other variables as experienced surgeons performed the tasks in quicker time, and junior surgeons improved in time to complete the task.

Endonasal/transsphenoidal surgery

Fourteen articles validated 13 skull base surgery simulators (Table 5). Additionally, two simulators are discussed elsewhere. The Fundamentals of Neurosurgery course involves an endoscopic nasal navigation module and is discussed in the training course section ²⁴. Described in the mixed simulator section is the perfusion-based cadaveric model, used to simulate a number of procedures including endonasal surgery with ICA injury and CSF leak repair ¹¹. In another article, the same model was described but evaluated specifically for endoscopic endonasal CSF leak repair among 9 residents ⁹⁰. Here the model gained favourable responses by survey, and thus achieved a LoE of 1, with no validity investigated.

A 3D-printed skull base model for transnasal endoscopic skull base surgery training demonstrated some content validity (rating 1) amongst 3 experienced neurosurgeons, and achieved a LoE of 1 as 10 Residents provided favourable responses concerning the use of the model for surgical education ⁹¹.

Gagliardi et al. used the skull base injectable tumour model to train the use of the Myriad tool for endoscopic skull base procedures ³⁸. In this study, endonasal resection of tumours was performed by 6 surgeons, who confirmed the educational characteristics of the model by questionnaire. Overall the model achieved an LoE of 1, as well as demonstrating limited content validity (rating 1).

An adult cadaver head perfusion model was used to simulate carotid injury control using muscle graft during endonasal surgery ⁴⁹. As blood loss was reduced in subsequent sessions by learners, a LoE of 2 was achieved by the model as better management was indicated. No validity was discussed for the technical skills component of this model.

Another study validated the Neuro-Endo Trainer (NET), a part task simulator based on CT scans of patients undergoing sellar-suprasellar-parasellar endoscopic endonasal surgery ⁹². The NET demonstrated level 1 relations to other variables as a statistical difference was noted between experts and novices in only a few tasks, and level 1 response processes as a Skills Assessment Scale was used. A LoE of 1 was awarded due to favourable responses post-training.

Endonasal Drilling was simulated in a bench model by Tai et al., and was tested for content amongst 8 neurosurgeons and residents. The model achieved a rating of 1, and a LoE of 1 also ⁹³.

Drilling skills were also improved in a different study when using a chicken egg and skull model ⁹⁴. Here, a rating 2 for relations to other variables as well as a LoE of 2 was achieved among 9 participants as a difference in area removed was found to improve with successive attempts and differed from experts.

A laser-sintered model was used to simulate ICA injury during endonasal surgery and demonstrated a LoE of 2 with an improvement in skills among 46 participants ⁵⁰. However, no validity was investigated.

Another group used a perfusion-based human cadaver model for ICA injury and demonstrated a LoE of 1 as it received favourable ratings by trainees ⁵¹. The same group attempted to demonstrate relations to other variables, however failed to do so as no difference was found between year of training or specialty ⁵². In this study however, an improvement in ability was demonstrated and the model was then awarded a LoE of 2.

A practical 3D-printed simulator for basic operational skills was evaluated among 18 participants ⁹⁵. A rating 2 for relations to other variables was allocated, as significant differences in drilling, curetting and aspirating scores were achieved. Further, 5 trainees demonstrated an improvement and thus a LoE of 2 was awarded to the model.

Another study assessed the content validity of 3D skull base models with pre-existing pathology ⁹⁶. A rating 1 was awarded for content among 15 ENT surgeons of varying experience, and a LoE of 1 was achieved also. A synthetic 3D cranial base model achieved favourable responses in successive sessions from self-assessment questionnaires and thus achieved a LoE of 1 ⁹⁷. No validity was investigated in this study however.

The NeuroTouch endoscopic endonasal module was used in a randomized study to assess whether simulator use would translate to superior outcomes in the operating theatre among 6 participants ⁹⁸. The simulator achieved a rating 2 for response processes as examiners were blinded, and a LoE of 5 was awarded as participants were evaluated over a 6 month period, and demonstrated superior outcomes compared to the untrained group.

Paediatric neurosurgery

Seven simulators and 1 training course was validated for paediatric neurosurgery training in 9 articles (Table 6). Additionally, discussed in the training program section is the fundamentals curriculum which included paediatric head models by the SNS^{66 99}.

The babyMARTYN model was used by 18 trainee neurosurgeons for a series of procedures including, posterior fossa haematoma evacuation, pterional craniotomy, fontanelle tapping, and EVD insertion¹⁰⁰. A LoE of 2 was achieved with improvement in post-training ratings and both content validity (rating 1) and response processes (rating 2) were achieved as the Physician Performance Diagnostic Inventory scale (PPDIS) was used.

A low-cost, synthetic and reusable simulator for endoscopic third ventriculostomy (ETV), called the SickKids simulator, demonstrated limited content validity among 4 neurosurgeons and 5 fellows, and achieved a LoE of 1²⁰. The SickKids simulator was then compared to The NeuroTouch simulator for ETV in a later study by Breimer et al.²¹. While both simulators demonstrated a LoE of 1 amongst 23 residents, and limited content validity (level=1) amongst 3 fellows, the authors concluded that the NeuroTouch was superior for anatomical learning whilst the bench model was superior for learning techniques and instrument familiarisation.

A full scale 3D adolescent hydrocephalus model for ETV training was evaluated by 17 subjects and demonstrated strong validity for response processes, internal structure and relations to other variables achieving a rating of 2 in each domain²². However, the model only achieved a LoE of 1.

The CURE Hydrocephalus and Spina Bifida fellowship combines anatomy review, web-based knowledge assessment, clinical exposure and an endoscopic simulation lab ¹⁰¹. Designed for low resource settings, this fellowship demonstrated improved academic output and clinical output resulting in level 2 consequences and a LoE of 5 as changes were demonstrated in various institutions after the fellowship.

Synthetic simulators for endoscope-assisted repair of metopic and sagittal craniosynostosis were evaluated by 25 participants by survey and demonstrated a rating 1 for content as well as a LoE of 1 ¹⁰².

A Prototype model for spinal detethering surgeries demonstrated a LoE of 1 as users regarded the experience as realistic, however, despite a correlation between user experience and assigned score, no mention of a significant difference was made and thus, only a rating of 0 for relations to other variables was awarded ¹⁰³.

A 3D Synthetic Model for numerous lumbar pathologies including tethered cord, fatty filum terminale, meningocele, myelomeningocele and lipomyelomeningocele was validated among 7 participants ¹⁰⁴. A LoE was not determined in the study, and the simulator failed to demonstrate a performance difference in skill levels based on pressure measurements. When participants were assessed by expert neurosurgeons however, a significant qualitative difference was found and thus a level 1 for relations to other variables was achieved.

Non-technical skills

An adapted Laerdal SimMan with cadaver head was used as part of a crisis management simulation involving dual neurosurgeon and anaesthetist participation (Table 4) ¹⁰⁵. Post-simulation surveys provided favourable responses to the simulation, demonstrating a LoE of 1, however little validity was demonstrated with level 0 achieved in relations to other variables as neurosurgeons were compared with anaesthetists with no significant difference.

A task and crisis simulator for vertebroplasty in an immersive operating theatre was rated well for training and realism, by both expert and junior neurosurgeons ¹⁰⁶. The VR-based simulator achieved a LoE of 1, and a rating 1 for content.

The earlier mentioned adult cadaveric head perfusion model by Ciporen et al., also assessed NTS with improved performance for situation awareness, decision making, communication, teamwork and leadership among 4 residents ⁴⁹. This earned the model a LoE of 2, and the simulator was also able to discriminate between experience of 6 residents thereby demonstrating rating 1 for relations to other variables.

Curriculum/Training Courses

Eight training courses were validated in 9 articles (Table 7). Additionally, 5 spinal surgery courses are discussed in the respective section.

The SBNS-accredited Neurosurgical Skills Workshop aimed to provide experience in positioning, burr hole creation, ventricular access, and flap formation to medical students

and foundation trainees with neurosurgical trainee mentors ¹⁰⁷. While no validity was determined, an improvement in knowledge was demonstrated and thus a LoE of 2 was awarded.

The SNS developed a fundamentals curriculum for early trainees which involved lectures and skills stations involving various bench models such as paediatric head models, lumbar torso and drain insertion kits, and plastic craniums ⁶⁶. Overall, the curriculum was well received by faculty neurosurgeons achieving a rating 2 for content, and a LoE of 2 as knowledge levels improved after the course. Participants were then contacted 6 months later and surveyed as to their knowledge retention and course effectiveness for improving patient care ⁹⁹. Thus the curriculum achieved a LoE of 3 as downstream effects was achieved.

The SNS boot camp was expanded to Bolivia, and involved didactic teaching with simulation-based training for both beginner and intermediate procedures including; ventriculostomy, lumbar puncture, craniotomy, and dural closure ²³. Whilst no validity was determined, the boot camp achieved a LoE of 1 due to favourable responses in a post-training survey.

The Fundamentals of Neurosurgery (FNS) involves a series of training modules including ventriculostomy, endoscopic nasal navigation, tumour debulking, hemostasis, and microdissection for technical skills acquisition in neurosurgical oncology ²⁴. These tasks were validated among an unspecified number of surgeons, who were deemed 'subject matter experts'. A rating 1 for content was thus awarded. No LoE was determined in the initial study.

The CNS developed a presigmoid approach model for drilling simulation, and an improvement in proficiency was found after repeated attempts at the simulator ¹⁰⁸. The model thus achieved a LoE of 2, however no validity was investigated.

Additionally, a trauma module involving didactic teaching followed by an ImmersiveTouch ventriculostomy simulation was evaluated by 12 residents ²⁵. Overall, the module achieved a rating 1 for response processes, as some evaluation measures were calculated by the VR simulator itself, and a rating 2 for relations to other variables as senior residents performed significantly better. The module also achieved a LoE of 2 as participants improved with use of the simulator and gained knowledge from the didactic component. Further, a craniotomy training model was incorporated into the trauma module and evaluated in a different study ¹⁰⁹. The model failed to distinguish between a junior and senior group for certain measures such as time taken, but did manage to for measures such as burr hole placement and thus achieved a rating 1 for relations to other variables. Also, a LoE of 2 was achieved as both technical skills and knowledge improved following the module.

A simulation-based curriculum with 68 exercises including spine surgery, basic skills, skull base models, and endovascular cases was evaluated by survey among 180 residents. The curriculum made use of sawbones models, the ImmersiveTouch platform and cadavers. Self-reported improvements in proficiency were significant, and thus the curriculum achieved a LoE of 1 ¹¹⁰.

Discussion

The present study highlights a great number of simulators within neurosurgery which have each demonstrated a variable extent of validity. No studies showed complete validity for all domains, and only 6 simulators or training programs were able to demonstrate the impact of using the simulator in the operating theatre and beyond, thus achieving levels of effectiveness greater than 2. Current data demonstrate a strong belief within the specialty that simulation tools could both supplement current training, and potentially improve patient outcomes ^{111 112}. The reality of such perceptions must be verified, and further studies are required to determine the 'translational outcomes' of many of the identified simulators.

With 96 simulators or training courses identified, neurosurgery employs more simulation-based training than many other specialities. Simulation in neurosurgery surpasses that of otolaryngology and orthopaedics, comparing well with urology, which may be regarded at the forefront of simulation-based training ^{113–115}. While an overlap between the specialties must be accounted for, as in the example of spine surgery, it is promising to see a great number of simulators that have been validated or have shown levels of effectiveness.

In particular, the creation of a Simulation Committee by the CNS, has done much to improve simulation-based training in neurosurgery ¹¹⁶. However, the downstream effects of such training have been less thoroughly investigated. Indeed, none of the CNS training courses reported translational outcomes in the operating theatre and thus had limited levels of effectiveness as demonstrated in the present study. Therefore, greater efforts are needed in this respect. The SNS is another group committed to implementing simulation-

based training, and notably launched a training camp in Bolivia. The CURE fellowship is another example of improving access to simulation-based training, as it provides subspecialty training in Uganda. The latter achieved the highest level of effectiveness in the review. Such programs have demonstrated the wider benefits of simulation-based training in resource-limited regions.

The use of modern definitions which recognise validity as a unitary construct enables a more rigorous assessment of the suitability of these models for training future neurosurgeons¹¹⁷. However, few studies identified actually used Messick's framework when validating the simulators. An example would be that many studies did not involve the input of experts and thus were limited in their determination of content validity, as outdated definitions were used. This is consistent with surgical simulation as a whole, in which only 6.6% of studies have been reported to use these definitions for validity assessment¹¹⁸. An emphasis must therefore be made for validation studies within neurosurgery to adopt modern validity definitions, thereby enabling a more informative appraisal of any simulators or training courses in development.

Despite a great number of simulators, there were very few for non-technical skills. Non-technical skills cover teamwork, decision making, communication and situational awareness. Indeed, demand for such training is increasing amongst neurosurgical trainees particularly when compared to senior tutors¹¹⁹. Thus, non-technical skills training will play a greater role in the future of neurosurgery, and development of further training modalities for this is necessary.

Full immersion simulation is widely regarded as an important modality for training, however the cost and maintenance of such facilities presents an obstacle to their accessibility and implementation ¹²⁰. An example of immersive simulation includes the perfusion-based cadaveric system by Zada et al. which was used in a mock operating theatre ¹¹. Virtual reality simulators such as the ImmersiveTouch and Neurotouch can provide a solution with ever increasing realism, including 3D technology to enable sounds of the operating theatre to be incorporated ¹²¹. However, the limitations of tactile feedback and interaction remain and thus VR remains suitable for initial training only. Overall, incorporating different modalities combined with immersive training is recommended.

The cost of simulation-based training is well demonstrated by the simulation curriculum developed by Gasco et al. ¹¹⁰. The authors mention the benefit of demonstrating translational outcomes of simulation-based training in obtaining financial support for such programs. Therefore, the benefits of investigating validity and translational outcomes are not unique to the simulator, but provide impetus for the use of simulation-based training as a whole. The advent of 3D-printing has enabled many low-cost models to be developed and such simulators have been validated ¹⁹³¹³⁴⁶². Ultimately, the importance of having multi-modal training, balanced with the need for both high- and low- cost options, will determine the progress of simulation-based training in neurosurgery.

Limitations

Neurosurgery within the United Kingdom involves training for spinal and paediatric procedures. Despite a rigorous search, some articles may have been missed as a result of this overlap with spinal orthopaedics and paediatric surgery. Furthermore, some

procedures are performed by interventional radiologists in the United Kingdom and are thus not part of the neurosurgical training program, as they are in other countries. An example would be cerebral angiography, and therefore certain simulators or training courses were omitted¹²².

Conclusion

Simulation training in neurosurgery is a progressing field with an increasing number of simulation-based models being developed. While some validation of these models has been performed, there is much progress to be made, with particularly few studies investigating translational outcomes to the operating theatre and beyond. The development of various training curricula is apparent, but again, greater efforts must be made to determine the translational benefit of these courses. Finally, utilising full immersion simulation, providing cost-effective training and developing simulators for non-technical skills is also warranted.

REFERENCES

1. Kerr B, O'Leary JP. The training of the surgeon: Dr. Halsted's greatest legacy. *Am Surg.* 1999. doi:10.3406/colan.1978.1208
2. Cobb MPH, Taekman JM, Zomorodi AR, Gonzalez LF, Turner DA. Simulation in Neurosurgery—A Brief Review and Commentary. *World Neurosurg.* 2016. doi:10.1016/j.wneu.2015.11.068
3. Akhigbe T, Zolnourian A, Bulters D. Mentoring models in neurosurgical training: Review of literature. *J Clin Neurosci Off J Neurosurg Soc Australas.* 2017;45:40-43. doi:10.1016/j.jocn.2017.07.025
4. Dawe SR, Windsor JA, Broeders JAJL, Cregan PC, Hewett PJ, Maddern GJ. A systematic review of surgical skills transfer after simulation-based training: laparoscopic cholecystectomy and endoscopy. *Ann Surg.* 2014;259(2):236-248. doi:10.1097/SLA.0000000000000245
5. Panic N, Leoncini E, De Belvis G, Ricciardi W, Boccia S. Evaluation of the endorsement of the preferred reporting items for systematic reviews and meta-analysis (PRISMA) statement on the quality of published systematic review and meta-analyses. *PLoS One.* 2013. doi:10.1371/journal.pone.0083138
6. Beckman TJ, Cook DA, Mandrekar JN. What is the validity evidence for assessments of clinical teaching? *J Gen Intern Med.* 2005;20(12):1159-1164. doi:10.1111/j.1525-1497.2005.0258.x
7. Messick S. Meaning and Values in Test Validation: The Science and Ethics of Assessment. *Educ Res.* 1989. doi:10.3102/0013189X018002005
8. McGaghie WC, Issenberg SB, Barsuk JH, Wayne DB. A critical review of simulation-based mastery learning with translational outcomes. *Med Educ.* 2014;48(4):375-385. doi:10.1111/medu.12391
9. Zymberg S, Vaz-Guimaraes Filho F, Lyra M. Neuroendoscopic training: presentation of a new real simulator. *Minim Invasive Neurosurg.* 2010;53(1):44-46. doi:10.1055/s-0029-1246169
10. Filho FVG, Coelho G, Cavalheiro S, Lyra M, Zymberg ST. Quality assessment of a new surgical simulator for neuroendoscopic training. *Neurosurg Focus.* 2011;30(4):E17. doi:10.3171/2011.2.focus10321
11. Zada G, Bakhsheshian J, Pham M, et al. Development of a Perfusion-Based Cadaveric Simulation Model Integrated into Neurosurgical Training: Feasibility Based On Reconstitution of Vascular and Cerebrospinal Fluid Systems. *Oper Neurosurg.* 2018;14(1):72-80. doi:10.1093/ons/oxp074
12. Craven C, Baxter D, Cooke M, et al. Development of a modelled anatomical replica for training young neurosurgeons. *Br J Neurosurg.* 2014;28(6):707-712. doi:10.3109/02688697.2014.913775
13. Shakur SF, Luciano CJ, Kania P, et al. Usefulness of a virtual reality percutaneous trigeminal rhizotomy simulator in neurosurgical training. *Clin Neurosurg.* 2015;11:420-425. doi:10.1227/NEU.00000000000000853
14. Okonogi S, Kondo K, Harada N, Masuda H, Nemoto M, Sugo N. Operative simulation of anterior clinoidectomy using a rapid prototyping model molded by a three-dimensional printer. *Acta Neurochir (Wien).* 2017;159(9):1619-1626. doi:10.1007/s00701-017-3202-4
15. Almeida DB, Hunhevicz S, Bordignon K, et al. A model for foramen ovale puncture training: Technical note. *Acta Neurochir (Wien).* 2006;148(8):881-883; discussion 883. doi:10.1007/s00701-006-0817-2
16. Waran V, Narayanan V, Karuppiah R, et al. Injecting realism in surgical training-initial simulation experience with custom 3D models. *J Surg Educ.* 2014;71(2):193-197. doi:10.1016/j.jsurg.2013.08.010
17. Clarke DB, Kureshi N, Hong M, Sadeghi M, D'Arcy RCN. Simulation-based training for burr hole surgery instrument recognition. *BMC Med Educ.* 2016;16:153. doi:10.1186/s12909-016-0669-2

18. Perin A, Prada FU, Moraldo M, et al. USim: A New Device and App for Case-Specific, Intraoperative Ultrasound Simulation and Rehearsal in Neurosurgery. A Preliminary Study. *Oper Neurosurg (Hagerstown, Md)*. 2018;14(5):572-578. doi:10.1093/ons/oxp144
19. Ploch CC, Mansi CSSA, Jayamohan J, Kuhl E. Using 3D Printing to Create Personalized Brain Models for Neurosurgical Training and Preoperative Planning. *World Neurosurg*. 2016;90:668-674. doi:10.1016/j.wneu.2016.02.081
20. Breimer GE, Haji FA, Cinalli G, Hoving EW, Drake JM. Validity Evidence for the Neuro-Endoscopic Ventriculostomy Assessment Tool (NEVAT). *Oper Neurosurg*. 2015;15(1):1. doi:10.1227/NEU.0000000000001158
21. Breimer GE, Haji FA, Bodani V, et al. Simulation-based Education for Endoscopic Third Ventriculostomy. *Oper Neurosurg*. 2016;13(1):1. doi:10.1227/NEU.0000000000001317
22. Weinstock P, Rehder R, Prabhu SP, Forbes PW, Roussin CJ, Cohen AR. Creation of a novel simulator for minimally invasive neurosurgery: fusion of 3D printing and special effects. *J Neurosurg Pediatr*. 2017;20(1):1-9. doi:10.3171/2017.1.PEDS16568
23. Ament JD, Kim T, Gold-Markel J, et al. Planning and Executing the Neurosurgery Boot Camp: The Bolivia Experience. *World Neurosurg*. 2017;104:407-410. doi:10.1016/j.wneu.2017.05.046
24. Choudhury N, Glinas-Phaneuf N, Delorme S, Del Maestro R. Fundamentals of Neurosurgery: Virtual Reality Tasks for Training and Evaluation of Technical Skills. *World Neurosurg*. 2013;80(5):e9-e19. doi:10.1016/j.wneu.2012.08.022
25. Schirmer CM, Elder JB, Roitberg B, Lobel DA. Virtual Reality-Based Simulation Training for Ventriculostomy. *Neurosurgery*. 2013;73(SUPPL. 4):S66-S73. doi:10.1227/NEU.0000000000000074
26. Perin A, Galbiati TF, Gambatesa E, et al. Filling the gap between the OR and virtual simulation: a European study on a basic neurosurgical procedure. *Acta Neurochir (Wien)*. 2018;160(11):2087-2097. doi:10.1007/s00701-018-3676-8
27. Banerjee PP, Luciano CJ, Lemole GM, Charbel FT, Oh MY. Accuracy of ventriculostomy catheter placement using a head- and hand-tracked high-resolution virtual reality simulator with haptic feedback. *J Neurosurg*. 2007;107(3):515-521. doi:10.3171/JNS-07/09/0515
28. Lemole M, Banerjee PP, Luciano C, Charbel F, Oh M. Virtual ventriculostomy with "shifted ventricle": neurosurgery resident surgical skill assessment using a high-fidelity haptic/graphic virtual reality simulator. *Neurol Res*. 2009;31(4):430-431. doi:10.1179/174313208x353695
29. Yudkowsky R, Luciano C, Banerjee P, et al. Practice on an augmented reality/haptic simulator and library of virtual brains improves residents' ability to perform a ventriculostomy. *Simul Healthc*. 2013;8(1):25-31. doi:10.1097/SIH.0b013e3182662c69
30. Dasgupta D, D'Antona L, Cat DA, et al. Simulation workshops as an adjunct to perioperative care bundles in the management of external ventricular drains: improving surgical technique and reducing infection. *J Neurosurg*. October 2018:1-5. doi:10.3171/2018.5.JNS172881
31. Bow H, He L, Raees MA, Pruthi S, Chitale R. Development and Implementation of an Inexpensive, Easily Producing, Time Efficient External Ventricular Drain Simulator Using 3-Dimensional Printing and Image Registration. *Oper Neurosurg (Hagerstown, Md)*. June 2018. doi:10.1093/ons/opy142
32. Waran V, Narayanan V, Karupiah R, et al. Neurosurgical Endoscopic Training via a Realistic 3-Dimensional Model With Pathology. *Simul Healthc J Soc Simul Healthc*. 2015;10(1):43-48. doi:10.1097/SIH.0000000000000060
33. Hooten KG, Lister JR, Lombard G, et al. Mixed Reality Ventriculostomy Simulation. *Neurosurgery*. 2014;10(4):576-581. doi:10.1227/NEU.0000000000000503
34. Russin J, Babiker H, Ryan J, Rangel-Castilla L, Frakes D, Nakaji P. Computational Fluid Dynamics to Evaluate the Management of a Giant Internal Carotid Artery Aneurysm. *World Neurosurg*. 2015;83(6):1057-1065. doi:10.1016/j.wneu.2014.12.038
35. Kirkman MA. Deliberate Practice, Domain-Specific Expertise, and Implications for Surgical

- Education in Current Climes. *J Surg Educ*. 2013;70(3):309-317. doi:10.1016/j.jsurg.2012.11.011
36. Tai BL, Rooney D, Stephenson F, et al. Development of a 3D-printed external ventricular drain placement simulator: technical note. *J Neurosurg*. 2015;123(4):1070-1076. doi:10.3171/2014.12.JNS141867
 37. Rooney DM, Tai BL, Sagher O, Shih AJ, Wilkinson DA, Savastano LE. Simulator and 2 tools: Validation of performance measures from a novel neurosurgery simulation model using the current Standards framework. *Surgery*. 2016;160(3):571-579. doi:10.1016/j.surg.2016.03.035
 38. Gagliardi F, Chau AM, Mortini P, Caputy AJ, Gragnaniello C. Skull Base Neuroendoscopic Training Model Using a Fibrous Injectable Tumor Polymer and the Nico Myriad. *J Craniofac Surg*. 2017;29(1):1. doi:10.1097/SCS.00000000000004042
 39. Mashiko T, Oguma H, Konno T, et al. Training of Intra-Axial Brain Tumor Resection Using a Self-Made Simple Device with Agar and Gelatin. *World Neurosurg*. 2018;109:e298-e304. doi:10.1016/j.wneu.2017.09.162
 40. Ashour AM, Elbabaa SK, Caputy AJ, Gragnaniello C. Navigation-Guided Endoscopic Intraventricular Injectable Tumor Model: Cadaveric Tumor Resection Model for Neurosurgical Training. *World Neurosurg*. 2016;96:261-266. doi:10.1016/j.wneu.2016.04.048
 41. Oliveira MM, Araujo AB, Nicolato A, et al. Face, Content, and Construct Validity of Brain Tumor Microsurgery Simulation Using a Human Placenta Model. *Oper Neurosurg*. 2016;12(1):61-67. doi:10.1227/NEU.0000000000001030
 42. Müns A, Mühl C, Haase R, et al. A neurosurgical phantom-based training system with ultrasound simulation. *Acta Neurochir (Wien)*. 2014;156(6):1237-1243. doi:10.1007/s00701-013-1918-3
 43. Müns A, Meixensberger J, Lindner D. Evaluation of a novel phantom-based neurosurgical training system. *Surg Neurol Int*. 2014;5(Supplement):173. doi:10.4103/2152-7806.146346
 44. Gélinas-Phaneuf N, Choudhury N, Al-Habib AR, et al. Assessing performance in brain tumor resection using a novel virtual reality simulator. *Int J Comput Assist Radiol Surg*. 2014;9(1):1-9. doi:10.1007/s11548-013-0905-8
 45. Holloway T, Lorsch ZS, Chary MA, et al. Operator experience determines performance in a simulated computer-based brain tumor resection task. *Int J Comput Assist Radiol Surg*. 2015;10(11):1853-1862. doi:10.1007/s11548-015-1160-y
 46. AlZhrani G, Alotaibi F, Azarnoush H, et al. Proficiency Performance Benchmarks for Removal of Simulated Brain Tumors Using a Virtual Reality Simulator NeuroTouch. *J Surg Educ*. 2015;72(4):685-696. doi:10.1016/j.jsurg.2014.12.014
 47. Alotaibi FE, AlZhrani GA, Mullah MAS, et al. Assessing Bimanual Performance in Brain Tumor Resection With NeuroTouch, a Virtual Reality Simulator. *Neurosurgery*. 2015;11 Suppl 2:1. doi:10.1227/NEU.0000000000000631
 48. Winkler-Schwartz A, Bajunaid K, Mullah MAS, et al. Bimanual Psychomotor Performance in Neurosurgical Resident Applicants Assessed Using NeuroTouch, a Virtual Reality Simulator. *J Surg Educ*. 2016;73(6):942-953. doi:10.1016/j.jsurg.2016.04.013
 49. Ciporen JN, Lucke-Wold B, Mendez G, Cameron WE, McCartney S. Endoscopic Management of Cavernous Carotid Surgical Complications: Evaluation of a Simulated Perfusion Model. *World Neurosurg*. 2017;98:388-396. doi:10.1016/j.wneu.2016.11.018
 50. Maza G, VanKoeveering KK, Yanez-Siller JC, et al. Surgical simulation of a catastrophic internal carotid artery injury: a laser-sintered model. *Int Forum Allergy Rhinol*. October 2018. doi:10.1002/alr.22178
 51. Pham M, Kale A, Marquez Y, et al. A Perfusion-based Human Cadaveric Model for Management of Carotid Artery Injury during Endoscopic Endonasal Skull Base Surgery. *J Neurol Surg Part B Skull Base*. 2014;75(05):309-313. doi:10.1055/s-0034-1372470
 52. Zada G, Bakhsheshian J, Pham M, et al. Development of a Perfusion-Based Cadaveric Simulation Model Integrated into Neurosurgical Training: Feasibility Based On Reconstitution

- of Vascular and Cerebrospinal Fluid Systems. *Oper Neurosurg*. 2018;14(1):72-80. doi:10.1093/ons/opx074
53. Mashiko T, Otani K, Kawano R, et al. Development of three-dimensional hollow elastic model for cerebral aneurysm clipping simulation enabling rapid and low cost prototyping. *World Neurosurg*. 2015;83(3):351-361. doi:10.1016/j.wneu.2013.10.032
 54. Wang L, Ye X, Hao Q, et al. Comparison of Two Three-Dimensional Printed Models of Complex Intracranial Aneurysms for Surgical Simulation. *World Neurosurg*. 2017;103:671-679. doi:10.1016/j.wneu.2017.04.098
 55. Wang L, Ye X, Hao Q, et al. Three-dimensional intracranial middle cerebral artery aneurysm models for aneurysm surgery and training. *J Clin Neurosci*. 2018;50:77-82. doi:10.1016/j.jocn.2018.01.074
 56. Alaraj A, Luciano CJ, Bailey DP, et al. Virtual Reality Cerebral Aneurysm Clipping Simulation With Real-Time Haptic Feedback. *Neurosurgery*. 2015;11 Suppl 2:1. doi:10.1227/NEU.0000000000000583
 57. Gmeiner M, Dirnberger J, Fenz W, et al. Virtual Cerebral Aneurysm Clipping with Real-Time Haptic Force Feedback in Neurosurgical Education. *World Neurosurg*. 2018;112:e313-e323. doi:10.1016/j.wneu.2018.01.042
 58. Belykh E, Miller EJ, Lei T, et al. Face, Content, and Construct Validity of an Aneurysm Clipping Model Using Human Placenta. *World Neurosurg*. 2017;105:952-960.e2. doi:10.1016/j.wneu.2017.06.045
 59. Oliveira MM, Wendling L, Malheiros JA, et al. Human Placenta Simulator for Intracranial–Intracranial Bypass: Vascular Anatomy and 5 Bypass Techniques. *World Neurosurg*. 2018;119:e694-e702. doi:10.1016/j.wneu.2018.07.246
 60. de Oliveira MMR, Ferrarez CE, Ramos TM, et al. Learning brain aneurysm microsurgical skills in a human placenta model: predictive validity. *J Neurosurg*. 2018;128(3):846-852. doi:10.3171/2016.10.JNS162083
 61. Aboud E, Aboud G, Al-Mefty O, et al. “Live cadavers” for training in the management of intraoperative aneurysmal rupture. *J Neurosurg*. 2015;123(5):1339-1346. doi:10.3171/2014.12.JNS141551
 62. Liu Y, Gao Q, Du S, et al. Fabrication of cerebral aneurysm simulator with a desktop 3D printer. *Sci Rep*. 2017;7:44301. doi:10.1038/srep44301
 63. Mashiko T, Kaneko N, Konno T, Otani K, Nagayama R, Watanabe E. Training in Cerebral Aneurysm Clipping Using Self-Made 3-Dimensional Models. *J Surg Educ*. 2017;74(4):681-689. doi:10.1016/j.jsurg.2016.12.010
 64. Cleary DR, Siler DA, Whitney N, Selden NR. A microcontroller-based simulation of dural venous sinus injury for neurosurgical training. *J Neurosurg*. 2018;128(5):1553-1559. doi:10.3171/2016.12.JNS162165
 65. Gasco J, Patel A, Luciano C, et al. A Novel Virtual Reality Simulation for Hemostasis in a Brain Surgical Cavity: Perceived Utility for Visuomotor Skills in Current and Aspiring Neurosurgery Residents. *World Neurosurg*. 2013;80(6):732-737. doi:10.1016/j.wneu.2013.09.040
 66. Selden NR, Origitano TC, Burchiel KJ, et al. A National Fundamentals Curriculum for Neurosurgery PGY1 Residents. *Neurosurgery*. 2012;70(4):971-981. doi:10.1227/NEU.0b013e31823d7a45
 67. Ghobrial GM, Anderson PA, Chitale R, Campbell PG, Lobel DA, Harrop J. Simulated Spinal Cerebrospinal Fluid Leak Repair. *Neurosurgery*. 2013;73(SUPPL. 4):S111-S115. doi:10.1227/NEU.0000000000000100
 68. Ghobrial GM, Balsara K, Maulucci CM, et al. Simulation Training Curricula for Neurosurgical Residents: Cervical Foraminotomy and Durotomy Repair Modules. *World Neurosurg*. 2015;84(3):751-755.e7. doi:10.1016/j.wneu.2015.04.056
 69. Zammar SG, Hamade YJ, Aoun RJN, et al. The Cognitive and Technical Skills Impact of the Congress of Neurological Surgeons Simulation Curriculum on Neurosurgical Trainees at the

- 2013 Neurological Society of India Meeting. *World Neurosurg.* 2015;83(4):419-423. doi:10.1016/j.wneu.2014.12.006
70. Ray WZ, Ganju A, Harrop JS, Hoh DJ. Developing an Anterior Cervical Discectomy and Fusion Simulator for Neurosurgical Resident Training. *Neurosurgery.* 2013;73(SUPPL. 4):S100-S106. doi:10.1227/NEU.0000000000000088
 71. Harrop J, Rezai AR, Hoh DJ, Ghobrial GM, Sharan A. Neurosurgical Training With a Novel Cervical Spine Simulator. *Neurosurgery.* 2013;73(SUPPL. 4):S94-S99. doi:10.1227/NEU.0000000000000103
 72. Coelho G, Defino HLA. The Role of Mixed Reality Simulation for Surgical Training in Spine. *Spine (Phila Pa 1976).* 2018;43(22):1609-1616. doi:10.1097/BRS.0000000000002856
 73. Ferguson D, Agyemang K, Barrett C, Mathieson C. A low cost dural closure simulation model for tomorrow's spinal neurosurgeons. *Br J Neurosurg.* November 2018:1-4. doi:10.1080/02688697.2018.1540765
 74. Podolsky DJ, Martin AR, Whyne CM, Massicotte EM, Hardisty MR, Ginsberg HJ. Exploring the Role of 3-Dimensional Simulation in Surgical Training. *J Spinal Disord Tech.* 2010;23(8):e70-e74. doi:10.1097/BSD.0b013e3181d345cb
 75. Walker JB, Perkins E, Harkey HL. A NOVEL SIMULATION MODEL FOR MINIMALLY INVASIVE SPINE SURGERY. *Oper Neurosurg.* 2009;65(6 SUPPL. 1):ons188-ons195. doi:10.1227/01.NEU.0000341534.82210.1B
 76. Chitale R, Ghobrial GM, Lobel D, Harrop J. Simulated Lumbar Minimally Invasive Surgery Educational Model With Didactic and Technical Components. *Neurosurgery.* 2013;73(SUPPL. 4):S107-S110. doi:10.1227/NEU.0000000000000091
 77. Sundar SJ, Healy AT, Kshetry VR, Mroz TE, Schlenk R, Benzel EC. A pilot study of the utility of a laboratory-based spinal fixation training program for neurosurgical residents. *J Neurosurg Spine.* 2016;24(5):850-856. doi:10.3171/2015.8.SPINE15119
 78. Luciano CJ, Banerjee PP, Bellotte B, et al. Learning Retention of Thoracic Pedicle Screw Placement Using a High-Resolution Augmented Reality Simulator With Haptic Feedback. *Oper Neurosurg.* 2011;69(SUPPL. 1):ons14-ons19. doi:10.1227/NEU.0b013e31821954ed
 79. Luciano CJ, Banerjee PP, Sorenson JM, et al. Percutaneous Spinal Fixation Simulation With Virtual Reality and Haptics. *Neurosurgery.* 2013;72(SUPPL. 1):A89-A96. doi:10.1227/NEU.0b013e3182750a8d
 80. Tanner G, Vojdani S, Komatsu DE, Barsi JM. Development of a Saw Bones Model for training pedicle screw placement in scoliosis. *BMC Res Notes.* 2017;10(1):678. doi:10.1186/s13104-017-3029-3
 81. Gasco J, Patel A, Ortega-Barnett J, et al. Virtual reality spine surgery simulation: an empirical study of its usefulness. *Neurol Res.* 2014. doi:10.1179/1743132814Y.0000000388
 82. Rambani R, Ward J, Viant W. Desktop-Based Computer-Assisted Orthopedic Training System for Spinal Surgery. *J Surg Educ.* 2014;71(6):805-809. doi:10.1016/j.jsurg.2014.04.012
 83. Park HJ, Wang C, Choi KH, Kim HN. Use of a life-size three-dimensional-printed spine model for pedicle screw instrumentation training. *J Orthop Surg Res.* 2018;13(1):86. doi:10.1186/s13018-018-0788-z
 84. Boody BS, Hashmi SZ, Rosenthal BD, et al. The Effectiveness of Bioskills Training for Simulated Lumbar Pedicle Screw Placement. *Glob Spine J.* 2018;8(6):557-562. doi:10.1177/2192568217743505
 85. Boody BS, Rosenthal BD, Jenkins TJ, Patel AA, Savage JW, Hsu WK. The Effectiveness of Bioskills Training for Simulated Open Lumbar Laminectomy. *Glob Spine J.* 2017;7(8):794-800. doi:10.1177/2192568217703337
 86. Adermann J, Geißler N, Bernal LE, Kotsch S, Korb W. Development and validation of an artificial wetlab training system for the lumbar discectomy. *Eur Spine J.* 2014;23(9):1978-1983. doi:10.1007/s00586-014-3257-3
 87. Babaei M, Ghahramani-Asl R, Sadoughi HR, Sardari D, Shahzadi S. Evaluation of

- Bremsstrahlung radiation dose in stereotactically radiocolloid therapy of cystic craniopharyngioma tumors with 32 P radio-colloid. *Australas Phys Eng Sci Med*. 2018;41(3):697-711. doi:10.1007/s13246-018-0665-4
88. Hou Y, Shi J, Lin Y, Chen H, Yuan W. Virtual surgery simulation versus traditional approaches in training of residents in cervical pedicle screw placement. *Arch Orthop Trauma Surg*. 2018;138(6):777-782. doi:10.1007/s00402-018-2906-0
 89. Xiang L, Zhou Y, Wang H, et al. Significance of preoperative planning simulator for junior surgeons' training of pedicle screw insertion. *J Spinal Disord Tech*. 2015;28(1):E25-E29. doi:10.1097/BSD.0000000000000138
 90. Christian EA, Bakhsheshian J, Strickland BA, et al. Perfusion-based human cadaveric specimen as a simulation training model in repairing cerebrospinal fluid leaks during endoscopic endonasal skull base surgery. *J Neurosurg*. 2018;129(3):792-796. doi:10.3171/2017.5.JNS162982
 91. Zheng J-P, Li C-Z, Chen G-Q, Song G-D, Zhang Y-Z. Three-Dimensional Printed Skull Base Simulation for Transnasal Endoscopic Surgical Training. *World Neurosurg*. 2018;111:e773-e782. doi:10.1016/j.wneu.2017.12.169
 92. Singh R, Baby B, Damodaran N, et al. Design and Validation of an Open-Source, Partial Task Trainer for Endonasal Neuro-Endoscopic Skills Development: Indian Experience. *World Neurosurg*. 2016;86:259-269. doi:10.1016/j.wneu.2015.09.045
 93. Tai BL, Wang AC, Joseph JR, et al. A physical simulator for endoscopic endonasal drilling techniques: technical note. *J Neurosurg*. 2016;124(3):811-816. doi:10.3171/2015.3.JNS1552
 94. Okuda T, Yamashita J, Fujita M, Yoshioka H, Tasaki T, Kato A. The chicken egg and skull model of endoscopic endonasal transsphenoidal surgery improves trainee drilling skills. *Acta Neurochir (Wien)*. 2014;156(7):1403-1407. doi:10.1007/s00701-014-2035-7
 95. Wen G, Cong ZX, Liu KD, et al. A practical 3D printed simulator for endoscopic endonasal transsphenoidal surgery to improve basic operational skills. *Child's Nerv Syst*. 2016;32(6):1109-1116. doi:10.1007/s00381-016-3051-0
 96. Narayanan V, Narayanan P, Rajagopalan R, et al. Endoscopic skull base training using 3D printed models with pre-existing pathology. *Eur Arch Oto-Rhino-Laryngology*. 2015;272(3):753-757. doi:10.1007/s00405-014-3300-3
 97. Muto J, Carrau RL, Oyama K, Otto BA, Prevedello DM. Training model for control of an internal carotid artery injury during transsphenoidal surgery. *Laryngoscope*. 2017;127(1):38-43. doi:10.1002/lary.26181
 98. Thawani JP, Ramayya AG, Abdullah KG, et al. Resident simulation training in endoscopic endonasal surgery utilizing haptic feedback technology. *J Clin Neurosci Off J Neurosurg Soc Australas*. 2016;34:112-116. doi:10.1016/j.jocn.2016.05.036
 99. Selden NR, Anderson VC, McCartney S, Origitano TC, Burchiel KJ, Barbaro NM. Society of Neurological Surgeons boot camp courses: knowledge retention and relevance of hands-on learning after 6 months of postgraduate year 1 training. *J Neurosurg*. 2013;119(3):796-802. doi:10.3171/2013.3.JNS122114
 100. Craven CL, Cooke M, Rangeley C, Alberti SJMM, Murphy M. Developing a pediatric neurosurgical training model. *J Neurosurg Pediatr*. 2018;21(3):329-335. doi:10.3171/2017.8.PEDS17287
 101. Dewan MC, Onen J, Bow H, Ssenyonga P, Howard C, Warf BC. Subspecialty pediatric neurosurgery training: a skill-based training model for neurosurgeons in low-resourced health systems. *Neurosurg Focus*. 2018;45(4):E2. doi:10.3171/2018.7.FOCUS18249
 102. Eastwood KW, Bodani VP, Haji FA, Looi T, Naguib HE, Drake JM. Development of synthetic simulators for endoscope-assisted repair of metopic and sagittal craniosynostosis. *J Neurosurg Pediatr*. 2018;22(2):128-136. doi:10.3171/2018.2.PEDS18121
 103. Bailey J, Nair K, Macuk A, et al. Design and Development of a Modular Medical Simulation Prototype for Pediatric Spinal Detethering Surgeries. *J Med Device*. 2013;7(2):020945.

- doi:10.1115/1.4024304
104. Mattei TA, Frank C, Bailey J, et al. Design of a synthetic simulator for pediatric lumbar spine pathologies. *J Neurosurg Pediatr.* 2013;12(2):192-201. doi:10.3171/2013.4.PEDS12540
 105. Ciporen J, Lucke-Wold B, Dogan A, Cetas J, Cameron W. Dual Endoscopic Endonasal Transsphenoidal and Precaruncular Transorbital Approaches for Clipping of the Cavernous Carotid Artery: A Cadaveric Simulation. *J Neurol Surg Part B Skull Base.* 2016;77(06):485-490. doi:10.1055/s-0036-1584094
 106. Wucherer P, Stefan P, Weidert S, Fallavollita P, Navab N. Task and crisis analysis during surgical training. *Int J Comput Assist Radiol Surg.* 2014;9(5):785-794. doi:10.1007/s11548-013-0970-z
 107. Kamel M, Fountain DM, May P, Ashpole R. Launching an SBNS-accredited neurosurgical skills workshop for medical students and foundation trainees. *Br J Neurosurg.* 2017;31(6):724-726. doi:10.1080/02688697.2017.1354123
 108. Jabbour P, Chalouhi N. Simulation-Based Neurosurgical Training for the Presigmoid Approach With a Physical Model. *Neurosurgery.* 2013;73(SUPPL. 4):S81-S84. doi:10.1227/NEU.0000000000000090
 109. Elsharkawy A, Lehečka M, Niemelä M, et al. Anatomic Risk Factors for Middle Cerebral Artery Aneurysm Rupture. *Neurosurgery.* 2013;73(5):825-837. doi:10.1227/NEU.0000000000000116
 110. Gasco J, Holbrook TJ, Patel A, et al. Neurosurgery simulation in residency training: feasibility, cost, and educational benefit. *Neurosurgery.* 2013;73 Suppl 1(SUPPL. 4):39-45. doi:10.1227/NEU.0000000000000102
 111. Ganju A, Aoun SG, Daou MR, et al. The Role of Simulation in Neurosurgical Education: A Survey of 99 United States Neurosurgery Program Directors. *World Neurosurg.* 2013;80(5):e1-e8. doi:10.1016/j.wneu.2012.11.066
 112. Haji FA, Dubrowski A, Drake J, de Ribaupierre S. Needs assessment for simulation training in neuroendoscopy: a Canadian national survey. *J Neurosurg.* 2013;118(2):250-257. doi:10.3171/2012.10.JNS12767
 113. Musbahi O, Aydin A, Al Omran Y, Skilbeck CJ, Ahmed K. Current Status of Simulation in Otolaryngology: A Systematic Review. *J Surg Educ.* 2017. doi:10.1016/j.jsurg.2016.09.007
 114. Morgan M, Aydin A, Salih A, Robati S, Ahmed K. Current Status of Simulation-based Training Tools in Orthopedic Surgery: A Systematic Review. *J Surg Educ.* 2017. doi:10.1016/j.jsurg.2017.01.005
 115. Aydin A, Shafi AMA, Shamim Khan M, Dasgupta P, Ahmed K. Current Status of Simulation and Training Models in Urological Surgery: A Systematic Review. *J Urol.* 2016. doi:10.1016/j.juro.2016.01.131
 116. Harrop J, Lobel DA, Bendok B, Sharan A, Rezai AR. Developing a neurosurgical simulation-based educational curriculum: an overview. *Neurosurgery.* 2013;73 Suppl 1:25-29. doi:10.1227/NEU.0000000000000101
 117. Noureldin YA, Lee JY, McDougall EM, Sweet RM. Competency-Based Training and Simulation: Making a "Valid" Argument. *J Endourol.* 2018. doi:10.1089/end.2017.0650
 118. Borgersen NJ, Naur TMH, Sørensen SMD, et al. Gathering Validity Evidence for Surgical Simulation: A Systematic Review. *Ann Surg.* 2018. doi:10.1097/SLA.0000000000002652
 119. Khan A, Doke T, Boeris D. Nurturing the Next Generation of Neurosurgeons: How Important Are Nontechnical Skills? *World Neurosurg.* 2018;120:e227-e233. doi:10.1016/j.wneu.2018.08.036
 120. Kassab E, Tun JK, Arora S, et al. "Blowing up the barriers" in surgical training: Exploring and validating the concept of distributed simulation. *Ann Surg.* 2011. doi:10.1097/SLA.0b013e318228944a
 121. Bernardo A. Virtual Reality and Simulation in Neurosurgical Training. *World Neurosurg.* 2017;106:1015-1029. doi:10.1016/j.wneu.2017.06.140
 122. Zammar SG, El Tecle NE, El Ahmadieh TY, et al. Impact of a Vascular Neurosurgery Simulation-

Based Course on Cognitive Knowledge and Technical Skills in European Neurosurgical Trainees. *World Neurosurg.* 2015;84(2):197-201. doi:10.1016/j.wneu.2014.12.001

Table 1: An overview of the available Neurosurgery simulation models described in the literature (2000–2018). Abbreviations: VR- virtual reality, AR- augmented reality, H- high, L- low, Y- yes, N- no

Name of Model (Institution/Manufacturer)	Fidelity	Availability	Type of Model	Describing Study
Mixed Simulators & General Procedures				
S.I.M.O.N.T. Neurosurgical Endotrainer (Discipline of Neurosurgery, Escola Paulista de Medicina da Universidade Federal de São Paulo, Brasil; Pro Delphus Company)	L	Y	Bench	Filho ¹⁰ Zymberg ⁹
Foramen Ovale Puncture Model (Instituto de Neurologia de Curitiba, Curitiba, Brazil)	L	Y	Bench	Almeida ¹⁵
Custom 3D Head Models with Medtronic Platform (Division of Neurosurgery, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia)	H	Y	Bench	Waran ¹⁶
ImmersiveTouch: Trigeminal Rhizotomy Simulator (Department of Neurosurgery, University of Illinois at Chicago, Chicago, Illinois)	H	Y	VR	Shakur ¹³
Perfusion-based Cadaveric Simulation Model (Department of Neurological Surgery, Keck School of Medicine, University of Southern California, Los Angeles, California)	H	Y	Cadaver	Zada ¹¹ Christian ⁹⁰
Modelled Anatomical Replica for Training Young Neurosurgeons ‘MARTYN’ (Royal College of Surgeons England)	L	Y	Bench	Craven ¹²
PeriopSim™ (Conquer Mobile)	L	Y	App	Clarke ¹⁷
USim (Camelot Biomedical Systems, Genova, Italy; Neurosurgery Department, Fondazione IRCCS Istituto Neurologico Nazionale “C.Besta”, Milan, Italy)	L	Y	Bench/App	Perin ¹⁸
Gelatin 3D Brain Model (Department of Mechanical Engineering, Stanford University, Stanford, California, USA)	L	Y	Bench	Ploch ¹⁹
Rapid Prototyped 3D Anterior Clinoidectomy Model (Department of Neurosurgery, Toho University Graduate School of Medicine, Tokyo, Japan)	L	Y	Bench	Okonogi ¹⁴
Ventriculostomy				
Low-cost Ventriculostomy simulator with V-PAT and OSATS assessment tools (Department of Learning Health Sciences, University of Michigan Medical School, Ann Arbor, Michigan, USA)	L	Y	Bench	Rooney ³⁷ Tai ³⁶
University of Florida Mixed Ventriculostomy Simulator (Department of Neurological Surgery, University of Florida, Gainesville, Florida, USA)	H	Y	MR	Hooten ³³

Medtronic StealthStation w/ Resin Head Model (Victor Horsley Department of Neurosurgery, The National Hospital for Neurology and Neurosurgery, Queen Square, University College London Hospitals NHS Foundation Trust, London, United Kingdom; Medtronic, Minneapolis, MN, USA)	H	Y	Bench	Kirkman ³⁵
ImmersiveTouch (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	H	Y	VR	Perin ²⁶
ImmersiveTouch: Virtual Brain Library (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA; Department of Medical Education, College of Medicine, University of Chicago, Chicago, USA)	H	Y	VR	Yudkowsky ²⁹
ImmersiveTouch: Ventriculostomy w/ Shifted Ventricle	H	Y	VR	Lemole ²⁸
ImmersiveTouch: Ventriculostomy Catheter Placement (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	H	Y	VR	Banerjee ²⁷
Hollow 3D Printed Head (Department of Neurosurgery, Vanderbilt University Medical Center, Nashville, Tennessee)	L	Y	Bench	Bow ³¹
Simulation Workshop w/ Rowena (Victor Horsley Department of Neurosurgery, National Hospital for Neurology, Queen Square; Neurodesign Ltd.)	L	Y	Bench	Dasgupta ³⁰
3D Hydrocephalus Model (University of Malaya, Kuala Lumpur, Malaysia)	H	N	Bench	Waran ³²
3D Cerebral Lateral Ventriculostomy Model (Division of Neurological Surgery Barrow Neurological Institute, Phoenix, Arizona, USA)	L	Y	Bench	Ryan ³⁴
Vascular Neurosurgery				
Whole 3D Cerebral Aneurysm Model (Department of Neurosurgery, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China)	L	Y	Bench	Wang ⁵⁴
Regional 3D Cerebral Aneurysm Model (Department of Neurosurgery, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China)	L	Y	Bench	Wang ⁵⁴
3D Whole & Regional Middle Cerebral Aneurysm Model (Department of Neurosurgery, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China)	L	Y	Bench	Wang ⁵⁵
3D Cerebral aneurysm Simulator (School of Mechanical Engineering, Zhejiang University, Hangzhou 310027, China)	L	Y	Bench	Liu ⁶²
Live Cadaver Model (Arkansas Neuroscience Institute, St. Vincent Health System, Little Rock, Arkansas, USA)	H	Y	Cadaver	Aboud ⁶¹
Human Placenta (Barrow Neurological Institute, St. Joseph's Hospital and Medical Center, Phoenix,	H	Y	Human	Belykh ⁵⁸ , de Oliveira ⁶⁰

Arizona)				Tissue	
(Microsurgical Laboratory, Department of Surgery, Federal University of Minas Gerais, Belo Horizonte, Minas Gerais, Brazil)					
Human Cadaver (Microsurgical Laboratory, Department of Surgery, Federal University of Minas Gerais, Belo Horizonte, Minas Gerais, Brazil)	H	Y	Cadaver	de Oliveira ⁶⁰	
Cerebral Aneurysm Clipping Training Program (Department of Neurosurgery, Jichi Medical University, Shimotsuke Tochigi, Japan)	L	Y	Bench	Mashiko ⁶³	
Virtual Cerebral Aneurysm Clipping with Real Time Haptic Force Feedback Model (Department of Neurosurgery, Kepler University Hospital, Linz, Austria)	H	Y	VR	Gmeiner ⁵⁷	
ImmersiveTouch: Aneurysm Clipping (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	H	Y	VR	Alaraj ⁵⁶	
3D Hollow Elastic Models (Medical Simulation Center, Jichi Medical University, Tochigi, Japan)	L	Y	Bench	Mashiko ⁵³	
ImmersiveTouch: Hemostasis Simulation (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	H	Y	VR	Gasco ⁶⁵	
Venous Sinus Injury and Air Embolus Surgical Simulation (Department of Neurological Surgery, Oregon Health & Science University, Portland, Oregon)	H	Y	Bench	Cleary ⁶⁴	
Human Placenta Model: Intracranial Bypass (Placentarium, Federal University of Minas Gerais, Belo Horizonte, Minas Gerais, Brazil; Lysterly Neurosurgical Simulation Laboratory, Baptist Medical Center, Florida, USA)	H	Y	Human Tissue	Oliveira ⁵⁹	
Tumour Resection					
Phantom-based Training System (Department of Neurosurgery, University Hospital Leipzig, Leipzig, Germany)	L	Y	Bench	Müns ⁴² Müns ⁴³	
Agar-Gelatin Tumour resection Models (Department of Neurosurgery, Jichi Medical University, Shimotsuke Tochigi, Japan)	L	Y	Bench	Mashiko ³⁹	
NeuroTouch: Glioblastoma Module (CAE Healthcare, Montreal, Quebec, Canada)	H	Y	VR	Holloway ⁴⁵	
NeuroTouch: Glioma Scenarios (CAE Healthcare, Montreal, Quebec, Canada)	H	Y	VR	Alzhrani ⁴⁶ , Alotaibi ⁴⁷ Winkler-Schwartz ⁴⁸	
NeuroTouch: Meningioma-like convexity tumor (CAE Healthcare, Montreal, Quebec, Canada)	H	Y	VR	Gélinas-Phaneuf ⁴⁴	
Navigation-Guided Endoscopic Intraventricular Injectable Tumor Model (Department of Neurosurgery, St. Louis University, St. Louis, Missouri, USA)	H	Y	Cadaver	Ashour ⁴⁰	
Human Placenta Model (Microsurgical Laboratory, Department of Surgery, Federal University of Minas	H	Y	Human	Oliveira ⁴¹	

Gerais, Belo Horizonte, Minas Gerais, Brazil)			Tissue	
Training Courses				
National Fundamentals Curriculum (Society of Neurological Surgeons)	H	Y	Bench	Selden ⁶⁶ , Selden ⁹⁹
Neurosurgery Boot Camp: Bolivia (Bolivian Society for Neurosurgery; Foundation for International Education in Neurological Surgery; Solidarity Bridge; University of Massachusetts)	H	Y	Bench	Ament ²³
Fundamentals of Neurosurgery (Department of Neurology and Neurosurgery, Montreal Neurological Institute and Hospital, Brain Tumour Research Centre, McGill University, Montréal, Quebec, Canada)	H	Y	VR	Choudhury ²⁴
Trauma Module w/ ImmersiveTouch: Ventriculostomy (Congress of Neurological Surgeons)	H	Y	VR	Schirmer ²⁵
Trauma Module w/ Physical Craniotomy Model (Congress of Neurological Surgeons)	L	Y	Bench	Lobel ¹⁰⁹
Presigmoid Approach Simulation Module (Congress of Neurological Surgeons)	L	Y	Bench	Jabbour ¹⁰⁸
SBNS-accredited Neurosurgical Skills Workshop (Society of British Neurological Surgeons; The Neurology and Neurosurgery Interest Group)	L	Y	Bench	Kamel ¹⁰⁷
Simulation Curriculum (Division of Neurosurgery; and Division of Epidemiology & Biostatistics, University of Texas Medical Branch, Galveston, Texas, USA)	H	Y	Bench/VR/Cadaver	Gasco ¹¹⁰
Durotomy Repair Module (Congress of Neurological Surgeons)	L	Y	Bench	Ghobrial ⁶⁷
Cervical Foraminotomy and Durotomy Repair Modules (Congress of Neurological Surgeons)	L	Y	Bench	Ghobrial ⁶⁸
Microanastomosis, Anterior Cervical Discectomy and Fusion (ACDF), Posterior Cervical Fusion (PCF), and Durotomy Repair Modules (Congress of Neurological Surgeons)	L	Y	Bench	Zammar ⁶⁹
Anterior Discectomy and Fusion Simulator (Congress of Neurological Surgeons)	L	Y	Bench	Ray ⁷⁰
Cervical Spine Simulator (Congress of Neurological Surgeons)	L	Y	Bench	Harrop ⁷¹
Spine Surgery				
Mixed Reality Spine Simulator (Paediatric Neurosurgery Center, Beneficencia Portuguesa Hospital, Sao Paulo, Brazil)	H	Y	Bench/VR	Coelho ⁷²
Low Cost Dural Closure Simulation Model (Department of Neurosurgery, Institute of Neurological Sciences, Glasgow, UK)	L	Y	Bench	Ferguson ⁷³
3D Software-based Pedicle Screw Simulator (Faculty of Medicine, Orthopaedic Biomechanics Laboratory, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada)	L	Y	VR	Podolsky ⁷⁴
Simulated Lumbar Minimally Invasive Surgery Education Model (Department of Neurosurgery, Thomas	L	Y	Bench	Chitale ⁷⁶

Jefferson University Hospital, Philadelphia, Pennsylvania; Mayo Clinic, Rochester, Minnesota, USA)				
Laboratory-based Spinal Fixation Training Program (Case Western Reserve University School of Medicine; Department of Neurological Surgery; Center for Spine Health, Neurological Institute, Cleveland, Ohio)	H	Y	Bench/Cadaver	Sundar ⁷⁷
Minimally Invasive Spine Surgery Simulator (Department of Neurosurgery, University of Mississippi Medical Center, Jackson, Mississippi, USA)	H	Y	Bench/Animal	Walker ⁷⁵
ImmersiveTouch: Percutaneous Spinal Needle Placement (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	H	Y	VR	Luciano ⁷⁹
ImmersiveTouch: Thoracic Pedicle Screw Placement (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	H	Y	VR	Luciano ⁷⁸
Saw Bones Scoliosis Model (Stony Brook University Hospital, Stony Brook, USA)	L	Y	Bench	Tanner ⁸⁰
Desktop-based Computer Assisted Orthopaedic Training System (Department of Computer Sciences, University of Hull, United Kingdom; Department of Orthopaedics, Leeds Teaching Hospital NHS Trust, Leeds, United Kingdom)	L	Y	VR	Rambani ⁸²
Life-Sized 3D Spine Model (Department of Orthopaedic Surgery, Kangnam Sacred Heart Hospital, Hallym University College of Medicine, Seoul, South Korea)	H	N	Bench	Park ⁸³
Bioskills Training Module: Lumbar Pedicle Screw (Northwestern Memorial Hospital, Chicago, Illinois, USA)	H	Y	Bench	Boody ⁸⁴
Bioskills Training Module: Lumbar Laminectomy (Northwestern Memorial Hospital, Chicago, Illinois, USA)	H	Y	Bench	Boody ⁸⁵
Artificial Wetlab Training System for Lumbar Discectomy (Krankenhaus Winsen, Orthopädie, Winsen, Germany)	H	Y	Bench	Adermann ⁸⁶
Cadaveric Spinal Surgery Simulation; Thiel, Crosado and formaldehyde (Health Education Yorkshire and the Humber, University of Leeds, Leeds, United Kingdom)	H	Y	Cadaver	Tomlinson ⁸⁷
Virtual Surgical Training System (Department of Orthopaedic Surgery, Changzheng Hospital, Second Military Medical University, Shanghai, China)	H	Y	VR	Hou ⁸⁸
ImmersiveTouch: Pedicle Screw Placement (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	H	Y	VR	Gasco ⁸¹
3D Patient-Specific Rendering: Pedicle Screw Insertion (Department of Orthopedics, General Hospital of Shenyang Military Area Command of Chinese PLA, Shenyang, Liaoning, China)	H	N	VR	Xiang ⁸⁹
Durotomy Repair Module (Congress of Neurological Surgeons)	L	Y	Bench	Ghobrial ⁶⁷
Cervical Foraminotomy and Durotomy Repair Modules (Congress of Neurological Surgeons)	L	Y	Bench	Ghobrial ⁶⁸

Microanastomosis, Anterior Cervical Discectomy and Fusion (ACDF), Posterior Cervical Fusion (PCF), and Durotomy Repair Modules (Congress of Neurological Surgeons)	L	Y	Bench	Zammar ⁶⁹
Anterior Discectomy and Fusion Simulator (Congress of Neurological Surgeons)	L	Y	Bench	Ray ⁷⁰
Cervical Spine Simulator (Congress of Neurological Surgeons)	L	Y	Bench	Harrop ⁷¹
Endoscopic Endonasal Transphenoidal Surgery				
Endoscopic Endonasal Drilling Model (Department of Learning Health Sciences, University of Michigan Medical School, Ann Arbor, Michigan, USA)	L	N	Bench	Tai ⁹³
Chicken Eggs and Skull Model (Department of Neurosurgery, Kinki University Faculty of Medicine, Osaka, Japan)	L	Y	Bench	Okuda ⁹⁴
Laser Sintered Model (Ohio State University, Columbus, Ohio, USA)	L	Y	Bench	Maza ⁵⁰
Perfusion-Based Human Cadaveric Simulation for ICA injury (Keck School of Medicine, Los Angeles, California, United States)	H	Y	Cadaver	Shen ⁵² Pham ⁵¹
Practical 3D Printed Simulator (Department of Neurosurgery, Jinling Hospital, School of Medicine, Nanjing University, China)	H	Y	Bench	Wen ⁹⁵
3D Endoscopic Skull base models w/ Pre-existing pathology (Division of Neurosurgery, Department of Surgery, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia)	L	N	Bench	Narayanan ⁹⁶
3D Cranial base model (Department of Neurological Surgery, The Ohio State University Wexner Medical Center, Columbus, Ohio, USA)	L	Y	Bench	Muto ⁹⁷
NeuroTouch Simulator: Endoscopic Endonasal Module (CAE Healthcare, Montreal, Quebec, Canada)	H	Y	VR	Thawani ⁹⁸
3D Printed Skull base for Transnasal endoscopic Skull base Surgery (Beijing Neurosurgical Institute, Beijing Tiantan Hospital, Capital Medical University, Beijing, China)	L	N	Bench	Zheng ⁹¹
Neuro-Endo-Trainer (NET) SkullBase-Task-GraspPickPlace (Centre for Biomedical Engineering, Indian Institute of Technology Delhi, New Delhi, India)	L	Y	Bench	Singh ⁹²
Perfusion-based Cadaveric Simulation Model: Endoscopic Endonasal CSF Leak Repair (Department of Neurological Surgery, Keck School of Medicine, University of Southern California, Los Angeles, California)	H	Y	Cadaver	Christian ⁹⁰
Skull Base Injectable Tumor Model (Macquarie Neurosurgery, Australian School of Advanced Medicine, Macquarie University, Sydney, Australia)	L	Y	Bench	Gagliardi ³⁸
Adult Cadaver Head Perfusion Model (Oregon Health & Science University Body Donation Program, Portland, Oregon)	H	Y	Cadaver	Ciporen ⁴⁹

Paediatric neurosurgery

baby Modeled Anatomical Replica for Training Young Neurosurgeons ‘babyMARTYN’ (Victor Horsley Department of Neurosurgery, National Hospital for Neurology and Neurosurgery, Queen Square)	L	Y	Bench	Craven ¹⁰⁰
SickKids brain simulator (Centre for Image Guided Innovation and Therapeutic Intervention, Toronto, Ontario, Canada)	L	Y	Bench	Breimer ²⁰ Breimer ²¹
NeuroTouch Simulator: ETV Module (CAE Healthcare, Montreal, Quebec, Canada)	H	Y	VR	Breimer ²¹
Full Scale Hydrocephalus Head Model (Harvard Medical School, Boston, Massachusetts and Division of Pediatric Neurosurgery, Johns Hopkins Hospital, Baltimore, Maryland, USA)	H	N	Bench	Weinstock ²²
Synthetic Endoscope-assisted Craniosynostosis Models (Center for Image-Guided Innovation and Therapeutic Innovation, The Hospital for Sick Children, Toronto)	H	Y	Bench	Eastwood ¹⁰²
CURE Hydrocephalus and Spina Bifida Fellowship (CURE Childrens Hospital of Uganda, Mbale, Uganda)	H	Y	Bench	Dewan ¹⁰¹
Prototype for Pediatric Spinal Detethering Surgeries (University of Illinois ,College of Medicine at Peoria, Peoria, IL, USA)	H	N	Bench	Bailey ¹⁰³
3D Synthetic Model for Pediatric Lumbar Spine Pathologies (Department of Neurosurgery, University of Illinois College of Medicine, USA)	L	Y	Bench	Mattei ¹⁰⁴

Non-Technical Skills

Crisis Management Simulation: Dual Neurosurgery and Anaesthesia Training Experience (Anesthesiology and Perioperative Medicine, Oregon Health and Science University, Portland, Oregon, United States; Department of Neurosurgery, West Virginia University School of Medicine, Morgantown, West Virginia, United States)	H	Y	Bench	Ciporen ¹⁰⁵
Adult Cadaver Head Perfusion Model (Oregon Health & Science University Body Donation Program, Portland, Oregon)	H	Y	Cadaver	Ciporen ⁴⁹
Task and Crisis Vertebroplasty Simulation (Technische Universitat Munchen, Munich, Germany)	H	Y	VR	Wucherer ¹⁰⁶

Table 2: Validation studies on training models for General Neurosurgery. Abbreviations: LoE- Level of Effectiveness, VR- virtual reality, MR – Mixed Reality.

Model (Institution / Manufacturer)	Type of Model	Study	Validation	n	Participants Demographics	LoE
Mixed Simulators & General Procedures						
S.I.M.O.N.T. Neurosurgical Endotrainer (Discipline of Neurosurgery, Escola Paulista de Medicina da Universidade Federal de São Paulo, Brasil; Pro Delphus Company)	Bench	Zymberg (2010) ⁹	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	37	9 Expert Neurosurgeons 28 Novice Neurosurgeons	1
	Bench	Filho (2011) ¹⁰	Content: 2 Response processes: N Internal structure: 2 Relations to other variables: 0 Consequences: N	22	9 Experts 13 Novices	2
Perfusion-based Cadaveric Simulation Model (Department of Neurological Surgery, Keck School of Medicine, University of Southern California, Los Angeles, California)	Cadaver	Zada (2018) ¹¹	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	52	Residents	1
Modelled Anatomical Replica for Training Young Neurosurgeons ‘MARTYN’ (Royal College of Surgeons England)	Bench	Craven (2014) ¹²	Content: 1 Response processes: N Internal structure N Relations to other variables: N Consequences: N	18	5 Experts 4 Intermediate 9 Novices	1
Percutaneous Trigeminal Rhizotomy Simulator (Department of Neurosurgery, University of Illinois at Chicago, Chicago, Illinois)	VR	Shakur (2015) ¹³	Content: N Response processes: N Internal structure: N Relations to other variables: 2 Consequences: N	71	27 Senior Residents 44 Junior Residents	0
Rapid Prototyped 3D Anterior Clinoidectomy Model (Department of Neurosurgery, Toho University Graduate School of Medicine, Tokyo, Japan)	Bench	Okonogi (2017) ¹⁴	Content: 1 Response processes: N Internal structure: N	10	Neurosurgeons	0

Foramen Ovale Puncture Model (Instituto de Neurologia de Curitiba, Curitiba, Brazil)	Bench	Almeida (2006) ¹⁵	Relations to other variables: N Consequences: N Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	5	Neurosurgeons	0
Custom 3D Head Models with Medtronic Platform (Division of Neurosurgery, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia)	Bench	Waran (2014) ¹⁶	Content: N Response processes: N Internal structure: N Relations to other variables: 1 Consequences: N	6	2 Neurosurgeons 4 Trainees	2
PeriopSim™: Instrument Trainer & Burr Hole Surgery (Conquer Mobile)	App	Clarke (2016) ¹⁷	Content: N Response processes: 2 Internal structure: N Relations to other variables: 0 Consequences: N	24	18 Residents 6 Expert Neurosurgeons	2
USim (Camelot Biomedical Systems, Genova, Italy; Neurosurgery Department, Fondazione IRCCS Istituto Neurologico Nazionale "C.Besta", Milan, Italy)	Bench/App	Perin (2018) ¹⁸	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	14	Residents	2
Gelatin 3D Brain Model (Department of Mechanical Engineering, Stanford University, Stanford, California, USA)	Bench	Ploch (2016) ¹⁹	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	10	Neurosurgeons Residents	1
Tumour Resection						
Phantom-based Training System (Department of Neurosurgery, University Hospital Leipzig, Leipzig, Germany)	Bench	Müns (2014) ⁴²	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	5	Residents	1

Agar-Gelatin Tumour resection Models (Department of Neurosurgery, Jichi Medical University, Shimotsuke Tochigi, Japan)	Bench	Müns (2014) ⁴³	Content: N Response processes: 2 Internal structure: N Relations to other variables: 1 Consequences: N	9	Residents	2
	Bench	Mashiko (2018) ³⁹	Content: 1 Response processes: N Internal structure: N Relations to other variables: 1 Consequences: N	9	4 Neurosurgeons 5 Residents	2
	VR	Gélinas-Phaneuf ⁴⁴	Content: N Response processes: N Internal structure: N Relations to other variables: 1 Consequences: N	72	44 Senior Residents 18 Junior Residents 10 Medical Students	1
	VR	Holloway (2015) ⁴⁵	Content: 1 Response processes: N Internal structure: N Relations to other variables: 2 Consequences: N	83	12 Residents 71 Medical Students	2
	VR	Alzhrani (2015) ⁴⁶	Content: N Response processes: 2 Internal structure: N Relations to other variables: 2 Consequences: N	33	17 Neurosurgeons 7 Senior Residents 9 Junior Residents	0
	VR	Alotaibi (2015) ⁴⁷	Content: 1 Response processes: 2 Internal structure: N Relations to other variables: 2 Consequences: N	18	6 Neurosurgeons 12 Residents	1
	VR	Winkler-Schwartz (2016) ⁴⁸	Content: N Response processes: 2 Internal structure: N Relations to other variables: 2	22	6 Residents 16 Medical Students	0

Navigation-Guided Endoscopic Intraventricular Injectable Tumor Model (Department of Neurosurgery, St. Louis University, St. Louis, Missouri, USA)	Cadaver	Ashour (2016) ⁴⁰	Consequences: N Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	n/a	Trainee Neurosurgeons	1
	Human Tissue	de Oliveira (2015) ⁴¹	Content: 1 Response processes: N Internal structure: N Relations to other variables: 1 Consequences: N	16	8 Neurosurgeons 8 Residents	1
Ventriculostomy						
Low-cost Ventriculostomy simulator (Department of Learning Health Sciences, University of Michigan Medical School, Ann Arbor, Michigan, USA)	Bench	Tai (2015) ³⁶	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	17	4 Faculty Neurosurgeons 12 residents 1 Fellow	1
	Bench	Rooney (2015) ³⁷	Content: 1 Response processes: 2 Internal structure: 2 Relations to other variables: 2 Consequences: 2	14	3 Expert Neurosurgeons 11 Novice Neurosurgeons	0
With V-PAT and OSATS assessment tools						
Medtronic StealthStation w/ Resin Head Model (Victor Horsley Department of Neurosurgery, The National Hospital for Neurology and Neurosurgery, Queen Square, University College London Hospitals NHS Foundation Trust, London, United Kingdom; Medtronic, Minneapolis, MN, USA)	Bench	Kirkman (2015) ³⁵	Content: 0 Response processes: 2 Internal structure: N Relations to other variables: N Consequences: N	31	10 Experienced 11 Moderate experience 10 Novice 2 Medical Students	2
3D cerebral lateral ventriculostomy model (Division of Neurological Surgery Barrow Neurological Institute, Phoenix, Arizona, USA)	Bench	Ryan (2015) ³⁴	Content: N Response processes: N Internal structure: N Relations to other variables: N	10	4 Residents 6 Students	1

			Consequences: N			
3D Hydrocephalus Model (University of Malaya, Kuala Lumpur, Malaysia)	Bench	Waran (2015) ³²	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	15	3 Neurosurgeons 12 Trainees	1
University of Florida Mixed Ventriculostomy Simulator (Department of Neurological Surgery, University of Florida, Gainesville, Florida, USA)	MR	Hooten (2014) ³³	Content: N Response processes: 2 Internal structure: N Relations to other variables: 2 Consequences: N	263	120 Senior Residents 71 Junior Residents 72 Interns	1
ImmersiveTouch: Ventriculostomy w/ Shifted Ventricle	VR	Lemole (2009) ²⁸	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	48	Residents	2
ImmersiveTouch: Ventriculostomy Catheter Placement (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	VR	Banerjee (2007) ²⁷	Content: N Response processes: N Internal structure: N Relations to other variables: 0 Consequences: N	78	Residents	0
ImmersiveTouch: Virtual Brain Library (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA; Department of Medical Education, College of Medicine, University of Chicago, Chicago, USA)	VR	Yudkowsky (2013) ²⁹	Content: N Response processes: N Internal structure: N Relations to other variables: 2 Consequences: N	16	Residents	3
Simulation Workshop w/ Rowena (Victor Horsley Department of Neurosurgery, National Hospital for Neurology, Queen Square; Neurodesign Ltd.)	Bench	Dasgupta (2018) ³⁰	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: 0	n/a	-	4

Hollow 3D Printed Head (Department of Neurosurgery, Vanderbilt University Medical Center, Nashville, Tennessee)	Bench	Bow (2018) ³¹	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: 0	11	3 Interns 8 Medical Students	2
ImmersiveTouch (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	VR	Perin (2018) ²⁶	Content: 2 Response processes: N Internal structure: N Relations to other variables: 2 Consequences: N	92	18 Neurosurgeons 25 Junior Residents 49 Senior Residents	1

Table 3: Validation studies on training models for Vascular neurosurgery. Abbreviations: LoE- Level of Effectiveness, VR- virtual reality,

Name of Model (Institution / Manufacturer)	Type of Model	Study	Validation	Participants		LoE
				n	Demographics	
Whole 3D Cerebral Aneurysm Model (Department of Neurosurgery, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China)	Bench	Wang (2017) ⁵⁴	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	n/a	Neurosurgeons	1
Regional 3D Cerebral Aneurysm Model (Department of Neurosurgery, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China)	Bench	Wang (2017) ⁵⁴	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	n/a	Neurosurgeons	1
3D Whole & Regional Middle Cerebral Aneurysm Model (Department of Neurosurgery, Beijing Tian Tan Hospital, Capital Medical University, Beijing, China)	Bench	Wang (2018) ⁵⁵	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	6	Residents	1
3D Cerebral aneurysm Simulator (School of Mechanical Engineering, Zhejiang University, Hangzhou 310027, China)	Bench	Liu (2017) ⁶²	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	10	4 Students 6 Neurosurgeons	1

Live Cadaver Model (Arkansas Neuroscience Institute, St. Vincent Health System, Little Rock, Arkansas, USA)	Cadaver	Aboud (2015) ⁶¹	Content: 2 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	91	27 Faculty 64 Participants	1
Human Placenta Aneurysm Model (Barrow Neurological Institute, St. Joseph's Hospital and Medical Center, Phoenix, Arizona)	Human Tissue	Belykh (2017) ⁵⁸	Content: 2 Response processes: 2 Internal structure: 2 Relations to other variables: 2 Consequences: N	30	4 Students 12 Residents 4 Fellows 10 Neurosurgeons	1
(Microsurgical Laboratory, Department of Surgery, Federal University of Minas Gerais, Belo Horizonte, Minas Gerais, Brazil)	Human Tissue	de Oliveira (2018) ⁶⁰	Content: 2 Response processes: 2 Internal structure: N Relations to other variables: N Consequences: N	21	12 Neurosurgeons 9 Residents	5
Human Cadaver (Microsurgical Laboratory, Department of Surgery, Federal University of Minas Gerais, Belo Horizonte, Minas Gerais, Brazil)	Cadaver	de Oliveira (2018) ⁶⁰	Content: 1 Response processes: 2 Internal structure: N Relations to other variables: N Consequences: N	21	12 Neurosurgeons 9 Residents	5
Cerebral Aneurysm Clipping Training Program (Department of Neurosurgery, Jichi Medical University, Shimotsuke Tochigi, Japan)	Bench	Mashiko (2017) ⁶³	Content: N Response processes: N Internal structure: N Relations to other variables: 1 Consequences: N	7	4 Residents 2 Junior Neurosurgeons 1 Senior Neurosurgeon	2

Virtual Cerebral Aneurysm Clipping with Real Time Haptic Force Feedback Model (Department of Neurosurgery, Kepler University Hospital, Linz, Austria)	Bench	Gmeiner (2018) ⁵⁷	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	18	4 Residents 14 Surgeons	1
ImmersiveTouch: Aneurysm Clipping (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	VR	Alaraj (2015) ⁵⁶	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	17	Residents	1
3D Hollow Elastic Models (Medical Simulation Center, Jichi Medical University, Tochigi, Japan)	Bench	Mashiko (2015) ⁵³	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	18	12 Neurosurgeons 6 Junior Neurosurgeons	1
Human Placenta Model: Intracranial Bypass (Placentarium, Federal University of Minas Gerais, Belo Horizonte, Minas Gerais, Brazil; Lyerly Neurosurgical Simulation Laboratory, Baptist Medical Center, Florida, USA)	Human Tissue	Oliveira (2018) ⁵⁹	Content: 1 Response processes: N Internal structure: 2 Relations to other variables: 2 Consequences: N	60	5 Expert Neurosurgeons 5 Novice Neurosurgeons 50 Health Professionals	0
Venous Sinus Injury and Air Embolus Surgical Simulation (Department of Neurological Surgery, Oregon Health & Science University, Portland, Oregon)	Bench	Cleary (2018) ⁶⁴	Content: N Response processes: 1 Internal structure: N Relations to other variables: 1 Consequences: N	12	1 Attending Physician 11 Residents	1
ImmersiveTouch: Hemostasis Simulation (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	VR	Gasco (2013) ⁶⁵	Content: N Response processes: N Internal structure: N Relations to other	54	14 Residents 20 Senior Medical Students 20 Junior Medical Students	1

variables: N
Consequences: N

Table 4: Validation studies on training models for Non-Technical Skills & Spine Surgery. Abbreviations: LoE- Level of Effectiveness, VR- virtual reality

Model (Institution / Manufacturer)	Type of Model	Study	Validation	n	Participants Demographics	LoE
Non-Technical Skills						
Crisis Management Simulation: Dual Neurosurgery and Anaesthesia Training Experience (Anesthesiology and Perioperative Medicine, Oregon Health and Science University, Portland, Oregon, United States; Department of Neurosurgery, West Virginia University School of Medicine, Morgantown, West Virginia, United States)	Bench	Ciporen (2018) ¹⁰⁵	Content: N Response processes: N Internal structure: N Relations to other variables: 0 Consequences: N	14	7 Neurosurgery Residents 7 Anaesthesia Residents	1
Adult Cadaver Head Perfusion Model (Oregon Health & Science University Body Donation Program, Portland, Oregon)	Cadaver	Ciporen (2017) ⁴⁹	Content: N Response processes: N Internal structure: N Relations to other variables: 1 Consequences: N	6	Senior Residents	2
Task and Crisis Vertebroplasty Simulation (Technische Universitat Munchen, Munich, Germany)	VR	Wucherer (2014) ¹⁰⁶	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	5	1 Expert Neurosurgeon 2 Senior Neurosurgeons 2 Junior Neurosurgeons	1
Spine Surgery						
Mixed Reality Spine Simulator (Paediatric Neurosurgery Center, Beneficencia Portuguesa Hospital, Sao Paulo, Brazil)	Bench/VR	Coelho (2018) ⁷²	Content: 2 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	16	Spine Surgeons	0
Durotomy Repair Module (Congress of Neurological Surgeons)	Bench	Ghobrial (2013) ⁶⁷	Content: N Response processes: N Internal structure: N Relations to other variables: N		1 Retired Neurosurgeon 4 Residents 1 Physician Assistant	2

Congress of Neurological Surgeons Cervical Foraminotomy and Durotomy Repair Modules (Congress of Neurological Surgeons)	Bench	Ghobrial (2015) ⁶⁸	Consequences: N Content: N Response processes: 2 Internal structure: N Relations to other variables: 0	20	Residents	2
Microanastomosis, Anterior Cervical Discectomy and Fusion (ACDF), Posterior Cervical Fusion (PCF), and Durotomy Repair Modules (Congress of Neurological Surgeons)	Bench	Zammar (2015) ⁶⁹	Consequences: N Content: N Response processes: 2 Internal structure: N Relations to other variables: N	20	Residents	2
Low Cost Dural Closure Simulation Model (Department of Neurosurgery, Institute of Neurological Sciences, Glasgow, UK)	Bench	Ferguson (2018) ⁷³	Consequences: N Content: N Response processes: 2 Internal structure: N Relations to other variables: N	15	2 Spine Surgeons Fellows 13 Specialty Trainees	2
3D Software-based Pedicle Screw Simulator (Faculty of Medicine, Orthopaedic Biomechanics Laboratory, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada)	VR	Podolsky (2010) ⁷⁴	Consequences: N Content: N Response processes: N Internal structure: N Relations to other variables: N	37	Trainees	1
Simulated Lumbar Minimally Invasive Surgery Education Model (Department of Neurosurgery, Thomas Jefferson University Hospital, Philadelphia, Pennsylvania; Mayo Clinic, Rochester, Minnesota, USA)	Bench	Chitale (2013) ⁷⁶	Consequences: N Content: N Response processes: N Internal structure: N Relations to other variables: N	8	Trainees	2
Anterior Discectomy and Fusion Simulator (Congress of Neurological Surgeons)	Bench	Ray (2013) ⁷⁰	Consequences: N Content: N Response processes: N Internal structure: N Relations to other variables: N	6	1 Neurosurgeon 3 Residents 2 Medical Students	2
Cervical Spine Simulator (Congress of Neurological Surgeons)	Bench	Harrop (2013) ⁷¹	Consequences: N Content: N Response processes: N Internal structure: N	11	1 Neurosurgeon 10 Trainees	2

Laboratory-based Spinal Fixation Training Program (Case Western Reserve University School of Medicine; Department of Neurological Surgery; Center for Spine Health, Neurological Institute, Cleveland, Ohio)	Bench/Cadaver	Sundar (2016) ⁷⁷	Relations to other variables: N Consequences: N Content: N Response processes: 2 Internal structure: N	10	8 Residents 2 Medical Students	2
Minimally Invasive Spine Surgery Simulator (Department of Neurosurgery, University of Mississippi Medical Center, Jackson, Mississippi, USA)	Bench/Animal	Walker (2009) ⁷⁵	Relations to other variables: N Consequences: N Content: N Response processes: N Internal structure: N	8	Residents	1
ImmersiveTouch: Percutaneous Spinal Needle Placement (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	VR	Luciano (2013) ⁷⁹	Relations to other variables: N Consequences: N Content: N Response processes: N Internal structure: N	63	Fellows Residents	2
ImmersiveTouch: Thoracic Pedicle Screw Placement (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	VR	Luciano (2011) ⁷⁸	Relations to other variables: N Consequences: N Content: N Response processes: N Internal structure: N	51	Fellows Residents	2
Saw Bones Scoliosis Model (Stony Brook University Hospital, Stony Brook, USA)	Bench	Tanner (2017) ⁸⁰	Relations to other variables: 1 Consequences: N Content: N Response processes: 2 Internal structure: N	20	10 Senior Residents 10 Junior Residents	0
Desktop-based Computer Assisted Orthopaedic Training System (Department of Computer Sciences, University of Hull, United Kingdom; Department of Orthopaedics, Leeds Teaching Hospital NHS Trust, Leeds, United Kingdom)	VR	Rambani (2014) ⁸²	Relations to other variables: N Consequences: N Content: N Response processes: 2 Internal structure: 0	12	Junior Trainees	2
Life-Sized 3D Spine Model (Department of Orthopaedic Surgery, Kangnam Sacred Heart Hospital, Hallym University College of Medicine, Seoul, South Korea)	Bench	Park (2018) ⁸³	Relations to other variables: N Consequences: N Content: N Response processes: N	2	Novice Surgeons	2

Korea)			Internal structure: N Relations to other variables: N Consequences: N			
Bioskills Training Module: Lumbar Pedicle Screw (Northwestern Memorial Hospital, Chicago, Illinois, USA)	Bench	Boody (2018) ⁸⁴	Content: N Response processes: 2 Internal structure: N Relations to other variables: N Consequences: N	19	Orthopaedic Residents Medical Students	2
Bioskills Training Module: Lumbar Laminectomy (Northwestern Memorial Hospital, Chicago, Illinois, USA)	Bench	Boody (2017) ⁸⁵	Content: N Response processes: 2 Internal structure: N Relations to other variables: N Consequences: N	20	Orthopaedic Residents Medical Students	2
Artificial Wetlab Training System for Lumbar Discectomy (Krankenhaus Winsen, Orthopadie, Winsen, Germany)	Bench	Adermann (2014) ⁸⁶	Content: 1 Response processes: 1 Internal structure: N Relations to other variables: N Consequences: N	12	Experienced Spine Surgeons	1
Cadaveric Spinal Surgery Simulation; Thiel, Crosado and formaldehyde (Health Education Yorkshire and the Humber, University of Leeds, Leeds, United Kingdom)	Cadaver	Tomlinson (2016) ⁸⁷	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	10	Fellows Consultants	1
Virtual Surgical Training System (Department of Orthopaedic Surgery, Changzheng Hospital, Second Military Medical University, Shanghai, China)	VR	Hou (2018) ⁸⁸	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	10	Novice Residents	2
ImmersiveTouch: Pedicle Screw Placement (ImmersiveTouch Inc., University of Illinois at Chicago Medical Center, Chicago, IL, USA)	VR	Gasco (2014) ⁸¹	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	26	Medical Students	2
3D Patient-Specific Rendering: Pedicle Screw Insertion (Department of	VR	Xiang (2015) ⁸⁹	Content: N	4	2 Experienced Spine	2

Orthopedics, General Hospital of Shenyang Military Area Command of
Chinese PLA, Shenyang, Liaoning, China)

Response processes: N
Internal structure: N
Relations to other variables: 1
Consequences: N

Surgeons
2 Junior Surgeons

Table 5: Validation studies on training models for Endoscopic Endonasal Transphenoidal Surgery . Abbreviations: LoE- Level of Effectiveness, VR- virtual reality.

Model (Institution / Manufacturer)	Type of Model	Study	Validation	n	Participants	LoE
					Demographics	
Endoscopic Endonasal Drilling Model (Department of Learning Health Sciences, University of Michigan Medical School, Ann Arbor, Michigan, USA)	Bench	Tai (2016) ⁹³	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	8	Neurosurgeons Residents	1
Chicken Eggs and Skull Model (Department of Neurosurgery, Kinki University Faculty of Medicine, Osaka, Japan)	Bench	Okuda (2014) ⁹⁴	Content: N Response processes: N Internal structure: N Relations to other variables: 2 Consequences: N	9	4 Experts 5 Residents	2
Laser Sintered Model (Ohio State University, Columbus, Ohio, USA)	Bench	Maza (2018) ⁵⁰	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	46	20 Otolaryngologists 26 Neurosurgeons	2
Perfusion-Based Human Cadaveric Simulation for ICA injury (Keck School of Medicine, Los Angeles, California, United States)	Cadaver	Shen (2018) ⁵²	Content: N Response processes: N Internal structure: N Relations to other variables: 0 Consequences: N	35	19 Neurosurgeons 16 Otolaryngologists	2
	Cadaver	Pham (2014) ⁵¹	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	n/a	Residents	1
Practical 3D Printed Simulator (Department of Neurosurgery, Jinling Hospital, School of Medicine, Nanjing University, China)	Bench	Wen (2016) ⁹⁵	Content: N Response processes: N Internal structure: N Relations to other variables: 2 Consequences: N	23	5 Expert Neurosurgeons 7 Assistant Neurosurgeons 6 Novice	2

					Neurosurgeons 5 Trainees	
3D Endoscopic Skull base models w/ Pre-existing pathology (Division of Neurosurgery, Department of Surgery, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia)	Bench	Narayanan (2015) ⁹⁶	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	15	ENT Surgeons	1
3D Cranial base model (Department of Neurological Surgery, The Ohio State University Wexner Medical Center, Columbus, Ohio, USA)	Bench	Muto (2017) ⁹⁷	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	5	Trainees	1
NeuroTouch Simulator: Endoscopic Endonasal Module (CAE Healthcare, Montreal, Quebec, Canada)	VR	Thawani (2016) ⁹⁸	Content: N Response processes: 2 Internal structure: N Relations to other variables: N Consequences: N	6	Residents	5
3D Printed Skull base for Transnasal endoscopic Skull base Surgery (Beijing Neurosurgical Institute, Beijing Tiantan Hospital, Capital Medical University, Beijing, China)	Bench	Zheng (2018) ⁹¹	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	13	3 Neurosurgeons 10 Residents	1
Neuro-Endo-Trainer (NET) SkullBase-Task-GraspPickPlace (Centre for Biomedical Engineering, Indian Institute of Technology Delhi, New Delhi, India)	Bench	Singh (2016) ⁹²	Content: 0 Response processes: 1 Internal structure: N Relations to other variables: N Consequences: N	61	4 Expert Neurosurgeons 19 Novice Neurosurgeons 38 Residents	1
Perfusion-based Cadaveric Simulation Model: Endoscopic Endonasal CSF Leak Repair (Department of Neurological Surgery, Keck School of Medicine, University of Southern California, Los Angeles, California)	Cadaver	Christian (2018) ⁹⁰	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	9	Residents	1
Skull Base Injectable Tumor Model (Macquarie Neurosurgery, Australian	Bench	Gagliardi	Content: 1	6	3 Residents	1

School of Advanced Medicine, Macquarie University, Sydney, Australia)		(2018) ³⁸	Response processes: N Internal structure: N Relations to other variables: N Consequences: N		3 Faculty Surgeons	
Adult Cadaver Head Perfusion Model (Oregon Health & Science University Body Donation Program, Portland, Oregon)	Cadaver	Ciporen (2017) ⁴⁹	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	6	Senior Residents	2

Table 6: Validation studies on training models for Paediatric neurosurgery . Abbreviations: LoE- Level of Evidence, LoR- Level of Recommendation, VR- virtual reality.

Name of Model (Institution / Manufacturer)	Type of Model	Study (Year)	Validation	n	Participants	LoE
					Demographics	
'babyMARTYN' (Victor Horsley Department of Neurosurgery, National Hospital for Neurology and Neurosurgery, Queen Square)	Bench	Craven (2018) ¹⁰⁰	Content: 1 Response processes: 2 Internal structure: N Relations to other variables: N Consequences: N	11	3 Neurosurgeons 8 Trainees	2
		Breimer (2015) ²⁰	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	25	16 Residents 5 Fellows 4 Neurosurgeons	1
SickKids brain simulator (Centre for Image Guided Innovation and Therapeutic Intervention, Toronto, Ontario, Canada)	Bench	Breimer (2017) ²¹	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	26	23 Residents 3 Fellows	1
NeuroTouch Simulator: ETV Module (CAE Healthcare, Montreal, Quebec, Canada)	VR	Breimer (2017) ²¹	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	26	23 Residents 3 Fellows	1
Full Scale Hydrocephalus Head Model (Harvard Medical School, Boston, Massachusetts and Division of Pediatric Neurosurgery, Johns Hopkins Hospital, Baltimore, Maryland, USA)	Bench	Weinstock (2017) ²²	Content: N Response processes: 2 Internal structure: 2 Relations to other	17	13 Residents 4 Fellows	1

CURE Hydrocephalus and Spina Bifida Fellowship (CURE Childrens Hospital of Uganda, Mbale, Uganda)	Bench	Dewan (2018) ¹⁰¹	variables: 2 Consequences: N Content: N Response processes: N Internal structure: N Relations to other variables: N	33	Fellows	5
Synthetic Endoscope-assisted Craniosynostosis Models (Center for Image-Guided Innovation and Therapeutic Innovation, The Hospital for Sick Children, Toronto)	Bench	Eastwood (2018) ¹⁰²	Consequences: 2 Content: 1 Response processes: N Internal structure: N Relations to other variables: N	25	5 Expert 9 Fellows 11 Residents	1
Prototype for Pediatric Spinal Detethering Surgeries (University of Illinois ,College of Medicine at Peoria, Peoria, IL, USA)	Bench	Bailey (2013) ¹⁰³	Consequences: N Content: 1 Response processes: N Internal structure: N Relations to other variables: N	12	1 Neurosurgeon 1 Fellow 6 Residents 4 Students	1
3D Synthetic Model for Pediatric Lumbar Spine Pathologies (Department of Neurosurgery, University of Illinois College of Medicine, USA)	Bench	Mattei (2013) ¹⁰⁴	Consequences: N Content: N Response processes: N Internal structure: N Relations to other variables: 1 Consequences: N	7	3 Experts 1 Medical Student 3 Residents	0

Table 7: Validation studies on training models for Training Courses. Abbreviations: LoE- Level of Effectiveness, VR- virtual reality.

Model (Institution / Manufacturer)	Type of Model	Study	Validation	n	Participants Demographics	LoE
Training Courses						
National Fundamentals Curriculum (Society of Neurological Surgeons)	Bench	Selden (2012) ⁶⁶	Content: 2 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	261	75 Neurosurgeons 186 Residents	2
		Selden (2017) ⁹⁹	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	164	Residents	3
Neurosurgery Boot Camp: Bolivia (Bolivian Society for Neurosurgery; Foundation for International Education in Neurological Surgery; Solidarity Bridge; University of Massachusetts)	Bench	Ament (2017) ²³	Content: N Response processes: N Internal structure: N Relations to other variables: N Consequences: N	29	5 Neurosurgeons 24 Residents	1
Fundamentals of Neurosurgery (Department of Neurology and Neurosurgery, Montreal Neurological Institute and Hospital, Brain Tumour Research Centre, McGill University, Montréal, Quebec, Canada)	VR	Choudhury (2013) ²⁴	Content: 1 Response processes: N Internal structure: N Relations to other variables: N Consequences: N	n/a	Subject Matter Experts	0
Trauma Module w/ ImmersiveTouch: Ventriculostomy (Congress of Neurological Surgeons)	VR	Schirmer (2013) ²⁵	Content: N Response processes: 1 Internal structure: N Relations to other variables: 2 Consequences: N	12	Residents	2
Trauma Module w/ Physical Craniotomy Model (Congress of Neurological Surgeons)	Bench	Lobel (2013) ¹⁰⁹	Content: N Response processes: N	14	Residents Medical Students	2

SBNS-accredited Neurosurgical Skills Workshop (Society of British Neurological Surgeons; The Neurology and Neurosurgery Interest Group)	Bench	Kamel (2017) ¹⁰⁷	Internal structure: N Relations to other variables: 1 Consequences: N Content: N Response processes: N	10		2
Presigmoid Approach Simulation Module (Congress of Neurological Surgeons)	Bench	Jabbour (2013) ¹⁰⁸	Internal structure: N Relations to other variables: N Consequences: N Content: N Response processes: N	9	Residents	2
Simulation Curriculum (Division of Neurosurgery; and Division of Epidemiology & Biostatistics, University of Texas Medical Branch, Galveston, Texas, USA)	Bench/V R/ Cadaver	Gasco (2013) ¹¹⁰	Internal structure: N Relations to other variables: N Consequences: N Content: N Response processes: N	130	Residents	1

PRISMA 2009 Flow Diagram

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit www.prisma-statement.org.

Supplementary Tables 1-2

[Click here to download Supplementary Material \(Video/Media Files\): Supplementary Tables 1 & 2 .docx](#)