King's Research Portal

DOI:

[10.1177/1359104518778327](https://doi.org/10.1177/1359104518778327)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Hannah, B., & Woolgar, M. (2018). Secondary trauma and compassion fatigue in foster carers. *Clinical Child Psychology and Psychiatry*, 23(4), 629-643. <https://doi.org/10.1177/1359104518778327>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Author Accepted Manuscript: Clinical Child Psychology & Psychiatry

Title: Secondary Trauma and Compassion Fatigue in Foster Carers

Dr Beatrice Hannah¹ and Dr Matt Woolgar¹²

¹King's College London, Institute of Psychiatry, Psychology and Neuroscience; ²South London and Maudsley NHS Foundation Trust, UK

Corresponding author: Dr Beatrice Hannah, King's College London, Institute of Psychiatry Psychology & Neuroscience, Department of Psychology, Denmark Hill, London, SE5 8AF
Email: Beatrice.hannah@kcl.ac.uk

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

This research was undertaken as part of a DClinPsy degree to Dr Beatrice Hannah

Acknowledgements

The authors would like to thank all of the local authorities who assisted with recruitment and to all of the foster carers who took the time to participate in the study. They also thank Dr Wendy Geraghty for her contributions to the project.

Abstract

Background: With the number of children being placed in the care of Local Authorities increasing, the need to recruit and retain foster carers is essential. Compassion fatigue is recognised as a negative effect for professionals working with adults and children who have experienced trauma. This study aims to confirm the proof of concept within foster carers and to explore the potential risks associated with intent to continue fostering, overall job satisfaction and psychological factors [avoidant coping] that could be targets for interventions.

Methods: 131 foster carers completed an online survey including self-report measures of compassion fatigue and associated risk factors.

Results: Results confirm the existence of compassion fatigue amongst foster carers with rates in line with previous studies on other professionals working with children. High compassion fatigue was associated with lower intent to continue fostering and lower job satisfaction. Avoidant cognitive styles of psychological inflexibility and thought suppression were associated with compassion fatigue.

Conclusions: The confirmation of compassion fatigue amongst foster carers and the potential risks to job retention are important findings for social care. The associations with avoidant cognitive styles have clinical implications for potential interventions.

Recommendations for further research and the limitations of this study are also discussed.

Keywords: Foster carers, Looked After Children, compassion fatigue, secondary trauma, burnout, job retention

According to the Department for Education (2016) in the UK, 69,540 children were in the care of local authorities on 31st March 2015, with 75% (52,050) of these children living with foster carers. With numbers of children being taken into care increasing, and social care being faced with a growing shortage of foster carers (Harber, Oakley, & Exchange, 2012; Rodger, Cummings, & Leschied, 2006), the need to recruit and retain foster carers is essential. Research shows that foster carers report high levels of stress related to their role (Murray, Tarren-Sweeney, & France, 2011; Vanschoonlandt & Vanderfaellie, 2013; Whenan, Oxlad, & Lushington, 2009). The typical stressors faced by foster carers include placement disruptions, allegations, difficulties with social services, and contact with birth families (Wilson, Sinclair, & Gibbs, 2000). High levels of strain are associated with a foster carers overall job satisfaction and intention to quit (Farmer, Lipscombe, & Moyers, 2005). In addition, a recent review highlights the impact of carer-wellbeing on overall placement stability (Rock, Michelson, Thomson, & Day, 2015). Limited training, feeling unsupported and low self-efficacy also appear to contribute to a foster carer's likelihood of leaving the profession (Denby, Rindfleisch, & Bean, 1999; Rhodes, Orme, & Buehler, 2001; Sinclair, Gibbs, & Wilson, 2004). Whilst placement stability is a priority for social care, research exploring the overall wellbeing of foster carers remains extremely limited.

The reasons children are placed in the care of a local authority are often related to potentially traumatising experiences of abuse and neglect (Department of Children, Schools and Families, 2009). Given the adverse circumstances usually associated with a child being taken into care, looked after children often present with high rates of behavioural difficulties and emotional distress (Cook et al., 2005; Greeson, Briggs, & Kisiel, 2011; Oswald, Fegert, & Goldbeck, 2010). It is well established that looked after children in the care of a local authority experience higher rates of psychiatric disorders (Ford, Vostanis, Meltzer, & Goodman, 2007;

McAuley & Davis, 2009; Meltzer, Gatward, Corbin, Goodman, & Ford, 2003; Richardson & Joughin, 2000) and increased rates of behavioural difficulties than children living with their biological parents (Burns et al., 2004; Clausen, Landsverk, Ganger, Chadwick, & Litrownik, 1998; Halfon, Mendonca, & Berkowitz, 1995; Sawyer, Carbone, Searle, & Robinson, 2007). With this in mind, the impact of caring for children presenting with a history of complex trauma, as well as behavioural and emotional difficulties understandably puts a strain on the foster carers with whom they are placed.

Psychological impact of working with traumatised individuals

A number of terms have been used to describe the psychological impact of working with traumatised individuals including burnout (Maslach, 1982; Maslach, Schaufeli, & Leiter, 2001), secondary trauma (Figley, 1995) and compassion fatigue (Bride, Radey, & Figley, 2007; Stamm, 2005).

- *Compassion Fatigue* is considered to be an over-arching term used to describe the emotional impact of working with traumatised individuals including secondary trauma and burnout (Figley, 2002, 2013; Stamm, 2010).
- *Secondary Trauma* is characterised by tension and preoccupation with the suffering of others as a result of working in trauma related contexts. Secondary trauma symptoms are similar to symptoms of Post-Traumatic Stress Disorder (PTSD) and include heightened states of arousal, intrusive thoughts or images, and avoidance of thoughts or reminders of the person's traumatic experiences (Figley, 1995).
- *Burnout* is the response that occurs as a result of prolonged exposure to demanding interpersonal situations and is characterised by emotional and physical exhaustion, depersonalisation, and reduced personal accomplishment

(Maslach et al., 2001). The symptoms of burnout tend to develop gradually over time but if ignored can progressively worsen (Figley, 1995).

- *Compassion Satisfaction* is described as the positive emotional response that results from helping others, including positive feelings about colleagues, contributions to the work setting, and doing something good for society (Phelps, Lloyd, Creamer, & Forbes, 2009; Stamm, 2002). Compassion satisfaction has been found to mitigate the effects of burnout and secondary trauma (Ray, Wong, White, & Heaslip, 2013; Wee & Myers, 2003).

The presence of compassion fatigue has been widely reported in various professionals working with traumatised individuals including nurses (Beck, 2011; Burtson & Stichler, 2010; Hooper, Craig, Janvrin, Wetsel, & Reimels, 2010); trauma counsellors (Carmel & Friedlander, 2009; Ghahramanlou & Brodbeck, 2000); domestic violence staff (Baird & Jenkins, 2003); and social workers of adult trauma victims (Boscarino, Figley, & Adams, 2004). There have been fewer studies investigating the impact of working with traumatised children, however there are indications of similar experiences of compassion fatigue in child protection workers (Conrad & Kellar-Guenther, 2006; Cornille & Meyers, 1999; Sprang, Craig, & Clark, 2011); forensic interviewers of abused children (Perron & Hiltz, 2006); and paediatric healthcare providers (Meadors, Lamson, Swanson, White, & Sire, 2010). The emotional impact of working with children may occur directly as a result of the challenges of managing behavioural difficulties or indirectly through hearing about a child's traumatic history (Meyers & Cornille, 2002).

Avoidance and trauma

Avoidance of trauma related thoughts and memories is central to the definition of compassion fatigue, and specifically secondary trauma (Figley, 1995). More broadly, avoidant coping has been found to be common in a range of psychological problems including

depression and anxiety disorders (Holahan et al., 2005; Thompson and Waltz, 2010; Tull et al., 2004). As such, this exploratory study aims to investigate the relationship of avoidant coping with compassion fatigue. Two cognitive processes that have been described as highly related to avoidance are psychological inflexibility and thought suppression. Psychological flexibility is described as “the ability to fully contact the present moment and the thoughts and feelings it contains without needless defence” (Hayes, Luoma, Bond, Masuda, & Lillis, 2006, p. 7). Thought suppression is the conscious attempt to stop thinking about a particular thought (Wegner, Schneider, Carter, & White, 1987). Research indicates a paradoxical effect of thought suppression and avoidance of unwanted internal experiences with continued avoidance over time actually causing maintenance or increased frequency of the unwanted experiences (Hayes & Wilson, 1996; Kumpula, Orcutt, Bardeen, & Varkovitzky, 2011; Marx & Sloan, 2005; Wegner et al., 1987).

The present study

This is a proof of concept study that aims to examine whether the model of compassion fatigue, described in other professionals, applies to this population. The study draws similarities from previous unpublished work by McLain (2008) and a recent study by Ottaway and Selwyn (2016), who investigated compassion fatigue in foster carers. This study expands on previous work by also considering the potential relationship with job retention and satisfaction as well as underlying avoidant mechanisms. The study examined the association between secondary trauma, burnout and compassion satisfaction on intent to continue fostering and overall job satisfaction, two outcomes that will have significant practical implications. Finally, the study explored the association of compassion fatigue with two cognitive processes, psychological inflexibility and thought suppression, which are known to be related to symptoms of avoidance in trauma.

Method

Participants

Participants were 131 local authority foster carers, foster carers from the independent sector and kinship carers approved as foster carers working in the United Kingdom. Inclusion criteria for participation in this study were that the foster carers were (a) registered foster carer for at least 6 months and (b) currently looking after at least one child placed by the Local Authority, or had done so within the last 6 months. An a priori power analysis was carried out to determine the minimum sample size required to detect a statistically significant relationship between compassion fatigue and job satisfaction and intent to continue fostering. Based on results of previous studies investigating compassion fatigue, with an effect size of $r=.33$ (Perron & Hiltz, 2006), a sample size of at least 102 was required with an alpha level set at 0.01 and power of 0.80. There is currently no comprehensive data on the demographic profile of foster carers in the UK, however limited data suggests that the current sample is of a similar age, gender and marital status, (McDermid, Holmes, Kirton, & Signoretta, 2012), although more ethnically diverse, which may be a reflection of the urban recruitment in the current sample.

Measures

Professional Quality of Life Questionnaire (ProQoL; Stamm, 2005): The ProQoL is a 30 item, 6-point Likert scale based on experiences in the last 30 days measuring three components: compassion satisfaction, burnout, and compassion fatigue, with each score measuring distinct theoretical concepts. This measure has good psychometric properties from a range of populations including social work and (Adams, Boscarino, & Figley, 2006; Potter et al.,

2010). The ProQol was scored using the method outlined in the ProQol Manual (Stamm, 2010) which includes cut-offs for high (75%) and low (25%) scores based on a databank of 1,289 cases created from multiple studies across various helping professions. For this study, Cronbach's Alpha indicates good internal reliability for all subtests: secondary trauma (.75), burnout (.80) and compassion satisfaction (.87).

Secondary Trauma Stress Scale (STSS; Bride, Robinson, Yegidis, & Figley, 2004): The STSS is a 17-item, 5-point Likert scale designed to assess the frequency of intrusion, avoidance and arousal symptoms over the past 7 days. A score of 38 or higher indicates secondary trauma. The STSS has demonstrated good validity and reliability in studies of social workers (Bride et al., 2004; Ting, Jacobson, Sanders, Bride, & Harrington, 2005). In relation to this study, the total STSS score showed good internal reliability (.94) as did the subtests of intrusions (.80), avoidance (.87) and arousal (.84).

Intent to continue and job satisfaction were measured using two items from a previous study investigating foster carer satisfaction and intent to continue (Denby et al., 1999). The first item, "all in all, how satisfied are you as a foster parent?" consisted of a four-point scale. To create binary categories, "Not at all satisfied" and "somewhat dissatisfied" were combined and "somewhat satisfied" and "very satisfied" to create binary categories of low and high satisfaction. The second item, "how likely is it that a year from now you will still be working as a foster carer?" consisted of a three-level scale of "Not at All Likely", "Somewhat Likely" or "Very Likely". The latter two scores were combined to create a high intent to continue score.

Acceptance and Action Questionnaire (AAQ-II; Bond et al., 2011): The AAQ-II consists of 7 items on a 7-point Likert scale. The scale reflects the single domain of psychological inflexibility with higher scores indicating greater psychological inflexibility. The psychometric

properties of this measure are well documented in non-clinical samples (Bond et al., 2011). In this study, internal reliability for this scale was high (.93).

The White Bear Suppression Inventory (WBSI; Wegner & Zanakos, 1994): The WBSI is a 15 item measure on a 5-point Likert scale measuring a person's tendency to suppress thoughts. This measure has been found to have good reliability and validity (Luciano et al., 2006) and the internal consistency in this study was high (.96).

Procedure

Foster carers were contacted via local authorities in and around London as well as the voluntary sector organisation TACT, the UK's largest Fostering and Adoption charity. Following consent from gatekeepers, foster carers were sent an email invitation to participate in the study containing the information sheet and a link to the online survey. Consent was implied by participants completing the survey. Once the survey had been completed, participants were given an opportunity to contact the researcher via email to claim a £5 voucher. Participants were able to remain anonymous and not claim the voucher should they wish.

Both the information sheet and the exit page of the survey contained information for accessing further support for participants who experienced any anxiety or distress as a result of completing the survey. The research was approved by King's College London Research Ethics Committee and consent forms and information sheets were available via the online survey. Data collection was conducted between October 2015 to January 2016.

Data Analysis Plan

The data from the survey was exported from the Bristol Online Survey into IBM SPSS Statistics 22 where it was then coded and analysed. Due to data not being normally distributed, the bootstrap function (5000 samples) was used to generate more accurate standard errors, confidence intervals and p-values (Good, 2005). The bootstrap function on independent sample

t-tests do not use the t-distribution and degrees of freedom to derive the p-value and confidence intervals and so these are not provided. Effect sizes for t-tests were defined as weak (0.2) moderate (0.5) or large (0.8) as recommended by (Cohen, 1992). For bivariate analyses, correlations were defined as weak ($r=.1$), moderate ($r=.3$) or strong ($r\geq.5$) based on recommendations by (Cohen, 1992).

Results

Descriptive statistics

The majority of participants were females (77.1%); over 40 years old (91.6%); married (66.4%); and had their own children (90.1%). 43.5% of the sample were White British and 49.6% had obtained higher education or professional qualifications. The majority of participants reported they were very likely to continue fostering (77.9%) and either somewhat (42.7%) or very (43.5%) satisfied in their jobs (see Table 1 for demographic information).

TABLE 1 HERE

Overall, 25.2% ($n=33$) of participants reported high levels of secondary trauma and 30.5% ($n=40$) reported high levels of burnout. 18.3% ($n=24$) of participants rated themselves as high for compassion satisfaction (see Table 2).

TABLE 2 HERE

The mean scores and standard deviations for the STSS can be found in Table 3. The STSS provides a specific clinical cut off of 38 and this study found that 19.8% ($n=26$) of participants were above the clinical cut off for secondary trauma.

TABLE 3 HERE

In line with the hypothesis, there was evidence supporting the presence of secondary trauma and burnout in foster carers. Further analysis indicated significant correlations between the two measures of secondary trauma (see Table 4). Furthermore, and in line with previous findings, compassion satisfaction was negatively associated with secondary trauma and burnout scales.

TABLE 4 HERE

Intent to continue fostering was significantly associated with secondary trauma (CI: 4.07 to 15.90, $p=.006$, $d=.70$), burnout (CI: 2.85 to 7.70, $p<.001$, $d=.88$), and compassion satisfaction (CI: -6.39 to -0.68, $p=.036$, $d=-.54$). Participants with low intent to continue fostering had significantly higher secondary trauma and burnout scores and significant lower compassion fatigue scores than those with high intent to continue fostering (see Figure 1).

FIGURE 1 HERE

Low job satisfaction was significantly associated with secondary trauma (CI: -17.92 to -2.57, $p=.035$, $d=0.73$), burnout (CI: -9.34 to -1.88, $p=.009$, $d=0.82$), and compassion satisfaction (CI: 1.26 to 9.43, $p=.023$, $d=0.77$). Participants with low job satisfaction had significantly higher secondary trauma and burnout scores, and significantly lower compassion satisfaction scores (see Figure 2).

FIGURE 2 HERE

Results indicate that psychological inflexibility was significantly and positively correlated to secondary trauma ($r=.349$, $p<.001$) and burnout ($r=.520$, $p<.001$). Thought suppression, as measured by the WBSI, was also positively associated with secondary trauma ($r=.597$, $p<.001$) as well as burnout ($r=.437$, $p<.001$). As predicted, higher levels of psychological inflexibility and increased thought suppression are related to higher secondary trauma and burnout scores (see Table 5).

TABLE 5 HERE**Discussion**

This study investigated the experiences of compassion fatigue in foster carers working in the UK. A sample of 131 foster carers completed an online survey to measure compassion fatigue and several related variables. The primary aim of the study was to establish whether the concept of compassion fatigue, the over-arching term to describe the effects of secondary trauma and burnout, occurs within this population. This study also examined the impact of compassion fatigue on risk factors related to intent to continue fostering and job satisfaction. This study also examined the associations between compassion fatigue and avoidant cognitive styles of psychological inflexibility and thought suppression.

Results indicate that 19.8% of participants were above the clinical cut off for secondary trauma based on the STSS and 25.2% presented as high risk for secondary trauma and 30.5% for burnout on the ProQol. Whilst STSS scores appear somewhat lower than previous studies on nurses (38%, Quinal et al., 2009; 33%,

Dominquez-Gomez & Rutledge, 2009), the results remain concerning. Additionally, results from the ProQol appear to be consistent with previous work with foster carers (Ottaway and Selwyn, 2016) as well as other professionals working with trauma victims (Bride et al., 2007; Stamm, 2005; Van Hook & Rothenberg, 2009). Compassion satisfaction was found to be negatively associated with compassion fatigue, in line with Stamm's (2010) model, supporting the idea that potential for mitigating effects of compassion satisfaction on compassion fatigue. Whilst similarities can be drawn from studies examining other caring professions, it should be noted here that fostering vulnerable children is unique and complex work unlike any other profession, and requires incredible emotional commitment. As such, all references to fostering as a job does not intend to simplify this important and at times complicated role. With this in mind, this study focuses primarily on compassion fatigue, but acknowledges there are range of stressors related to this role that have not been measured in this study (e.g. interactions with the local authority, issues around intimacy and decision making or personal stressors) which may also impact a carers' job satisfaction and intent to continue fostering.

Local authorities are currently facing a significant challenge in recruiting and retaining sufficient numbers of foster carers to take on the growing numbers of looked after children in the UK. The high rates of mental health difficulties and behavioural challenges in looked after children are well documented and these are known to contribute to the success of a placement. However, the experience of the foster carer themselves is under researched and this study aimed to explore the relationship between compassion fatigue with job satisfaction and foster carer's intent to continue fostering. Foster carers with lower job satisfaction and intent to continue had significantly higher compassion fatigue scores and this could be a target for intervention as discussed below. Additionally, compassion satisfaction scores were higher for foster carers with higher intent to continue and job satisfaction. The report of physical harm at

48% appears high and may well contribute to the level of secondary trauma in this sample. However, the data was collected as a single item as part of the demographic data and as such we would not claim that it is necessarily a precise representation.

Bivariate analyses found strong correlations between compassion fatigue and psychological inflexibility and thought suppression. Given the paradoxical effect of avoidant cognitive styles, foster carers with higher psychological inflexibility and a tendency to suppress thoughts may be at increased risk for compassion fatigue. Similarly, the negative emotional impact of their work may be exacerbated by a predisposition to engage in avoidant styles, i.e., be bidirectional (see below).

This is an initial study to establish proof of concept and the relation to some key variables. As such, the sample is relatively small and diverse in terms of the type of foster placement, and recruitment centring in and around one major urban region, all of which may impact the representativeness. One potential limitation is that the content of the survey may have influenced the type of respondent and affected the return rate. One possibility is that those experiencing higher strain and possibly at greater risk of compassion fatigue did not feel able to take the time to complete the survey and the current results underestimate the overall levels of compassion fatigue amongst foster carers. On the other hand, the survey content may have appealed to those experiencing high levels of strain as a means of reflecting experiences. Qualitative studies may be more able to investigate the potential influences driving participation and responses.

As a cross sectional study, we are unable to describe the direction of effects. Further research should be conducted to examine the direct effect compassion fatigue may have on intention to leave. Additionally, the outcomes for job retention and satisfaction were measured as binaries, derived from ordinal scales and although this has been done in other studies (e.g.

Denby et al., 1999), the relationships between satisfaction and retention are likely to be more nuanced than captured in these binary constructs. Future studies would benefit from more in depth investigation of these constructs, especially, if there was an aim to predict risk of leaving the profession based on current satisfaction. Conducting a longitudinal study, with repeated measures of compassion fatigue, may be particularly valuable in assessing the trajectory of foster carers' job retention in relation to their compassion fatigue levels. In terms of the risk factors investigated in this study, the relationship between compassion fatigue and other variables may well be bi-directional.

Clinical implications

The current study provides important evidence of the presence of compassion fatigue in foster carers which is relevant to those managing and supporting the foster carers in their work. The association between compassion fatigue and job satisfaction and intent to continue provides further support for increased awareness and support for foster carers as symptoms may have negative implications to placement stability and overall retention. Formal training on compassion fatigue and related risk factors, as well as encouraging services to screen for symptoms may be beneficial. This study has shown the ProQol and STSS are two reliable measures that could be used for screening symptoms of compassion fatigue.

In addition, the associations between avoidant cognitive styles and compassion fatigue have implications on further interventions that aim to reduce thought suppression and increase psychological flexibility, as a way to target possible mechanisms underlying compassion fatigue. Acceptance and Commitment Therapy (ACT; Hayes, Strosahl, & Wilson, 1999) is a psychological therapy that promotes a non-judgmental willingness to experience aversive events whilst simultaneously moving toward valued life goals, and an intervention incorporating an ACT framework could be useful for foster carers. Interventions incorporating mindfulness

(which may address avoidant coping styles, Shapiro et al., 2006) may also be worth consideration, although further research is required to explore their efficacy in relation to compassion fatigue.

Conclusions

This study is the first to describe rates of compassion fatigue in foster carers with approximately one quarter of participants reporting high levels of secondary trauma and burnout. This study also replicates previous findings of the association between compassion satisfaction and compassion fatigue. Higher levels of compassion fatigue were also associated with lower job satisfaction and lower intent to continue fostering. In order to promote placement stability and foster carer retention, services should pay attention to the wellbeing of foster carers particularly regarding recognition of compassion fatigue. Findings confirm an association between avoidant cognitive styles and compassion fatigue. Possible ways to target this could be to consider interventions based on increased psychological flexibility such as Acceptance and Commitment Therapy (ACT). Clinical implications are discussed in terms of direct support for foster carers through reflective practice and clinical interventions targeting cognitive styles such as ACT or mindfulness to promote foster carer wellbeing. Further research is required to explore the efficacy of these interventions and the wellbeing of foster carers generally.

References

- Adams, R. E., Boscarino, J. A., & Figley, C. R. (2006). Compassion fatigue and psychological distress among social workers: a validation study. *American Journal of Orthopsychiatry*, 76(1), 103-108.
- Baird, S., & Jenkins, S. R. (2003). Vicarious traumatization, secondary traumatic stress, and burnout in sexual assault and domestic violence agency staff. *Violence and victims*, 18(1), 71-86.
- Beck, C. T. (2011). Secondary traumatic stress in nurses: a systematic review. *Archives of psychiatric nursing*, 25, 1-10.
- Bond, F. W., Hayes, S. C., Baer, R. A., Carpenter, K. M., Guenole, N., Orcutt, H. K., Waltz, T., & Zettle, R. D. (2011). Preliminary psychometric properties of the Acceptance and Action Questionnaire-II: A revised measure of psychological inflexibility and experiential avoidance. *Behavior Therapy*, 42(4), 676-688.
- Boscarino, J. A., Figley, C. R., & Adams, R. E. (2004). Compassion fatigue following the September 11 terrorist attacks: a study of secondary trauma among New York City social workers. *International Journal of Emergency Mental Health*, 6, 57-66.
- Bride, B. E., Radey, M., & Figley, C. R. (2007). Measuring compassion fatigue. *Clinical Social Work Journal*, 35(3), 155-163.
- Bride, B. E., Robinson, M. M., Yegidis, B., & Figley, C. R. (2004). Development and validation of the secondary traumatic stress scale. *Research on social work practice*, 14(1), 27-35.
- Burns, B. J., Phillips, S. D., Wagner, H. R., Barth, R. P., Kolko, D. J., Campbell, Y., & Landsverk, J. (2004). Mental health need and access to mental health services by youths involved with child welfare: A national survey. *Journal of the American Academy of Child & Adolescent Psychiatry*, 43(8), 960-970.

- Burtson, P. L., & Stichler, J. F. (2010). Nursing work environment and nurse caring: relationship among motivational factors. *Journal of Advanced Nursing*, 66(8), 1819-1831.
- Carmel, M. J. S., & Friedlander, M. L. (2009). The relation of secondary traumatization to therapists' perceptions of the working alliance with clients who commit sexual abuse. *Journal of Counseling Psychology*, 56(3), 461-467.
- Clausen, J. M., Landsverk, J., Ganger, W., Chadwick, D., & Litrownik, A. (1998). Mental health problems of children in foster care. *Journal of Child and Family Studies*, 7(3), 283-296.
- Cohen, J. (1992). A power primer. *Psychological bulletin*, 112(1), 155-159.
- Conrad, D., & Kellar-Guenther, Y. (2006). Compassion fatigue, burnout, and compassion satisfaction among Colorado child protection workers. *Child abuse & neglect*, 30(10), 1081-1080.
- Cook, A., Spinazzola, J., Ford, J., Lanktree, C., Blaustein, M., Cloitre, M., & Van der Kolk, B. (2005). Complex trauma. *Psychiatric annals*, 35(5), 390-398.
- Cornille, T., & Meyers, T. (1999). Secondary traumatic stress among child protective service workers: Prevalence, severity and predictive factors. *Traumatology*, 5(1), 1-17.
- Denby, R., Rindfleisch, N., & Bean, G. (1999). Predictors of foster parents' satisfaction and intent to continue to foster. *Child abuse & neglect*, 23(3), 287-303.
- Department for Education. (2016). *Outcomes for children looked after by LAs: 31 March 2015*. UK: Crown Copyright.
- Department of Children Schools and Families. (2009). *Annual Report*. Crown Copyright.
- Dominguez-Gomez, E., & Rutledge, D. N. (2009). Prevalence of secondary traumatic stress among emergency nurses. *Journal of Emergency Nursing*, 35(3), 199-204.
- Farmer, E., Lipscombe, J., & Moyers, S. (2005). Foster carer strain and its impact on parenting and placement outcomes for adolescents. *British Journal of Social Work*, 35(2), 237-253.

- Figley, C. R. (1995). Compassion fatigue: Toward a new understanding of the costs of caring. In B. Stamm (Ed.), *Secondary Traumatic Stress: Self-care issues for clinicians, researchers and educators* (2 ed.). Lutherville, MD: Sidran Press.
- Figley, C. R. (2002). Compassion fatigue: Psychotherapists' chronic lack of self care. *Journal of clinical psychology*, 58(11), 1433-1441.
- Figley, C. R. (2013). *Compassion fatigue: Coping with secondary traumatic stress disorder in those who treat the traumatized*: Routledge.
- Ford, T., Vostanis, P., Meltzer, H., & Goodman, R. (2007). Psychiatric disorder among British children looked after by local authorities: comparison with children living in private households. *The British Journal of Psychiatry*, 190(4), 319-325.
- Ghahramanlou, M., & Brodbeck, C. (2000). Predictors of secondary trauma in sexual assault trauma counselors. *International Journal of Emergency Mental Health*, 2(4), 229-240.
- Good, P. (2005). *Permutation, Parametric, and Bootstrap Tests of Hypotheses* (3 ed.): Springer-Verlag New York.
- Greeson, J., Briggs, E., & Kisiel, C. (2011). Complex trauma and mental health in children and adolescents placed in foster care: Findings from the National Child Traumatic Stress Network. *Child Welfare*, 90(6), 91-108.
- Halfon, N., Mendonca, A., & Berkowitz, G. (1995). Health status of children in foster care: The experience of the Center for the Vulnerable Child. *Archives of Pediatrics & Adolescent Medicine*, 149(4), 386-392.
- Harber, A., Oakley, M., & Exchange, P. (2012). Fostering Aspirations: Reforming the foster care system in England and Wales. Retrieved from <http://www.policyexchange.org.uk/images/publications/fostering%20ambitions%20-%20jan%2012.pdf>

- Hayes, S., & Wilson, K. (1996). Experiential avoidance and behavioral disorders: A functional dimensional approach to diagnosis and treatment. *Journal of consulting and clinical psychology, 64*(6), 1152-1168.
- Hayes, S. C., Luoma, J. B., Bond, F. W., Masuda, A., & Lillis, J. (2006). Acceptance and commitment therapy: Model, processes and outcomes. *Behaviour research and therapy, 44*(1), 1-25.
- Hayes, S. C., Strosahl, K. D., & Wilson, K. G. (1999). *Acceptance and commitment therapy*. New York: Guilford Press.
- Holahan, C. J., Moos, R. H., Holahan, C. K., Brennan, P. L., & Schutte, K. K. (2005). Stress generation, avoidance coping, and depressive symptoms: a 10-year model. *Journal of consulting and clinical psychology, 73*(4), 658.
- Hooper, C., Craig, J., Janvrin, D. R., Wetsel, M. A., & Reimels, E. (2010). Compassion satisfaction, burnout, and compassion fatigue among emergency nurses compared with nurses in other selected inpatient specialties. *Journal of Emergency Nursing, 36*(5), 420-427.
- Kumpula, M. J., Orcutt, H. K., Bardeen, J. R., & Varkovitzky, R. L. (2011). Peritraumatic dissociation and experiential avoidance as prospective predictors of posttraumatic stress symptoms. *Journal of abnormal psychology, 120*(3), 617-627.
- Luciano, J. V., Belloch, A., Algarabel, S., Tomás, J. M., Morillo, C., & Lucero, M. (2006). Confirmatory factor analysis of the White Bear Suppression Inventory and the Thought Control Questionnaire: A comparison of alternative models. *European Journal of Psychological Assessment, 22*(4), 250-258.
- Marx, B. P., & Sloan, D. M. (2005). Peritraumatic dissociation and experiential avoidance as

predictors of posttraumatic stress symptomatology. *Behaviour research and therapy*, 43(5), 569-583.

Maslach, C. (1982). *Burnout: The cost of caring*. Englewood Cliffs, NJ: Prentice Hall.

Maslach, C., Schaufeli, W. B., & Leiter, M. P. (2001). Job burnout. *Annual review of psychology*, 52(1), 397-422.

McAuley, C., & Davis, T. (2009). Emotional well-being and mental health of looked after children in England. *Child & Family Social Work*, 14, 147-155.

McDermid, S., Holmes, L., Kirton, D., & Signoretta, P. (2012). The demographic characteristics of foster carers in the UK: Motivations, barriers and messages for recruitment and retention. Loughborough University Childhood Wellbeing Research Centre.

McLain, K. B. (2008). The impact of burnout, compassion fatigue, and compassion satisfaction on foster parenting. State University of New York at Buffalo.

Meadors, P., Lamson, A., Swanson, M., White, M., & Sire, N. (2010). Secondary traumatization in pediatric healthcare providers: Compassion fatigue, burnout, and secondary traumatic stress. *Journal of Death and Dying*, 60(2), 103-128.

Meltzer, H., Gatward, R., Corbin, T., Goodman, R., & Ford, T. (2003). *The mental health of young people looked after by local authorities in England*: HM Stationery Office.

Meyers, T., & Cornille, T. (2002). The trauma of working with traumatized children. In C. R. Figley (Ed.), *Treating Compassion Fatigue* (pp. 39-55). New York: Brunner-Routledge.

Murray, L., Tarren-Sweeney, M., & France, K. (2011). Foster carer perceptions of support and training in the context of high burden of care. *Child & Family Social Work*, 16, 149-158.

Oswold, J., Fegert, S., & Goldbeck, L. (2010). Post-Traumatic Stress Symptoms in Foster Children Following Maltreatment and Neglect. *Verhaltenstherapie*, 20(1), 37-44.

- Ottaway, H., & Selwyn, J. (2016). "No-one told us it was going to be like this": Compassion fatigue and foster carers summary report. Fostering Attachments Ltd.
- Perron, B. E., & Hiltz, B. S. (2006). Burnout and secondary trauma among forensic interviewers of abused children. *Child and Adolescent Social Work Journal*, 23(2), 216-234.
- Phelps, A., Lloyd, D., Creamer, M., & Forbes, D. (2009). Caring for carers in the aftermath of trauma. *Journal of aggression, maltreatment & trauma*, 18(3), 313-330.
- Potter, P., Deshields, T., Divanbeigi, J., Berger, J., Cipriano, D., Norris, L., & Olsen, S. (2010). Compassion fatigue and burnout: prevalence among oncology nurses. *Clinical journal of oncology nursing*, 14(5), 56-62.
- Quinal, L., Harford, S., & Rutledge, D. N. (2009). Secondary traumatic stress in oncology staff. *Cancer nursing*, 32(4)
- Ray, S., Wong, C., White, D., & Heaslip, K. (2013). Compassion satisfaction, compassion fatigue, work life conditions, and burnout among frontline mental health care professionals. *Traumatology*, 19(4), 255-267.
- Rhodes, K., Orme, J., & Buehler, C. (2001). A comparison of family foster parents who quit, consider quitting, and plan to continue fostering. *Social service review*, 75(1), 84-114.
- Richardson, J., & Joughin, C. (2000). *The mental health needs of looked after children*: RCPsych Publications.
- Rock, S., Michelson, D., Thomson, S., & Day, C. (2015). Understanding foster placement instability for looked after children: A systematic review and narrative synthesis of quantitative and qualitative evidence. *British Journal of Social Work*, 45(1), 177-203.
- Rodger, S., Cummings, A., & Leschied, A. W. (2006). Who is caring for our most vulnerable children?: The motivation to foster in child welfare. *Child abuse & neglect*, 30(10), 1129-1142.

- Sawyer, M., Carbone, J., Searle, A., & Robinson, P. (2007). The mental health and wellbeing of children and adolescents in home-based foster care. *Medical Journal of Australia*, 186(4), 181-184.
- Shapiro, S. L., Carlson, L. E., Astin, J. A., & Freedman, B. (2006). Mechanisms of mindfulness. *Journal of clinical psychology*, 62(3), 373-386.
- Sinclair, I., Gibbs, I., & Wilson, K. (2004). *Foster Carers: Why they stay and why they leave*: Jessica Kingsley Publishers.
- Sprang, G., Craig, C., & Clark, J. (2011). Secondary traumatic stress and burnout in child welfare workers: A comparative analysis of occupational distress across professional groups. *Child Welfare*, 90(6), 149-168.
- Stamm, B. (2010). The concise ProQOL manual. *Pocatello, ID: ProQOL. org*.
- Stamm, B. H. (2002). Measuring compassion satisfaction as well as fatigue: Developmental history of the Compassion Satisfaction and Fatigue Test.
- Stamm, B. H. (2005). *The ProQOL Manual: The Professional Quality of Life Scale: Compassion satisfaction, burnout & compassion fatigue/secondary trauma scales*. (Vol. Available at: www.isu.edu/~bhstamm). Baltimore, MD: Sidran Press.
- Thompson, B. L., & Waltz, J. (2010). Mindfulness and experiential avoidance as predictors of posttraumatic stress disorder avoidance symptom severity. *Journal of anxiety disorders*, 24(4), 409-415.
- Ting, L., Jacobson, J. M., Sanders, S., Bride, B. E., & Harrington, D. (2005). The secondary traumatic stress scale (STSS) confirmatory factor analyses with a national sample of mental health social workers. *Journal of Human Behavior in the Social Environment*, 11(3-4), 177-194.
- Tull, M. T., Gratz, K. L., Salters, K., & Roemer, L. (2004). The role of experiential avoidance in

posttraumatic stress symptoms and symptoms of depression, anxiety, and somatization.

The Journal of nervous and mental disease, 192(11), 754-761.

Van Hook, M. P., & Rothenberg, M. (2009). Quality of life and compassion satisfaction/fatigue and burnout in child welfare workers: A study of the child welfare workers in community based care organizations in central Florida. *Social Work and Christianity*, 36(1), 36.

Vanschoonlandt, F., & Vanderfaeillie, J. (2013). Parenting stress and parenting behavior among foster mothers of foster children with externalizing problems. *Children and Youth Services Review*, 35(10), 1742-1750.

Wee, D., & Myers, D. (2003). Compassion satisfaction, compassion fatigue, and critical incident stress management. *International Journal of Emergency Mental Health*, 5, 33-37.

Wegner, D. M., Schneider, D. J., Carter, S. R., & White, T. L. (1987). Paradoxical effects of thought suppression. *Journal of personality and social psychology*, 53(1), 5.

Wegner, D. M., & Zanakos, S. (1994). Chronic thought suppression. *Journal of personality*, 62(4), 615-640.

Whenan, R., Oxlad, M., & Lushington, K. (2009). Factors associated with foster carer well-being, satisfaction and intention to continue providing out-of-home care. *Children and Youth Services Review*.

Wilson, K., Sinclair, I., & Gibbs, I. (2000). The trouble with foster care: The impact of stressful 'events' on foster carers. *British Journal of Social Work*, 30(2), 193-209.

TABLES AND FIGURES

Figure 1: Mean scores and standard errors for low and high intent to continue fostering

Figure 2: Mean scores and standard errors for low and high job satisfaction

Table 1: Demographic Information

Variable		N	%
Gender	Female	101	77.1
	Male	30	22.9
Age	18-30	3	2.3
	31-40	8	6.1
	41-50	44	33.6
	51-60	60	45.8
	60+	16	12.2
Ethnicity	White British	57	43.5
	White Irish	1	0.8
	Any other White background	1	0.8
	Mixed White and Asian	1	0.8
	Asian or Asian British Indian	3	2.3
	Asian or Asian British Pakistani	1	0.8
	Asian or Asian British Bangladeshi	2	1.5
	Black or Black British Caribbean	7	5.3
	Black or Black British African	1	0.8
	Other Ethnic Groups: Any other ethnic group	1	0.8
Education	Missing	56	42.7
	Higher Education & professional/vocational equivalents	65	49.6
	A levels, vocational level 3 and equivalents	25	19.1
	GCSE/O Level grade A*-C, vocational level 2 and equivalents	27	20.6
	Other qualifications: level unknown	7	5.3
Relationship	No qualifications	7	5.3
	Single, never married	14	10.7
	Married	87	66.4
	Cohabiting	7	5.3
	Widowed	2	1.5
	Divorced	18	13.7
Children	Separated	3	2.3
	Have children of their own	118	90.1
Employer	Local Authority	110	84.0
	Private	20	15.3
	Both	1	0.8
Number of years fostering	Less than 1 year	6	4.6
	1-5 years	48	36.6
	5-10 years	32	24.4
	10-15 years	24	18.3
	15-20 years	8	6.1
	More than 20 years	13	9.9
Type of placement offered	Long-term and Short-term	50	38.2
	Long-term only	29	22.1
	Short-term only (includes respite and emergency care)	44	33.6

	Kinship	4	3.1
	Other	4	3.1
Typical length of placement offered	Short term (less than two weeks)	5	3.8
	Medium term (more than two weeks but less than one year)	36	27.5
	Long term (more than one year)	89	67.9
Child with disability	Currently caring for child with a physical and/or learning disability	39	29.8
Physical harm	Been physically hurt or threatened by a young person in their care	63	48.1
Mental health services	Looked after a child who required CAMHS involvement	82	62.6
Intent to continue	Not at all likely to continue	8	6.1
	Somewhat likely to continue	21	16.0
	Very likely to continue	102	77.9
Job Satisfaction	Not at all satisfied	5	3.8
	Somewhat dissatisfied	13	3.9
	Somewhat satisfied	56	42.7
	Very satisfied	57	43.5

Table 2: Rates of secondary trauma, burnout and compassion satisfaction from the ProQol

	Secondary Trauma	Burnout	Compassion Satisfaction
High (>75th percentile)	33 (25.2%)	40 (30.5%)	24 (18.3%)
Moderate	67 (51.1%)	62 (47.3%)	71 (54.2%)
Low (<25th percentile)	31 (23.7%)	29 (22.1%)	36 (27.5%)

Table 3: Mean scores and standard deviations from STSS

	Mean (SD)
STSS Total	30.58 (12.03)
Intrusions*	9.36 (3.75)
Avoidance	9.47 (4.06)
Arousal	11.78 (5.02)

*missing data from 1 participant

Table 4: Bivariate correlation of ProQol and STSS measures

	ProQol		
	Secondary Trauma	Burnout	Compassion Satisfaction
STSS Total	.707**	.676**	-.376**
Intrusions	.619**	.526**	-.256**
Avoidance	.669**	.681**	-.395**
Arousal	.692**	.669**	-.392**

** . Correlation is significant at the 0.01 level (2-tailed).

Table 5: Bivariate correlations of secondary trauma, intrusions, avoidance and arousal (from STSS) and burnout (from ProQol) with psychological inflexibility (AAQ-II) and thought suppression (WBSI)

	AAQ-II	WBSI
Secondary Trauma	.349**	.597**
Intrusions	.279**	.558**
Avoidance	.344**	.563**
Arousal	.351**	.556**
Burnout	.520**	.437**

** . Correlation is significant at the 0.01 level (2-tailed).