King's Research Portal

DOI:

[10.1016/j.freeradbiomed.2019.09.029](https://doi.org/10.1016/j.freeradbiomed.2019.09.029)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Knock, G. A. (2019). NADPH oxidase in the vasculature: Expression, regulation and signalling pathways; role in normal cardiovascular physiology and its dysregulation in hypertension. *Free Radical Biology and Medicine*, 145, 385-427. <https://doi.org/10.1016/j.freeradbiomed.2019.09.029>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

NADPH oxidase in the vasculature: expression, regulation and signalling pathways; role in normal cardiovascular physiology and its dysregulation in hypertension

Dr Greg Knock, PhD

Author for correspondence:

Dr Greg Knock
Senior Lecturer
Dpt. of Inflammation Biology
School of Immunology & Microbial Science
Faculty of Life Science & Medicine
King's College London

Table of contents

1. Introduction
2. NADPH oxidase isoform expression and regulation in the vasculature
 - 2.1. NOX isoforms
 - 2.2. NOX1 and NOX2 'Canonical' activation pathway.
 - 2.3. Other 'non-canonical' regulatory sub-units and NOX chaperones.
 - 2.4. Regulation of NOX4 in the vasculature.
 - 2.5. Regulation of NOX5 in the vasculature.
 - 2.6. Compartmentalisation of NOX isoforms and ROS metabolism.
3. Proximal targets for NOX-derived ROS in the vasculature.
 - 3.1. Protein tyrosine phosphatases.
 - 3.2. Receptor and non-receptor tyrosine kinases.
 - 3.3. Protein kinase C.
4. Role of NOX in VSM contraction.
 - 4.1. Pro-contractile actions of exogenous ROS.
 - 4.2. NOX/ROS-dependent Ca^{2+} influx pathways.
 - 4.3. NOX/ROS-dependent Ca^{2+} store handling.
 - 4.4. NOX/ROS-dependent, Ca^{2+} -independent contractile signalling.
 - 4.5. Role of NOX in GPCR-induced contraction.
 - 4.6. Role of NOX in myogenic, pressure/stretch-induced contraction.
5. Role of NOX in VSM relaxation.
 - 5.1. Activation of Endothelial NOX.
 - 5.2. Cross-talk between NOX and eNOS in the endothelium.
 - 5.3. Role of NOX-derived ROS in endothelial hyperpolarisation.
 - 5.4. NOX-derived H_2O_2 as a diffusible hyperpolarising factor.
 - 5.5. Pro-relaxant targets for ROS in VSM.
6. Role of NOX-derived ROS in hypertension.
 - 6.1. Expression of NOX isoforms/subunits in hypertension.
 - 6.2. Impaired endothelium-dependent relaxation and enhanced contractility.
 - 6.3. Vascular remodelling in hypertension.
 - 6.4. Effects of selective NOX inhibition or over-expression in experimental hypertension.
7. NOX in the pulmonary circulation.
 - 7.1. Differences between pulmonary and systemic arteries.
 - 7.2. Role for NOX-derived ROS in HPV.
 - 7.3. Role of NOX in pulmonary hypertension.
8. Summary and conclusions.

Figure legends

Tables

List of abbreviations

4EBP	Eukaryotic translation initiation factor 4E-binding protein
5-HT	5-hydroxytryptamine
16 α -HOE	16 α -hydroxyestrone
Ang II	Angiotensin II
ASK-1	Apoptosis signal-regulating kinase-1
BCL6	B-cell lymphoma 6
BH ₄	Tetrahydrobiopterin
BK _{Ca}	Large conductance Ca ²⁺ activated K ⁺ channel
BMPR2	Bone morphogenetic protein receptor-2
[Ca ²⁺] _i	Intracellular Ca ²⁺ concentration
cADPR	Cyclic ADP-ribose
CAMK-II	Ca ²⁺ -calmodulin activated kinase-II
CPI-17	C-kinase potentiated protein phosphatase-1 Inhibitor-17
CSK	c-Src kinase
DAG	Diacylglycerol
DCF	2,7-dichlorofluorescein
DHE	Dihydroethidium
DOCA salt rat	Deoxycorticosterone acetate salt rat
DPI	Diphenyleneiodonium
EBP50	Ezrin-radixin-moesin binding phosphoprotein 50
EDHF	Endothelium-derived hyperpolarising factor
EGFR	Epidermal growth factor receptor
eNOS	Endothelial nitric oxide synthase
ET-1	Endothelin-1
FAK	Focal adhesion kinase
GPCR	G-protein coupled receptor
GRB2	Growth-factor receptor bound protein-2
GSH	Reduced glutathione
H ₂ O ₂	Hydrogen peroxide
HETE	Hydroxyeicosatetraenoic acid
HIF-1 α	Hypoxia-inducible factor-1 α
HPV	Hypoxic pulmonary vasoconstriction
HSP27	Heat-shock protein 27
HSP90	Heat-shock protein 90
IGFR	Insulin-like growth factor receptor
IK _{Ca}	Intermediate conductance Ca ²⁺ -activated K ⁺ channel
K _{IR}	Inward rectifier K ⁺ channel
ILK	Integrin-linked kinase
IP ₃	Inositol trisphosphate
IP ₃ R	IP ₃ receptor
JAK2	Janus kinase-2
K _V	Voltage-gated K ⁺ channel
LMW-PTP	Low molecular weight protein tyrosine phosphatase
LVGCC	L-type voltage-gated Ca ²⁺ channels
MAP	Mean arterial pressure
MCT	Monocrotaline
MLC ₂₀	Myosin-light chain-20
MLCK	Myosin light-chain kinase
MLCP	Myosin light-chain phosphatase

mTOR	Mammalian target of rapamycin
MYPT1	Myosin phosphatase targeting subunit-1
NAADP	Nicotinic acid adenine dinucleotide phosphate
NF- κ B	Nuclear factor kappa-light-chain-enhancer of activated B cells
NO $\cdot$	Nitric oxide
NOX	Nicotinamide adenine dinucleotide phosphate (NADPH) oxidase
NRTK	Non-receptor tyrosine kinase
ONOO $^-$	Peroxynitrite
PDGF	Platelet-derived growth factor
PDGFR	Platelet-derived growth factor receptor
PDI	Protein disulphide isomerase
PECAM-1	Platelet-endothelial cell adhesion molecule-1
PH	Pulmonary hypertension
PAH	Pulmonary arterial hypertension
PIP ₂	Phosphatidyl inositol bisphosphate
PIP ₃	Phosphatidyl inositol trisphosphate
PI3K	Phosphatidyl inositol-3 kinase
pKa	Dissociation constant for ionisation (of amino acid side-chains)
PKC	Protein kinase C
PKG	Protein kinase G
PLA ₂	Phospholipases A ₂
PLC	Phospholipases C
PLD	Phospholipase D
Poldip2	Polymerase delta interacting protein 2
PPAR γ	Peroxisome proliferator-activated receptor gamma
PTP	Protein tyrosine phosphatase
PTP1B	Tyrosine protein phosphatase non-receptors type 1
PTP ϵ M	Tyrosine protein phosphatase- ϵ M
PTEN	Phosphatase and tensin homolog
PYK2	Protein tyrosine kinase 2 β
RacGEF	Rac-specific guanine nucleotide exchange factor
RhoGEF	Rho-specific guanine nucleotide exchange factor
ROCK	Rho-activated kinase
ROS	Reactive oxygen species
RTK	Receptor tyrosine kinase
RyR	Ryanodine sensitive channel
S1P	Sphingosine-1 phosphate
SERCA	Sarcoplasmic reticulum Ca ²⁺ ATPase
SHP2	Src-homology region-2 domain-containing phosphatase-2
SHR	Spontaneously hypertensive rat
SK _{Ca}	Small conductance Ca ²⁺ -activated K ⁺ channel
SOD	Superoxide dismutase
STAT	Signal transducer and activator of transcription
TGF- β	Transforming growth factor- β
TRPA1	Transient receptor potential ankyrin-1
TRPC1	Transient receptor potential canonical-1
TRPC6	Transient receptor potential canonical-6
TSP-1	Thrombospondin-1
TNF- α	Tumor necrosis factor- α
VEGFR	Vascular endothelial growth factor receptor

VSM	Vascular smooth muscle
VSMCs	Vascular smooth muscle cells
WKY	Wistar-Kyoto rat

Abstract

The last 20-25 years have seen an explosion of interest in the role of NADPH oxidase (NOX) in cardiovascular function and disease. In vascular smooth muscle and endothelium, NOX generates reactive oxygen species (ROS) that act as second messengers, contributing to the control of normal vascular function. NOX activity is altered in response to a variety of stimuli, including G-protein coupled receptor agonists, growth-factors, perfusion pressure, flow and hypoxia. NOX-derived ROS are involved in smooth muscle constriction, endothelium-dependent relaxation and smooth muscle growth, proliferation and migration, thus contributing to the fine-tuning of blood flow, arterial wall thickness and vascular resistance. Through reversible oxidative modification of target proteins, ROS regulate the activity of protein tyrosine phosphatases, kinases, G proteins, ion channels, cytoskeletal proteins and transcription factors. There is now considerable, but somewhat contradictory evidence that NOX contributes to the pathogenesis of hypertension through oxidative stress. Specific NOX isoforms have been implicated in endothelial dysfunction, hyper-contractility and vascular remodelling in various animal models of hypertension, pulmonary hypertension and pulmonary arterial hypertension, but also have potential protective effects, particularly NOX4. This review explores the multiplicity of NOX function in the healthy vasculature and the evidence for and against targeting NOX for antihypertensive therapy.

1. Introduction

Vascular smooth muscle (VSM) function is regulated by a highly complex network of intracellular signalling pathways incorporating cell surface receptors, G proteins, ion channels/transporters, kinases/phosphatases, cytoskeletal proteins/adaptors, transcription factors, all under the regulation of endocrine, neuronal, paracrine/autocrine, metabolic and mechanical stimuli. For these stimuli to contribute effectively to vascular function, they must be able to induce the production of second messenger molecules and these must be able to influence the function of specific cellular targets in a pre-determined spacio-temporal manner, and these effects must be readily reversible. Established second messenger molecules include cytosolic Ca^{2+} , cyclic nucleotides, phospholipids and more recently, reactive oxygen species (ROS). ROS can be defined as any electrophilic oxygen containing species, but in living cells, the primary species are superoxide ($\text{O}_2^{\cdot-}$) and its dismutation product hydrogen peroxide (H_2O_2), which are produced through one or two electron reduction of molecular oxygen, respectively. Electron transfer is catalysed by members of a large family of oxidoreductase enzymes. For many of these, including complex I and III of the mitochondrial electron transfer chain, uncoupled endothelial nitric oxide synthase (eNOS), xanthine oxidase, cyclooxygenases and lipoxygenases, ROS production is considered to be secondary to the principle function of the enzyme, at least under normal physiological conditions [1-7]. However, the main source of ROS in the vasculature (including in VSM and endothelium), is from the family of nicotinamide adenine dinucleotide phosphate (NADPH) oxidases (NOX), for which their sole function is the production of superoxide and/or H_2O_2 [8-10]. In phagocytes the function of NOX is to generate an 'oxidative burst' the purpose of which is to kill and destroy engulfed pathogens. In non-phagocytic cells, including VSM and endothelium however, it is now known that NOX-derived ROS act as essential second messenger molecules contributing to the regulation of normal cellular function and their production is tightly regulated. Only during 'oxidative stress', a key feature of cardiovascular disease, are ROS produced in excess and considered harmful. Of the seven known isoforms of NOX, four are expressed in VSM and endothelium, namely NOX1, 2, 4, and 5.

Experimental and theoretical analysis of the chemical properties of superoxide and H_2O_2 in aqueous or hydrophobic environments suggests that both of these ROS species can modify protein function through direct oxidation of the sulfhydryl side chains (-SH) of cysteine residues [11, 12], resulting in conformational changes that may enhance or inhibit protein activity. Single electron oxidation (sulfenylation; R-SOH) is readily reversible through the combined action of thioredoxins, glutathione peroxidases and reduced glutathione (GSH). Additional oxidation of -SOH (sulfinylation; R-SO₂H) is also reversible through the action of sulfiredoxin-1, but further oxidation of -SO₂H (sulfonylation; R-SO₃H) is considered irreversible [13-15]. Cysteine oxidation may also result in reversible formation of intra-molecular or inter-molecular disulphide bonds that stabilize a particular tertiary or quaternary structure, again to either promote or inhibit function. Under normal conditions, cysteine oxidation is highly selective, being dependent on the dissociation constant (pKa) of the sulfhydryl group in question and therefore its propensity to donate electrons. In addition to direct oxidation by ROS, cysteine residues may be oxidised indirectly by oxidised peroxyredoxin (PRX) or nitrosylated (R-SNO) by the reactive nitrogen species, nitric oxide ($\text{NO}^{\cdot}$) [16, 17]. Nitrosylation requires a prior single

electron oxidation ($R-S\cdot$) and is reversible through the action of thioredoxin-1 and GSNO-reductase [17].

Thioredoxin may also catalyse trans-nitrosylation of cysteine by nitroso-glutathione (GSNO) [17], or cysteine may be directly glutathionylated by oxidised glutathione (R-SSG). Recent work has also suggested that ROS-dependent oxidation of methionine (R-SCHO) may also influence protein and vascular function and this is reversible through the action of methionine sulfoxide reductase [18, 19]. Cysteine, lysine and histidine residues may also be modified through the formation of Michael adducts with lipid peroxides ($LOO\cdot$) such as 4-hydroxynonenal, generated by oxidation of poly-unsaturated fatty acids. Being lipids, this latter form of modification may be of particular importance for the oxidative regulation of membrane bound proteins, such as ion channels [20].

Superoxide spontaneously reacts with $NO\cdot$, to form peroxynitrite ($ONOO\cdot$). $ONOO\cdot$ also modifies protein function by nitrating tyrosine residues ($R-ONO$), via reaction with CO_2 and formation of the intermediate nitrogen dioxide and carbonate anion radicals, $NO_2\cdot$ and $CO_3\cdot^-$ [17]. Nitration influences protein function by preventing or mimicking phosphorylation [21]. In the presence of free cytosolic Fe^{2+} , H_2O_2 is converted to the very powerful oxidant hydroxyl radical ($OH\cdot$) which oxidises proteins indiscriminately, regardless of pKa, thus is unlikely to be of physiological relevance. Key targets for reversible oxidative modification by NOX-derived ROS in VSM function include a wide range of cell surface and intracellular proteins, including phosphatases, kinases and other cytosolic enzymes such as G-proteins and adapter proteins, ion channels and transporters and transcription factors.

This review will provide an update on expression, compartmentalisation, regulation and function of the different NOX isoforms expressed in the vasculature. It will detail the complex signalling networks both upstream of NOX activation and downstream of ROS production. It will report on how NOX activity is regulated by factors that influence VSM function, including vasoconstrictors, vasodilators, pressure/stretch, flow/shear-stress, mitogenic stimuli and hypoxia. Principle ROS effector proteins and downstream signalling pathways relevant to normal VSM function (contraction/relaxation) and the influence of endothelial factors on that function ($NO\cdot$, endothelium-derived hyperpolarising factor, EDHF) will be examined. Also included is a summary of how the expression and function of specific NOX isoforms is altered in hypertension, with often conflicting evidence for and against their importance to the pathogenesis of the disease. The role of NOX4 is particularly contentious. Finally, there will be a discussion of NOX function in the pulmonary circulation, how it differs from systemic, its importance to the pulmonary vasoconstriction response to acute hypoxia and its potential role in pulmonary hypertension.

2. NADPH oxidase isoform expression and regulation in the vasculature

2.1. NOX isoforms. The first identified and best characterised of the NADPH oxidases is the membrane-spanning flavocytochrome b558, a heterodimer composed of gp91^{phox} (now known as NOX2), and p22^{phox}. gp91^{phox} is composed of six transmembrane α -helices connected by five loops, three of which are extracellular and the other two intracellular, while p22^{phox} is composed of two transmembrane α -helices with cytosolic tails at both termini [22-24]. b558 generates superoxide by

the single electron reduction of two molecules of oxygen, the two electrons being transferred one at a time from a single molecule of NADPH at a dehydrogenase domain via FAD and two internally bound haem units [25]. This enzyme was identified as being responsible for the phagocytic respiratory burst in leucocytes [22], but soon after found to be expressed at considerably lower but functionally significant levels in both vascular smooth muscle cells (VSMCs) [26] and endothelium [27]. NOX2 requires specific stimuli for full activation, and collectively the intracellular loops of gp91^{phox} and p22^{phox} interact with cytosolic regulatory subunits p47^{phox}, p67^{phox} and p40^{phox} following a sequence of signalling events resulting in their translocation from the cytosol where they reside as a non-active heterotrimer [28].

The essential steps in the activation of the phagocytic oxidase begin with activation of protein kinase C (PKC) and PKC-dependent phosphorylation of the cytosolic regulatory subunit p47^{phox}. This phosphorylation triggers cytosolic to membrane translocation of p47^{phox}, together with other cytosolic regulatory subunits p67^{phox} and p40^{phox}, their association with gp91^{phox} and p22^{phox} and partial oxidase activation [29-31]. p67^{phox} is thought to contribute to the dehydrogenation of NADPH and the transfer of e⁻ to FAD [30, 32-34]. Full oxidase activity also requires phosphatidylinositol-3 kinase (PI3K) and the small G proteins Rac1 or Rac2. PI3K is required mainly because p47^{phox} and p40^{phox} also possesses phosphoinositide binding sites [35-37], while Rac1 binds the dehydrogenase domain of gp91^{phox} and p67^{phox} and is required for complete assembly and full activation of the oxidase [22, 30, 38]. gp91^{phox} also appears to require direct phosphorylation by PKC during its activation [39]. As detailed below, vascular NOX2 is activated in a similar way, although the specifics of how PKC, PI3K and Rac are activated and subcellular distribution of subunits may vary.

There are now known to be seven isoforms of the catalytic subunit of NOX (NOX 1-5 and DUOX1 & 2), of which NOX 1, 2, 4 & 5 are expressed in the vasculature [23]. The relative abundance of NOX1 and NOX2 appears to vary depending on the cell type, size of vessel and vascular bed, as well as species and physiological/pathophysiological conditions [23, 26, 40-45]. Collectively, the literature suggests that both are expressed in both endothelium and VSMCs but NOX2 appears to be the dominant subtype in endothelium, while NOX1 predominates in VSMCs. NOX4 is ubiquitously expressed at high levels in both VSMCs and endothelium of humans and rodents [23, 45-53] and in one report, is 100-fold more abundant in endothelium than NOX2 [53]. NOX5 is highly expressed in the human vasculature, but curiously absent from rodents [54-57]. All other NOX/phox subunits expressed in phagocytes are also present in VSMCs and endothelium [23, 26, 58-60]. As in phagocytes, p22^{phox} is essential for G-protein coupled receptor (GPCR)-induced NOX1/2 activity in the vasculature [61-63]. As with NOX2, p47^{phox}, p67^{phox}, p40^{phox} and Rac also participate in the activation of NOX1 [64], but vascular cells and tissues expressing NOX1 also express homologues of p47^{phox} and p67^{phox}, NOXO1 and NOXA1, respectively [23](discussed further in section 2.3).

Low levels of preassembled NOX1 or NOX2 [41, 60] together with active NOX4 [65] or NOX5 [66] probably contribute to basal ROS levels in unstimulated cells, and under these conditions, the rate limiting step for basal NOX-derived superoxide is probably the availability of cytosolic NADPH [67].

This 'basal' ROS undoubtedly contributes to normal resting cellular functions such as basal eNOS activity and resting smooth muscle tone, as well as making a contribution to ROS-dependent protein-folding in the endoplasmic reticulum [46, 51, 68-70]. However, NOX isoforms respond to external stimuli by generating brief bursts of ROS in the short term or sustained increases over longer periods. Over the short to medium terms (seconds to several minutes), enhanced NOX activity requires an initial stimulus which, through a series of signalling events, results in full activity of the enzyme. The exact nature of the stimulus, activation cascades and their effects depend largely on the NOX isoform being examined, its location in the cell and in which cell type it is expressed. As expected, acute stimuli for the production of ROS in VSM and/or endothelium include GPCR, flow/shear-stress, pressure/stretch and hypoxia, but also some growth factor and cytokine receptors (Table 1).

2.2. NOX1 and NOX2 'Canonical' activation pathway. GPCR agonists are the most frequently studied short term stimuli for NOX1 and NOX2 activity in the vasculature. The most extensively studied of the vascular GPCR agonists for NOX1 or NOX2 is angiotensin II (Ang II), acting via the type 1 angiotensin receptor (AT1R) to induce vasoconstriction, gene expression, growth, proliferation and migration [26, 40, 43, 45, 55, 71-108], but there are many others known to activate NOX in VSMC or endothelium through similar if not identical signalling pathways. These include endothelin-1 (ET-1) [55, 84, 109-114], ATP/purinergic agonists [64, 115-118], urotensin II [119-121], thrombin [62, 122-130], vasopressin [131], catecholamines (especially, alpha agonists) [131-137], 5-hydroxytryptamine (5-HT) [138], arachidonic acid and various eicosanoids, including prostaglandins, hydroxyeicosatetraenoic acids (HETEs) [6, 58, 139-146] and sphingolipids [147-150]. Here, common pathways for the activation of NOX in both VSM and endothelium are described. Additional activation by mechanical forces in the vasculature, most notably pressure/stretch in VSM and flow/shear-stress in endothelium, are discussed later.

The multiple steps required to be taken to achieve full and complete activation of NOX1 or NOX2 in vascular tissue are similar to those that apply to the phagocytic oxidase in leukocytes and, as illustrated in Figure 1, can be divided into two phases: (1) a first phase, corresponding to an initial rapid, often transient elevation in ROS production, and (2) a second phase, corresponding to a sustained longer lasting increase in ROS production. The most important step in the first phase is the phosphorylation and translocation of p47^{phox} to the membrane, together with recruitment of p67^{phox} and p40^{phox} and their association with NOX1/2 and p22^{phox}. For this to occur, the initial stimulus must be capable of activating phospholipases C, D or A₂ (PLC, PLD, PLA₂) and the kinases PKC and/or Src, which then phosphorylate p47^{phox}. The requisite step in the second phase is the recruitment of Rac to the partly assembled enzyme. Activation of Rac and complete assembly of the oxidase appears to require further ROS-dependent activation of Src and activation of PI3K, often via transactivation of a receptor tyrosine kinase (RTK), usually epidermal growth factor receptor (EGFR). Each of these key steps are outlined below.

Phospholipase-dependent activation of PKC is an important first step in assembly of vascular NOX1 or NOX2 by GPCR agonists acting via Gq, although which phospholipase and which lipid products are considered most important for this activation appears to vary. Many GPCR expressed by VSMCs are

coupled to Gq and the activation of PLC, which generates inositol trisphosphate (IP₃) and diacylglycerol (DAG). Conventional PKC isozymes are activated by Ca²⁺ released from IP₃-sensitive stores and both conventional and novel isozymes are activated by DAG. Despite this, relatively few studies have directly shown NOX activation to be PLC-dependent in the vasculature. These include NOX2-dependent ROS production induced by Ang II in renal preglomerular VSMCs [133] and NOX1-dependent ROS production induced by sphingosylphosphorylcholine in mesenteric and pulmonary artery [148], sphingosine-1 phosphate (S1P) in rat aorta [147], advanced glycosylation end-products in retinal capillaries [151] or by thrombospondin-1 (TSP-1) in mouse aorta [152]. Alternatively, there is substantial evidence for a role of the PLD- or PLA₂-derived lipids phosphatidic acid, arachidonic acid or lysophosphatidylcholine, in activation of NOX by GPCR. PLD and PLA₂ are activated by Gβγ subunits [153, 154], while PLD activation in response to Ang II and other GPCR agonists also requires the small GTPase ARF6 [90, 108, 155]. In response to Ang II in VSMC, these lipid metabolites are likely to be activating PKC, since as outlined below, PKC-dependent serine phosphorylation of p47^{phox} is a prerequisite for p47^{phox} translocation to the membrane and its association with NOX1/2 and p22^{phox}, although which PKC isozyme is specifically shown to be PLD or PLA₂-dependent in this context is rarely specified [79, 80, 100, 156, 157]. Various stimuli in endothelium, including thrombin, low density lipoprotein and tumor necrosis factor-α (TNF-α) appear to couple PLA₂-derived arachidonic acid metabolites to NOX activity [124, 158-160], and two of these stimuli are likely to involve PKC as the target of those metabolites [158, 159].

There is strong evidence supporting a necessary role for PKC in NOX activation in VSM of multiple vascular beds and in some endothelial preparations, as demonstrated by the blocking effect of PKC inhibitors on NOX activity and ROS production in response to various stimuli including direct PKC activation by phorbol esters [67, 140, 161, 162]. These include Ang II in mouse and human coronary, human subcutaneous artery, rat aortic VSMCs and mouse renal preglomerular arterioles [77, 80, 100, 133, 152, 163], TSP-1 in mouse aorta [152], TNF-α or platelet-derived growth factor (PDGF) in human pulmonary endothelium and rat aortic VSMCs [158, 163], U46619 or sphingosylphosphorylcholine in bovine coronary, rat mesenteric or bovine aortic endothelium [6, 58, 140, 148] and direct PKC activators in rat or bovine aorta or bovine coronary artery [140, 156, 164]. Hypoxia is also capable of activating NOX via PKC [165] and is almost certainly doing so via upstream stimulation of mitochondrial ROS production (section 7.2), since PKC is also ROS-sensitive (section 3.3). Multiple studies have used p47^{phox}^{-/-} mice, siRNA or inhibitory peptides that target p47^{phox} interaction with p22^{phox}, gp91^{phox} and/or NOX1 at the membrane to show the dependence of the initial stimulated ROS production on p47^{phox} in the vasculature [72, 93, 97, 104, 118, 132, 140, 144, 148, 152, 158, 165-168]. As in phagocytes, several studies have confirmed that the main role of PKC in vascular NOX activation is to serine phosphorylate p47^{phox}, triggering translocation of p47^{phox} and p67^{phox} to the membrane and their association with NOX and p22^{phox} [6, 26, 58, 97, 118, 152, 158, 162, 169]. In addition, some studies have implicated specific PKC isozymes in vascular NOX activation, namely PKC-α [162], PKC-β [163, 170, 171], PKC-δ [169], PKC-ε [148, 165] or PKC-ζ [6, 108, 158]. There is also evidence for a role for phosphorylation of p22^{phox}, p67^{phox} and p40^{phox} by a range of serine/threonine kinases in non-vascular

cells [22, 30, 172], much of which may also be relevant in the vasculature but is yet to be demonstrated.

Interestingly, NOX activity in VSMCs and endothelium, unlike that in leukocytes, also appears to be dependent on tyrosine kinase activity, particularly the non-receptor tyrosine kinase Src, or other closely related kinases. Src is strongly involved in both pressor and vascular remodelling responses to Ang II and a range of other stimuli [101] and is strongly associated with ROS signalling in the vasculature eg. [173, 174] and reviewed in [175-177]. Multiple studies have used Src inhibitors, siRNA or Src^{-/-} mouse to show the necessity of Src activity for either NOX1 or NOX2 activation in various vascular preparations, including bovine coronary artery VSMCs [140], rat or mouse mesenteric and pulmonary arteries [101, 148], renal preglomerular VSMCs [133], rat or mouse aortic VSMCs [77, 101], human coronary or subcutaneous arteries [82, 178] or human pulmonary endothelium [179]. Like PKC, activated Src can trigger p47^{phox} translocation and initial NOX1/2 activation by direct tyrosine phosphorylation of p47^{phox} [82, 140, 144, 179] or indirectly via phosphorylation and activation of PKC [173] which also phosphorylates p47^{phox}, as described above. Interestingly, Src may also be activated by PKC-dependent phosphorylation [133, 140, 180, 181], resulting in positive feedback between the two kinases. This association between the two kinases is potentially initiated by basal ROS since both kinases are ROS sensitive [176, 177] (sections 3.2 and 3.3). A number of additional pathways have been suggested for the activation of Src by GPCR, including direct activation by Gβγ subunits [182, 183]. Translocation of cytosolic subunits and NOX activation may also involve a degree of actin cytoskeleton re-organisation, via Src and/or PKC-dependent phosphorylation of actin-associated proteins. One study has suggested that before PKC-δ can phosphorylate and trigger the translocation of p47^{phox} to the plasma membrane it must first phosphorylate smooth muscle-22α thus freeing itself from the actin cytoskeleton [169]. Others show in human VSMCs and endothelium that Src phosphorylates cortactin, facilitating phosphorylation of p47^{phox} and its translocation to the membrane [85, 184, 185]. A third study has also suggested that translocation of p47^{phox} and activation of vascular NOX1 is facilitated by binding of p47^{phox} to ezrin-radixin-moesin binding phosphoprotein 50 [89].

Once the holoenzyme is assembled at the plasma membrane, full activation of vascular NOX1 or NOX2 also requires association of the catalytic subunit and p67^{phox} with activated Rac1 or Rac2, as described above for the phagocytic oxidase. This has been shown for basal ROS and ROS production induced by GPCR agonists and growth factors in mouse, rodent and human VSMCs and bovine endothelium [62, 71, 77, 90, 144, 178, 186, 187]. Being a small G-protein, Rac requires activation by Rac-specific guanine nucleotide exchange factors (RacGEFs). RacGEFs in turn require phospholipid binding and/or tyrosine phosphorylation for their activation. RacGEF tyrosine phosphorylation is Src-dependent, either direct or via intermediary activation of other non-receptor tyrosine kinases focal adhesion kinase (FAK), protein tyrosine kinase 2β (PYK2) or Janus kinase-2 (JAK2) [188-192]. In response to GPCR stimulation, Src also indirectly contributes to activation of the lipid kinase PI3K via phosphorylation and transactivation of growth factor receptors (particularly EGFR) [77, 125, 130, 145, 149, 193-195]. Subsequently, PI3K-derived phosphatidyl inositol biphosphate (PIP₂) and/or phosphatidyl inositol

trisphosphate (PIP₃) promote translocation and activation of Rac1 and complete NOX complex assembly at the membrane [77, 96, 196]. PIP₃ binds and activates multiple RacGEFs [197-199], while in neutrophils, but not yet shown in VSM or endothelium, PIP₂ and PIP₃ bind and provoke conformational changes in p47^{phox} and p40^{phox} thus promoting/stabilising NOX2 assembly [36, 37]. PIP₂ and PIP₃ production is also supported by concomitant inhibition of the lipid phosphatase PTEN (phosphatase and tensin homolog) by the same stimuli that activate PI3K [142], perhaps by the initial NOX-derived ROS [196, 200] (see also section 3.1).

Interestingly, a number of studies have shown that growth factor receptor transactivation and PI3K activation in vascular cells is also NOX-dependent [62, 149, 150]. A popular explanation for this is that partial activation of NOX generates sufficient ROS to further activate Src, promoting docking to and transactivation of the growth factor receptor and that the NOX/Src/EGFR/PI3K/Rac/NOX signalling axis constitutes multiple nested positive feedback loops between NOX1/2-derived ROS and ROS dependent signalling (particularly Src and PKC) [42, 77, 140]. Indeed, exogenously applied H₂O₂ stimulates NOX activity in lysed VSMC membrane fractions [165]. It has been suggested that this feedback loop also forms the basis of the late sustained phase of ROS production which persists for at least 30 min after the initial GPCR-induced stimulus in VSMCs, following an EGFR-independent transient rapid phase [77]. Finally, GPCR-induced EGFR transactivation in VSMCs may also occur through direct oxidative activation of EGFR [200], ROS-dependent matrix metalloprotease activation, and epidermal growth factor shedding [130, 136, 201] or via oxidative inhibition of the tyrosine phosphatase Src-homology region-2 domain-containing phosphatase-2 (SHP2) [121, 200]. These are all parallel pathways which can probably operate independently of Src. Of relevance to downstream signalling, activated growth factor receptors also act as docking sites for the assembly of growth and proliferation signalling complexes, including growth-factor receptor bound protein-2 (GRB2) and PI3K/AKT (section 6.3).

2.3. Other ‘non-canonical’ regulatory sub-units and NOX chaperones. NOX1 can form functional oxidase units at the membrane with NOXO1 instead of p47^{phox} [202] and/or with NOXA1 instead of p67^{phox} [126, 202, 203]. Unlike p47^{phox}, NOXO1 lacks the auto-inhibitory domain and PKC binding site, so does not require phosphorylation for its membrane targeting and association with NOX1/p22^{phox}, perhaps conferring a greater degree of constitutive activity to the oxidase than with p47^{phox} [202], although NOX1 presumably still requires PI3K and Rac1 for full activity. Alternatively, NOX1 may also be activated by a combination of p47^{phox} and NOXA1 [126, 203], indeed one study suggests that while human endothelium expresses p67^{phox}, it is completely absent from rodent aortic VSMCs and replaced by NOXA1 [203], and it has also been noted that NOXA1 expression in VSMC is lower than that of other cytoplasmic activators of NOX, thus introducing a possible rate limiting step for NOX1 activation, but suggesting regulation of NOXA1 expression may be a key way of altering ROS production in the long term [126]. Alternatively, in VSMCs, physical association between NOXA1 and NOX1 may be facilitated by direct PKC-β-mediated phosphorylation of NOX1 [163]. NOX2 may also be activated by NOXA1, but with lower affinity/efficacy than by p67^{phox} [204]. NOX1/2 expression and activity are also positively regulated by protein disulphide isomerase (PDI) and heat-shock protein 90 (HSP90) in both

endothelium and VSM [205-207]. PDI forms a ROS-induced intermolecular disulphide bond with p47^{phox} which is apparently necessary for NOX1 enzyme complex stability [208].

2.4. Regulation of NOX4 in the vasculature. NOX4 is expressed in both endothelium and VSM and according to some reports, is the most highly expressed NOX in both cell types [209, 210]. Like NOX1 and NOX2, NOX4 also possesses a dehydrogenase domain that requires FAD for the transfer of electrons to O₂ [211], but it differs in that it usually makes H₂O₂ instead of superoxide, presumably by the direct transfer of two electrons to each O₂ molecule rather than one to each of two O₂ molecules [45, 212-215], but may instead make superoxide in some intracellular compartments [212, 216]. Despite some reports that NOX4 preferentially uses NADH over NADPH as an electron donor [216, 217], other studies in isolated membrane preparations demonstrated a similar selectivity for NADPH as for other NOX isoforms [212]. Like NOX1 and NOX2, the catalytic subunit of NOX4 associates with p22^{phox} and this binding is necessary for catalytic activity [213, 214, 216, 218, 219], but unlike NOX1 or NOX2, it does not require p47^{phox}, p67^{phox}, NOX01, NOXA1, p40^{phox} or HSP90 for its activation [205, 213], suggesting that it exhibits a high degree of constitutive activity and makes a major contribution to basal ROS production in the vasculature, particularly basal H₂O₂ [45, 46, 65, 211-213, 215, 216, 220].

Detailed characterisation of signalling pathways for the acute activation of NOX4 is scarce, but there is certainly sufficient evidence to support the suggestion that it responds acutely to flow/shear-stress in endothelium [52] (section 5.1), GPCR and growth factor stimuli in endothelium [215, 221-223], and various GPCR agonists in VSM [45, 131, 224-226] (Table 2). Some studies suggest that NOX4 activity is supported by arachidonic acid [210, 224] the production of which is stimulated by some GPCR, but they disagree over whether NOX4 activation by this lipid is also dependent on Rac [223, 224, 227], as is the case with NOX1 and NOX2, or is entirely independent of it [210, 213, 222]. There is also disagreement over whether or not NOX4 responds to activators of PKC and therefore whether it requires serine/threonine phosphorylation for full activity [210, 216]. There is as yet no evidence that NOX4 activity is regulated by tyrosine phosphorylation. The only clearly defined cytosolic regulatory protein for NOX4 is polymerase delta interacting protein 2 (Poldip2), which is necessary for NOX4-mediated H₂O₂ production in VSMC [226, 228-230] and is a known actin and integrin-associated protein [49]. These relatively simple interactions are summarised in Figure 2. NOX4 is also expressed in subcellular organelles, such as the sarcoplasmic reticulum, but whether it can respond to acute stimuli at this location is unknown (see below).

2.5. Regulation of NOX5 in the vasculature. NOX5 is similar to other NOX isoforms in that it has six transmembrane domains, two hemes and a dehydrogenase domain that binds FAD and NADPH, but it is fully functional as an oxidase in the absence of both p22^{phox} and any cytosolic regulatory subunits [231, 232], except perhaps for HSP90 which appears to be essential for the long-term stability of the enzyme complex [205, 233], perhaps by antagonising the enzymatic instability induced by caveolin in endothelial caveolae [234]. The primary site of acute regulation of NOX5 is an N-terminal cytosolic domain containing four EF hands that bind free Ca²⁺, as well as Ca²⁺-calmodulin binding domains, that collectively relieve oxidase auto-inhibition [235, 236]. As a result, NOX5-derived ROS production is

enhanced by stimuli that raise intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) in endothelium and VSM, such as thrombin, Ang II, ET-1, U46619 and bradykinin, acting both via voltage-dependent extracellular Ca^{2+} influx and sarcoplasmic reticulum Ca^{2+} -induced Ca^{2+} release [55, 66, 232, 237]. NOX5 is therefore essentially constitutively active but with a 'basal' activity that is dependent on $[\text{Ca}^{2+}]_i$ [66]. However, in addition to this basal activity, the sensitivity of vascular NOX5 to increases in $[\text{Ca}^{2+}]_i$ may also be enhanced via phosphorylation by kinases such as PKC- α [238, 239], Ca^{2+} -calmodulin activated kinase-II (CAMK-II) [240], ERK-MAP kinase [241] or c-Abl [242], but inhibited by the Ca^{2+} -independent PKC- ϵ [239]. These various routes for the activation of NOX5 are illustrated in Figure 2. Since H_2O_2 elevates $[\text{Ca}^{2+}]_i$ (sections 4.2 and 4.3), NOX5 activation may also contribute to a positive feedback loop between multiple NOX isoforms [242], although this has yet to be demonstrated in vascular cells.

2.6. Compartmentalisation of NOX isoforms and ROS metabolism. Since ROS are generally highly reactive species, it has been suggested that for them to operate as effective selective second messenger molecules, the site of ROS production should be localized near to or in the same sub-cellular compartments as the target proteins or even be in direct contact with those target proteins in the same compartment. There is a growing body of evidence to support this [15, 48, 243-246]. As described above, all NOX isoforms expressed in the vasculature are located in the plasma membrane, but most are likely concentrated in or around specific structures on the cell surface. In VSM, plasma membrane NOX4 preferentially localises with vinculin and clustered integrins [48, 234], where presumably Poldip2 couples NOX4-derived ROS to activation of FAK, integrin-linked kinase (ILK) and RhoA at focal adhesions [49, 226, 229, 230, 247]. This may suggest a possible role for NOX4 in pressure/stretch-induced myogenic contraction (section 4.6) and VSM migration and proliferation (section 6.3). In endothelium, NOX1 and 2 are found in caveolae where they will associate with and be activated by the 'mechanosome' of the endothelial flow/shear-stress sensor. NOX5 is also located in caveolae, but is not known to be activated by flow. NOX4 is sensitive to flow but unlike NOX1, 2 and 5, it is not located in the caveolae of endothelial cells and therefore caveolin cannot influence its activity in the short term, but it does negatively regulate its expression [234]. In caveolae, NOX1, 2 and 5 are maintained in a resting state by caveolin-1 through disruption of the enzyme complex until this is relieved by HSP90 [205, 233]. NOX4 is presumably located elsewhere on the surface of endothelial cells, but to which structures it binds is unknown. The role of endothelial cell surface NOXs in sensing flow is discussed in more depth later in this review (section 5.1).

Due to the orientation of NOX catalytic subunits in the plasma membrane, they will be producing primarily extracellular ROS. However, NOX4-derived H_2O_2 is free to diffuse through the plasma membrane to have intracellular actions, and this may be facilitated by cell surface aquaporins, widely expressed in the cardiovascular system [248-250] while for the negatively charged superoxide generated by NOX1, 2 or 5 to also have intracellular actions, transport must be facilitated by plasma membrane anion channels, including the chloride channel CLIC-3 [251] and the volume sensitive channel LRRC8A [252], or it must first be dismutated to H_2O_2 by extracellular superoxide dismutase (SOD) [15, 253]. Alternatively, it is now fairly well established that functional NOX enzymes are also

located within various subcellular organelles, including the endosomes, sarcoplasmic reticulum, nucleus and mitochondria.

It is well known that growth factor receptors, cytokine receptors and GPCR are endocytosed upon ligand binding or following transactivation in the case of RTKs, either to be recycled back to the plasma membrane or sent to lysosomes for degradation. However, it is also now known that key signalling events occur in the endosome following receptor activation [243, 252]. It has been suggested that this includes activation of NOX1 in VSM [130, 252, 254, 255], and for activation of NOX2 in endothelium [256, 257]. This appears to occur following endocytosis of the activated NOX enzyme together with the activating receptor. For NOX1 and NOX2, this may be facilitated by p40^{phox}-dependent interaction with phosphoinositides in the endosomal membrane [36] or NOX-dependent phosphorylation of dynamin or caveolin-1 [252]. NOX4 and NOX5 are also found in endosomes and therefore may also contribute to endosomal RTK or GPCR signalling [242, 258]. Also endocytosed are several associated activator and effector proteins, such as Src, Rac and tyrosine phosphatases [245, 255, 258, 259]. Endosomes also contain SOD1, so superoxide can be dismutated to H₂O₂, enabling both NOX-derived ROS species to contribute to cell signalling in the endosome as a 'redoxosome' [15, 245, 246, 254, 257, 258, 260]. Alternatively, there is evidence that endosomal superoxide may be released into the cytosol through volume-sensitive anion channels (LRRC8) [252] or CIC-3 [254, 255, 261] and of course, H₂O₂ is also free to diffuse into the cytosol.

NOX4 is also co-expressed with p22^{phox} in the sarcoplasmic reticulum of endothelium and VSM [46, 51, 53, 219, 225] and probably generates ROS within the lumen of the organelle [219]. The H₂O₂ produced here is also free to diffuse out or exit the compartment via aquaporins [15], but specific functions for NOX-derived ROS within the endoplasmic reticulum have been considered. In particular, it may contribute to the oxidising environment required for normal disulphide-dependent protein folding [15, 68] or contribute to the oxidative modification of Ca²⁺ release channel function (section 4.3). NOX5 is also expressed in the endoplasmic reticulum of endothelium [232], but its functions there are unknown. A functional splice variant of NOX4 lacking transmembrane domains but which still associates with p22^{phox} has been detected in the nucleus of VSMCs where it may directly contribute to oxidative regulation of gene expression [216, 262, 263]. NOX4 is also reportedly expressed in the mitochondria of cardiac myocytes [264, 265], but whether it is also expressed there in VSM or endothelium and what its function may be is unclear. The mitochondria also generates superoxide through leakage of electrons from the inner membrane electron transport chain [1, 2] and this is of particular functional importance in vascular responses to hypoxia, where it may be coupled to plasma membrane or endosomal NOX activity (sections 7.2 and 7.3). Movement of ROS from within mitochondria and sarcoplasmic reticulum into the cytosol is also facilitated by aquaporins and CIC-3 (or equivalent) [15].

The multiple feedforward and positive feedback loops between ROS and upstream signalling pathways derived from NOX1 or NOX2 will ensure the response to a stimulus is rapid and amplifiable, but for this to be of any use as a controllable cell signal and to contribute to the highly compartmentalised

nature of the response there must also be an 'off-switch'. For feedback involving Src and PKC it has been suggested that this may arise out of the fact that as well as being activated by ROS, the two kinases are also inactivated by ROS, but the latter perhaps only at higher concentrations [176, 244, 266]. Apart from this, cells possess powerful antioxidant enzyme systems capable of limiting the accumulation and diffusion of any ROS generated and to some extent these too are compartmentalised. If superoxide does not react with target proteins or with NO $\cdot$ to produce ONOO $\cdot$, then it will eventually be dismutated to H $_2$ O $_2$. Superoxide dismutase is present in the cytosol and endosomes (SOD1) [260], mitochondria (SOD1, MnSOD/SOD2) and extracellular space (EC-SOD) [15]. While H $_2$ O $_2$ is more freely diffusible and less reactive than superoxide, its ultimate complete reduction to H $_2$ O is catalysed by cytosolic catalase, or by peroxiredoxins or glutathione peroxidases located in the cytosol, mitochondria and endoplasmic reticulum [15]. The action of glutathione peroxidase is supported by the main cellular REDOX buffer, reduced glutathione (GSH). Oxidised glutathione (GSSG) is re-reduced to GSH by glutathione reductase utilising NADPH, which is itself generated in the pentose phosphate pathway, the ultimate source of cellular reducing equivalents. The compartmentalisation and metabolism of ROS are illustrated in Figure 3.

3. Proximal targets for NOX-derived ROS in the vasculature.

A wide range of signalling proteins involved in vascular smooth muscle function have been identified as being sensitive to ROS and often particularly sensitive to NOX-derived ROS. For many of these, the target protein is probably not directly sensitive to cysteine oxidation but is responding to ROS-induced modulation of an upstream 'proximal ROS target'. Downstream effector pathways include multiple protein kinases, ion channels, guanine nucleotide exchange factors for small G-proteins of the Rho family, cytoskeletal proteins and transcription factors. Their roles in NOX-dependent vascular function will be detailed in subsequent sections. This section will focus on essential proximal ROS sensors, protein tyrosine phosphatases (PTPs) and non-receptor tyrosine kinases (NRTKs), as well as PKC and CAMK-II.

Tyrosine phosphorylation has long been known as an essential early signalling event in response to growth factors and cytokines, since growth factor receptors are themselves tyrosine kinases and cytokine receptors directly activate non-receptor tyrosine kinases as an essential early step in their signalling cascades [21, 191, 267, 268], but more recently it has also been shown that vascular smooth muscle pro-contractile stimuli, GPCR agonists in general, stretch in many vascular beds and hypoxia in the pulmonary circulation, are also associated with enhanced total cellular tyrosine phosphorylation in the vasculature [21, 103, 269-273] and reviewed in: [176, 177]. Enhancements in cellular tyrosine phosphorylation induced by either growth or contractile stimuli in VSM are inhibited by antioxidants [267, 274, 275] or SOD and/or catalase [267, 276] and altered by inhibition/over-expression of NOX [103, 259, 274, 277, 278]. Similarly, exogenous ROS cause an increase in tyrosine phosphorylation in vascular tissues [103, 174, 191, 273, 279-283]. Net tyrosine phosphorylation and thus relative activity of the phosphorylated target proteins will always be determined by the relative opposing activities of tyrosine phosphatases and kinases [284]. For example, tyrosine phosphatase inhibitors generally induce constriction in vascular smooth muscle, whereas tyrosine kinase inhibitors induce relaxation

[82, 273, 279, 283]. As discussed in previous sections, all the stimuli mentioned above induce NOX-derived ROS production in the vascular wall, therefore the observed NOX-dependent effects of ROS on net tyrosine phosphorylation are most likely occurring via the combined effects of ROS-induced PTP inhibition and ROS-induced NRTK activation. As discussed below, there is a direct link between the induced NOX activity and the stimulation of downstream contractile and mitogenic effector pathways.

3.1. Protein tyrosine phosphatases. Tyrosine phosphatases, dual-specific phosphatases and the lipid phosphatase PTEN share a unique signature motif HC(X)₅R(S/T) containing a nucleophilic cysteine which confers an unusually low pKa for ionization of 4-5 compared to the typical pKa of nonreactive cysteines of 8.5 in most other proteins [285]. Oxidation usually results in the formation of an intramolecular disulphide or sulfenamide bond which inhibits phosphatase activity by preventing docking to protein targets and relief of auto-inhibition. Like most serine threonine phosphatases, reduced PTPs (under low ROS conditions) do not require external stimuli for their activation and will be constitutively active when bound to target phospho-proteins, but will be progressively inactivated when cells are stimulated to increase ROS production, [196, 244, 286-290]. In some cases, as with PTEN, the phosphatase is protected from oxidation by PRX, but oxidation of PRX results in disulphide formation and inactivation of the phosphatase [16]. This oxidative inhibition is readily reversible through the action of glutathione peroxidase or thioredoxin [268, 288, 290].

Key tyrosine phosphatases important in the control of VSM or endothelial function that are sensitive to NOX-derived ROS include SHP2 [121, 291], PTP1B (tyrosine protein phosphatase non-receptors type 1) [51, 276], protein tyrosine phosphatase- ϵ M (PTP ϵ M) and low molecular weight protein tyrosine phosphatase (LMW-PTP) [292, 293]. All of these are key negative regulators of growth factor receptors and the signalling complexes that are activated by them. SHP2 is particularly important in suppressing activity of EGFR and platelet-derived growth factor receptor (PDGFR), by dephosphorylating phosphotyrosines essential for the docking and activation of PI3K and GRB2. This may occur in endosomes co-expressing growth factor receptors and NOX1 or NOX4 and contributes to the feedback between RTK transactivation, PI3K activity and NOX activation [258]. PTP1B negatively regulates vascular endothelial growth factor receptor (VEGFR) signalling in endothelium [276] and appears to co-localise with NOX4 and EGFR in the endoplasmic reticulum where oxidative inhibition of the phosphatase promotes EGFR trafficking [51]. There is also some evidence that dual-specific tyrosine/threonine phosphatases are ROS-sensitive. This class of phosphatase is responsible for the de-phosphorylation and inactivation of MAP kinases and the oxidative inactivation of MAP kinase phosphatase-1 contributes to MAP kinase activation in VSM [294]. As mentioned above, oxidative inhibition of PTEN contributes to NOX1/2 activation by supporting PI3K-derived PIP₂ and PIP₃ production [196], although this has yet to be directly demonstrated in vascular tissues. PIP₃ is also required for the activation of AKT, thus PTEN is a possible route through which ROS may promote VSM growth. Obviously, phosphatases will oppose the actions of kinases by dephosphorylating downstream target proteins, such as ion channels, guanine nucleotide exchange factors and other kinases, but PTPs also de-

phosphorylate tyrosine residues on kinases themselves and these phospho-tyrosines may be inhibitory or stimulatory of kinase activity. Indeed most tyrosine kinases possess both (See below).

3.2. Receptor and non-receptor tyrosine kinases. There are two major families of tyrosine kinase, the receptor tyrosine kinases (RTK) and the non-receptor tyrosine kinases (NRTK). A feature shared by all tyrosine kinases is that they require auto-phosphorylation for full kinase activity. Most growth factor receptors are RTKs, and auto-phosphorylation is triggered by agonist binding and receptor dimerization or by transactivation involving phosphorylation by another kinase, usually an NRTK such as Src, in response to other stimuli, usually GPCR. This generates phospho-tyrosine docking sites for multiple downstream signalling pathways, including PI3K, GRB2 and PLC- γ . There are multiple families of NRTK, including the Src family kinases (notably Src, Fyn & Yes in vascular tissues), FAK, the closely related PYK2 and Janus kinase (especially JAK2). These kinases are important for mediating growth factor and cytokine-induced gene expression and growth-related signalling but are also key regulators of smooth muscle electrical excitability, intracellular Ca^{2+} and actin cytoskeleton dynamics, and therefore contractility. All of the above have, in one way or another, been implicated in ROS-dependent signalling in the vasculature and are either upstream or downstream of NOX-dependent ROS production, and make a major contribution to the control of smooth muscle tone induced by GPCR agonists [133, 174, 176, 177, 272, 295, 296], stretch [194] and hypoxia [174, 271, 297] (section 7.2), or the control of VSM proliferation [95, 298-300] (section 6.3).

Src (the principle member of the Src-family kinases) is one of a handful of protein kinases known to be directly ROS-sensitive, although the majority of evidence for this stems from extensive work in non-vascular cells and tissues. Src possesses two key phospho-tyrosine residues, an inhibitory one on the C-terminal tail (Y527 in humans) and a stimulatory one on the kinase activation loop (Y418 in humans). When Y527 is phosphorylated by c-Src kinase (CSK), internal SH2 binding promotes folding of the kinase, hiding the active site, and disruption of this internal binding by de-phosphorylation of Y527, or PKC-directed serine phosphorylation of the SH2 domain, triggers auto-phosphorylation of Y418, fully activating the kinase [301, 302]. Src also contains several reactive cysteine residues, including one in the SH2 domain and one in the kinase domain, oxidation of which induces hyper-phosphorylation of Y418 [303], and/or intermolecular disulfide bridge formation that may prevent internal Y527/SH2 binding, thus prolonging kinase activity [304]. Oxidation of these two cysteine residues and Src activation occurs in response to NOX activation [305]. ROS may also module Src activity indirectly via oxidative inhibition of CSK and PTPs that de-phosphorylate Y418 (stimulatory of Src) or of PTPs that dephosphorylate Y527 (inhibitory of Src). The former may ensure a rapid response to and positive feedback to NOX activation, while the latter may contribute to a timely termination of the signal [301, 306-308]. In line with Src activity generally having a positive relationship with NOX1/2-derived ROS generation, there is evidence that exogenous ROS activate Src and that VSM contractile or proliferative stimuli induce ROS-dependent or NOX-dependent Src oxidation and auto-phosphorylation [125, 173, 174, 259, 303, 309-312], and stimuli that activate NOX-dependent contractile responses and contractions induced by exogenous ROS are both suppressed by inhibitors of Src family kinases [148, 174, 177, 311].

Several other NRTKs, including FAK, PYK2 and JAK2, are ROS sensitive in the vasculature, but unlike Src, are probably not directly ROS-sensitive. The likeliest route is downstream of Src, which phosphorylates these kinases, triggering stimulatory auto-phosphorylation [145, 309, 313-315]. Alternatively, it may occur downstream of oxidative inhibition of a PTP that de-phosphorylates their activation loop [189, 291], or phosphorylation by a ROS sensitive PKC [316]. Either way, they are activated by GPCR agonists and other contractile, migratory or mitogenic stimuli in a manner dependent on NOX and endogenous ROS production or by exogenous ROS [109, 110, 126, 189, 191, 229, 309, 317, 318]. A couple of these studies have implicated specific NOX isoforms in activation of NRTK: JAK2 is stimulated by PDGF via NOX5-derived ROS [317] while FAK is activated by PDGF at focal attachments in a NOX4/Poldip2 dependent manner [229] and subsequently phosphorylates multiple focal adhesion associated proteins [319]. As discussed earlier, transactivation of growth factor receptors such as EGFR, PDGFR and insulin-like growth factor receptor (IGFR) is generally considered to be indirectly NOX and ROS-sensitive because the NRTK (usually Src) that initially phosphorylates the receptor in response to NOX activation is itself ROS-sensitive, as discussed above [125, 173, 194, 309]. As far as can be determined, only one study also suggests that EGFR transactivation may also involve direct sulfenylation of the EGFR kinase domain [200].

3.3. Protein kinase C and Ca²⁺-calmodulin-activated kinase II. PKC is a family of 12 cytosolic serine/threonine kinases and like Src, it appears to be both upstream of NOX activation and a directly ROS-sensitive NOX effector. Six PKC isozymes of three classes are expressed in vascular tissues: conventional (PKC- α , - β_1 , - β_2), novel (PKC- δ and - ϵ) and atypical (PKC- ζ) [320, 321]. All PKC isozymes are inactive when maintained in a folded configuration due to a regulatory domain containing an auto-inhibitory pseudo-substrate, which prevents access of the catalytic domain to substrate proteins. Unfolding and release of the pseudo-substrate is normally induced by Ca²⁺ (α and β only), DAG (α , β δ and ϵ only), other lipid mediators (all isozymes) and by tyrosine phosphorylation (all isozymes). Src is a key NRTK activator of PKC; reviewed in: [173, 176, 321]. Essential to the release of the pseudo-substrate are multiple cysteine residues in the regulatory domain and these are also highly sensitive to intramolecular disulphide bridge formation [322]. Cysteine oxidation removes the requirement for Ca²⁺, or lipid mediators, automatically triggering unfolding and activation of the kinase [176, 321]. There is also evidence that cysteine residues on the catalytic domain of PKC may also be sensitive to oxidation, but that this oxidation inhibits kinase activity [323]. However, this appears to require higher concentrations of ROS than oxidation of regulatory domain cysteine residues, so may contribute to the termination of the positive feedback between PKC and NOX once sufficient ROS have been generated [176, 323] (section 2.6). These biochemical studies were all performed in non-vascular systems, but are supported by studies in vascular tissues showing that PKC undergoes cytosolic to membrane translocation and is activated in response to exogenous ROS in VSM or endothelium [324-327] and that inhibition of specific PKC isozymes inhibits ROS-induced or ROS-dependent GPCR mediated contraction and associated signalling events in isolated arteries [148, 283, 325-327]. CAMK-II is another kinase that is potentially both upstream (specifically of NOX5) and downstream of ROS. This kinase is normally activated by Ca²⁺-calmodulin, binding of which triggers autophosphorylation at

Thr287, but is also activated by oxidation of two methionine residues (Met281/282). This oxidative activation is maintained independently of Ca^{2+} , but has the same effect on kinase activity [19, 328].

4. Role of NOX in VSM contraction.

VSM constriction negatively influences tissue blood flow while positively influencing vascular resistance and perfusion pressure. Thus, in order to match blood flow to the metabolic demands of the tissue, prevent tissue damage due to over-perfusion, contribute to thermoregulation and inflammation, or to regulate total peripheral resistance and MAP, VSM tone is under the control of circulating hormones (eg Ang II), peripheral neurotransmitters (noradrenaline, acetylcholine), paracrine and autocrine factors (bradykinin, histamine etc) haemodynamic forces (pressure/stretch, flow/shear-stress) as well as products of metabolism and PO_2 . Interestingly, NOX and the two principle ROS, superoxide and H_2O_2 , contribute to both contraction and relaxation responses in the vasculature, depending on the vascular beds and the nature of the stimulus. This has in part been investigated by testing the effects of application of exogenous H_2O_2 and artificially generated intracellular superoxide with xanthine/xanthine oxidase or cell permeable quinones such as LY83583 or menadione [329, 330] (Table 3). Despite being considered somewhat of a blunt instrument due to the use of potentially supra-physiological concentrations and the loss of subcellular compartmentalisation [244], the results of such experiments are largely supported by pharmacological and genetic manipulation of specific NOX isoforms as well as the use of non-specific antioxidants and highly selective antioxidant enzymes (SOD and catalase). There are several effector pathways through which ROS influence VSM contraction and relaxation, acting on both the VSM and the endothelium. The simplest is by a shift in the balance between the availability of eNOS-derived $\text{NO}\cdot$ and NOX-derived superoxide in favour of superoxide and/or a change in the ability of SOD to remove superoxide. This results in more superoxide chemically reacting with and 'scavenging' $\text{NO}\cdot$, generating the poor vasodilator $\text{ONOO}\cdot$, thus impairing endothelium-dependent relaxation and enhancing vascular tone [99, 102, 152, 331-333]. However, this mechanism by no means fully accounts for the observed effects of superoxide in the vasculature because inhibiting or ablating eNOS does not abolish the effect [137, 311, 334-336]. The alternatives are more complex. As discussed throughout this section and illustrated in Figures 4 and 5, ROS can induce phosphorylation of myosin-light chain-20 (MLC_{20}), a necessary step in actin-myosin cross-bridge cycling and contractile force generation, either by directly influencing VSM $[\text{Ca}^{2+}]_i$, which induces phosphorylation of MLC_{20} by myosin light-chain kinase (MLCK), or by inhibition of myosin light-chain phosphatase (MLCP). There is also some evidence for a role in MLC_{20} -independent effects on actin dynamics. In the following sections, evidence for the role of specific NOX isoforms in mediating contractile and relaxation stimuli will be discussed. Please note that many of the studies cited in the following sections have made use of pharmacological agents designed as specific inhibitors of NOX, such as apocynin, DPI and VAS2870, but which have subsequently been queried as non-specific flavoprotein inhibitors or even as general antioxidants rather than NOX inhibitors, particularly apocynin [337, 338]. Nevertheless, such studies are included here because they are by and large supported by parallel studies utilising specific NOX isoform/regulatory subunit siRNA, over-expression and knock-out/knock-in mouse models, as well as newer NOX inhibitors and antioxidant enzymes. Conclusions drawn from these studies are also corroborated by the parallel use of ROS measurement

and/or exogenous ROS, often in the same articles. Please see Tables 2 & 4 for an overview of the relevant literature in this regard.

4.1. Pro-contractile actions of exogenous ROS. In the absence of pre-constriction, superoxide directly constricts rat aorta [339] and rat pulmonary artery [311, 340], but not mesenteric or femoral artery [311]. In the absence of pre-constriction, H₂O₂ directly constricts mouse and rat aorta [131, 341-344], bovine coronary artery [194], canine cerebral artery [283], mouse carotid [342] and rat or rabbit pulmonary artery [146, 326, 340, 343, 343, 345-347], but may or may not directly constrict rat mesenteric artery [341, 341-343, 348, 348]. In various mammalian systemic arterial preparations pre-constricted with GPCR agonists or myogenic tone, both superoxide and H₂O₂ appear to exert mixed, concentration-dependent, contraction/relaxation effects [311, 325, 327, 334, 343, 349, 350], although the dominant effect of H₂O₂ tends to be relaxation (section 5). Superoxide consistently enhances GPCR-induced pre-constriction in pulmonary artery [174, 311, 351] (section 7.1.). Alternatively, in various mammalian systemic vascular preparations, both H₂O₂ and superoxide consistently enhance sub-maximal contraction induced by KCl [148, 341, 342, 344, 350-354], because KCl removes the influence of ROS on K⁺ channels (as explained further in section 5.5). Note that in some studies, superoxide generating systems X/XO or LY83583 reportedly cause contraction that is largely suppressed by SOD or only partly suppressed by catalase, suggesting that the contractile effect of superoxide does not necessarily require conversion to H₂O₂ [311, 339, 340, 351]. It is also worth highlighting that vasoconstriction in various vascular beds induced by exogenous ROS is inhibited by non-specific tyrosine kinase inhibitors [283, 341, 346], specific blockers of Src family kinases [174, 311, 352], and PKC inhibitors [283, 325, 326], suggesting that these proximal kinases contribute to ROS-induced VSM contraction either as direct ROS-sensors or downstream of PTP inhibition. There is evidence supporting a role for NOX-derived ROS, as well as Src and/or PKC, in all three of the main VSM contraction effector pathways (an increase in [Ca²⁺]_i, inhibition of MLCP and actin polymerisation).

4.2. NOX/ROS-dependent Ca²⁺ influx pathways. Smooth muscle contraction is primarily dependent on phosphorylation of MLC₂₀ by the Ca²⁺-calmodulin-dependent MLCK. Exogenous H₂O₂ and superoxide applied in the absence of pre-constrictor agents induce increases in [Ca²⁺]_i in aorta and cerebral or mesenteric arteries [131, 283, 325, 348], while in pulmonary artery, H₂O₂ does [326], but superoxide does not [311]. However, superoxide does enhance GPCR-induced Ca²⁺ responses in pulmonary artery [351]. A major source of this ROS-induced [Ca²⁺]_i response appears to be influx via L-type voltage-gated Ca²⁺ channels (LVGCC), particularly in systemic vessels. This is based partly on the inhibitory effects of specific LVGCC antagonists on exogenous ROS-induced [Ca²⁺]_i and contraction in aorta and cerebral, mesenteric and renal arteries [283, 325, 327, 344, 348, 354, 355] and the enhancement of whole-cell L-type Ca²⁺ currents in mesenteric artery VSMCs by superoxide [148] and in cerebral artery by H₂O₂ [105]. NOX-derived endogenous ROS also contribute to GPCR-induced [Ca²⁺]_i influx and contraction in part via enhanced opening of LVGCC. For example, in bovine pulmonary artery VSMCs, the U46619-induced [Ca²⁺]_i increase is partially sensitive to inhibition of either NOX1/2 or LVGCC [58], Ang II-induces apocynin or catalase-sensitive Ca²⁺ sparklets and contraction in rat

cerebral artery [105, 325] and sphingosylphosphorylcholine induces antioxidant-sensitive voltage-gated Ca^{2+} current in mesenteric artery VSMCs in association with p47^{phox} -dependent contraction [148]. In addition, mouse mesenteric artery expressing human NOX5 exhibit enhanced antioxidant-sensitive contraction induced by depolarisation with sub-maximal KCl [66]. It is not clear whether cell surface NOX4 regulates Ca^{2+} channel activity in VSM but it does activate transient receptor potential canonical-6 (TRPC6) in podocytes in response to Ang II stimulation [356].

There is conflicting evidence over the influence of ROS on LVGCC activity through direct cysteine oxidation or S-nitrosylation, some suggesting that it is stimulatory [357, 358] and others that it is inhibitory [359-362], perhaps depending on which region of the channel is targeted. Alternatively, ROS-dependent stimulation of LVGCC Ca^{2+} influx may be indirect, either via oxidative activation of protein kinases that activate the channel by phosphorylation, or via prior action on other membrane channels that depolarises the plasma membrane, thus opening LVGCC. VSM LVGCC open probability is increased by both serine/threonine and tyrosine phosphorylation, most probably PKC, CAMK-II and/or Src-mediated, because the pore-forming subunit of LVGCC ($\text{Cav}1.2$) is a phosphorylation target for all three kinases [325, 358, 363-365]. PKC- α contributes to Ang II-induced NOX-dependent induction of LVGCC Ca^{2+} sparklet activity and contraction in rat cerebral artery [325]. CAMK-II enhances H_2O_2 induced $\text{Cav}1.2$ current in ventricular myocytes [358]. While there is no direct evidence that Src phosphorylates $\text{Cav}1.2$ in response to ROS, it is a reasonable assumption, considering it is itself a key ROS target in VSM. Conceivably, ROS could depolarise VSM by either inhibiting K^+ channel activity or enhancing the activity of non-selective cation channels. One study suggests that LVGCC were being opened in response to ROS-dependent voltage-gated K^+ channel (K_v) inhibition and subsequent depolarisation [58]. However this response may be specific to the pulmonary vasculature (section 7.2), while a number of other studies in systemic vessels suggests that K_v channel activity in VSM is enhanced by ROS, contributing to a pro-relaxant effect [350, 351, 366] (section 5.5). Alternatively, ROS are more likely to depolarise VSM through the opening of Ca^{2+} -activated Cl^- channels or TRP channels, which themselves also conduct Ca^{2+} [327]. In aortic VSMCs, GPCR agonists induce inward Ca^{2+} currents and contraction that are dependent on both TRPC6 and NOX (probably 1 or 2) and these effects are mimicked by exogenous H_2O_2 [131, 367]. In rat pulmonary artery, H_2O_2 enhances TRPC1/Stim1/Orai1-dependent Ca^{2+} influx [368], and Ca^{2+} -sensing receptor/TRPC6-dependent Ca^{2+} influx [369], while in renal afferent arterioles superoxide activates Ca^{2+} -activated Cl^- channels to enhance myogenic tone [327]. As with LVGCC, ROS are likely to be activating these channels indirectly via intracellular signalling, perhaps dependent on PLC/DAG and/or PKC [327, 367, 370]. There is also some evidence, albeit primarily from cardiac muscle, that the plasma membrane Ca^{2+} ATPase is inhibited by ROS, a pro-contractile effect [13], while the $\text{Na}^+/\text{Ca}^{2+}$ exchanger may be activated, opposing this effect [13, 371].

4.3. NOX/ROS-dependent Ca^{2+} store handling. Most VSM GPCR are coupled to G_q to stimulate production of IP_3 , so will trigger Ca^{2+} release from the sarcoplasmic reticulum via the IP_3 receptor (IP_3R). However, they may also open IP_3R indirectly via NOX-derived ROS, since both exogenous and NOX-derived superoxide in VSM and endothelium appear to directly sensitize IP_3R to IP_3 , increasing

the rate of stimulated Ca^{2+} release [372, 373]. Alternatively, H_2O_2 -induced contractile responses in aorta and pulmonary artery are partially dependent on sarcoplasmic reticulum Ca^{2+} release via ryanodine sensitive channels (RyR) [326, 344]. In addition, many GPCR agonists induce RyR-dependent Ca^{2+} release, usually secondary to nearby Ca^{2+} influx or release by other sarcoplasmic reticulum channels such as the Two-Pore channel or IP_3R (Ca^{2+} -induced Ca^{2+} release). Of the three subtypes (RyR1, 2 and 3), RyR1 is probably pro-relaxant in systemic blood vessels ([374], but may be pro-contractile in others [326], whereas RyR2 is universally considered pro-contractile [375]. All three subtypes are known to be directly ROS sensitive, containing multiple reactive cysteine residues that are important for channel activity [13, 376]. However, the functional significance of this has not been demonstrated in VSM. Instead Ang II-induced, RyR-mediated, sarcoplasmic reticulum Ca^{2+} release in renal afferent arterioles and pulmonary artery is indirectly dependent on NOX2 and also perhaps NOX4, via ROS-dependent production of CD38-derived cyclic ADP-ribose (cADPR) and nicotinic acid adenine dinucleotide phosphate (NAADP), activators of the RyR and Two-Pore channel, respectively [377-379]. In addition, vascular RyR2 may also be activated by ROS-induced dissociation of the inhibitory FKBP12.6 protein [375]. Ca^{2+} is pumped back into the sarcoplasmic reticulum by the sarcoplasmic reticulum Ca^{2+} ATPase (SERCA), and there is evidence that SERCA activity is suppressed by direct irreversible cysteine oxidation [13, 380], thus prolonging increases in $[\text{Ca}^{2+}]_i$. Recent evidence suggest that this oxidation of SERCA, albeit in cardiac muscle, may be mediated by NOX4-derived ROS in the sarcoplasmic reticulum [381]. It is not known whether NOX5 expressed in the sarcoplasmic reticulum contributes the regulation of Ca^{2+} homeostasis. The role of NOX-derived ROS in VSM Ca^{2+} influx and Ca^{2+} release signalling pathways is summarised in Figure 4.

4.4. NOX/ROS-dependent, Ca^{2+} -independent contractile signalling. In parallel with Ca^{2+} -dependent activation of MLCK, there is also concomitant inhibition of MLCP which further enhances MLC_{20} phosphorylation and contraction independently of a change in $[\text{Ca}^{2+}]_i$, a process that is often termed ' Ca^{2+} -sensitization' [382]. In line with this, exogenous ROS induce contraction in permeabilised, Ca^{2+} -clamped rat pulmonary artery [311, 326, 343, 351], and a high concentration of superoxide also does so in non-permeabilised arteries without raising $[\text{Ca}^{2+}]_i$ [311]. The main signalling pathway in VSM for inhibition of MLCP is phosphorylation of myosin phosphatase targeting subunit-1 (MYPT1) by Rho-activated kinase (ROCK) [382, 383]. ROCK is a serine threonine kinase that exists in a folded inactive form in the cytosol until activated by the monomeric G-protein RhoA [384]. Prior to this, RhoA is activated by the exchange of bound GDP for bound GTP, a process usually catalysed by Rho-specific guanine nucleotide exchange factors (RhoGEFs).

Contractions elicited by exogenous superoxide in rat aorta and pulmonary artery are blocked by inhibition of ROCK and are independent of dismutation to H_2O_2 [114, 311, 339, 351]. Sub-cellular translocation of ROCK, ROCK-mediated phosphorylation of MYPT1 and subsequent MLC_{20} phosphorylation, are all induced by exogenous superoxide [176, 311, 339, 351]. Although there is no evidence that Rho-kinase is directly ROS-sensitive, exogenous superoxide causes cytosolic to membrane translocation of RhoA in rat aorta and pulmonary artery [174, 176, 339]. This may be because RhoA is directly ROS sensitive, as it contains reactive cysteine residues that when oxidised

mimic the effect of RhoGEF binding, lowering the affinity for guanine nucleotides and triggering exchange of GDP for GTP [385, 386]. Alternatively, ROS may activate RhoA and ROCK upstream of RhoGEF activation, since superoxide induced contraction in rat pulmonary artery is suppressed by RhoGEF inhibition [174]. Such a pathway probably involves tyrosine phosphorylation of the RhoGEF by ROS-sensitive NRTKs. Src is a prime candidate, being directly ROS-sensitive and important for GPCR and superoxide-induced RhoGEF translocation, RhoA translocation and/or MYPT1 phosphorylation in rat afferent arterioles and pulmonary artery [106, 133, 174, 272, 311]. Src is also known to activate RhoGEFs indirectly via other tyrosine kinases FAK, PYK2 or JAK2 [145, 313, 314]. Once activated, RhoA also directly promotes actin polymerisation and, via ROCK and LIM kinase, inhibits the actin severing protein cofilin [387]. There are two main physiological contractile stimuli that may activate RhoGEFs and subsequently activate RhoA and ROCK via NOX-derived ROS. GPCR directly activate RhoGEFs via $G_{12/13}$, but also activate NOX1 or NOX2, and the superoxide produced will activate Src, PYK2 and JAK2 to promote RhoGEF activation [174, 295], as described above. Stretch induces integrin engagement with the extracellular matrix, FAK auto-phosphorylation and subsequent RhoGEF phosphorylation [270]. This 'outside-in' FAK activation is also associated with activation of NOX4/Poldip2 [229]. One study shows that NOX4/Poldip2-dependent H_2O_2 production induced by integrin engagement results in direct cysteine oxidation of filamentous actin and this promotes subsequent maturation of focal attachments [49]. Another study has shown that NOX4 derived ROS oxidise integrin $\alpha_7\beta_1$ and that this induces extracellular matrix binding [388] and potentially additional activation of FAK, suggesting a two-way relationship between integrin engagement and NOX4 activation which may strengthen the transmission of contractile force [389]. This response to stretch is of particular importance to the role of NOX in myogenic contraction (section 4.6). In line with the importance of RhoA/ROCK signalling in GPCR-induced VSM contraction and the role of NOX in activating ROCK, GPCR-induced translocation/activation of RhoGEFs/RhoA and/or MYPT1/MLC₂₀ phosphorylation in rat afferent arterioles, and mesenteric, tail and pulmonary arteries, are inhibited by non-specific antioxidants, SOD-mimetics or NOX blockers [106, 114, 133, 135, 174].

Unlike superoxide, some reports suggest that exogenous H_2O_2 induces contraction in a number of vascular beds that is insensitive to ROCK inhibition and is not associated with RhoA or ROCK translocation or phosphorylation of MYPT1, and is even independent of MLC₂₀ phosphorylation [311, 326, 343, 346, 347]. Indeed H_2O_2 actually suppresses basal MLC₂₀ phosphorylation at the main MLCK target site (Ser19) [326, 343]. Instead, in permeabilised, Ca^{2+} -clamped tissues H_2O_2 causes a contraction that is sensitive to inhibition of PKC [326, 343]. In parallel to ROCK-dependent MLCP inhibition via MYPT1, PKC, which is itself directly ROS sensitive [176, 321, 322], also inhibits MLCP through phosphorylation of the inhibitory protein, C-kinase potentiated protein phosphatase-1 Inhibitor-17 (CPI-17) [135, 382, 390]. In addition, there are two further routes through which ROS, with the involvement of PKC, could influence contractile force, and both involve actin polymerisation. Firstly, there is heat-shock protein-27 (HSP27) a chaperone which prevents actin polymerisation, unless phosphorylated and inhibited by p38 MAP-kinase via MAP-Kinase activated protein kinase [315]. Secondly, there is caldesmon, which inhibits actin-myosin interaction unless it is phosphorylated and inhibited by ERK MAP-kinase [391]. ERK and p38 MAP-kinases are both indirectly activated by ROS via

upstream PKC-mediated MEK phosphorylation [156, 194, 392]. MEK is also activated via oxidative inhibition of upstream inhibitory PTPs [268], oxidative activation of Ras [393] or Src-dependent RTK transactivation and recruitment of GRB2, Ras and Raf. In line with these signalling pathways being pro-contractile in VSM, constrictions induced by exogenous H₂O₂ or NOX-dependent GPCR agonists or stretch are inhibited by blockers of either p38 or ERK MAP-kinases [103, 194, 283, 341, 352-354, 394, 395]. ROS-dependent, Ca²⁺-independent contractile signalling pathways are summarised in Figure 5.

4.5. Role of NOX in GPCR-induced contraction. Having established the pro-contractile effects of ROS in VSM and the signalling pathways involved, this review now addresses the relevance of NOX-derived ROS to physiological vasoconstrictor stimuli, starting with GPCR agonists. These include agents involved in the control of mean arterial blood pressure and central control of regional blood flow (Ang II, noradrenaline) or local control of blood flow (ET-1, adenosine, 5-HT, prostaglandins, sphingolipids etc), all of which are known to activate NOX and stimulate ROS production (Table 1). Complementary to this, a variety of studies in isolated arteries from various vascular beds have utilised non-selective antioxidants, pharmacological manipulation of NOX and antioxidant enzymes, to examine the role of NOX in VSM contraction (Table 2). For example, Ang II in isolated aorta, mesenteric, cerebral and pulmonary arteries and renal afferent arterioles [89, 99, 104, 105, 133, 137, 325, 335, 377, 378], catecholamines in aorta, tail artery and renal arteries [131, 135, 137, 226], ET-1 in mesenteric and pulmonary arteries [66, 114], eicosanoids in mesenteric, coronary and pulmonary arteries [66, 140, 146, 148, 174], adenosine in aorta [116] and 5-HT in pulmonary artery [138], all cause constriction in part via NOX and ROS. A couple of these studies have also examined the role of NOX-derived ROS in renal haemodynamics *in vivo*. Infusion of an antioxidant or NOX inhibitor enhances baseline renal blood flow [137]. Such infusion also reverses the inhibition of renal blood flow by Ang II or noradrenaline and antagonises the concomitant increase in renal vascular resistance [133, 137]. Through the use of knockout mice or specific inhibitors, these responses have been attributed to NOX1 or NOX2 (p47^{phox}^{-/-}) [104], NOX1 alone (NOX1^{-/-}) [89, 138, 148], NOX2 alone (gp91ds-tat, NOX2^{-/-}) [133, 140], or Poldip2/NOX4 (Poldip2^{+/-}) [226]. The role of NOX in Ang II induced vasoconstriction is of particular importance to the study of NOX involvement in hypertension (section 6).

4.6. Role of NOX in myogenic, pressure/stretch-induced contraction. In a number of vascular beds, arterioles constrict in response to increased perfusion pressure (myogenic tone) in order to maintain adequate organ blood flow despite changes in MAP and prevent tissue damage through excessive flow. This is important in the cerebral [325, 396] and renal circulations [397], but has been studied most extensively in the renal afferent arteriole, where NOX-derived ROS play an important role. In the perfused pressurised afferent arteriole, endogenous superoxide production is increased by raising perfusion pressure [104, 230, 327, 334] (Table 1) and the response is suppressed in p47^{phox}^{-/-} mice, suggesting the involvement of NOX1 or NOX2 [104, 230] or in Poldip2^{+/-} mice suggesting the additional involvement of NOX4 [230] (Table 2). There is limited evidence that stretch activates NOX in other vascular beds. Myogenic tone is of marginal importance in coronary arteries, and pulmonary arteries do not generate force in response to stretch alone, but passive stretch does influence force generated in response to other stimuli. For example, depolarisation induced contraction in bovine pulmonary

and coronary arteries is enhanced by previous application of supra-physiological stretch and this enhancement appears to be dependent on NOX-derived ROS [194, 353].

Exactly how NOX1 or NOX2 are activated by stretch in VSM is unknown, but it is reasonable to speculate that a key step is the integrin-mediated activation of Src and FAK following stretch-induced binding of integrins with the extracellular matrix, the so-called 'outside-in' activation pathway, perhaps via interaction with transmembrane PTPs [398]. This may also involve subsequent activation of PKC. Indeed, one study in femoral artery suggests that stretch-induced ROS production requires PKC-dependent phosphorylation of p47^{phox} [162]. Like GPCR-induced responses, myogenic contraction involves both Ca²⁺ influx (primarily via LVGCC) and RhoA/Rho-kinase mediated Ca²⁺ sensitization [325, 327, 355, 399] and NOX-derived ROS will contribute to these, perhaps via stretch-induced Src- and PKC-dependent LVGCC phosphorylation [363-365] or RhoGEF phosphorylation [145, 313, 314]. An important role for NOX1 or NOX2-derived superoxide in myogenic contraction is suggested by the inhibitory effects of antioxidants, pharmacological NOX inhibitors and p47^{phox} knockout in isolated perfused renal afferent arterioles [99, 104, 230, 325, 334]. A pathway for NOX4 activation by integrin engagement which may contribute to myogenic tone can be inferred from the known association between Poldip2/NOX4, integrins, FAK and the actin cytoskeleton, and its role in integrin-induced VSM focal adhesion turnover [47, 49, 107, 228, 229] (Figure 5, and see also section 6.3 on the role of NOX in VSM migration). However, contrary to this and contrary to the proposed role for NOX4 in Ca²⁺-independent contraction via actin polymerisation, NOX4-derived H₂O₂ appears to be antagonistic of NOX1- or NOX2-derived superoxide and inhibitory of myogenic tone, rather than contributing to it. Note that afferent arteriole myogenic tone is also inhibited by over-expressing SOD and suppressed with PEG-SOD or enhanced in SOD^{-/-} mice, strongly suggesting that superoxide, not H₂O₂, is the specific ROS mediating this response [99, 230, 334]. Conversely, inhibiting NOX4-derived H₂O₂ with PEG-catalase or in Poldip2^{+/-} mice has the opposite effect, enhancing myogenic tone, indicating a pro-relaxant effect of H₂O₂ [230, 327]. Consistent with this, exogenous superoxide enhances myogenic tone in cerebral arteries and renal afferent arterioles [325, 327] while exogenous H₂O₂ has a mixed dose-dependent effect on myogenic tone in renal afferent arterioles, enhancing the response at low concentrations but causing relaxation at higher concentrations [334]. The pro-relaxant properties of ROS, and H₂O₂ in particular, are discussed further in the next section.

5. Role of NOX in VSM relaxation.

Exogenous superoxide, as generated extracellularly by xanthine/xanthine oxidase or intracellularly by LY83583 or menadione, relaxes rat aorta and mesenteric and femoral arteries pre-stimulated with vasoconstrictor GPCR agonists [311, 333, 351] and suppresses rat cerebral myogenic auto-regulation *in vivo* [396]. Superoxide may do this independently of its dismutation to H₂O₂ [311], but exogenous H₂O₂ also clearly causes relaxation in a range of systemic vascular preparations pre-stimulated with a range of vasoconstrictor GPCR agonists [336, 350, 354, 400-407] or pressure-induced myogenic tone [327, 334, 354, 408-411] and increases canine coronary blood flow *in vivo* [336] (Table 3). Thus, perhaps paradoxically, as well as mediating vasoconstriction, endogenous ROS also mediate vasodilation, often in the same vascular beds. This section discusses the evidence for a vasodilatory

action of NOX-derived ROS in response to normal physiological vasodilatory stimuli, focussing on the regulation of its production, its role in modulating endothelial function and its target pro-relaxant effector pathways in the smooth muscle.

5.1. Activation of Endothelial NOX. Physiological stimuli that trigger VSM relaxation and increase blood flow include vasodilatory GPCR agonists such as acetylcholine (from parasympathetic efferent nerves), histamine and bradykinin (local inflammatory mediators) and others, as well as flow/shear-stress. A response to shear-stress is important in some vascular beds for accommodating increased flow in circumstances where cardiac output and MAP are increased, such as muscle during exercise. Conversely, in tissues where relatively constant flow is desirable such as brain and kidney, flow-induced dilation balances and is balanced by pressure-induced myogenic contraction [230, 334, 412]. GPCR-induced and flow-induced relaxation are both endothelium-dependent and not wholly dependent on eNOS, therefore the ROS contributing to this relaxation are likely to be endothelial in origin and may act in parallel with or even influence eNOS and EDHF. As in VSM, NOX isoforms 1, 2, 4 and 5 are all expressed in the vascular endothelium [53, 54, 215, 234, 413-415] and are likely sources of the ROS contributing to or influencing these relaxation responses. Indeed, acetylcholine, histamine, bradykinin and flow all induce NOX-derived ROS production in vascular tissues and cultured endothelial cells [237, 402, 405, 408, 414, 416-418, 418-421] (Table 1). Vascular relaxation responses in isolated arteries and cerebral blood flow are improved by non-selective NOX inhibitors [421, 422], and several studies have examined the role of specific endothelial NOX isoforms in endothelium-dependent relaxation (Table 2). So far, the only unambiguous evidence for a pro-relaxant isoform is for NOX4. Femoral artery relaxation responses are suppressed in NOX4 knockout [52] while coronary perfusion pressure is depressed by over-expression of NOX4 [215]. Conversely, deletion of NOX1 appears to improve endothelium-dependent relaxation in isolated aorta or pressurised mesenteric arteries [423, 424] while expression of NOX5 in mouse vasculature appears to suppress it [66]. For NOX2, there is conflicting evidence. Some studies using selective inhibition of NOX2 or NOX2 knockout suggest that NOX2-derived superoxide antagonises endothelium-dependent relaxation [418, 425, 425], while other suggest that it supports endothelium-dependent relaxation in response to both flow and acetylcholine [418]. As discussed below, these differences may relate to the different pathways through which the four NOX isoforms respond to stimuli acting on the endothelium and the nature of their effects within the endothelium, but perhaps most significantly, the fact that NOX4 produces H₂O₂, not superoxide.

Activation of NOX1, NOX2 and NOX5 isoforms by GPCR agonists in endothelium is essentially the same as in VSM and has already been discussed, while the exact mechanism of NOX4 activation by GPCR in endothelium remains unclear (section 2). NOX activation by laminar flow/ shear-stress on the other hand is specific to the endothelium because only the endothelium is in direct contact with flowing blood. Endothelial cells possess a mechanical flow sensor or 'mechanosome', a protein complex composed of platelet-endothelial cell adhesion molecule-1 (PECAM-1) and the adapter protein VE-cadherin, associated with caveolin in caveolae. In response to changes in flow, this complex transactivates the type-2 VEGFR which then acts as a platform for the activation of PI3K [426], PKC- δ

[427], Src [428] and Rac1 [429], all of which contribute to the assembly and activation of NOX1 and NOX2 (see Figure 1 for more specific details). Secondly, VEGFR activation also triggers closure of K_{ATP} channels, similarly associated with caveolae [430, 431]. It is believed that the resultant endothelial depolarisation opens T-type Ca^{2+} channels, thus raising endothelial $[Ca^{2+}]_i$ [432, 433]. This initial increase in $[Ca^{2+}]_i$, as well as having downstream actions, may activate PKC- α or - β , further contributing to NOX1/2 activation. It may also directly activate NOX5, although this has not been demonstrated experimentally. In the endothelium, caveolin-1 appears to constitutively suppress NOX1 and NOX2 activity in the short term and suppress their expression in the long term [48, 234, 430, 431] and it is likely that HSP90 relieves this suppression by displacing caveolin-1 from the oxidase catalytic subunit [234], in a manner similar to that previously described for the activation of eNOS [434]. However, it should be noted that several earlier publications on the relationship between NOX and caveolin (not cited here) have since been retracted, so later studies should perhaps be interpreted with caution. NOX4 is also activated by sustained laminar shear-stress in endothelium, possibly explaining the high baseline levels of NOX4-derived endothelial H_2O_2 *in vivo* [291]. However, it is unclear how laminar shear-stress activates NOX4 since it is not located in caveolae [23] and the activation mechanism must differ from that described for VSM because integrins are not expressed on the apical surface of endothelium. Once activated, endothelial NOX generates H_2O_2 (NOX4) or superoxide (NOX1, 2 and 5) within the cell, where it contributes to the regulation of vascular tone through three main pathways: interaction with eNOS to regulate $NO\cdot$ production, both positively (H_2O_2) and negatively (superoxide), opening of endothelial K^+ channels as part of an endothelium-derived hyperpolarisation response, and diffusion (H_2O_2 only) to the VSM to cause relaxation directly, primarily via opening of VSM K^+ channels.

5.2. Cross-talk between NOX and eNOS in the endothelium. A key mediator of VSM relaxation is endothelium-derived $NO\cdot$, produced by eNOS through the transfer of an electron from NADPH to L-arginine in the presence of tetrahydrobiopterin (BH_4) [435]. The initial trigger for eNOS dimerization and activation is usually an increase in $[Ca^{2+}]_i$, either via sarcoplasmic reticulum Ca^{2+} release in response to GPCR agonist binding (eg acetylcholine, bradykinin and adenosine), or Ca^{2+} influx triggered by laminar shear-stress, as described above. $NO\cdot$ diffuses to the smooth muscle to cause relaxation, primarily via the activation of guanylate cyclase, production of cGMP and activation of protein kinase G (PKG), resulting in opening of K^+ channels. Classically, endothelial or smooth muscle derived superoxide is considered a major opponent of $NO\cdot$ mediated relaxation because it will chemically react with $NO\cdot$, generating peroxynitrite ($ONOO^-$) which is a relatively weak vasodilator [362, 436]. This is supported by the fact that applying or over-expressing SOD supports $NO\cdot$ mediated relaxation, while knocking out SOD has the opposite effect [99, 437]. Also, superoxide has the potential to trigger uncoupling of eNOS by oxidising BH_4 , thus forcing eNOS to make more superoxide instead of $NO\cdot$ [413, 435]. On the other hand, endogenous SOD normally dismutates much of the endothelial superoxide into H_2O_2 , which is pro-relaxant [402, 404, 410, 438], and $NO\cdot$ may indirectly support H_2O_2 production over that of superoxide through selective inhibitory S-nitrosylation of p47^{phox} (NOX1, 2) and of NOX5 [439, 440], but leaving NOX4-derived H_2O_2 unaffected [52, 440]. There is also evidence that endothelial ROS more directly promote eNOS activity and $NO\cdot$ production, with either NOX2-derived

superoxide [414, 418, 420] or NOX4-derived H₂O₂ [209, 291, 441], activating several protein kinases, including CAMK-II, p38 MAPK and FAK, either sequentially or in parallel, resulting in eNOS phosphorylation and further enhanced NO $\cdot$ production [209, 291, 442]. Interestingly, it is also apparent that sustained periods of physiological laminar pulsatile shear-stress also suppress gp91^{phox}/NOX2 expression, thus reducing basal superoxide production [414, 443]. It also upregulates eNOS expression [444], while at the same time it promotes NO $\cdot$ dependent down-regulation of NOX2 expression [420].

5.3. Role of NOX-derived ROS in endothelial hyperpolarisation. The second pathway through which ROS can promote endothelium-dependent relaxation from within the endothelium is by opening the intermediate and small conductance Ca²⁺-activated K⁺ channels (IK_{Ca}/SK_{Ca}) on the endothelium to trigger hyperpolarisation [405, 408], which then spreads to the VSM via myo-endothelial gap junctions [406, 445] or by the effluxed K⁺ diffusing to the VSM and activating VSM inward rectifier K⁺ channel (K_{IR}) and Na⁺/K⁺ATPase, which then hyperpolarises the VSM [446, 447]. There is little evidence that IK_{Ca} or SK_{Ca} are directly ROS-sensitive, but they are activated by an increase in endothelial [Ca²⁺]_i which can be induced by H₂O₂. Endothelial H₂O₂ may elevate [Ca²⁺]_i by promoting Ca²⁺ release from the sarcoplasmic reticulum or by triggering endothelial Ca²⁺ influx, and there is evidence for both. As in VSM, endothelial Ca²⁺ release occurs through a combination of IP₃R and RyR efflux pathways [448, 449] and, also as in VSM, these may be enhanced by H₂O₂ [225, 450]. This idea is supported by a study in isolated rabbit iliac artery, where exogenous H₂O₂ enhances the release of endothelial sarcoplasmic reticulum Ca²⁺ and relaxation triggered by pharmacological SERCA inhibition, and the enhanced relaxation is suppressed by blockers of IK_{Ca} / SK_{Ca} and myo-endothelial gap junctions [406]. Alternatively, the endothelial transient receptor potential ankyrin-1 (TRPA1) is also directly ROS-sensitive [451, 452]. TRPA1 has a large conductance and may itself facilitate sufficient endothelial Ca²⁺ influx to activate co-localised SK_{Ca} and IK_{Ca} on the endothelial basolateral surface [453-456]. One additional study provides evidence supporting a role for NOX-derived H₂O₂ in endothelial hyperpolarisation. Here, enhanced ACh-induced relaxation induced by endothelium-selective over-expression of NOX4 in isolated aorta was normalised by catalase, KCl and blockers of SK_{Ca} and IK_{Ca}, suggesting that the extra H₂O₂ generated was acting within the endothelium to promote endothelial hyper-polarisation [215]. Therefore, H₂O₂ may play an accessory role in endothelium-dependent relaxation as a promotor of endothelial-VSM electrical coupling.

5.4. NOX-derived H₂O₂ as a diffusible hyperpolarising factor. EDHF was originally defined to describe an unknown factor or factors that may be responsible for the component of endothelium-dependent hyperpolarising vasodilation not due to NO $\cdot$ or prostacyclin [457, 458]. We now know that EDHF probably comprises no single factor but a range of diffusible factors or processes coupling endothelial hyperpolarisation to smooth muscle hyperpolarisation and relaxation [446, 459]. In addition to contributing to endothelial hyperpolarisation and electrical coupling between endothelium as described above, H₂O₂ may itself be one of these proposed factors that diffuses to the VSM and causes hyperpolarisation [410]. Flow/shear-stress and vasodilatory GPCR agonists such as acetylcholine, bradykinin, histamine, substance P and adenosine induce hyperpolarisation and dilation of a wide

range of isolated resistance-sized systemic arteries/arterioles from various mammalian species that is suppressed when SOD is inhibited or knocked out or when catalase is applied, and is restored by applying SOD or SOD mimetics [52, 102, 117, 139, 215, 400-402, 404-406, 408, 410, 411, 418, 421, 425, 437, 438, 460-463]. Thus, it is presumably H_2O_2 not superoxide, that plays the major role in relaxation responses to physiological vasodilator stimuli, even though exogenous superoxide is also capable of relaxing VSM. This may be partly because superoxide scavenges $NO\cdot$ but H_2O_2 does not, and partly because H_2O_2 is more freely diffusible and membrane permeable than superoxide. In many of the studies implicating H_2O_2 as an EDHF, the principle source of endothelial H_2O_2 is not clearly defined, although a role for NOX can often be inferred. Two studies have directly linked NOX to EDHF, suggesting that either inhibition of NOX2 or knockout of NOX4 will remove the PEG-catalase inhibitable component of adenosine-induced relaxation of isolated rat cerebral arterioles [117] or of flow-induced femoral dilation *in vivo* [52]. One alternative source is uncoupled eNOS [402, 461].

5.5. Pro-relaxant targets for ROS in VSM. In arteries pre-stimulated with constrictor GPCR agonist or myogenic tone, the EDHF component of relaxation induced by vasodilatory GPCR agonists or flow after blockade of eNOS or prostacyclin production, is inhibited by a depolarising KCl solution to remove the influence of K^+ channels on VSM membrane potential [402, 405, 411], suggesting that EDHF ultimately relaxes the VSM by opening K^+ channels, triggering hyperpolarisation, closing LVGCC and lowering $[Ca^{2+}]_i$. In similarly pre-constricted vessels, the relaxation induced by either H_2O_2 or superoxide is also inhibited, or even turned into a contractile response by KCl, supporting the hypothesis that H_2O_2 is an EDHF [148, 336, 341, 342, 350-352, 354, 402, 405, 407, 408, 411].

VSM express a range of K^+ channels including, most notably, a number of different K_v channels and the large conductance Ca^{2+} activated K^+ channel (BK_{Ca}). However, there appears to be conflicting evidence over which of these two types of channel are the most important in mediating VSM hyperpolarisation responses to physiological vasodilatory stimuli. Certainly there is conflicting evidence over which is important in mediating the hyperpolarising and pro-relaxant effects of exogenous H_2O_2 or superoxide. Some studies show that superoxide or H_2O_2 -induced relaxations of GPCR-mediated or myogenic contractions in coronary and mesenteric arteries or renal afferent arterioles are prevented by selective pharmacological inhibition of K_v channels (eg. 4-aminopyridine, 4-AP, linopirdine), but not of BK_{Ca} channels (eg. TEA, iberiotoxin) [327, 336, 350, 351, 366, 407, 464], or that H_2O_2 enhances coronary blood flow in dogs *in vivo* and this is blocked by 4-AP [336]. In line with this, at the single cell or single channel level, a few studies show that exogenous superoxide (10 μ M LY83583) and somewhat high concentrations of H_2O_2 (1-10mM) both activate 4-AP sensitive K_v currents in isolated coronary and mesenteric VSMCs [350, 351, 366, 407]. K_v channels expressed in VSM may be directly ROS sensitive since the current enhancement by H_2O_2 in coronary arterial VSMCs is prevented by thiol alkylation [366] and $K_v2.1$ is modified by direct S-glutathionylation (oxidised glutathione) in mesenteric artery VSMCs to increase its voltage sensitivity [407], or it may be indirect via MAP-kinase dependent phosphorylation [409].

Curiously, despite all the evidence described above that K_V channels are preferentially activated by ROS and contribute to ROS mediated relaxation, when examining hyperpolarisation and VSM relaxation induced by physiological vasodilatory stimuli acting via endogenous H_2O_2 , the majority of evidence points towards BK_{Ca} as the likeliest target channel because acetylcholine and flow both induced relaxation in coronary or mesenteric arteries that is suppressed by inhibitors of BK_{Ca} channels (TEA, TBA, iberiotoxin, paxilline) [402, 403, 410, 411, 460, 462, 462]. This is also supported by additional studies showing that exogenous superoxide (X/XO) and physiological concentrations of H_2O_2 (1-100 μ M), as well as perfusate from mesenteric arteries containing endogenous H_2O_2 , activate single channel BK_{Ca} currents in canine cerebral artery or human coronary artery VSMCs [396, 403, 410, 460]. Unlike for K_V channels, there is limited evidence for direct oxidative activation of BK_{Ca} . Although the human BK_{Ca} channel reportedly possesses a disulphide switch in non-vascular cells [465], there is no evidence to support its direct oxidative modification in vascular cells, indeed physiological concentrations of exogenous H_2O_2 (<100 μ M) only effectively activates BK_{Ca} in cell-attached patch-clamp configuration, indicating the requirement of intracellular signalling [403, 410]. It is more likely that it requires oxidative activation of kinases that phosphorylate and open these channels, particularly $PKGI\alpha$, either directly (independently of cGMP) [327, 400, 401, 403, 466], or via oxidative activation of guanylate cyclase which generates cGMP, the main activator of PKG [467, 468]. In addition to opening K^+ channels, there is some evidence that under certain circumstances, ROS and/or $NO\cdot$ may promote relaxation by inhibiting L-type Ca^{2+} channels through direct thiol sulfonylation or S-nitrosylation [359-362]. However, it is unclear how this mechanism can operate and be regulated alongside the better characterised indirect opening of L-type Ca^{2+} channels by ROS-dependent phosphorylation or membrane depolarisation (section 4.2). Finally, $[Ca^{2+}]_i$ may be lowered to promote relaxation through glutathionylation and activation of VSM SERCA [380]. All of the interactions between endothelial superoxide, H_2O_2 and $NO\cdot$, their contribution to EDHF and VSM relaxation and the effector pathways are summarised in Figure 6.

A big question remains though: how ROS can have opposing effects, contraction and relaxation, in the same tissue or even the same cell within that tissue, and how the final net response is obtained. This presumably relates to the relative sensitivity of the various ROS effector pathways to ROS and therefore the concentration of ROS applied, but also to the relative expression levels of components of those pro-contractile and pro-relaxant pathways and their possible co-localisation with NOX in the endothelium and VSM. Differential effects of superoxide vs. H_2O_2 within the cell may also be important, and therefore the relative activities and distributions of SOD and catalase. Clearly, the nature of the pre-constriction influences the outcome, not only by altering the balance between LVGCC and K^+ channel activity but also the contributions of VSM sarcoplasmic reticulum Ca^{2+} release and Ca^{2+} sensitization pathways, both of which are also ROS sensitive but minimally influenced by membrane potential or $NO\cdot$.

6. Role of NOX-derived ROS in hypertension.

Hypertension is a persistent increase in baseline MAP, which if uncorrected will result in cardiac remodelling and ultimate heart failure. There are various causes, including over-activation of the

renin-angiotensin-aldosterone system and sympathetic nervous system, various genetic factors and environmental/life-style factors, such as a high salt diet, excess body weight and smoking [469-472]. Regardless of cause, disease progression involves several underlying mechanisms, including inflammation, smooth muscle hyper-contraction coupled with impaired endothelial function, smooth muscle proliferation and migration. A common feature of all these processes and the resultant hypertension is oxidative stress, as indicated by elevated levels of malondialdehyde and F2 isoprostanes, resulting from excessive ROS production, reduced NO $\cdot$ availability and impaired antioxidant defences [113, 415, 470, 472-478]. A clear causative link between oxidative stress and hypertension remains a matter of debate, especially in humans where there also remains uncertainty over the therapeutic benefit of antioxidants [469, 470, 472, 474, 476, 477, 479-482]. However, such a link is well characterised in multiple animal models of hypertension, such as the spontaneously hypertensive rat (SHR), renovascular hypertension, deoxycorticosterone acetate (DOCA) salt rat, obese rat and the Ang II-infused rat or mouse. These have been used to examine the mechanisms underlying oxidative stress and its role in the pathophysiology of hypertension. Throughout earlier sections of this review are detailed the various cellular target signalling pathways for NOX-derived ROS that contribute to the normal functioning of the vasculature, both pro-contractile and pro-relaxant. In this section will be outlined evidence for how these pathways are altered in oxidative stress to promote a hyper-contraction state, how excess ROS induce vascular remodelling through stimulation of VSM growth, proliferation and migration and fibrosis, and briefly, the role of NOX/ROS in vascular inflammation.

6.1. Expression of NOX isoforms/subunits in hypertension. Individually, NOX isoforms may contribute to VSM hyper-contraction or remodelling in response to hypertensive stimuli through altered activity, expression or both [80]. NOX1 and 2 are both upregulated in SHR compared to Wistar-Kyoto control (WKY) rats [415, 483-485]. The regulation of subunit NOX expression by long-term treatment of vascular cells/tissues with mitogenic GPCR agonists and growth factors/cytokines has been reviewed extensively elsewhere [23, 474, 486]. Examples from rat or mouse models include expression of NOX1, NOX2, p22^{phox}, p47^{phox} and p67^{phox} in VSM being enhanced by Ang II infusion [26, 40, 45, 82, 423, 423, 473, 483, 487-491], NOX1 being enhanced by PDGF [40], NOX1 by stretch [492] and NOXA1 by TNF- α [126]. There is some disagreement over whether NOX4 is also upregulated in SHR, with several studies saying it is [70, 483, 485, 493, 494] and some saying it is not [415]. Regulation of NOX4 expression by mitogenic and inflammatory stimuli in hypertension is also contentious. Several reports suggest NOX4 is upregulated by GPCR agonists and inflammatory mediators in VSM [47, 120, 263, 423, 489, 494, 495] while a few others suggest it is down-regulated by Ang II and PDGF [40, 45, 473]. However, a recent study suggests that, either in SHR or during Ang II infusion, expression of NOX4 in VSM is enhanced, and this occurs by the suppression of B-cell lymphoma 6 (BCL6) [494]. BCL6 itself normally suppresses NOX4 expression, reduces oxidative stress and impairs aortic wall thickening [494]. NOX5 expression is enhanced by Ang II, ET-1 and inflammatory mediators such as TNF- α in human VSMCs *in vitro* [57, 496]. It is not clear whether native NOX5 expression is altered in VSM in hypertensive humans but it is upregulated in renal proximal tubule cells [497]. There is limited evidence that NOX polymorphisms are associated with hypertension risk in humans [470]. p22^{phox} polymorphisms show

either a protective association or an increased risk [498] and seven identified polymorphisms in NOX5 result in reduced or absent enzyme activity [499]. See Figure 7 for a summary of all the factors influencing NOX expression in hypertension.

6.2. Impaired NO--dependent relaxation and enhanced contractility. In SHR, amplitude of phenylephrine-induced contraction in isolated aorta is enhanced while acetylcholine induced relaxation is impaired compared to WKY controls and these differences are normalised by antioxidants and NOX inhibitors (tiron, tempol, apocynin, VAS2870) or catalase [415, 423, 425, 489]. If there is an imbalance between ROS production and antioxidant defences, then excess superoxide in the vasculature will have two immediate effects. Firstly, it will react with NO· in an uncontrolled way, resulting in the production of excess ONOO⁻ and secondly it will trigger uncoupling of eNOS through oxidation of BH₄, forcing eNOS to make superoxide instead of NO·. Excessive ROS, particularly hydroxyl radical, may also directly inactivate guanylate cyclase [484, 500]. Hypertensive agents such as Ang II and TXA₂ promote eNOS uncoupling [5, 6, 66, 414, 422, 435, 501]. The degree to which this eNOS uncoupling impairs agonist-induced relaxation will also depend on how much of the superoxide produced is dismutated to H₂O₂ in the endothelium, and this will depend on the expression and location of SOD, which may or may not be down-regulated [99, 422]. The net effect will be impaired NO· mediated relaxation via K⁺ channel closure and VSM depolarisation [425]. However, these pathways do not solely account for the hyper-contractile phenotype. Hypertension is also associated with a dysregulation of intracellular Ca²⁺ in the VSM [70, 502-504] and this is influenced by VSM oxidant stress [13, 70, 376]. There is evidence for both enhanced Ca²⁺ influx through LVGCC [503-507] and altered sarcoplasmic reticulum Ca²⁺ handling [506, 508, 509], but it perhaps remains to be clarified which of all the various ROS-dependent Ca²⁺ signalling pathways described in this review are predominantly upregulated in hypertension. For example, one study shows in mesenteric arteries from SHR, that H₂O₂ induces a greater LVGCC-mediated Ca²⁺ influx than in WKY control rats [348], while others suggests that one consequence of oxidative stress on [Ca²⁺]_i is that excessive cytosolic superoxide results in irreversible inhibition of smooth muscle SERCA, preventing the normal pro-relaxant, NO--mediated S-glutathionylation of the ATPase [380] (section 5.5).

Hypertension is also associated with an increased activity of the RhoA/Rho-kinase signalling pathway and this also mediates both enhanced VSM contractile responses and VSM proliferation/migration [510-514]. As discussed earlier in this review, RhoA is both directly [385, 386] and indirectly ROS sensitive, so this enhanced activity could be NOX-dependent, perhaps via an upstream ROS-sensitive tyrosine kinase. For example, an Ang II-induced increase in MAP and aortic remodelling is suppressed in smooth muscle JAK2 knockout mice and part of the effect is ROS-dependent [192]. In carotid artery from SHR, enhanced agonist-induced contraction is inhibited by antioxidants and NOX inhibition while exogenous H₂O₂ causes a greater degree of Rho-kinase-dependent contraction than in WKY rats [515]. Antioxidants such as the putative SOD mimetic tempol and vitamins C and E lower blood pressure in various rodent models of hypertension [133, 516-520]. This effect is also accompanied by corrected endothelial dysfunction, particularly improved NO· activity and reduced agonist evoked ROS production, and normalised blood vessel contractility and vascular resistance [519]. SOD itself has a

similar restorative effect on endothelium-dependent relaxation [102, 425] and blood pressure [488], while knocking out SOD1 enhances the transient phase of Ang II-induced hypertension in mice, suggesting a specific role for superoxide in mediating the initial pressor response [99]. Taken as a whole, these data suggest that a key feature of hypertension is indeed a shift in the balance between pro-contractile and pro-relaxant effects of ROS contributing towards a hyper-contractile phenotype (Figure 7).

6.3. Vascular remodelling in hypertension. Vascular remodelling, resulting in thickening of the smooth muscle, narrowing of the lumen and increased vessel stiffness, is a key feature of hypertension, either as a primary cause of elevated MAP or as a secondary response to prolonged periods of elevated perfusion pressure. Either way, ROS are essential components of the signalling pathways involved in this process (Figures 7 & 8). For VSM proliferation, various stimuli and proximal ROS-sensitive kinases, including Src, CAMK-II and PKC, converge on ERK MAP-kinase via growth-factor transactivation and Ras [521], or via phosphorylation of Raf and/or MEK [522, 523] or via PYK2/JAK2 [524]. Once phosphorylated, ERK translocates to the nucleus to induce gene expression and cell division and ROCK may supporting this translocation [525]. In addition, there is the JAK/STAT pathway, a common route for inflammatory responses and gene expression. JAK2 is principally induced by cytokine receptor activation, but it is also activated by GPCRs to phosphorylate signal transducer and activator of transcription (STAT), triggering its translocation to the nucleus to regulate gene expression. JAK2 also stimulates ERK phosphorylation [526], possibly downstream of ROS-activated Src family kinases [188, 527], PKC- δ [170] or PYK2 [524]. PYK2 is activated by Ca^{2+} , often in response to induced sarcoplasmic reticulum Ca^{2+} release, triggered by IP_3 generated by PLC- β (GPCR coupled) or PLC- γ (RTK-coupled). p38 MAP-kinase is also activated by MEK in response to various mitogenic GPCR or RTK stimuli via NOX-derived ROS or by TNF- α via apoptosis signal-regulating kinase-1 (ASK-1) in VSM endosomes. The latter requires NOX1-dependent oxidative activation of ASK-1 [248, 252]. Finally, PI3K is a key stimulator of cell growth and cell survival, acting via AKT, mammalian target of rapamycin (mTOR), S6-kinase and eukaryotic translation initiation factor 4E-binding protein (4EBP) [528].

Cell migration requires a dynamic cell polarisation involving co-ordinated cycles of actin depolymerisation and polymerisation at the trailing and leading edges. This is dependent on activation of small G-proteins RhoA, Rac and cdc42, as well as FAK, interaction with ILK and cross-talk with extracellular matrix [107, 529]. ROS-activated CAMK-II may facilitate VSMC migration by activating the matrix metalloprotease MMP9 [328, 530]. At the leading edge of the migrating cell, actin depolymerisation is required for the formation of new actin fibres and this is achieved through the ROS-dependent de-phosphorylation and activation of cofilin by the PTP slingshot-1L following NOX4-dependent oxidation of 14-3-3 [531, 532]. Rho and Rac are often activated by migratory stimuli via tyrosine phosphorylation of their respective guanine nucleotide exchange factors, by Src, FAK, PYK2 or JAK2 and once activated they collectively promote actin reorganisation [145, 189, 190, 192, 313, 314]. Poldip2 may facilitate activation of RhoA and focal adhesions via NOX4-derived ROS activating associated RhoGEFs [247], perhaps via ROS-induced tyrosine phosphorylation. CAMK-II also promoted

migration via Rac [533]. As well as responding to inflammatory mediators [534], VSM NOX-derived ROS also positively influence vascular inflammation in hypertension. One study suggests that vascular oxidant stress is sensed by resident dendritic cells through the formation of isoketal protein adducts, leading to T-cell recruitment and activation and the release of cytokines that promote fibrosis and arterial stiffening [535].

FAK is a common point for cross-talk between GPCRs, RTKs and integrins. The 'inside-out' FAK activation pathway involves GPCR mediated NOX1 or NOX2-dependent activation of FAK, probably via Src [314], subsequent dissociation of ILK from clustered integrins, paxillin phosphorylation and RhoA activation, promoting migration through actin polymerisation and focal adhesion turnover at the leading edge [107, 529, 536-538]. There may then be secondary further integrin-dependent activation of NOX, possibly NOX4, following GPCR-induced FAK-activation, involving integrin $\alpha 1\beta 1$ [107]. FAK may also promote VSM proliferation and migration in response to enhanced perfusion pressure, because it will be activated by 'outside-in' stretch-induced integrin engagement [539]. FAK is also coupled to NOX4/Poldip2 activation and in combination they are capable of subsequently activating both Src and RhoA [49, 226, 229, 296, 539-541] and acts as an alternative platform for GRB2 docking and Ras/Raf/MEK/ERK signalling and PI3K/AKT activation, thus promoting both proliferation and migration of VSMCs in response to elevated pressure [107, 529, 539, 541]. Activation of FAK by stretch may also be ROS-dependent considering that NOX4 is located at focal attachments and potentially influences integrin-dependent signalling by oxidising integrin $\alpha 7\beta 1$ [49, 388]. FAK activation by NOX4 at focal adhesions probably involves oxidative inhibition of FAK phosphatases LMW-PTP or SHP2 [189, 542, 543]. Phosphorylation of Src, FAK and PYK2 are all increased in VSM of SHR [485, 544], while Ang II-induced ERK phosphorylation and impairment of vascular function are normalised by Src inhibition [101, 544].

In VSMCs, exogenous H₂O₂ induces transactivation of EGFR, PDGFR or IGFR, GRB2/Ras/Raf assembly and subsequent ERK phosphorylation [103, 173, 267, 274, 309, 545] and AKT activation [309]. Phosphorylation of EGFR, PDGFR and IGFR are also all enhanced in VSMC of SHR [485]. Evidence supports a role for Src or Fyn, PKC- δ , PYK2 and JAK2 in mediating these effects [173, 527, 546] and Src auto-phosphorylation is also enhanced in VSMC of SHR [485]. Knocking out SOD has a similar effect to that of exogenous ROS, promoting ERK and JAK/STAT phosphorylation and proliferation [547]. In addition to the proliferative, hypertrophic and migratory pathways mediated by ERK, JAK/STAT, PI3K/AKT and FAK/RhoA another important element of VSM hypertensive remodelling is nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B). NF- κ B is normally involved in inflammatory and cell survival responses to cellular stress such as cytokines and ROS [255, 534, 548] but becomes over-activated under oxidative stress, and contributes to the stimulation of VSM proliferation. NF- κ B is activated by these stimuli through the degradation of an inhibitory I- κ B protein and Src may contribute to ROS-dependent NF- κ B activation in VSM by phosphorylating I- κ B alpha [259, 549]. Alternatively, NF- κ B may be activated via p38 MAP-kinase downstream of TNF- α receptor-1 stimulation via TRAF2 and ASK-1 [252, 550]. The NF- κ B signalling pathway is opposed by nuclear factor erythroid 2-related factor 2 (Nrf2) which plays a cyto-protective role under oxidative stress by

inducing expression of anti-oxidant enzymes [548]. Nrf2 translocation to the nucleus is promoted by oxidative inhibition of KEAP-1 which degrades cytosolic Nrf2 [551].

In vivo, one would expect an early key response to oxidative stress to be excessive oxidative inhibition of PTPs and over-stimulation of NRTKs resulting in an overall enhancement of tyrosine phosphorylation, which would have consequences for both contractility and VSM growth/proliferation and migration [21]. One recent study suggests that coupled with ER stress, oxidative stress may induce VSM proliferation through irreversible hyper-oxidation (sulfinic and sulfonic acid) of PTPs [70]. Antioxidants also suppress many of the above described signalling pathways and processes associated with hypertension, including NRTK phosphorylation [110, 538, 546], growth factor receptor transactivation [103, 121, 136, 147, 149, 194, 274], AKT activation [109, 136, 150], ERK phosphorylation [64, 103, 109, 110, 136, 147, 149, 156, 194, 267, 274] and VSM hypertrophy [96, 552], proliferation [118, 121, 187, 267, 317, 485] and migration [100, 222, 229, 267, 538], in response to a range of GPCR agonists, growth factors and stretch (Table 4, Figures 7 & 8). A few studies highlight the importance of Src in hypertension, both as an upstream regulator of NOX and as a proximal ROS-sensitive kinase important in contraction and proliferation, by showing that oxidative stress and impaired cardiovascular function induced by Ang II infusion or in SHR vs. WKY rats are all improved by treatment with a Src inhibitor [82, 101, 485].

In animal models of hypertension, or in VSMC treated with hypertensive agents, the broad spectrum NOX/flavoprotein inhibitors diphenyleneiodonium (DPI) and apocynin inhibit growth-factor transactivation [103, 121, 136, 147, 149, 194, 316], ERK phosphorylation [64, 103, 109, 110, 121, 136, 147, 149, 156, 194, 316], AKT activation [109, 126], proliferation [79, 80, 107, 118, 121, 126, 127], hypertrophy [79, 80, 96, 109, 136, 147] and migration [100, 147, 178, 222]. Apocynin also normalises blood pressure and vasoconstrictor responses in spontaneously hypertensive rats [553, 554], DOCA salt rat [555] and dopamine D2 deficient mice [556]. A more recently developed inhibitor, VAS2870 also normalises MAP and improves acetylcholine-induced relaxation in isolated aorta [415]. While originally thought to be NOX-specific, DPI, apocynin and VAS2870 are considered as broad spectrum flavoprotein inhibitors, potentially acting on other sources of ROS or as direct antioxidants [474]. However, a new class of dual specific NOX inhibitors has been developed, including the NOX1/4 dual inhibitor GKT137831. This may be effective against hypertension induced intermittent hypoxia [557] and prevents hypertensive cardiac remodelling in rats [558], but does not significantly lower MAP in rats with hypertension induced by aortic banding [558]. GKT137831 has been deemed safe for human use and has progressed to clinical trials as a NOX blocker, but not yet for hypertension [474, 559]. Thus, the role for specific NOX isoforms in oxidative stress and its effects on the vascular wall in hypertension remains an avenue of active research.

6.4. Effects of selective NOX inhibition or over-expression in experimental hypertension. Several studies have examined the effects of knockout, siRNA depletion or over-expression of NOX isoforms or regulatory subunits on blood pressure, associated functional abnormalities and signalling in various mouse or rat models of hypertension. Ang II-induced hypertension and associated increases in ROS

production, ERK phosphorylation, proliferation and migration are blunted in p47^{phox}^{-/-} mice [41, 72, 72, 104, 166] or by p22^{phox} siRNA or antisense [61, 157, 560], while the hypertrophic response is enhanced by over-expressing p22^{phox} [552]. p22^{phox}^{-/-} or p22^{phox} siRNA also reduce MAP in salt-loaded stroke-prone SHR [561] or Ang II-induced hypertension [61, 157]. Individually, there are multiple, but often conflicting reports over which NOX isoforms, if any, predominate in experimental hypertension and associated signalling.

There is disagreement over whether or not NOX1 contributes to baseline MAP in Ang II-infused mice [423, 490] and disagreement over whether or not the sustained phase of Ang II-induced hypertension is suppressed in NOX1^{-/-} mice, with at least two saying it is [423, 490], and at least two saying it isn't [277, 562]. However, one study suggests that over-expressing NOX1 specifically in the VSM enhances Ang II-induced hypertension [520]. The Ang II pressor response in SHR is also blunted by NOX1 siRNA [483]. *In vitro*, the impaired Ach-induced relaxation in arteries from Ang II-induced hypertensive mice is restored by knocking out NOX1 [423]. There is also disagreement over whether NOX1^{-/-} suppresses *in vivo* aortic wall thickening in Ang II-induced hypertension, one showing a reduction [490] and one showing no effect [423]. While *in vitro*, Ang II-induced aortic VSMC proliferation is inhibited by NOX1 siRNA [107], GPCR-induced transactivation of EGFR in human and mouse VSMCs is NOX1-dependent [125, 178], Ang II-induced aortic VSM hypertrophy is inhibited by NOX1 siRNA [89] and over-expressing NOX1 enhances this Ang II-induced hypertrophy [520]. There is some evidence that NOX1 is important in VSM migration. The increased activation of Src and FAK in kidneys of renal hypertensive mice is prevented by knocking out NOX1 [277], migration of human coronary artery VSMCs is NOX1-dependent [178] and Ang II-induced aortic VSMC FAK phosphorylation, ILK activation and migration are all inhibited by NOX1 siRNA [107].

There is also disagreement over whether NOX2 is an essential mediator of Ang II-induced hypertension in mice. Several studies suggest that NOX2 knockout lowers basal MAP and the acute phase of Ang II-induced hypertension, but not the sustained increase in MAP [277, 562, 563], while others show that the NOX2 inhibitor gp91ds-tat effectively suppresses Ang II-induced hypertension [98, 133]. In SHR, the acute Ang II-induced pressor response and myogenic contraction responses in renal afferent arterioles are blunted by NOX2 siRNA [483, 564]. There is also some evidence that NOX2-derived ROS negatively regulates endothelium-dependent relaxation. Ach-induced vasodilation is impaired in isolated aorta from mice with renovascular hypertension but not in NOX2^{-/-} mice or in aorta from hypertensive mice treated with gp91ds-tat [425], while over-expressing endothelial NOX2 in mice potentiates the hypertensive effects of Ang II infusion without changing basal MAP [565]. An inhibitory role for NOX2 in the control of endothelium-dependent relaxation is supported by the observation that NO-dependent, flow-induced dilation is enhanced in humans with hereditary NOX2 deficiency [501]. NOX2 knockout also prevents Ang II infusion-induced aortic wall thickening [563] and NOX2 siRNA inhibits Ang II-induced aortic VSMC proliferation *in vitro* [107].

There are two opposing schools of thought about the role of NOX4 in the pathophysiology of hypertension, perhaps because it only generates H₂O₂ [45, 212-215], which is generally pro-relaxant

in the normotensive resistance vasculature, contributing to normal endothelial function [52, 209, 215, 291, 327, 422, 441, 444], but on the other hand its expression is upregulated in experimental hypertension [70, 423, 493] and by inflammatory mediators (eg TNF- α) in human aortic endothelium [263]. One theory is that like other NOX isoforms, NOX4 makes a direct contribution to oxidative stress and therefore to the pathogenesis of hypertension. Two studies show that NOX4^{-/-} or poldip2^{+/-} suppress Ang II infusion-induced increases in pulse pressure or MAP in mice [226, 566] while having no effect on baseline MAP or cardiac remodelling [566]. Yet another study suggests that NOX4 does contribute to pressure overload-induced cardiac dysfunction [264]. NOX4 may be pro-contractile in hypertension, as suggested by a study on the hypertensive effects of chronic intermittent hypoxia in rats. This was associated with an increased VSM expression of RhoA and Rho-kinase which was normalised by the selective NOX1/4 inhibitor GKT137831 [557]. Whether this increased expression of RhoA and Rho-kinase equates to an enhanced Ca²⁺ sensitizing contractile response to vasoconstrictor agents via MLCP inhibition, thus opposing the normal pro-relaxant effects of NOX4-derived H₂O₂, remains to be determined. A number of studies support a role for NOX4 in hypertensive vascular remodelling. Inhibiting NOX1/4 normalises mesenteric VSMC proliferation in SHR [70] and NOX4 appears to be the primary NOX isoform required for IGF-induced VSMC migration [222]. NOX4 may support proliferation via oxidative inhibition of SHP2 in association with PDGFR activation [258] and most likely supports VSMC migration in association with Poldip2, upstream of RhoA/FAK/integrin-mediated focal adhesion turnover and stress fiber formation [107, 228, 529, 536, 537].

The other theory is that upregulation of NOX4 in hypertension may be protective, particularly in the endothelium. Selectively over-expressing endothelial NOX4 reduces baseline MAP and prevents the initial transient phase of the hypertensive response to Ang II infusion but does not influence aortic wall thickness [215]. NOX4 is also upregulated by pathological oscillatory shear-stress or chronic flow overload in endothelium, but to a greater extent than is NOX2, thus potentially shifting the balance in ROS species away from pro-contractile superoxide towards pro-relaxant H₂O₂ [215, 327, 422, 444]. This is supported by the possible blood pressure lowering effect of VSM PKG activation by H₂O₂, probably derived from endothelial NOX4 [400]. In contrast to the endothelium-selective knockout described in [215], total NOX4 knockout has no effect on baseline MAP or the Ang II-infusion induced increase in MAP [52], but it enhances Ang II-induced aortic hypertrophy, impairs ACh-induced relaxation and potentiates Ang II induced expression of inflammatory cytokines, strongly suggesting a protective anti-hypertensive role for NOX4-derived H₂O₂ [52]. It is also suggested that NOX4-derived H₂O₂ is important in maintaining VSM differentiated state by promoting expression of contractile proteins and inhibiting proliferation, as opposed to NOX1 which is pro-proliferative [47, 567]. NOX4 may be anti-hypertensive and cardio-protective in part via induction of Nrf2 activity in VSM and heart [52, 568]. The idea of this protective NOX4/ Nrf2 axis is supported by the additional observation that prolonged superoxide generation by NOX1/2/5 (but not NOX4) generally results in activation of the pro-inflammatory and pro-proliferative transcription regulator NF κ B [534, 548] (Figure 7). Importantly, if this protective effect of NOX4 holds true for human hypertension it may suggest potential detrimental side-effects or even exacerbation of symptoms with antioxidant therapy as a treatment for the disease.

NOX5 appears to be pro-proliferative in human VSMC and endothelium, acting via activation of MAP-kinases and JAK2 [57, 317]. However, a thorough analysis of a potential role for NOX5 in hypertension is hampered the fact that rodents do not possess this NOX isoform. A few recent studies have addressed this by examining the effects of artificially expressing human NOX5 in mice, often in specific cell types. Expressing NOX5 in mouse VSM enhances basal superoxide production, enhances U46619 and ET-1 induced contraction and suppresses ACh-mediated relaxation in isolated mesenteric artery, but curiously does not significantly affect mesenteric artery media/lumen ratio or vessel stiffness or Ang II infusion-induced hypertension [66]. Thus VSM NOX5 may be involved in the control of tissue blood flow but not in the control of MAP. However, a number of recent studies suggest that NOX5 may be important in human hypertension indirectly by contributing to kidney dysfunction. When expressed specifically in renal podocytes of mice it impairs glomerular barrier function and raises systolic blood pressure [569] and when expressed in mouse smooth muscle and mesangial cells it accelerates the development of diabetic nephropathy [570]. Thirdly, a study in human renal proximal tubule cells shows that in tubule cells from patients with essential hypertension NOX5 expression was several fold higher while NOX1, NOX2 and NOX4 were unaffected [497]. Furthermore, cells from hypertensive patients exhibited reduced sensitivity to dopaminergic agonists, correlating with impaired suppression of tubular ROS production and sodium transport, implicating renal NOX5-derived ROS in the pathogenesis of essential hypertension [497].

Although roles for individual NOX isoforms in specific aspects of hypertension pathology, such as endothelial dysfunction or VSM proliferation have been suggested, there remain incompatibly opposing views over which specific isoforms of NOX are most important in determining the resultant increased MAP. Considering the general consensus over the effectiveness of antioxidants and broad spectrum NOX inhibitors at lowering MAP, improving endothelial function and associated remodelling in various animal models, and bearing in mind that the same knockout mouse often produces opposing results in the same model of hypertension, it is difficult to explain the discrepancies. It may well be that NOX4 does indeed exert both protective and pathogenic effects in hypertension, but still there are discrepancies. One can only speculate about subtle differences in experimental approach, such as doses of Ang II, duration of infusion, or methods used to measure the various functional and anatomical changes. Results with NOX5 are fewer but more consistent, if mice expressing human NOX5 are to be taken as valid models of human disease.

7. NOX in the pulmonary circulation. As discussed in previous sections, in the systemic circulation as a whole we can make a number of generalisations about the role of NOX-derived ROS in the control of vascular tone: on the one hand, superoxide and H₂O₂ produced within the VSM contribute to agonist-induced contraction and myogenic tone, while on the other, ROS produced by the endothelium contribute to agonist or flow-induced relaxation, primarily mediated by endothelial H₂O₂. Functionally, the pulmonary circulation differs from the systemic circulation in that it must respond to increases in cardiac output with vasodilation, such as during exercise, to accommodate the increased flow. It is therefore a low resistance/low pressure system, exhibiting no flow-autoregulation and

minimal or absent myogenic contraction. While isolated arteries do constrict in response to prostaglandins, 5-HT, ET-1 and sphingolipids, the pulmonary vasculature operates largely independently of circulating vasoactive hormones and the autonomic nervous system, but does exhibit endothelium-dependent relaxation, largely mediated by NO $\cdot$, but with an additional contribution of EDHF, which is relatively poorly characterised in this vascular bed [571-573]. Flow-induced dilation in the lung is believed to be achieved primarily through passive distension and recruitment of arterioles, although endothelium-dependent dilation may also contribute [291]. A perhaps unique feature of the pulmonary circulation is the fact that arteries constrict in response to hypoxia (hypoxic pulmonary vasoconstriction, HPV), as opposed to most systemic vessels, which dilate [574]. This contributes to ventilation/perfusion matching in healthy lungs, whereby regional hypoxia triggers constriction in that region, directing blood flow away from poorly ventilated areas to better ventilated areas of the lung, thus maximising blood oxygenation. This section will outline some of the key differences in the way pulmonary arteries respond to ROS, in comparison with systemic vessels, and the potential role of NOX in those differences.

7.1. Differences between pulmonary and systemic arteries. Within the context of this review, one may ask: are there fundamental differences in the balance between pro-contractile and pro-relaxant effects of NOX-derived ROS in pulmonary vs. systemic arteries? In terms of relaxation, very few studies have examined the role of ROS in endothelial function in pulmonary arteries and these appear to disagree on the key points. An earlier study, comparing isolated cat cerebral and pulmonary arteries, suggests that acetylcholine-induced relaxation is partly suppressed by blocking endothelial SOD in both vessel types but is suppressed by the addition of catalase in cerebral arteries only, suggesting an NO $\cdot$ scavenging contractile effect of superoxide in both vessels, but that a relaxant effect of endothelial H $_2$ O $_2$ is present in cerebral but absent from pulmonary [437]. In contrast, two recent studies have suggested a role for endothelial H $_2$ O $_2$ in mediating an EDHF response in the pulmonary circulation, one showing that shear-stress activates eNOS via endothelial NOX4-derived H $_2$ O $_2$ [291] and another that bradykinin-induced relaxation is suppressed by catalase, making it H $_2$ O $_2$ -dependent [573].

As has already been noted in previous sections, the use of exogenous ROS is not a perfect tool for elucidating the various functions of endogenous ROS, but nevertheless, there are apparent differences in the way that pulmonary and systemic arteries respond to exogenously applied ROS, suggesting that perhaps the key differences between vascular beds lie in the effector pathways. The majority of studies in pulmonary arteries show that they constrict in response to exogenous H $_2$ O $_2$, as well as to superoxide generated by X/XO or LY83583, whether it is applied in the presence of pre-constriction or not and regardless of concentration, whereas in systemic vessels, exogenous superoxide may have a mixed contraction/relaxation effect, depending on concentration, and H $_2$ O $_2$ will usually relax pre-constricted systemic vessels, regardless of concentration [291, 573]. It is also worth reiterating that direct constriction induced by exogenous ROS, in the absence of any pre-constriction, is common in pulmonary but very rare in systemic vessels, particularly resistance sized vessels (Table 3). If effector pathways are different or responding to ROS differently in pulmonary arteries, what may these

differences be? When comparing the effects of exogenous superoxide on pulmonary and mesenteric arteries, a key difference is that the Rho-kinase signalling pathway responds to superoxide in pulmonary but not mesenteric arteries. This was illustrated by LY83583 inducing contraction in α -toxin permeabilised pulmonary but not mesenteric arteries and enhancing Rho-kinase dependent MYPT1 phosphorylation in pulmonary but not mesenteric arteries, even though GPCR-induced contraction is sensitive to Rho-kinase inhibition in both pulmonary and mesenteric arteries [351]. Alternatively, there may be major differences in the way ion channels, particularly K^+ channels, respond to ROS. These contractile signalling pathways and their influence by ROS in the pulmonary circulation are discussed further in relation to the HPV response.

7.2. Role for NOX-derived ROS in HPV. In the pulmonary artery, there is almost universal agreement that ROS are essential mediators of HPV, but there are a number of theories about how ROS production or the effects of ROS on VSM function during acute hypoxia could be different in pulmonary arteries compared to systemic, and some of these include a role for NOX in that difference. Initial studies suggested NOX as the main source of ROS during HPV [575, 576] and that both basal and stimulated NOX-derived ROS production is several fold higher in pulmonary artery VSM compared to coronary artery VSM, for example [67]. Other studies showed that while HPV is preserved in gp91^{phox}^{-/-} mice [577, 578] and unaffected by NOX2 siRNA [579], hypoxia induces translocation of p47^{phox} in pulmonary VSM and acute hypoxia-induced H₂O₂ production and HPV are partially suppressed in p47^{phox}^{-/-} mice [165, 578] or by NOX4 siRNA [579]. Thus ROS derived from NOX4, with a possible contribution from NOX1, and none from NOX2, may be important in mediating HPV. Considering NOX as a possible ROS sensor in HPV, it was suggested that while levels of NOX4 expression in the two tissues were not different [67], higher expression of glucose-6-phosphate dehydrogenase in pulmonary compared to coronary VSM could support a greater availability of pentose phosphate pathway-derived NADPH and NADH during acute hypoxia, resulting in relatively greater NADPH- and NADH-dependent NOX-derived ROS production and downstream pro-contractile signalling [580]. However, it must now be acknowledged that the primary hypoxic sensor and leading source of any change in ROS during acute hypoxia almost certainly originates in the mitochondria. There are two key theories in this regard. The first theory is that acute hypoxia reduces the production of mitochondrial ROS, resulting in reduced K_v1.5 and/or K_v2.1 current amplitude, depolarisation and enhanced Ca²⁺ influx [581]. This theory is consistent with the widely observed activation of VSM K_v currents by ROS in various vascular beds (section 5.5), including pulmonary [351], but does not explain the apparent dependence of hypoxic pulmonary vasoconstriction on Rho-kinase mediated Ca²⁺ sensitization [271, 582, 583], RyR mediated Ca²⁺ release [375, 584, 585] and store-operated Ca²⁺ influx [368, 586]. The second, and perhaps now more widely accepted theory is that hypoxia *increases* the production of mitochondrial ROS [1, 2], permitting the also widely observed ROS-mediated activation of Rho-kinase, sarcoplasmic reticulum Ca²⁺ release and TRP channel mediated Ca²⁺ influx (sections 4.2-4.4). A recent study on the effects of cross-transplantation of mitochondria between pulmonary and femoral VSMCs supports this, showing that acute hypoxia increases ROS production in native VSMCs from both vessel types but does so to a greater extent in pulmonary and that mitochondrial cross-transplantation fundamentally reverses the differences between pulmonary and femoral ROS

production, $[Ca^{2+}]_i$ responses and contraction in response to hypoxia [584]. Nevertheless, the theory that hypoxia increases ROS production is also compatible with a secondary but potentially important role for NOX-derived ROS in HPV, using PKC as a link between the two sources of ROS. This was first suggested in perfused rabbit lungs where hypoxia appeared to stimulate PKC-dependent NOX activation [587]. It was then shown that PKC- ϵ in particular is activated by mitochondrial ROS during acute hypoxia in pulmonary VSM, but not in mesenteric, and that PKC- ϵ was required for HPV [588]. The connection with NOX was then made by showing that acute hypoxia enhanced p47^{phox} dependent ROS production, presumably from NOX1, because NOX2 was not expressed, and that this ROS production and HPV were suppressed in PKC- $\epsilon^{-/-}$ mice [165], indicating that PKC- ϵ activates NOX1 in response to mitochondrial ROS by phosphorylating p47^{phox}, as also described for PKC- δ and PKC- ζ (section 2.2). In contrast, this study also showed that hypoxia did not activate either PKC- ϵ or NOX in mesenteric arteries [165]. Interestingly, Src, another kinase known to be important in the assembly and activation of NOX1 or 2, is also activated by acute hypoxia via mitochondrial ROS [174]. There are as yet no reports of an involvement of NOX5 in HPV. As an additional effector pathway for ROS induced contraction during HPV, K^+ channel *inhibition* by hypoxia-induced, NOX4-derived ROS has been suggested [589], perhaps via ROS-induced channel degradation [590], as had been previously claimed for GPCR-induced contraction downstream of NOX-derived ROS in pulmonary artery [58, 146]. However, this is a relatively infrequently reported phenomenon in a wider context, with the majority of studies on K_v channels showing them to be activated by ROS, as discussed earlier in this review. It remains to be determined whether the inhibition of K_v currents by ROS is pulmonary-specific. A proposed model for the role of NOX in HPV is summarised in Figure 9.

7.3. Role of NOX in pulmonary hypertension. Pulmonary hypertension (PH) is defined as a mean pulmonary arterial pressure in excess of 25mmHg or in excess of 20 mmHg if there are other hemodynamic abnormalities present. PH has a number of causes, including idiopathic (genetic and pharmacologically induced, also known as pulmonary arterial hypertension, PAH)); and PH due to lung disease or chronic hypoxia [591]. There are other categories, but these two are the most widely studied, having approximate correlates in animal models. Common features of all types of PH are impaired endothelium-dependent relaxation, contractile hyper-responsiveness and vascular remodelling. As with systemic hypertension, oxidative stress is a central feature of the underlying pathophysiology of PH/PAH [59, 210, 234, 589, 592-599] and antioxidants or drugs that target specific sources of ROS, such as NOX, have been suggested as potential therapies for PH [600-604].

Research on enhanced pulmonary VSM contractility in PH has focussed on the mechanisms of elevated $[Ca^{2+}]_i$ and the role of RhoA/Rho-kinase, and as far as chronic hypoxia-induced PH is concerned many of the pathways implicated in normal physiological HPV, such as K_v channel inhibition, increased RyR-mediated Ca^{2+} release, increased LVGCC and TRPC-mediated Ca^{2+} influx and increased RhoA/Rho-kinase activity, are also implicated in PH [114, 605, 605-616], as well as NOX-dependent impairment of endothelium-dependent relaxation [594, 617]. Some of these studies suggest that many of the signalling proteins that respond to NOX-derived ROS in response to acute hypoxia are upregulated in chronic hypoxia-induced PH or monocrotaline (MCT)-induced PAH (a model of idiopathic PAH). These

include L-type (Cav1.2) and T-type (Cav 3.2) Ca²⁺ channels [610], RyR2 [611], TRPC1 and TRPC6 [612], Ca²⁺-activated chloride channel (TMEM16A) [613] and RhoA and Rho-kinase [605]. In addition, in PASM from patients with idiopathic PAH, expression of the Ca²⁺ sensing receptor is increased and this has been linked to increased Ca²⁺ influx via STIM/Orai/TRPC channels [614]. Conversely, expression of K⁺ channels, including Kv1.5 and KCNK3 (TASK-1) are down-regulated by chronic hypoxia and by mutations in bone morphogenetic protein receptor-2 (BMPR2) a common cause of idiopathic PAH in humans [618-621]. Vascular remodelling in PH includes hyper-proliferation and migration of VSMC, endothelium and fibroblasts and muscularisation of normally non-muscular arterioles and, although oxidative stress is implicated in the pathogenesis of this, there appears to have been relatively little targeted research into the ROS-sensitive signalling pathways involved. What can be gleaned however is that the main pathways that operate in systemic hypertension, such as PTP oxidation [597], growth-factor transactivation [622], Nrf2 suppression [597], and kinases Src [623], PYK2 [624, 625], JAK2 [626], FAK [269], ERK [210, 525, 593, 625, 627] and PI3K/AKT [628, 629] all play major roles in PH/PAH. What may differ is the role of individual NOX isoforms and how their expression and activity is altered to promote remodelling and hyper-contraction. A number of pathways exist through which expression of NOX and associated signalling proteins may be altered in PH/PAH. These include hypoxia-inducible factor-1 α (HIF-1 α) because it is known to be activated by mitochondrial ROS [606, 612, 618, 619, 630, 631], the production of which is increased by hypoxia [1, 2]. Activation of HIF-1 α by hypoxia may occur via the intermediary ROS-dependent activation of Src [632]. Also, as discussed further below, important roles for TSP-1 [633], NF κ B [625, 633] and peroxisome proliferator-activated receptor gamma (PPAR γ) [625, 633] in regulating NOX expression and associated pulmonary vascular remodelling have also been suggested.

Even though one study suggest that NOX1 is poorly expressed in PASM from large pulmonary artery from human donors [210], NOX1 expression is several-fold higher in PASM from patients with idiopathic PAH, particularly females and is further increased by treatment with either the estrogen metabolite 16 α -hydroxyestrone (16 α -HOE) or 5-HT, both of which have been associated with the pathogenesis of PAH in humans [59, 597, 634]. NOX1 expression is also upregulated by chronic hypoxia in human pulmonary artery fibroblasts [635]. Only a handful of studies have examined the role of NOX1 in animal models of PH/PAH, with occasional conflicting results. Chronic hypoxia induces pulmonary NOX1 expression in piglets [594, 617], but it is not altered in new-born lambs with PH induced by antenatal ligation of the ductus arteriosus [598]. In rats, chronic hypoxia appears not to alter NOX1 expression in pulmonary artery [589, 636] whereas MCT induces a 10-fold increase [593]. In the latter model, NOX1 siRNA normalises MCT-induced PASM ERK phosphorylation, proliferation and migration [593]. NOX1 knockout in mice suppresses the increased right ventricular systolic pressure and prevents the right ventricular hypertrophy and vascular remodelling induced by chronic hypoxia [59]. NOX1^{-/-} also suppresses oxidant stress and PASM proliferation and reverses suppression of Nrf2 activity induced by 5-HT [597]. Curiously, another study shows the opposite of the above, wherein NOX1^{-/-} induces PH under normoxic conditions [637].

NOX2 is also reportedly poorly expressed in PASMC of large pulmonary arteries from human donors [210], but unlike NOX1, PH in new-born lambs induced by antenatal ligation of the ductus arteriosus is associated with increased NOX2 expression [598]. In one study in rats, PH induced by both MCT and chronic hypoxia is associated with increased expression of NOX2 in pulmonary arterioles and in the endothelium, and it is suggested that this increase is due to a concomitant decrease in expression of caveolin-1 [234]. Alternatively, other studies in rats show that pulmonary NOX2 expression is not significantly influenced by chronic hypoxia [589, 596, 636] or MCT [593, 638], but that NOX2 activity may instead be increased [596]. Either an increased expression or an increased activity of NOX2 would be consistent with the inhibitory effect of gp91ds-tat on Ca^{2+} influx-independent contraction in pulmonary artery from chronic hypoxic rats [596]. It would also be consistent with the suppressing effect of NOX2 knockout on pulmonary constrictor hyper-responsiveness, PDGFR and AKT activation, vascular remodelling, increased right ventricular pressure and right heart hypertrophy induced by chronic hypoxia in mice [599, 639]. Another study in mice shows that chronic hypoxia does increase pulmonary NOX2 expression and suggests that this increase and the associated vascular remodelling are dependent on mitochondrial ROS [631].

In PASMC from idiopathic PAH patients, expression of NOX4 was increased 2.5 fold compared to matched controls [59, 629]. In cultured human PASMC, NOX4 expression is increased by chronic hypoxia and this is mediated in part by autocrine induction of IGF binding protein-3 [592, 635] or by upregulation of TSP-1 expression, increased NF κ B activity and suppression of PPAR γ (a negative regulator of proliferation) [625, 633]. In this instance, PYK2 appears to be upstream of NF κ B activation and the downstream ERK phosphorylation required for proliferation [625]. NOX4 expression in human pulmonary artery is also increased by another key mediator of VSMC proliferation, transforming growth factor- β (TGF- β) [52, 210]. In new-born lambs with PH induced by ductus arteriosus ligation, NOX4 and p22^{phox} expression is increased [598], and also in this model, cyclical stretch, which may exert a stronger influence on the vessel wall during PH, induces NOX4 expression via an increase in mitochondrial ROS production and induction of NF κ B [640]. In rats, a number of studies show that pulmonary NOX4 expression is upregulated by chronic hypoxia [234, 589, 595, 629, 636], MCT [234, 595, 641], another model of spontaneous PH [595] and by human BMPR2 activation [642], while another study finds no significant effect of MCT on pulmonary NOX4 expression in rats [593]. Also in contrast to rat studies, in piglets there is apparently no effect of chronic hypoxia on pulmonary NOX4 expression [594]. In mice, pulmonary NOX4 expression is increased by chronic hypoxia [631, 643], chronic intermittent hypoxia (CIH) [639] and MCT [638]. In one of these studies, the enhanced NOX4 expression and the associated PASMC proliferation required hypoxic induction of HIF-1 α [643], while in another, it and the associated vascular remodelling were also dependent on mitochondrial ROS, suggesting a link between mitochondrial ROS and cytosolic NOX4-derived ROS in PH [631], a parallel mechanism to that suggested for the acute HPV response [165, 588].

Of all the NOX isoforms expressed in the pulmonary vasculature, the greatest amount of work has been done on the role of NOX4 in the mechanisms of vascular remodelling associated with PH/PAH. Human PASMC proliferation induced by chronic hypoxia is suppressed by NOX4 siRNA [592, 627] and

the dual specific inhibitor GKT137831 [644]. GKT137831 also inhibits proliferation of these cells stimulated by the estrogen metabolite 16 α -OHE and does so more effectively in PASMC from PAH patients [59]. NOX4 siRNA also inhibits proliferation induced by TGF- β in human PASMC [210, 592], by cyclical stretch in PASMC from new-born lambs with PH induced by ductus arteriosus ligation [598, 640] and by thrombin and human BMP2 in rat PASMC [642, 645]. Putative NOX4-specific inhibitors VCC588646 and VCC202273 also partially prevent pulmonary vascular remodelling induced by MCT in rat lung [595]. Interestingly, in this last study, pulmonary NOX4 was found to be expressed primarily in adventitial fibroblasts and there is emerging evidence for its contribute to PH through promoting fibrosis and arterial stiffening [595, 646]. In terms of downstream signalling, NOX4 contributes to the enhanced mTOR, S6K and AKT phosphorylation, cell survival and proliferation observed in PASMC from PAH patients [629], and it appears to contribute to human PASMC proliferation in response to chronic hypoxia in part via suppression of PPAR γ expression, suggesting a reciprocal negative relationship between NOX4 and PPAR γ [627]. In rat PASMC, NOX4 appears to act via protein stabilization of HIF-2 α and induction of plasminogen activator inhibitor-1 [645], while in PASMC from new-born lambs with PH induced by ductus arteriosus ligation it acts via NF κ B activation [598]. In models of PH/PAH, there appears to be no evidence for NOX4 contributing to pulmonary VSMC migration, although one may expect this based on its known association with integrins, FAK/ILK and migratory stimuli in systemic hypertension [107]. There is also evidence that NOX4 participates in aberrant elevated [Ca²⁺]_i in PH by suppressing PASMC K_v1.5 current [589] and by enhancing expression of TRPC1 and TRPC6 [642]. In stark contrast to the above, two studies in mice show little involvement of NOX4 in PH, despite evidence for NOX4 upregulation by chronic hypoxia [631, 643], one showing that NOX4^{-/-} only reduces the chronic hypoxia-induced increase in right ventricular systolic pressure but has no effect on vascular remodelling [59], and the other showing no effect of NOX4^{-/-} on chronic hypoxia-induced increases in pulmonary artery pressure or vascular remodelling [647].

As with systemic hypertension, this data provides conflicting evidence over the importance of NOX to PH/PAH, particularly to the associated vascular remodelling. The data possibly suggests that there are species-dependent differences over which NOX isoform may be most important in PH, perhaps NOX4 and NOX1 in human and rat, NOX1 and NOX2 in mice, with NOX2 appearing of minimal importance in rats and humans and NOX4 possibly of minimal importance in mice. However, evidence also conflicts between studies using the same model in the same species, so differences may be explained by the use of different output measures, time points and drug delivery methods or different approaches to data collection. Interestingly, there is very little evidence of a protective effect of NOX4 in PH, unlike in systemic hypertension. This is in line with NOX4 being purely pro-contractile in the healthy pulmonary circulation and primarily pro-relaxant in the systemic. However, one recent study raises doubts about this too, suggesting that H₂O₂, probably derived (although not proven) from NOX4, exerts a protective effect in parallel with the damaging effects described above [648]. This study presented evidence for an enhanced oxidative activation of PKG1 α in lung tissues from humans with PH and an animal model of chronic hypoxia, and that this enhanced activity of PKG1 α was protective via both enhanced VSM relaxation and suppression of vascular remodelling. Clearly more work needs to be done to clarify the importance of NOX to the pathophysiology of PH and PAH and if possible

determine whether there are specific NOX isoforms that may be most suitable as potential therapeutic targets. The potential role of NOX in the pathogenesis of PH/PAH is summarised in Figure 10.

8. Summary and future directions.

There exists a highly complex network of tightly regulated signalling pathways both upstream and downstream of NOX-derived ROS production that operates in subtly different ways for different NOX isoforms in the VSM and endothelium of different vascular beds. They respond to a variety of hormonal, chemical and mechanical stimuli, to produce a range of physiological responses in the healthy vasculature. This includes VSM contraction, endothelium-dependent relaxation, VSM gene expression, growth, proliferation and migration. Although much has been done to characterise the roles of specific NOX isoforms in these functions, more work is needed to clarify the importance of compartmentalisation, particularly of NOX4 and NOX5 which are both expressed in sarcoplasmic reticulum and NOX4 in mitochondria and nucleus. There is extensive but inconclusive, and occasional contradictory evidence that expression and activity of NOX isoforms and associated signalling pathways are altered in various animal models of hypertension, pulmonary hypertension and pulmonary arterial hypertension. A role for NOX-derived ROS in the vascular inflammation that underlies many cardiovascular diseases, as well as the role of adventitial NOX, requires further investigation. For appropriate therapies to be devised and tested for use in humans, further work is needed to clarify the relative importance of specific NOX isoforms in these diseases, particularly NOX4 which appears to exert both detrimental and protective effects in hypertension and possibly in PH/PAH as well. Thus, whether general antioxidants or non-selective and highly selective NOX inhibitors prove beneficial in the treatment of hypertension in humans may depend on the relative effects of suppressing pathological and protective ROS function as well as their contributions to normal vascular homeostasis.

The diagram illustrates the signaling pathways for NOX1/2 and NOXO1/NOXA1 activation, divided into Phase one and Phase two.

Phase one: GPCR (G-protein coupled receptor) is activated by ligand A, leading to the activation of Gα_q, Arf6, and Gβγ. Gα_q activates PLC, which produces IP₃ and DAG. IP₃ releases Ca²⁺, which activates PKC α/β. Arf6 activates PLA₂, which produces AA, and PLD, which produces LPC and PA. Gβγ activates PKC-δ/ε/ζ. PKC-δ/ε/ζ, DAG, and Ca²⁺ all contribute to the activation of Src. Src is also activated by ROS. Activated Src leads to the translocation of NOX1/2 and NOXO1/NOXA1 to the membrane, where they are phosphorylated (P47^{phox}/NOXO1, P67^{phox}/NOXA1, and p40^{phox}) and associated with EBP50 and Cortactin.

Phase two: The activated NOX1/2 and NOXO1/NOXA1 complex produces superoxide (O₂^{•-}). This superoxide is converted to hydrogen peroxide (H₂O₂) by the enzyme MMP. H₂O₂ then activates the EGFR pathway. HB-EGF binds to EGFR, leading to the recruitment of PTEN, SHP2, PI3K, and Src. PI3K produces PIP₂ and PIP₃. PIP₂ is converted to PIP₃ by SHP2. PIP₃ recruits RacGEFs, which activate Rac. Rac then activates PKCβ. PKCβ, along with RacGEFs and Rac, leads to the activation of NOX1/2 and NOXO1/NOXA1. The activated NOX1/2 and NOXO1/NOXA1 complex produces superoxide (O₂^{•-}), which is converted to H₂O₂ by MMP. H₂O₂ then activates the EGFR pathway, leading to the activation of Src and PI3K, which in turn activate Rac and PKCβ, leading to the activation of NOX1/2 and NOXO1/NOXA1.

Figure 1. Activation of vascular NADPH oxidase-1 and -2 (NOX1/2) by G-protein coupled receptor (GPCR) stimulation. Inactive NOX1/2 exist in cell membranes coupled with p22^{phox}. Their activation follows a complex series of steps which can be divided into two phases. **Phase one:** Partial assembly and partial activation of the enzyme complex require translocation of regulatory subunits p47^{phox}, p67^{phox} and p40^{phox} from the cytosol to the membrane. This is achieved through phosphorylation of p47^{phox}, and possibly also of p67^{phox} and p40^{phox}, by multiple protein kinase C (PKC) isozymes and Src. PKC- δ may first need to phosphorylate SM_{22 α} before it can phosphorylate p47^{phox}. NOXO1 may substitute for p47^{phox} and NOXA1 may substitute for p67^{phox}. Src also phosphorylates cortactin, releasing it from the regulatory subunits, thus also supporting translocation. Thirdly, translocation is also facilitated through interaction with ezrin-radixin-moesin binding phosphoprotein 50 (EBP50). PKC isozymes are activated by GPCR through multiple pathways. Gq activates phospholipase C (PLC) which makes inositol trisphosphate (IP₃) and diacylglycerol (DAG). IP₃ triggers Ca²⁺ release from the sarcoplasmic reticulum to activate PKC- α or - β . DAG also is required for activation of PKC- α , - β , - δ and - ϵ . G $\beta\gamma$ subunits activate phospholipase A₂ (PLA₂) and phospholipase D (PLD), while PLD also requires the small G-protein ARF6. The products of these, arachidonic acid (AA), lysophosphatidylcholine (LPC) or phosphatidic acid (PA), also activate PKC- δ , - ϵ and - ζ . Src is activated directly by G $\beta\gamma$. Alternatively, PKC and Src are capable of activating each other by phosphorylation and are both activated by other sources of ROS. **Phase two:** Full, complete activation of the oxidase requires the binding of active Rac to p67^{phox} and the catalytic subunit. Rac activation requires prior activation of guanine nucleotide exchange factors (RacGEFs) and these are activated by PIP₃ and tyrosine phosphorylation by Src, FAK, PYP2 or JAK2. PIP₃ is generated by phosphatidylinositol-3 kinase (PI3K) which is primarily activated following transactivation of the epidermal growth-factor receptor (EGFR) by Src. Low level ROS produced by the partially active NOX are able to diffuse into the cytosol via anion channels or aquaporins and may be sufficient to further activate Src as well as inhibit the tyrosine phosphatase SHP2 (inactivates PI3K) and the lipid phosphatase PTEN (degrades PIP₃ and PIP₂). EGFR may also be activated by epidermal growth-factor (EGF) generated by a ROS-activated matrix metalloprotease. Finally, full activity of the oxidase is promoted by further phosphorylation by PKC- β , binding of PIP₂ and PIP₃ and stabilization by heat-shock protein 90 (HSP90) and protein disulphide isomerase (PDI). PDI may form an intermolecular disulphide bond with p47^{phox}. Open arrowheads indicate movement or production of a substance. Closed arrowheads indicate an effect or response.

Figure 2

Figure 2. Activation of vascular NADPH oxidase-4 (NOX4) and -5 (NOX5) in vascular cells. NOX4 exists in cell membranes as a stable complex with p22^{phox} and at the cell surface requires the additional binding of polymerase delta interacting protein 2 (Poldip2), which is associated with integrins and talin (not shown) at focal attachments in vascular smooth muscle. This may be sufficient for enzyme activity and H₂O₂ production, but activity may also be stimulated further by G-protein-coupled receptor (GPCR)-induced or receptor tyrosine kinase (RTK)-induced production of arachidonic acid (AA), activation of Rac or activation of protein kinase C (PKC). NOX4 is also present in the sarcoplasmic reticulum, although it is not clear whether or how it responds to cell surface receptor stimulation from this location or is simply constitutively active. NOX5 exists in cell membranes and, in endothelium, is associated with caveolin, which exerts an inhibitory effect on catalytic activity which in turn may be relieved by heat-shock protein 90 (HSP90). NOX5 is primarily activated acutely by an increase in intracellular Ca²⁺ concentration induced by GPCR, primarily via Ca²⁺ release from the sarcoplasmic reticulum (SR), which it senses via four 'EF hands' and Ca²⁺-calmodulin (CaM) binding sites on a long intracellular auto-inhibitory domain. Sensitivity of NOX5 to Ca²⁺ may also be enhanced by phosphorylation of the catalytic subunit by PKC-α, Ca²⁺-calmodulin-activated kinase II (CAMK-II) or c-Abl but may be inhibited by phosphorylation by PKC-ε. ECM = extracellular matrix. Open arrowheads indicate movement or production of a substance. Closed arrowheads indicate an effect or response.

Figure 3

Figure 3. Compartmentalisation and metabolism of NADPH oxidase (NOX) isoforms. All NOX isoforms are expressed at the cell surface, but in endothelium NOX1, 2 and 5 are concentrated in caveolae (associated with the flow sensing 'mechanosome' (M)), while in vascular smooth muscle NOX4 is concentrated at focal attachments in association with polymerase delta interacting protein-2 (Poldip2). Superoxide produced at the cell surface in response to G-protein coupled receptor (GPCR) stimulation is extracellular, but can enter the cell through anion channels (AC, either chloride channel-3 or LRRC8) or be dismutated by extracellular superoxide dismutase (EC-SOD) into H_2O_2 which can enter the cell via aquaporins. Alternatively, all NOX isoforms may be endocytosed with activating receptors (eg. receptor tyrosine kinases (RTK) or TNF- α receptors (TNFR)). ROS are produced within the endosome to influence effector pathways there or can enter the cytosol through anion channels. NOX4 is highly expressed in the sarcoplasmic reticulum (SR) where it will influence protein folding and possibly Ca^{2+} handling. H_2O_2 is free to leave the SR via aquaporins. NOX5 is also expressed in the SR and NOX4 is also expressed in the mitochondria and nucleus. In the nucleus, NOX4 may contribute to the regulation of gene expression, while the function of NOX5 in the SR and NOX4 in the mitochondria is unknown. The main source of mitochondrial ROS is most likely complexes I and III of the electron transport chain. These are also free to enter the cytosol through aquaporins and possibly also anion channels. Superoxide and H_2O_2 oxidise cysteine (-SOH) and methionine (-SCHO, not shown) in target proteins, either directly or using oxidised PRX as an intermediate. Oxidation of adjacent cysteine residues may trigger formation of a disulphide bridge. Nitric oxide ($NO\cdot$) modifies cysteine by nitrosylation (-SNO). Superoxide and $NO\cdot$ combine to form peroxynitrite ($ONOO^-$) which modifies proteins through nitration of tyrosine residues (-ONO). Glutathione peroxidase (GPX) uses the main intracellular reductant reduced glutathione (GSH) to reduce H_2O_2 , producing oxidised glutathione (GSSG). GSSG may also modify proteins through glutathionylation (-SSG). Antioxidant enzymes are widely expressed in the cytosol and intracellular organelles where they will limit the accumulation of ROS and contribute to the compartmentalisation of ROS signalling. Superoxide is dismutated by SOD1 in the cytosol, endosomes and mitochondrial outer membrane and by SOD2 in the mitochondrial matrix. H_2O_2 is reduced by cytosolic catalase (Cat) and GPX, by GPX in the mitochondria and by GPX and PRX in the SR. GSSG is re-reduced by glutathione reductase, using NADPH (not shown). Open arrowheads indicate movement or production of a substance. Closed arrowheads indicate an effect or response.

Figure 4

Figure 4. Role of NADPH oxidase (NOX) in vascular smooth muscle (VSM) Ca²⁺ signalling. G-protein coupled receptor (GPCR) stimulation activates phospholipase C (PLC, via G_q) and NOX1 or NOX2 on the cell surface. PLC generates inositol trisphosphate (IP₃) and diacylglycerol (DAG). IP₃ triggers Ca²⁺ release from the sarcoplasmic reticulum via IP₃R while DAG activates protein kinase C (PKC). NOX-derived ROS induce extracellular Ca²⁺ influx and Ca²⁺ release from the sarcoplasmic reticulum via multiple pathways. ROS stimulate Ca²⁺-activated chloride channels (Cl_{Ca}) and TRPC1/6/Orai, the latter probably via stimulation of PKC-dependent phosphorylation. Opening of these channels depolarises the cell membrane, thus opening L-type voltage-gated Ca²⁺ channels (LVGCC). ROS may further enhance LVGCC opening through channel phosphorylation by Src, PKC or CAMK-II, or by direct cysteine oxidation, but whether the latter is stimulatory or inhibitory is unclear. ROS promote Ca²⁺ release from the sarcoplasmic reticulum by enhancing the sensitivity of IP₃R to IP₃ and by activating the ryanodine receptor (RyR). RyR is also activated by ROS indirectly by displacing the inhibitory protein FKBP from RyR and by stimulating CD38, which makes cyclic ADP-ribose (cADPR) and nicotinic acid adenine dinucleotide phosphate (NAADP). cADPR further stimulates RyR while NAADP stimulates Ca²⁺ release via the Two-Pore channel (TPC). ROS also help maintain an elevated [Ca²⁺]_i by inhibiting Ca²⁺ extrusion via the plasma membrane Ca²⁺ ATPase (PMCA) and re-uptake into the sarcoplasmic reticulum via the sarco-endoplasmic Ca²⁺ ATPase (SERCA). Finally, ROS also may stimulate the Na⁺/Ca²⁺ exchanger, but this may oppose the other effects by extruding Ca²⁺. NOX4 is expressed in the sarcoplasmic reticulum of VSM, but details of its contributions to Ca²⁺ homeostasis are unknown. Open arrowheads indicate movement or production of a substance. Closed arrowheads indicate an effect or response.

Figure 5

Figure 5. Role of NADPH oxidase (NOX) in Ca^{2+} -independent pro-contractile signalling in vascular smooth muscle. Contractile force may be enhanced without an increase in $[\text{Ca}^{2+}]_i$ through two main pathways: inhibition of myosin light-chain phosphatase (MLCP) and actin polymerisation. MLCP is negatively regulated by Rho-kinase (ROCK) through inhibitory phosphorylation of myosin phosphatase regulatory subunit-1 (MYPT1). ROCK is activated by RhoA. RhoA is either activated directly by superoxide or by Rho guanine nucleotide exchange factors (RhoGEFs). RhoGEFs are activated by G-protein coupled receptors (GPCR) either directly via $\text{G}_{12/13}$, or indirectly by NOX1- or NOX2-dependent tyrosine phosphorylation by Src, PYK2 or JAK2. RhoGEFs are also activated by stretch via integrin engagement with extracellular matrix (ECM) and subsequent activation of focal adhesion kinase (FAK), in association with H_2O_2 produced by NOX4 coupled to polymerase delta interacting protein-2 (Poldip2). RhoA promotes actin polymerisation either directly or indirectly by promoting ROCK and LIM kinase (LIMK)-induced inhibition of the actin severing protein cofilin. Additional routes for the promotion of actin polymerisation are by direct oxidation of actin by NOX4-derived H_2O_2 , inhibition of the actin chaperone heat-shock protein 27 (HSP27) by p38 MAP-kinase and MAP-kinase activated protein kinase (MAPKAP) and ERK MAP-kinase-mediated inhibition of caldesmon which inhibits actin-myosin cross-bridge cycling. p38 and ERK MAP-kinases are both activation by ROS via MEK. NOX-derived H_2O_2 activates MEK via Src-dependent transactivation of receptor tyrosine kinases (RTK) and subsequent recruitment of growth-factor receptor binding protein-2 (GRB2), Ras and Raf, by direct oxidation of Ras, by oxidative inhibition of a MEK phosphatase or via protein kinase C (PKC) directed phosphorylation. PKC also inhibits MLCP by activating the inhibitory accessory protein C-kinase potentiated protein phosphatase-1 Inhibitor-17 (CPI-17). Open arrowheads indicate movement or production of a substance. Closed arrowheads indicate an effect or response.

Figure 6

Figure 6. Role of NADPH oxidase (NOX) in endothelium-dependent relaxation of the vasculature.

NOX1, 2, 4 and 5 are all expressed in the endothelium and 1, 2 and 5 are present in the caveolae. All isoforms are activated by pro-relaxant G-protein coupled receptor (GPCR) agonists and by flow via the endothelial flow 'mechanosome'. In caveolae, this consists of VEGFR, VE-cadherin and PECAM-1 and is associated with closure of K_{ATP} and opening of T-type Ca^{2+} channels (not shown). Caveolin-1 acts as a negative regulator of NOX1, 2 and 5 but this inhibition can be relieved by HSP90. The mechanism of NOX4 activation by GPCR and the exact nature of the NOX4 flow sensor are unclear. NOX4 generates H_2O_2 while NOX1/2/5 make superoxide, some of which is dismutated to H_2O_2 by SOD in the endothelium. GPCR also raise endothelial $[Ca^{2+}]_i$ by triggering release from the sarcoplasmic reticulum via IP_3R and RyR, and this Ca^{2+} stimulates eNOS to make $NO\cdot$, which diffuses to the vascular smooth muscle (VSM) to activate guanylate cyclase (GC). GC makes cGMP which activates protein kinase G (PKG). PKG phosphorylates and activates large conductance Ca^{2+} -activated K^+ channels (BK_{Ca}) to hyperpolarise the VSM, lower $[Ca^{2+}]_i$ and induce relaxation by closing L-type voltage-gated Ca^{2+} channels (LVGCC). Endothelial ROS, principally from NOX4, further promote eNOS activity by activating kinases which phosphorylate eNOS and by further raising endothelial $[Ca^{2+}]_i$ by enhancing Ca^{2+} release via RyR and Ca^{2+} influx via TRPA1 channels. VSM Endothelial Ca^{2+} (enhanced by H_2O_2) also activates the small and intermediate conductance Ca^{2+} -activated K^+ channels (SK_{Ca} and IK_{Ca}). The extruded K^+ diffuses to the VSM and activates the Na^+/K^+ ATPase and inward rectifier (K_{IR}) channels to hyperpolarise the VSM, while endothelial hyperpolarisation may also spread to the VSM via myo-endothelial gap

junctions (MEGJ). NOX4-derived H_2O_2 and any dismutated superoxide from NOX1, 2 or 5 also acts as a direct hyperpolarising factor. It diffuses to the VSM where it directly activates BK_{Ca} and voltage-gated K^+ channels (K_v), further activates GC and PKG and activates MAPK to further stimulate K_v channels, thus further hyperpolarising the VSM. Superoxide derived from endothelial NOX1, 2 and 5 partly antagonises eNOS by oxidising BH_4 , thus uncoupling eNOS, forcing it to make more superoxide. Conversely, $\text{NO}^\bullet$ also limits superoxide production by inhibiting the activity of NOX1, 2 and 5. $\text{NO}^\bullet$ and superoxide react to make ONOO^- . In the VSM, this promotes formation of oxidised glutathione (GSSG) which may activate SERCA, enhancing the rate of VSM Ca^{2+} uptake into the sarcoplasmic reticulum, thus promoting relaxation. H_2O_2 may also directly inhibit LVGCC. VEGFR = vascular endothelial growth-factor receptor, PECAM-1 = platelet-endothelial cell adhesion molecule-1, HSP90 = heat-shock protein 90, SOD = superoxide dismutase, EC-SOD = extracellular SOD, IP_3R = inositol trisphosphate receptor, RyR = ryanodine receptor channel. Open arrowheads indicate movement or production of a substance. Closed arrowheads indicate an effect or response.

Figure 7

Figure 7. Proposed role of vascular NADPH oxidase (NOX) in the pathogenesis of hypertension.

Multiple mitogenic and pro-contractile stimuli (Ang II, ET-1, PDGF, TNF- α) induce an increased vascular expression of NOX1, NOX2, regulatory subunits p22^{phox} (p22), p47^{phox} (p47), p67^{phox} (p67) and NOXA1, and NOX5. Vascular NOX1 and NOX2 expression are also increased in the spontaneously hypertensive rat (SHR). NOX5 activity may be suppressed by polymorphisms in humans, while activity of p22^{phox} may be increased (not shown). NOX4 expression is upregulated by inflammatory mediators (eg TNF- α) and mechanical forces associated with cardiovascular disease: oscillatory flow in endothelium (EC) and elevated perfusion pressure in vascular smooth muscle (VSM). Most studies agree that NOX4 expression is enhanced in SHR and in Ang II-induced hypertension, and this may be via inhibition of BCL6 expression, thus relieving BCL6-induced suppression of NOX4 expression. NOX4 expression may also be enhanced by chronic intermittent hypoxia (CIH). Collectively, the result of enhanced NOX expression is the production of excess superoxide ($O_2^{\cdot-}$, NOX1, NOX2 and NOX5) and H_2O_2 (NOX4), resulting in oxidant stress. Accumulation of superoxide may be exacerbated by reduced expression or activity of superoxide dismutase (SOD). Oxidant stress impairs endothelium-dependent relaxation through the combined effects of NO $\cdot$ scavenging and eNOS uncoupling. Excess superoxide also impairs guanylate cyclase (GC) activity. Vascular hyper-contraction is further exacerbated by elevated VSM $[Ca^{2+}]_i$, through suppression of SERCA activity, enhanced Ca^{2+} release from the sarcoplasmic reticulum and elevated Ca^{2+} channel activity, and through increased expression and activity of RhoA and Rho-kinase (ROCK). Vascular remodelling occurs through the combined action of ROS-stimulated VSMC migration (mediated by FAK, ILK, integrins and RhoA/ROCK), proliferation (primarily ERK MAP-kinase mediated, MAPK), increased survival (PI3K and AKT mediated) and fibrosis, coupled to inflammation. These processes are promoted by oxidative inhibition of protein tyrosine phosphatases (PTP) and enhanced activity of non-receptor tyrosine kinases (NRTK) and receptor tyrosine kinases (RTK). The NRTKs FAK, Src, PYK2 and JAK2 as well as protein kinase C (PKC) are all essential co-ordinators of vascular remodelling (see Figure 8 for more details). They also promote hyper-contraction via RhoA/ROCK. VSM proliferation and fibrosis is also supported by oxidative activation of the transcription regulator nuclear factor kappa-light-chain-enhancer of activated B cells (NF κ B). Opposing the 'oxidant stress' effects of enhanced NOX4 expression, excess H_2O_2 may also have a protective effect in hypertension, via activation of nuclear factor erythroid 2-related factor 2 (Nrf2), which may act to improve endothelial function and maintain a differentiated VSM phenotype. TNF- α = tumor necrosis factor- α , PDGF = platelet derived growth factor. Open arrowheads indicate movement or production of a substance. Closed arrowheads indicate an effect or response.

Figure 8

Figure 8. Role of NADPH oxidase (NOX) in vascular smooth muscle growth, proliferation and migration in response to G-protein coupled receptor activation and stretch. NOX1, 2 or 5 in the plasma membrane or endosomes are activated by G-protein coupled receptor (GPCR) agonists. The ROS generated produce parallel inhibition of multiple protein tyrosine phosphatases (PTPs), and activation of multiple non-receptor tyrosine kinases (Src, PYK2 and JAK2), CAMK-II and PKC. One consequence of this is Src-mediated transactivation of receptor tyrosine kinases (RTKs) which act as platforms for the parallel activation of PI3K, Ras and PLC- γ . PI3K activates AKT and mTOR and downstream protein synthesis and cell survival responses via S6K and 4EBP. Via GRB2, Ras activates Raf, MEK and ERK MAP-kinase (ERK) to promote cell division and proliferation. Alternatively, Raf and MEK activation occur via CAMK-II-mediated or PKC/PYK2/JAK2-mediated phosphorylation. Nuclear translocation of ERK is supported by Rho-kinase (ROCK). PLC- γ triggers Ca^{2+} release from the sarcoplasmic reticulum, which activates PYK2 and then JAK2. JAK2 also supports proliferation via STAT and ERK. In addition, Src promotes activity of the pro-proliferative transcription regulator NF κ B via inhibition of I κ B. p38 MAP-kinase (p38) is also involved in stimulating proliferation downstream of MEK or TRAF2/ASK, and contributes to the activation of NF κ B, particularly in response to TNF- α receptor activation. Both p38 activation pathways are NOX-dependent (not shown). As a braking mechanism to this, ROS also promote activity of the cyto-protective transcriptional regulator Nrf2 via degradation of the inhibitory KEAP1. Via Src, GPCR-induced ROS may also activate FAK at focal adhesions. This may lead to 'inside-out' activation of integrins and activation of NOX4. Alternatively, stretch may activate integrins leading to 'outside-in' activation of FAK and NOX4. Either way, activated FAK also acts as a platform for the GRB2/Ras proliferation pathway and the PI3K/AKT growth and cell survival pathway, while FAK and NOX4-derived ROS will promote actin polymerisation/de-polymerisation, focal adhesion turnover and migration via ILK/paxillin and RhoA, Rac and cofilin activity. CAMK-II also promotes migration via Rac or MMP9. NOX4 in the mitochondria or nucleus and NOX4 and NOX5 in the sarcoplasmic reticulum may also contribute to these responses through as yet uncharacterised pathways (not shown). CAMK-II = Ca^{2+} -calmodulin activated kinase-II. PYK2 = protein tyrosine kinase-2 β . JAK2 = Janus kinase-2. MMP9 = matrix metalloprotease-9. PKC = protein kinase C. PI3K = phosphatidylinositol-3 kinase. mTOR = mammalian target of rapamycin. S6K = S6-kinase. 4EBP = eukaryotic translation initiation factor 4E-binding protein. GRB2 = growth-factor receptor bound protein-2. PLC- γ = phospholipase C- γ . STAT = signal transducer and activator of transcription. NF κ B = nuclear factor kappa-light-chain-enhancer of activated B cells. ASK = apoptosis signal-regulating kinase 1. I κ B = inhibitor of κ B. Nrf2 = nuclear factor erythroid 2-related factor 2. Poldip2 = polymerase delta interacting protein-2. FAK = focal adhesion kinase. ILK = integrin-linked kinase. RhoGEFs = Rho guanine nucleotide exchange factors. S1L = Slingshot-1L. Double-headed arrows indicate a bi-directional activation pathway. Open arrowheads indicate movement or production of a substance. Closed arrowheads indicate an effect or response.

Figure 9

Figure 9. Proposed model for the role of NADPH oxidase (NOX) in hypoxic pulmonary vasoconstriction (HPV). Hypoxia may preferentially activate NOX4 by increasing the availability of cytosolic NADPH and NADH and/or by triggering an increased production of ROS from the mitochondrial electron transport chain (ETC) which activate PKC- ϵ and/or Src to phosphorylate regulatory subunits of NOX1 (PKC and Src) and directly phosphorylate NOX4 (PKC only). NOX-derived ROS may then raise $[Ca^{2+}]_i$ through a combination of K_v channel inhibition, TRPC/Orai store-operated Ca^{2+} channel opening and subsequent L-type voltage-gated Ca^{2+} channel opening (LVGCC). TRPC channel opening may be further stimulated via ROS-dependent activation of the Ca^{2+} -sensing receptor (CaSR). ROS also trigger Ca^{2+} release from the sarcoplasmic reticulum through ryanodine receptor channels (RyR) via enhanced production of cyclic ADP-ribose (cADPR). It is not known whether NOX4 in the sarcoplasmic reticulum contributes to this. Ca^{2+} activates myosin light-chain kinase (MLCK) to phosphorylate myosin light-chain-20 (MLC₂₀) and trigger contraction. In parallel to an increase in $[Ca^{2+}]_i$, ROS may also inhibit myosin light-chain phosphatase (MLCP) activity to further enhance MLC₂₀ phosphorylation. This most likely occurs via Rho-kinase-dependent inhibitory phosphorylation of MYPT-1, and ROS most likely activate Rho-kinase via direct activation of RhoA or via Src-dependent phosphorylation of Rho guanine nucleotide exchange factors (RhoGEFs). Open arrowheads indicate movement or production of a substance. Closed arrowheads indicate an effect or response.

Figure 10

Figure 10. Proposed role of vascular smooth muscle NADPH oxidase (NOX) in the pathogenesis of pulmonary hypertension (PH) and/or pulmonary arterial hypertension (PAH). Multiple mitogenic and pro-contractile stimuli induce the increased expression of NOX1, NOX2 and/or NOX4. Chronic hypoxia and cyclical stretch act via mitochondrial ROS, the induction of HIF-1 α and NF κ B and suppression of PPAR γ . Oxidant stress probably induces pulmonary arterial hyper-contractility through the combined effects of reduced NO $\cdot$ bioavailability, elevated [Ca²⁺]_i, through suppression of K⁺ channel expression and elevated Ca²⁺ channel expression, and enhanced RhoA/Rho-kinase expression and activity. Vascular remodelling occurs through the combined action of ROS-stimulated VSMC migration, proliferation, increased survival and fibrosis. These processes are promoted and co-ordinated by oxidative inhibition of protein tyrosine phosphatase (PTP) activity and enhanced activity of non-receptor tyrosine kinases (NRTK) and receptor tyrosine kinases (RTK). Counter to the pathogenic effects of oxidant stress, there may also be an underlying protective effect via activation of protein kinase G-1 α (PKG1 α) which may promote relaxation and suppress remodelling. MCT = monocrotaline, CIH = chronic intermittent hypoxia, 16 α -HOE = 16 α -hydroxyestrone, 5-HT = 5-hydroxytryptamine, BMPR2 = bone morphogenetic protein receptor-2, TSP-1 = thrombospondin-1, IGFBP3 = IGF binding protein-3, HIF-1 α = hypoxia-inducible factor-1 α , HIF-2 α = hypoxia-inducible factor-2 α , NF κ B = nuclear factor kappa-light-chain-enhancer of activated B cells, PPAR γ = peroxisome proliferator-activated receptor gamma, Nrf2 = nuclear factor erythroid 2-related factor 2, Cav-1 = caveolin-1, ROCK = Rho-kinase, Open arrowheads indicate movement or production of a substance. Closed arrowheads indicate an effect or response.

Table 1. Effects of vasoactive and mitogenic stimuli on ROS production in vascular tissues and cells.

DHE = dihydroethidium, DCF = 2,7-dichlorofluorescein. Note that in many of the publications cited here DHE, LO-12 and Lucigenin are nominally used as selective probes for superoxide, while DCF and Amplex Red are nominally assumed to be selective for extracellular and intracellular H₂O₂ respectively. However, for some probes the accuracy and validity of this assumed selectivity may depend on the sample preparation and detection methods use [649]. VSMCs = vascular smooth muscle cells.

stimulus	Species and vascular bed	Indicator(s)	Effect(s)	reference
Intact Tissue				
Angiotensin II	Mouse aorta Mouse renal afferent arterioles Rat and mouse mesenteric artery Human internal mammary artery	Lucigenin DHE LO-12 cytochrome C reduction	Increased superoxide	[89, 94, 98, 101, 101, 104, 106]
	Rat cerebral artery Mouse renal afferent arterioles Rat renal afferent arterioles Human subcutaneous arteries	Tempo-9AC H2DCFDA/DCF	Increased H ₂ O ₂	[82, 99, 105, 325, 377]
Prostanoids	Mouse aorta Mouse mesenteric artery Bovine coronary artery Rat pulmonary artery	L-012	Increased superoxide	[66, 140, 140, 174]
sphingosylphosphoryl choline	Mouse Mesenteric	C-DCFH/DA	Increased H ₂ O ₂	[148]
Other GPCR agonists	Rat tail artery Human coronary artery Rat or mouse pulmonary artery Rat carotid artery	Lucigenin DHE	Increased superoxide	[113, 114, 135, 138, 178]
Perfusion pressure & stretch	Mouse afferent arterioles Rat femoral artery Bovine coronary artery Bovine pulmonary artery	DHE Lucigenin ethidium bromide	Increased superoxide	[162, 194, 334, 353]
	Mouse afferent arterioles	DCF	Increased H ₂ O ₂	[230, 327]
Flow/ shear-stress	Mouse cerebral arteries (endothelium in situ) Human coronary arterioles (endothelium in situ)	DCF	Increased H ₂ O ₂	[408, 418]
Flow overload Flow reversal	Porcine carotid artery	Electron paramagnetic resonance LO-12	Increased superoxide	[161, 422]
hypoxia	Mouse and porcine pulmonary artery	Cytochrome C reduction assay Lucigenin	Increased superoxide	[165, 650]
	Porcine, mouse and rat pulmonary artery	DCF	Increased H ₂ O ₂	[588, 589, 650]
stimulus	Species and vascular bed	indicator	effect	reference
VSMCs				
Angiotensin II	Mouse and rat Aortic VSMC Rat renal afferent arteriole VSMCs Human resistance vessel VSMCs Bovine pulmonary artery VSMCs	DHE, Electron spin resonance spectroscopy Lucigenin Cytochrome C reduction	Increased superoxide	[72, 75, 80, 90, 95, 108, 133, 157, 163, 169]
	Rat aortic VSMCs Human aortic VSMC Human resistance vessel VSMCs Human coronary artery SMCs Human VSMC cell line	DCF Amplex red	Increased H ₂ O ₂	[26, 45, 77, 79, 93, 96, 100, 102, 107, 170]
Prostanoids and eicosanoids	Bovine pulmonary artery SMCs Rat aortic VSMCs Mouse and bovine pulmonary VSMC	Cytochrome C reduction assay DHE lucigenin	Increased superoxide	[58, 142, 147-149]
	Rat aortic VSMCs Rat pulmonary VSMC	DCF	Increased H ₂ O ₂	[145, 146, 651]

Other GPCR agonists	Rat and mouse aortic VSMCs Rat cerebral artery SMCs	Lucigenin DHE L012	Increased superoxide	[109, 117, 125, 126, 152]
	Rat and mouse aortic VSMCs Human coronary artery myocytes	DCF	Increased H ₂ O ₂	[116, 121, 130, 136, 225]
Cytokines and growth-factors	Mouse, rat and human aortic VSMC	Lucigenin, DHE, LO-12 H2FF-BSA (endosomal specific) Electron spin resonance	Increased superoxide	[126, 163, 222, 229, 252, 255]
Cytokines and growth-factors	Mouse and rat aortic VSMC Human pulmonary VSMC	DCF Homovanillic acid assay FAT-HyPer (focal adhesion targeted H ₂ O ₂ probe)	Increased H ₂ O ₂	[130, 163, 187, 203, 210, 267]
stimulus	Species and vascular bed	indicator	effect	reference
Endothelium				
Angiotensin II	Mouse aortic ECs Human skin microvascular ECs Human coronary microvascular ECs	DHE Electron spin resonance lucigenin	Increased superoxide	[55, 72, 97]
Vasodilator agonists (acetylcholine, histamine, bradykinin)	Mouse mesenteric ECs Rat femoral artery ECs Human umbilical vein ECs Human and bovine aortic ECs Bovine pulmonary ECs Porcine aortic ECs	DCF	Increased H ₂ O ₂	[237, 402, 405, 416, 417]
Other GPCR agonists	HMEC-1 (cell line) Human skin microvascular ECs	lucigenin DHE	Increased superoxide	[55, 124, 144, 232]
Laminar shear-stress	Rat basilar artery ECs Bovine aortic ECs	DHE Cytochrome C assay Electron spin resonance	Increased superoxide	[291, 419-421]
	Bovine aortic ECs	DCF, pHyper	Increased H ₂ O ₂	[414]
Laminar shear-stress (prolonged)	Mouse aortic ECs	DHE, electron spin resonance	Decreased superoxide	[443]
oscillatory shear-stress or cessation of flow	Human umbilical vein ECs Mouse and bovine aortic ECs	DHE electron spin resonance	Increased superoxide	[414, 419, 431, 443, 652]
	Mouse lung capillary ECs Mouse pulmonary artery ECs Mouse and bovine aortic ECs	DCF Amplex red	Increased H ₂ O ₂	[431, 432, 444]
	Mouse and human pulmonary ECs Human umbilical vein ECs	Fe ²⁺ oxidation xylenol orange Amplex Red	Increased H ₂ O ₂	[52, 158, 221, 223]

Table 2. Effects of antioxidants and NOX inhibition on vascular smooth muscle contraction responses to GPCR agonists, pressure and hypoxia and on relaxation responses to GPCR agonists and flow/shear stress. Note that drugs commonly used as non-selective NOX inhibitors, particularly diphenyleneiodonium (DPI) and apocynin are no longer considered selective enough for NOX and may indeed simply be acting as antioxidants [337], so any results with these drugs should be supported with the use of more selective drugs or specific NOX isoform siRNA or knockout. NaC = N-acetyl cysteine. SOD = superoxide dismutase.

Stimulus	Species and vascular bed	Inhibitor and effect	Target	Inferred role of Target	Reference
Pro-contractile GPCR agonists					
Angiotensin II-induced contraction	Rabbit aorta Rat cerebral Mouse mesenteric Mouse renal afferent arterioles Rat pulmonary artery	Tiron, Tempol, PEG-catalase, SOD-tg, apocynin DPI p47 ^{phox} -/- NOX1 siRNA Gp91ds-tat All inhibit contraction	ROS NOX1 NOX2	ROS, NO1 and NOX2 = Pro-contractile	[89, 99, 104, 105, 133, 325, 335, 377, 378]
Phenylephrine-induced contraction	Mouse aorta Rat Tail artery	Tiron, NaC, apocynin, VAS2870 catalase, Poldip2 ^{-/-} all inhibit contraction	ROS NOX4	ROS and NOX4 = Pro-contractile	[131, 135, 226]
ET-1-induced contraction	Rat pulmonary artery Mouse mesenteric artery	Tiron N-Acetyl cysteine Both inhibit contraction	ROS	ROS = Pro-contractile	[66, 114]
Sphingosine phosphorylcholine-induced Contraction	Rat mesenteric artery	Tempol, VAS2870 -inhibit p47 ^{phox} -/- -inhibits but gp91 ^{phox} -/- has no effect	ROS NOX1	NOX1 = Pro-contractile but NOX2 is not	[148]
U46619-induced contraction	Bovine coronary artery Mouse mesenteric artery Rat pulmonary artery	NaC PEG-catalase, Tempol, ebselen, apocynin, DPI gp91-ds-tat, NOX2 ^{-/-} , p47 ^{phox} -/- All inhibit contraction	ROS NOX2	ROS and NOX2 = Pro-contractile	[66, 140, 146, 174]
Adenosine-induced contraction	Mouse aorta	Apocynin, PEG-SOD/Cat All inhibit contraction	ROS NOX	NOX = Pro-contractile	[116]
5-HT-induced contraction	pulmonary (mouse)	SOD, apocynin Both inhibit contraction but gp91 ^{phox} -/- has no effect	Superoxide Not NOX2	Superoxide = Pro-contractile Not NOX2-derived	[138]
Stretch/pressure					
Myogenic contraction	Mouse renal afferent arterioles	PEG-SOD, tempol, ebselen, apocynin, p47 ^{phox} -/- All inhibit contraction	superoxide NOX1 or 2	NOX1/2 = Pro-contractile	[99, 104, 230, 327, 334]
		POLDIP2 ^{-/-} . PEG-catalase Both inhibit contraction	NOX4, H ₂ O ₂	NOX4-derived H ₂ O ₂ = Pro-contractile	[230, 327]
Stretch induced contraction	Bovine coronary artery Bovine pulmonary artery	Apocynin, DPI, ebselen All inhibit contraction	ROS NOX	NOX = Pro-contractile	[194, 353]
Hypoxia					
Hypoxic pulmonary vasoconstriction	Mouse pulmonary artery	Apocynin, p47 ^{-/-} P47 translocation inhibitor	NOX, ROS	NOX contributes to HPV	[165]
	Porcine and guinea-pig pulmonary artery	SOD inhibits DETCA (SOD inhibitor) potentiates	superoxide	Superoxide contributes to HPV	[650, 653]
	Mouse and bovine pulmonary artery	gp91 ^{phox} -/- no effect p47 ^{phox} -/- partial inhibition	NOX1	NOX1 but not NOX2 contributes to HPV	[578]
		NOX2 siRNA –no effect NOX4 siRNA –partial inhibition	NOX2 NOX4	NOX4 contributes to HPV bit NOX2 does not	[579]
Stimulus	Species and vascular bed	Inhibitor and effect	Target	Inferred role of Target	Reference

Pro-relaxant GPCR agonists					
Acetylcholine-induced relaxation	Rat aorta Mouse mesenteric artery Cat cerebral artery Dog coronary artery Rat femoral artery	SOD restores impaired relaxation Catalase, eNOS ^{-/-} inhibit relaxation	H ₂ O ₂	H ₂ O ₂ = Pro-relaxant (eNOS-derived) H ₂ O ₂ = Pro-relaxant	[102, 400, 402, 405, 437, 462]
	Mouse mesenteric artery	NOX1 ^{-/-} improved relaxation	NOX1, ROS	NOX1 = Pro-contractile	[424]
	Aorta from renovascular hypertensive mice	NOX2 ^{-/-} , gp91ds-tat, Tiron, PEG-SOD All restore impaired relaxation	NOX2	NOX2 = pro-contractile	[425]
	Perfused cerebral artery from mouse	PEG-cat/gp91ds-tat Both inhibit relaxation	NOX2/ H ₂ O ₂	NOX2 = Pro-relaxant	[418]
	Mouse aorta and perfused heart	Endothelium-specific NOX4 over-expression enhances relaxation and reduces perfusion pressure	NOX4, H ₂ O ₂ (not superoxide)	NOX4 = Pro-relaxant	[215]
	Mouse mesenteric artery	Human NOX5 expression in VSM Inhibits relaxation	NOX5	NOX5 = Pro-contractile	[66]
Bradykinin-induced relaxation	Perfused mouse mesenteric artery Human mesenteric artery	SOD ^{-/-} inhibits increased flow)Catalase inhibits hyper-polarisation and relaxation	H ₂ O ₂	H ₂ O ₂ = Pro-relaxant	[404, 438]
Adenosine-induced relaxation	Rat cerebral artery	PEG-SOD, PEG-Cat	ROS	ROS = pro-relaxant	[117]
Histamine-induced relaxation	Mouse aorta and perfused heart	EC-specific NOX4 over-expression enhances relaxation response and reduces perfusion pressure	H ₂ O ₂	NOX4 = Pro-relaxant H ₂ O ₂ = Pro-relaxant	[215]
Flow/ shear-stress					
Flow/ shear-stress Induced relaxation	Human coronary artery Rat mesenteric artery Human coronary arterioles Mouse cerebral arteries	Catalase, PEG-catalase gp91ds-tat All inhibit relaxation	H ₂ O ₂	H ₂ O ₂ = pro-relaxant NOX2 = pro-relaxant	[408, 410, 411, 418]

Table 3. Effects of exogenous reactive oxygen species (ROS) on vascular tone. Pro-contractile and pro-relaxant effects of H₂O₂ or superoxide (generated extracellularly with xanthine/xanthine oxidase (X/XO) or intracellularly with LY83583) on arterial preparations *in vitro* or blood flow *in vivo*.

Pro-contractile effects of ROS					
ROS stimulus	Pre-constriction?	Vascular bed	Bath conc.	effect	reference
Superoxide (X/XO, LY83583)	None	Rat aorta and pulmonary artery	1-400μM	Causes contraction	[311, 339, 340]
	PGF _{2α} U46619/L-NAME	Rat pulmonary artery, mesenteric artery and femoral artery	1-10μM	Enhances pre-constriction	[174, 311, 351]
	Pressure/myogenic	Mouse renal afferent arteriole Rat cerebral artery	0.1-250μM	Enhances pre-constriction	[325, 327]
	KCl	Rat pulmonary and mesenteric artery	1-10μM	Enhances pre-constriction	[148, 351]
	Permeabilised, pCa 6.9	Rat pulmonary artery	10μM	Enhances pre-constriction	[311]
H ₂ O ₂	None	Rat and mouse aorta Rat mesenteric artery Bovine coronary artery Canine cerebral artery SHR carotid artery Mouse carotid artery Rat, rabbit & porcine pulmonary artery	1μM – 10mM	Causes contraction	[131, 146, 194, 283, 326, 340-342, 342, 343, 343-348, 515]
	phenylephrine	Rat mesenteric artery	10-100μM	Enhances pre-constriction	[350]
	Pressure/myogenic	Rat mesenteric artery	100μM	Enhances pre-constriction	[343]
	KCl	Mouse aorta Mouse mesenteric artery Mouse carotid artery Bovine pulmonary artery	1μM-1mM	Enhances pre-constriction	[341, 341, 342, 342, 342, 352-354]
Pro-Relaxant Effects of ROS					
ROS stimulus	Pre-constriction?	Vascular bed	Bath conc.	effect	reference
Superoxide (LY83583, menadione, X/XO)	PGF _{2α} U46619/L-NAME Phenylephrine	Rat mesenteric and femoral artery Rat aorta	10-50μM	Causes relaxation	[311, 333, 351]
	Myogenic	Cerebral circulation <i>in vivo</i> (rat)	200μM	Suppresses myogenic auto-regulation	[396]
H ₂ O ₂	U46619 PGF _{2α} ET-1 Noradrenaline Phenylephrine	Mouse, rat and human mesenteric artery Porcine coronary artery Human coronary arteriole Rat femoral artery Rabbit iliac artery	1μM-10mM	Causes relaxation	[336, 350, 354, 400-407]
	Pressure	Coronary circulation <i>in vivo</i>	1-300μM	Increases blood flow	[336]
	Myogenic	Mouse renal afferent arteriole Human coronary arterioles Mouse mesenteric artery Rat coronary artery	1μM-1mM 25μM	Causes relaxation	[327, 334, 354, 408-411]
	Hypoxia	Guinea-pig pulmonary artery	10-300μM	Causes relaxation	[653]

Table 4. Effects of antioxidants and NOX inhibition on VSM growth/proliferative/migration & associated signalling in response to mitogenic and hypertensive stimuli. DPI = diphenyleneiodonium. NaC = N-acetyl cysteine. SOD = superoxide dismutase.

stimulus	Species and vascular bed	Inhibitor and effect	Target	Inferred role of target	reference
Proliferation					
Angiotensin II induced proliferation	Mouse and rat aortic VSMC Human resistance artery VSMCs	Apocynin, DPI P22 ^{phox} siRNA NOX1 siRNA NOX1-/- NOX2 siRNA All inhibit proliferation	NOX1/2	NOX1 and NOX2 = pro-proliferative	[79, 107, 490, 560]
	Aorta (mouse)	NOX4 over-expression Does NOT influence proliferation	NOX4/ H ₂ O ₂	NOX4 is not pro-proliferative	[215]
Other GPCR agonists (Thrombin, ATP, Urotensin II, 5-HT) Induce proliferation	Rat and human aortic VSMC Rat pulmonary VSMC	NaC DPI, apocynin NOXA1 shRNA NOX1-/- NOX4 siRNA All inhibit proliferation	NOX NOXA1	NOXA1 = pro-proliferative NOX1 = pro-proliferative NOX4 = pro-proliferative	[118, 121, 126, 127, 597, 645]
Growth factors (PDGF, TGF-β) Induce proliferation	Rat and human aortic VSMCs Human pulmonary VSMC	NaC, catalase NOX4 siRNA NOX5 siRNA All inhibit proliferation	ROS NOX5	NOX4 = pro-proliferative NOX5 = pro-proliferative	[187, 210, 267, 317]
Chronic hypoxia Induced VSMC proliferation and/or vascular remodelling	Mouse, human Pulmonary vasculature or VSMCs	NOX1-/- NOX2-/- NOX4 siRNA GKT137831 All inhibit pulmonary media thickening or induce apoptosis	NOX1 NOX2 NOX4	NOX1, NOX2 and NOX4 promote pulmonary vascular remodelling and PH	[59, 592, 599, 635, 644] [639]
	Mouse pulmonary vasculature	NOX4-/- No effect on pulmonary vascular remodelling	NOX4	NOX4 NOT involved in vascular remodelling in PH	[59, 647]
Monocrotaline induced VSMC proliferation	Rat pulmonary VSMC	NOX1 siRNA VCC588646 and VCC202273 All inhibit proliferation	NOX1 NOX4	NOX1 and NOX4 promote VSMC proliferation and remodelling	[593, 595]
Other models of PAH Induced VSMC proliferation	Human pulmonary VSMC Lamb pulmonary VSMC Rat pulmonary VSMC	NOX4 siRNA GKT137831 Inhibit proliferation	NOX4	NOX4 = pro-proliferative	[59, 210, 629, 642]
RTK transactivation and ERK phosphorylation					
Angiotensin II	Rat aortic VSMCs	Nac, Tiron, α-tocopherol, DPI P22 ^{phox} siRNA All inhibit EGFR transactivation and ERK phosphorylation	NOX, ROS	NOX = pro-proliferative	[90, 95, 103, 274, 560]
Other GPCR agonists ET-1, thrombin, phenylephrine, Lyso-phosphatidyl choline, Sphingosine 1-phosphate, urotensin II Induced EGFR transactivation and/or ERK phosphorylation	Rat and mouse aortic VSMCs	NAC, Tiron catalase, DPI, apocynin dom neg p47 NOX1-/- and NOX1 siRNA All inhibit EGFR transactivation and ERK phosphorylation	ROS NOX1	NOX1 = pro-proliferative	[64, 109, 110, 121, 125, 130, 136, 147, 149, 156]
Growth factors (PDGF, TGF-β) Induce ERK phosphorylation	Rat aortic VSMCs Human pulmonary VSMC	Catalase or NaC NOX4 siRNA Inhibit ERK phosphorylation	NOX4	NOX4 = pro-proliferative	[210, 267]
stretch	Bovine coronary artery	Ebselen, DPI, apocynin Inhibit EGFR transactivation and ERK phosphorylation	ROS/NOX	NOX = pro-proliferative	[194]
VSM Hypertrophy					
Angiotensin II Induced VSMC hypertrophy	Rat and mouse aortic VSMCs	NaC, DPI, p22 ^{phox} antisense NOX1 siRNA All inhibit hypertrophy	NOX1, 2 or 4	NOX1/2/4 promote growth	[89, 96, 157, 169]

		p22 ^{phox} overexpression in VSMC Enhances hypertrophy	ROS/NOX1/2 /4	NOX1/2/4 promote growth	[552]
PI3K/AKT signalling and protein synthesis					
Angiotensin II	Human resistance artery VSMCs Rat aortic VSMCs	DPI NOX1 siRNA Inhibit AKT phosphorylation and protein synthesis	NOX1	NOX1 promotes growth	[79, 107]
Other GPCR agonists ET-1, thrombin, phenylephrine, Sphingosine 1-phosphate	Rat aortic VSMCs	NAC, ebselen, Apocynin, DPI NOXA1 shRNA Inhibit PI3K activation, AKT activation and protein synthesis	NOX1	NOX1 and NOXA1 promote growth	[109, 126, 136, 150]
VSMC Migration					
Angiotensin II Induced VSMC migration	Human coronary artery SMCs Rat aortic VSMCs	NAC, DPI P22 ^{phox} siRNA NOX1 siRNA Inhibit migration	NOX1	NOX1 stimulates migration	[100, 107, 560]
Other GPCR agonists (ET-1, thrombin, Sphingosine 1-phosphate, urokinase) Induced VSMC migration	Rat aortic VSMCs Human coronary artery	DPI, apocynin NOX1-/- (inhibits migration)	NOX1	NOX1 stimulates migration	[125, 147, 178]
Growth factors (PDGF, IGF, TNF- α) Induce VSMC migration	Rat and mouse aortic VSMCs	Catalase, NAC NOX1-/- NOX4 siRNA, POLDIP2 siRNA (inhibits focal adhesion turnover and migration)	NOX1 NOX4	NOX1 and NOX4 stimulate migration	[163, 222, 229, 267]

Reference List

- [1] Waypa G.B., Chandel N.S., Schumacker P.T. Model for hypoxic pulmonary vasoconstriction involving mitochondrial oxygen sensing. *Circ Res* ;**88**:1259-66;2001
- [2] Waypa G.B., Marks J.D., Guzy R., Mungai P.T., Schriewer J., Dokic D. et al. Hypoxia triggers subcellular compartmental redox signaling in vascular smooth muscle cells. *Circ Res* ;**106**:526-35;2010
- [3] Al G., I, Khoo N.K., Knaus U.G., Griendling K.K., Touyz R.M., Thannickal V.J. et al. Oxidases and peroxidases in cardiovascular and lung disease: new concepts in reactive oxygen species signaling. *Free Radic Biol Med* ;**51**:1271-88;2011
- [4] Murphy M.P. How mitochondria produce reactive oxygen species. *Biochem J* ;**417**:1-13;2009
- [5] Dikalova A.E., Gongora M.C., Harrison D.G., Lambeth J.D., Dikalov S., Griendling K.K. Upregulation of Nox1 in vascular smooth muscle leads to impaired endothelium-dependent relaxation via eNOS uncoupling. *Am J Physiol Heart Circ Physiol* ;**299**:H673-H679;2010
- [6] Zhang M., Song P., Xu J., Zou M.H. Activation of NAD(P)H oxidases by thromboxane A2 receptor uncouples endothelial nitric oxide synthase. *Arterioscler Thromb Vasc Biol* ;**31**:125-32;2011
- [7] Tejero J., Shiva S., Gladwin M.T. Sources of Vascular Nitric Oxide and Reactive Oxygen Species and Their Regulation. *Physiol Rev* ;**99**:311-79;2019
- [8] Griendling K.K., Ushio-Fukai M. NADH/NADPH Oxidase and Vascular Function. *Trends Cardiovasc Med* ;**7**:301-7;1997
- [9] Bedard K., Krause K.H. The NOX family of ROS-generating NADPH oxidases: physiology and pathophysiology. *Physiol Rev* ;**87**:245-313;2007
- [10] Bedard K., Jaquet V., Krause K.H. NOX5: from basic biology to signaling and disease. *Free Radic Biol Med* ;**52**:725-34;2012
- [11] Cardey B., Enescu M. Cysteine oxidation by the superoxide radical: a theoretical study. *Chemphyschem* ;**10**:1642-8;2009
- [12] Luo D., Smith S.W., Anderson B.D. Kinetics and mechanism of the reaction of cysteine and hydrogen peroxide in aqueous solution. *J Pharm Sci* ;**94**:304-16;2005
- [13] Hidalgo C., Donoso P. Crosstalk between calcium and redox signaling: from molecular mechanisms to health implications. *Antioxid Redox Signal* ;**10**:1275-312;2008
- [14] Griendling K.K., Touyz R.M., Zweier J.L., Dikalov S., Chilian W., Chen Y.R. et al. Measurement of Reactive Oxygen Species, Reactive Nitrogen Species, and Redox-Dependent Signaling in the Cardiovascular System: A Scientific Statement From the American Heart Association. *Circ Res* ;**119**:e39-e75;2016
- [15] Yoboue E.D., Sitia R., Simmen T. Redox crosstalk at endoplasmic reticulum (ER) membrane contact sites (MCS) uses toxic waste to deliver messages. *Cell Death Dis* ;**9**:331;2018

- [16] Kim Y., Jang H.H. Role of Cytosolic 2-Cys Prx1 and Prx2 in Redox Signaling. *Antioxidants (Basel)* ;**8**;2019
- [17] Moller M.N., Rios N., Trujillo M., Radi R., Denicola A., Alvarez B. Detection and quantification of nitric oxide-derived oxidants in biological systems. *J Biol Chem* ;2019
- [18] Sjoberg B., Foley S., Cardey B., Fromm M., Enescu M. Methionine oxidation by hydrogen peroxide in peptides and proteins: A theoretical and Raman spectroscopy study. *J Photochem Photobiol B* ;**188**:95-9;2018
- [19] Erickson J.R., Joiner M.L., Guan X., Kutschke W., Yang J., Oddis C.V. et al. A dynamic pathway for calcium-independent activation of CaMKII by methionine oxidation. *Cell* ;**133**:462-74;2008
- [20] Trevisani M., Siemens J., Materazzi S., Bautista D.M., Nassini R., Campi B. et al. 4-Hydroxynonenal, an endogenous aldehyde, causes pain and neurogenic inflammation through activation of the irritant receptor TRPA1. *Proc Natl Acad Sci U S A* ;**104**:13519-24;2007
- [21] Monteiro H.P., Arai R.J., Travassos L.R. Protein tyrosine phosphorylation and protein tyrosine nitration in redox signaling. *Antioxid Redox Signal* ;**10**:843-89;2008
- [22] Babior B.M. NADPH oxidase: an update. *Blood* ;**93**:1464-76;1999
- [23] Lassegue B., San M.A., Griending K.K. Biochemistry, physiology, and pathophysiology of NADPH oxidases in the cardiovascular system. *Circ Res* ;**110**:1364-90;2012
- [24] Santillo M., Colantuoni A., Mondola P., Guida B., Damiano S. NOX signaling in molecular cardiovascular mechanisms involved in the blood pressure homeostasis. *Front Physiol* ;**6**:194;2015
- [25] Biberstine-Kinkade K.J., DeLeo F.R., Epstein R.I., LeRoy B.A., Nauseef W.M., Dinanuer M.C. Heme-ligating histidines in flavocytochrome b(558): identification of specific histidines in gp91(phox). *J Biol Chem* ;**276**:31105-12;2001
- [26] Touyz R.M., Chen X., Tabet F., Yao G., He G., Quinn M.T. et al. Expression of a functionally active gp91phox-containing neutrophil-type NAD(P)H oxidase in smooth muscle cells from human resistance arteries: regulation by angiotensin II. *Circ Res* ;**90**:1205-13;2002
- [27] Meyer J.W., Holland J.A., Ziegler L.M., Chang M.M., Beebe G., Schmitt M.E. Identification of a functional leukocyte-type NADPH oxidase in human endothelial cells :a potential atherogenic source of reactive oxygen species. *Endothelium* ;**7**:11-22;1999
- [28] Lapouge K., Smith S.J., Groemping Y., Rittinger K. Architecture of the p40-p47-p67phox complex in the resting state of the NADPH oxidase. A central role for p67phox. *J Biol Chem* ;**277**:10121-8;2002
- [29] Fontayne A., Dang P.M., Gougerot-Pocidalo M.A., El-Benna J. Phosphorylation of p47phox sites by PKC alpha, beta II, delta, and zeta: effect on binding to p22phox and on NADPH oxidase activation. *Biochemistry* ;**41**:7743-50;2002

- [30] Belambri S.A., Rolas L., Raad H., Hurtado-Nedelec M., Dang P.M., El-Benna J. NADPH oxidase activation in neutrophils: Role of the phosphorylation of its subunits. *Eur J Clin Invest* ;e12951;2018
- [31] Kuribayashi F., Nunoi H., Wakamatsu K., Tsunawaki S., Sato K., Ito T. et al. The adaptor protein p40(phox) as a positive regulator of the superoxide-producing phagocyte oxidase. *EMBO J* ;**21**:6312-20;2002
- [32] Mizuki K., Takeya R., Kuribayashi F., Nobuhisa I., Kohda D., Nunoi H. et al. A region C-terminal to the proline-rich core of p47phox regulates activation of the phagocyte NADPH oxidase by interacting with the C-terminal SH3 domain of p67phox. *Arch Biochem Biophys* ;**444**:185-94;2005
- [33] Dahan I., Smith S.M., Pick E. A Cys-Gly-Cys triad in the dehydrogenase region of Nox2 plays a key role in the interaction with p67phox. *J Leukoc Biol* ;**98**:859-74;2015
- [34] Dang P.M., Babior B.M., Smith R.M. NADPH dehydrogenase activity of p67PHOX, a cytosolic subunit of the leukocyte NADPH oxidase. *Biochemistry* ;**38**:5746-53;1999
- [35] Ago T., Kuribayashi F., Hiroaki H., Takeya R., Ito T., Kohda D. et al. Phosphorylation of p47phox directs phox homology domain from SH3 domain toward phosphoinositides, leading to phagocyte NADPH oxidase activation. *Proc Natl Acad Sci U S A* ;**100**:4474-9;2003
- [36] Kanai F., Liu H., Field S.J., Akbary H., Matsuo T., Brown G.E. et al. The PX domains of p47phox and p40phox bind to lipid products of PI(3)K. *Nat Cell Biol* ;**3**:675-8;2001
- [37] Bagaitkar J., Barbu E.A., Perez-Zapata L.J., Austin A., Huang G., Pallat S. et al. PI(3)P-p40phox binding regulates NADPH oxidase activation in mouse macrophages and magnitude of inflammatory responses in vivo. *J Leukoc Biol* ;**101**:449-57;2017
- [38] Kao Y.Y., Gianni D., Bohl B., Taylor R.M., Bokoch G.M. Identification of a conserved Rac-binding site on NADPH oxidases supports a direct GTPase regulatory mechanism. *J Biol Chem* ;**283**:12736-46;2008
- [39] Raad H., Paclet M.H., Boussetta T., Kroviarski Y., Morel F., Quinn M.T. et al. Regulation of the phagocyte NADPH oxidase activity: phosphorylation of gp91phox/NOX2 by protein kinase C enhances its diaphorase activity and binding to Rac2, p67phox, and p47phox. *FASEB J* ;**23**:1011-22;2009
- [40] Lassegue B., Sorescu D., Szocs K., Yin Q., Akers M., Zhang Y. et al. Novel gp91(phox) homologues in vascular smooth muscle cells : nox1 mediates angiotensin II-induced superoxide formation and redox-sensitive signaling pathways. *Circ Res* ;**88**:888-94;2001
- [41] Lassegue B., Clempus R.E. Vascular NAD(P)H oxidases: specific features, expression, and regulation. *Am J Physiol Regul Integr Comp Physiol* ;**285**:R277-R297;2003
- [42] Lyle A.N., Griendling K.K. Modulation of vascular smooth muscle signaling by reactive oxygen species. *Physiology (Bethesda)* ;**21**:269-80;2006
- [43] Nguyen Dinh C.A., Montezano A.C., Burger D., Touyz R.M. Angiotensin II, NADPH oxidase, and redox signaling in the vasculature. *Antioxid Redox Signal* ;**19**:1110-20;2013

- [44] Souza H.P., Laurindo F.R., Ziegelstein R.C., Berlowitz C.O., Zweier J.L. Vascular NAD(P)H oxidase is distinct from the phagocytic enzyme and modulates vascular reactivity control. *Am J Physiol Heart Circ Physiol* ;**280**:H658-H667;2001
- [45] Dikalov S.I., Dikalova A.E., Bikineyeva A.T., Schmidt H.H., Harrison D.G., Griendling K.K. Distinct roles of Nox1 and Nox4 in basal and angiotensin II-stimulated superoxide and hydrogen peroxide production. *Free Radic Biol Med* ;**45**:1340-51;2008
- [46] Petry A., Djordjevic T., Weitnauer M., Kietzmann T., Hess J., Gorlach A. NOX2 and NOX4 mediate proliferative response in endothelial cells. *Antioxid Redox Signal* ;**8**:1473-84;2006
- [47] Clempus R.E., Sorescu D., Dikalova A.E., Pounkova L., Jo P., Sorescu G.P. et al. Nox4 is required for maintenance of the differentiated vascular smooth muscle cell phenotype. *Arterioscler Thromb Vasc Biol* ;**27**:42-8;2007
- [48] Hilenski L.L., Clempus R.E., Quinn M.T., Lambeth J.D., Griendling K.K. Distinct subcellular localizations of Nox1 and Nox4 in vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol* ;**24**:677-83;2004
- [49] Vukelic S., Xu Q., Seidel-Rogol B., Faidley E.A., Dikalova A.E., Hilenski L.L. et al. NOX4 (NADPH Oxidase 4) and Poldip2 (Polymerase delta-Interacting Protein 2) Induce Filamentous Actin Oxidation and Promote Its Interaction With Vinculin During Integrin-Mediated Cell Adhesion. *Arterioscler Thromb Vasc Biol* ;**38**:2423-34;2018
- [50] Wingler K., Wunsch S., Kreutz R., Rothermund L., Paul M., Schmidt H.H. Upregulation of the vascular NAD(P)H-oxidase isoforms Nox1 and Nox4 by the renin-angiotensin system in vitro and in vivo. *Free Radic Biol Med* ;**31**:1456-64;2001
- [51] Chen K., Kirber M.T., Xiao H., Yang Y., Keaney J.F., Jr. Regulation of ROS signal transduction by NADPH oxidase 4 localization. *J Cell Biol* ;**181**:1129-39;2008
- [52] Schroder K., Zhang M., Benkhoff S., Mieth A., Pliquett R., Kosowski J. et al. Nox4 is a protective reactive oxygen species generating vascular NADPH oxidase. *Circ Res* ;**110**:1217-25;2012
- [53] van Buul J.D., Fernandez-Borja M., Anthony E.C., Hordijk P.L. Expression and localization of NOX2 and NOX4 in primary human endothelial cells. *Antioxid Redox Signal* ;**7**:308-17;2005
- [54] Guzik T.J., Chen W., Gongora M.C., Guzik B., Lob H.E., Mangalat D. et al. Calcium-dependent NOX5 nicotinamide adenine dinucleotide phosphate oxidase contributes to vascular oxidative stress in human coronary artery disease. *J Am Coll Cardiol* ;**52**:1803-9;2008
- [55] Montezano A.C., Burger D., Paravicini T.M., Chignalia A.Z., Yusuf H., Almasri M. et al. Nicotinamide adenine dinucleotide phosphate reduced oxidase 5 (Nox5) regulation by angiotensin II and endothelin-1 is mediated via calcium/calmodulin-dependent, rac-1-independent pathways in human endothelial cells. *Circ Res* ;**106**:1363-73;2010
- [56] Montezano A.C., Tsiropoulou S., Dulak-Lis M., Harvey A., Camargo L.L., Touyz R.M. Redox signaling, Nox5 and vascular remodeling in hypertension. *Curr Opin Nephrol Hypertens* ;**24**:425-33;2015

- [57] Pandey D., Patel A., Patel V., Chen F., Qian J., Wang Y. et al. Expression and functional significance of NADPH oxidase 5 (Nox5) and its splice variants in human blood vessels. *Am J Physiol Heart Circ Physiol* ;**302**:H1919-H1928;2012
- [58] Chakraborti S., Chowdhury A., Kar P., Das P., Shaikh S., Roy S. et al. Role of protein kinase C in NADPH oxidase derived O₂⁻-mediated regulation of KV-LVOCC axis under U46619 induced increase in [Ca²⁺]_i in pulmonary smooth muscle cells. *Arch Biochem Biophys* ;**487**:123-30;2009
- [59] Hood K.Y., Montezano A.C., Harvey A.P., Nilsen M., MacLean M.R., Touyz R.M. Nicotinamide Adenine Dinucleotide Phosphate Oxidase-Mediated Redox Signaling and Vascular Remodeling by 16alpha-Hydroxyestrone in Human Pulmonary Artery Cells: Implications in Pulmonary Arterial Hypertension. *Hypertension* ;**68**:796-808;2016
- [60] Li J.M., Shah A.M. Intracellular localization and preassembly of the NADPH oxidase complex in cultured endothelial cells. *J Biol Chem* ;**277**:19952-60;2002
- [61] Modlinger P., Chabrashvili T., Gill P.S., Mendonca M., Harrison D.G., Griendling K.K. et al. RNA silencing in vivo reveals role of p22phox in rat angiotensin slow pressor response. *Hypertension* ;**47**:238-44;2006
- [62] Herkert O., Diebold I., Brandes R.P., Hess J., Busse R., Gorlach A. NADPH oxidase mediates tissue factor-dependent surface procoagulant activity by thrombin in human vascular smooth muscle cells. *Circulation* ;**105**:2030-6;2002
- [63] Hanna I.R., Hilenski L.L., Dikalova A., Taniyama Y., Dikalov S., Lyle A. et al. Functional association of nox1 with p22phox in vascular smooth muscle cells. *Free Radic Biol Med* ;**37**:1542-9;2004
- [64] Schuchardt M., Prufer J., Prufer N., Wiedon A., Huang T., Chebli M. et al. The endothelium-derived contracting factor uridine adenosine tetraphosphate induces P2Y(2)-mediated pro-inflammatory signaling by monocyte chemoattractant protein-1 formation. *J Mol Med (Berl)* ;**89**:799-810;2011
- [65] Ellmark S.H., Dusting G.J., Fui M.N., Guzzo-Pernell N., Drummond G.R. The contribution of Nox4 to NADPH oxidase activity in mouse vascular smooth muscle. *Cardiovasc Res* ;**65**:495-504;2005
- [66] Montezano A.C., De Lucca C.L., Persson P., Rios F.J., Harvey A.P., Anagnostopoulou A. et al. NADPH Oxidase 5 Is a Pro-Contractile Nox Isoform and a Point of Cross-Talk for Calcium and Redox Signaling-Implications in Vascular Function. *J Am Heart Assoc* ;**7**;2018
- [67] Gupte S.A., Kaminski P.M., Floyd B., Agarwal R., Ali N., Ahmad M. et al. Cytosolic NADPH may regulate differences in basal Nox oxidase-derived superoxide generation in bovine coronary and pulmonary arteries. *Am J Physiol Heart Circ Physiol* ;**288**:H13-H21;2005
- [68] Santos C.X., Nabeebaccus A.A., Shah A.M., Camargo L.L., Filho S.V., Lopes L.R. Endoplasmic reticulum stress and Nox-mediated reactive oxygen species signaling in the peripheral vasculature: potential role in hypertension. *Antioxid Redox Signal* ;**20**:121-34;2014
- [69] Amanso A.M., Debbas V., Laurindo F.R. Proteasome inhibition represses unfolded protein response and Nox4, sensitizing vascular cells to endoplasmic reticulum stress-induced death. *PLoS One* ;**6**:e14591;2011

- [70] Camargo L.L., Harvey A.P., Rios F.J., Tsiropoulou S., Da Silva R.N.O., Cao Z. et al. Vascular Nox (NADPH Oxidase) Compartmentalization, Protein Hyperoxidation, and Endoplasmic Reticulum Stress Response in Hypertension. *Hypertension* ;**72**:235-46;2018
- [71] Wassmann S., Laufs U., Baumer A.T., Muller K., Konkol C., Sauer H. et al. Inhibition of geranylgeranylation reduces angiotensin II-mediated free radical production in vascular smooth muscle cells: involvement of angiotensin AT1 receptor expression and Rac1 GTPase. *Mol Pharmacol* ;**59**:646-54;2001
- [72] Landmesser U., Cai H., Dikalov S., McCann L., Hwang J., Jo H. et al. Role of p47(phox) in vascular oxidative stress and hypertension caused by angiotensin II. *Hypertension* ;**40**:511-5;2002
- [73] Cruzado M.C., Risler N.R., Miatello R.M., Yao G., Schiffrin E.L., Touyz R.M. Vascular smooth muscle cell NAD(P)H oxidase activity during the development of hypertension: Effect of angiotensin II and role of insulinlike growth factor-1 receptor transactivation. *Am J Hypertens* ;**18**:81-7;2005
- [74] Garrido A.M., Griendling K.K. NADPH oxidases and angiotensin II receptor signaling. *Mol Cell Endocrinol* ;**302**:148-58;2009
- [75] Griendling K.K., Minieri C.A., Ollerenshaw J.D., Alexander R.W. Angiotensin II stimulates NADH and NADPH oxidase activity in cultured vascular smooth muscle cells. *Circ Res* ;**74**:1141-8;1994
- [76] Griendling K.K., Ushio-Fukai M., Lassegue B., Alexander R.W. Angiotensin II signaling in vascular smooth muscle. New concepts. *Hypertension* ;**29**:366-73;1997
- [77] Seshiah P.N., Weber D.S., Rocic P., Valppu L., Taniyama Y., Griendling K.K. Angiotensin II stimulation of NAD(P)H oxidase activity: upstream mediators. *Circ Res* ;**91**:406-13;2002
- [78] Tabet F., Schiffrin E.L., Callera G.E., He Y., Yao G., Ostman A. et al. Redox-sensitive signaling by angiotensin II involves oxidative inactivation and blunted phosphorylation of protein tyrosine phosphatase SHP-2 in vascular smooth muscle cells from SHR. *Circ Res* ;**103**:149-58;2008
- [79] Touyz R.M., Schiffrin E.L. Ang II-stimulated superoxide production is mediated via phospholipase D in human vascular smooth muscle cells. *Hypertension* ;**34**:976-82;1999
- [80] Touyz R.M., Schiffrin E.L. Increased generation of superoxide by angiotensin II in smooth muscle cells from resistance arteries of hypertensive patients: role of phospholipase D-dependent NAD(P)H oxidase-sensitive pathways. *J Hypertens* ;**19**:1245-54;2001
- [81] Touyz R.M., Tabet F., Schiffrin E.L. Redox-dependent signalling by angiotensin II and vascular remodelling in hypertension. *Clin Exp Pharmacol Physiol* ;**30**:860-6;2003
- [82] Touyz R.M., Yao G., Schiffrin E.L. c-Src induces phosphorylation and translocation of p47phox: role in superoxide generation by angiotensin II in human vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol* ;**23**:981-7;2003
- [83] Touyz R.M. Reactive oxygen species and angiotensin II signaling in vascular cells -- implications in cardiovascular disease. *Braz J Med Biol Res* ;**37**:1263-73;2004

- [84] Touyz R.M., Yao G., Viel E., Amiri F., Schiffrin E.L. Angiotensin II and endothelin-1 regulate MAP kinases through different redox-dependent mechanisms in human vascular smooth muscle cells. *J Hypertens* ;**22**:1141-9;2004
- [85] Touyz R.M., Yao G., Quinn M.T., Pagano P.J., Schiffrin E.L. p47phox associates with the cytoskeleton through cortactin in human vascular smooth muscle cells: role in NAD(P)H oxidase regulation by angiotensin II. *Arterioscler Thromb Vasc Biol* ;**25**:512-8;2005
- [86] Ushio-Fukai M., Zafari A.M., Fukui T., Ishizaka N., Griendling K.K. p22phox is a critical component of the superoxide-generating NADH/NADPH oxidase system and regulates angiotensin II-induced hypertrophy in vascular smooth muscle cells. *J Biol Chem* ;**271**:23317-21;1996
- [87] Viridis A., Neves M.F., Amiri F., Touyz R.M., Schiffrin E.L. Role of NAD(P)H oxidase on vascular alterations in angiotensin II-infused mice. *J Hypertens* ;**22**:535-42;2004
- [88] Viridis A., Duranti E., Taddei S. Oxidative Stress and Vascular Damage in Hypertension: Role of Angiotensin II. *Int J Hypertens* ;**2011**:916310;2011
- [89] Al G., I, Meijles D.N., Mutchler S., Zhang Q., Sahoo S., Gorelova A. et al. Binding of EBP50 to Nox organizing subunit p47phox is pivotal to cellular reactive species generation and altered vascular phenotype. *Proc Natl Acad Sci U S A* ;**113**:E5308-E5317;2016
- [90] Bourmoum M., Charles R., Claing A. The GTPase ARF6 Controls ROS Production to Mediate Angiotensin II-Induced Vascular Smooth Muscle Cell Proliferation. *PLoS One* ;**11**:e0148097;2016
- [91] Hashad A.M., Sancho M., Brett S.E., Welsh D.G. Reactive Oxygen Species Mediate the Suppression of Arterial Smooth Muscle T-type Ca(2+) Channels by Angiotensin II. *Sci Rep* ;**8**:3445;2018
- [92] Zhang X., Yang J., Yu X., Cheng S., Gan H., Xia Y. Angiotensin II-Induced Early and Late Inflammatory Responses Through NOXs and MAPK Pathways. *Inflammation* ;**40**:154-65;2017
- [93] Dechend R., Viedt C., Muller D.N., Ugele B., Brandes R.P., Wallukat G. et al. AT1 receptor agonistic antibodies from preeclamptic patients stimulate NADPH oxidase. *Circulation* ;**107**:1632-9;2003
- [94] Berry C., Hamilton C.A., Brosnan M.J., Magill F.G., Berg G.A., McMurray J.J. et al. Investigation into the sources of superoxide in human blood vessels: angiotensin II increases superoxide production in human internal mammary arteries. *Circulation* ;**101**:2206-12;2000
- [95] Touyz R.M., Cruzado M., Tabet F., Yao G., Salomon S., Schiffrin E.L. Redox-dependent MAP kinase signaling by Ang II in vascular smooth muscle cells: role of receptor tyrosine kinase transactivation. *Can J Physiol Pharmacol* ;**81**:159-67;2003
- [96] Lee H., Ham S.A., Kim M.Y., Kim J.H., Paek K.S., Kang E.S. et al. Activation of PPARdelta counteracts angiotensin II-induced ROS generation by inhibiting rac1 translocation in vascular smooth muscle cells. *Free Radic Res* ;**46**:912-9;2012
- [97] Li J.M., Shah A.M. Mechanism of endothelial cell NADPH oxidase activation by angiotensin II. Role of the p47phox subunit. *J Biol Chem* ;**278**:12094-100;2003

- [98] Rey F.E., Cifuentes M.E., Kiarash A., Quinn M.T., Pagano P.J. Novel competitive inhibitor of NAD(P)H oxidase assembly attenuates vascular O₂(-) and systolic blood pressure in mice. *Circ Res* ;**89**:408-14;2001
- [99] Carlstrom M., Lai E.Y., Ma Z., Steege A., Patzak A., Eriksson U.J. et al. Superoxide dismutase 1 limits renal microvascular remodeling and attenuates arteriole and blood pressure responses to angiotensin II via modulation of nitric oxide bioavailability. *Hypertension* ;**56**:907-13;2010
- [100] Yasunari K., Maeda K., Nakamura M., Yoshikawa J. Pressure promotes angiotensin II--mediated migration of human coronary smooth muscle cells through increase in oxidative stress. *Hypertension* ;**39**:433-7;2002
- [101] Callera G.E., Antunes T.T., He Y., Montezano A.C., Yogi A., Savoia C. et al. c-Src Inhibition Improves Cardiovascular Function but not Remodeling or Fibrosis in Angiotensin II-Induced Hypertension. *Hypertension* ;**68**:1179-90;2016
- [102] Rajagopalan S., Kurz S., Munzel T., Tarpey M., Freeman B.A., Griendling K.K. et al. Angiotensin II-mediated hypertension in the rat increases vascular superoxide production via membrane NADH/NADPH oxidase activation. Contribution to alterations of vasomotor tone. *J Clin Invest* ;**97**:1916-23;1996
- [103] Frank G.D., Eguchi S., Yamakawa T., Tanaka S., Inagami T., Motley E.D. Involvement of reactive oxygen species in the activation of tyrosine kinase and extracellular signal-regulated kinase by angiotensin II. *Endocrinology* ;**141**:3120-6;2000
- [104] Lai E.Y., Solis G., Luo Z., Carlstrom M., Sandberg K., Holland S. et al. p47(phox) is required for afferent arteriolar contractile responses to angiotensin II and perfusion pressure in mice. *Hypertension* ;**59**:415-20;2012
- [105] Chaplin N.L., Amberg G.C. Hydrogen peroxide mediates oxidant-dependent stimulation of arterial smooth muscle L-type calcium channels. *Am J Physiol Cell Physiol* ;**302**:C1382-C1393;2012
- [106] Montezano A.C., Callera G.E., Yogi A., He Y., Tostes R.C., He G. et al. Aldosterone and angiotensin II synergistically stimulate migration in vascular smooth muscle cells through c-Src-regulated redox-sensitive RhoA pathways. *Arterioscler Thromb Vasc Biol* ;**28**:1511-8;2008
- [107] Moraes J.A., Frony A.C., Dias A.M., Renovato-Martins M., Rodrigues G., Marcinkiewicz C. et al. Alpha1beta1 and integrin-linked kinase interact and modulate angiotensin II effects in vascular smooth muscle cells. *Atherosclerosis* ;**243**:477-85;2015
- [108] Chakraborti S., Sarkar J., Chakraborti T. Role of PLD-PKCzeta signaling axis in p47phox phosphorylation for activation of NADPH oxidase by angiotensin II in pulmonary artery smooth muscle cells. *Cell Biol Int* ;**43**:678-94;2019
- [109] Daou G.B., Srivastava A.K. Reactive oxygen species mediate Endothelin-1-induced activation of ERK1/2, PKB, and Pyk2 signaling, as well as protein synthesis, in vascular smooth muscle cells. *Free Radic Biol Med* ;**37**:208-15;2004

- [110] Banes-Berceli A.K., Ogobi S., Tawfik A., Patel B., Shirley A., Pollock D.M. et al. Endothelin-1 activation of JAK2 in vascular smooth muscle cells involves NAD(P)H oxidase-derived reactive oxygen species. *Vascul Pharmacol* ;**43**:310-9;2005
- [111] Wedgwood S., Dettman R.W., Black S.M. ET-1 stimulates pulmonary arterial smooth muscle cell proliferation via induction of reactive oxygen species. *Am J Physiol Lung Cell Mol Physiol* ;**281**:L1058-L1067;2001
- [112] Meyer M.R., Barton M., Prossnitz E.R. Functional heterogeneity of NADPH oxidase-mediated contractions to endothelin with vascular aging. *Life Sci* ;**118**:226-31;2014
- [113] Li L., Fink G.D., Watts S.W., Northcott C.A., Galligan J.J., Pagano P.J. et al. Endothelin-1 increases vascular superoxide via endothelin(A)-NADPH oxidase pathway in low-renin hypertension. *Circulation* ;**107**:1053-8;2003
- [114] Jernigan N.L., Walker B.R., Resta T.C. Reactive oxygen species mediate RhoA/Rho kinase-induced Ca²⁺ sensitization in pulmonary vascular smooth muscle following chronic hypoxia. *Am J Physiol Lung Cell Mol Physiol* ;**295**:L515-L529;2008
- [115] Carlstrom M., Lai E.Y., Ma Z., Patzak A., Brown R.D., Persson A.E. Role of NOX2 in the regulation of afferent arteriole responsiveness. *Am J Physiol Regul Integr Comp Physiol* ;**296**:R72-R79;2009
- [116] El-Awady M.S., Ansari H.R., Fil D., Tilley S.L., Mustafa S.J. NADPH oxidase pathway is involved in aortic contraction induced by A3 adenosine receptor in mice. *J Pharmacol Exp Ther* ;**338**:711-7;2011
- [117] Gebremedhin D., Weinberger B., Lourim D., Harder D.R. Adenosine can mediate its actions through generation of reactive oxygen species. *J Cereb Blood Flow Metab* ;**30**:1777-90;2010
- [118] Lee I.T., Lin C.C., Wang C.H., Cherng W.J., Wang J.S., Yang C.M. ATP stimulates PGE(2)/cyclin D1-dependent VSMCs proliferation via STAT3 activation: role of PKCs-dependent NADPH oxidase/ROS generation. *Biochem Pharmacol* ;**85**:954-64;2013
- [119] Diebold I., Petry A., Burger M., Hess J., Gorlach A. NOX4 mediates activation of FoxO3a and matrix metalloproteinase-2 expression by urotensin-II. *Mol Biol Cell* ;**22**:4424-34;2011
- [120] Djordjevic T., BelAiba R.S., Bonello S., Pfeilschifter J., Hess J., Gorlach A. Human urotensin II is a novel activator of NADPH oxidase in human pulmonary artery smooth muscle cells. *Arterioscler Thromb Vasc Biol* ;**25**:519-25;2005
- [121] Tsai C.S., Loh S.H., Liu J.C., Lin J.W., Chen Y.L., Chen C.H. et al. Urotensin II-induced endothelin-1 expression and cell proliferation via epidermal growth factor receptor transactivation in rat aortic smooth muscle cells. *Atherosclerosis* ;**206**:86-94;2009
- [122] Brandes R.P., Viedt C., Nguyen K., Beer S., Kreuzer J., Busse R. et al. Thrombin-induced MCP-1 expression involves activation of the p22phox-containing NADPH oxidase in human vascular smooth muscle cells. *Thromb Haemost* ;**85**:1104-10;2001
- [123] Gorlach A., Diebold I., Schini-Kerth V.B., Berchner-Pfannschmidt U., Roth U., Brandes R.P. et al. Thrombin activates the hypoxia-inducible factor-1 signaling pathway in vascular smooth muscle cells: Role of the p22(phox)-containing NADPH oxidase. *Circ Res* ;**89**:47-54;2001

- [124] Holland J.A., Meyer J.W., Chang M.M., O'Donnell R.W., Johnson D.K., Ziegler L.M. Thrombin stimulated reactive oxygen species production in cultured human endothelial cells. *Endothelium* ;**6**:113-21;1998
- [125] Jagadeesha D.K., Takapoo M., Banfi B., Bhalla R.C., Miller F.J., Jr. Nox1 transactivation of epidermal growth factor receptor promotes N-cadherin shedding and smooth muscle cell migration. *Cardiovasc Res* ;**93**:406-13;2012
- [126] Niu X.L., Madamanchi N.R., Vendrov A.E., Tchivilev I., Rojas M., Madamanchi C. et al. Nox activator 1: a potential target for modulation of vascular reactive oxygen species in atherosclerotic arteries. *Circulation* ;**121**:549-59;2010
- [127] Patterson C., Ruef J., Madamanchi N.R., Barry-Lane P., Hu Z., Horaist C. et al. Stimulation of a vascular smooth muscle cell NAD(P)H oxidase by thrombin. Evidence that p47(phox) may participate in forming this oxidase in vitro and in vivo. *J Biol Chem* ;**274**:19814-22;1999
- [128] Vendrov A.E., Madamanchi N.R., Niu X.L., Molnar K.C., Runge M., Szyndralewicz C. et al. NADPH oxidases regulate CD44 and hyaluronic acid expression in thrombin-treated vascular smooth muscle cells and in atherosclerosis. *J Biol Chem* ;**285**:26545-57;2010
- [129] Wang Z., Castresana M.R., Newman W.H. Reactive oxygen species-sensitive p38 MAPK controls thrombin-induced migration of vascular smooth muscle cells. *J Mol Cell Cardiol* ;**36**:49-56;2004
- [130] Miller F.J., Jr., Chu X., Stanic B., Tian X., Sharma R.V., Davisson R.L. et al. A differential role for endocytosis in receptor-mediated activation of Nox1. *Antioxid Redox Signal* ;**12**:583-93;2010
- [131] Ding Y., Winters A., Ding M., Graham S., Akopova I., Muallem S. et al. Reactive oxygen species-mediated TRPC6 protein activation in vascular myocytes, a mechanism for vasoconstrictor-regulated vascular tone. *J Biol Chem* ;**286**:31799-809;2011
- [132] Bleeke T., Zhang H., Madamanchi N., Patterson C., Faber J.E. Catecholamine-induced vascular wall growth is dependent on generation of reactive oxygen species. *Circ Res* ;**94**:37-45;2004
- [133] Jackson E.K., Gillespie D.G., Zhu C., Ren J., Zacharia L.C., Mi Z. Alpha2-adrenoceptors enhance angiotensin II-induced renal vasoconstriction: role for NADPH oxidase and RhoA. *Hypertension* ;**51**:719-26;2008
- [134] Nishio E., Watanabe Y. The involvement of reactive oxygen species and arachidonic acid in alpha 1-adrenoceptor-induced smooth muscle cell proliferation and migration. *Br J Pharmacol* ;**121**:665-70;1997
- [135] Tsai M.H., Jiang M.J. Reactive oxygen species are involved in regulating alpha1-adrenoceptor-activated vascular smooth muscle contraction. *J Biomed Sci* ;**17**:67;2010
- [136] Zhang H., Chalothorn D., Jackson L.F., Lee D.C., Faber J.E. Transactivation of epidermal growth factor receptor mediates catecholamine-induced growth of vascular smooth muscle. *Circ Res* ;**95**:989-97;2004

- [137] Just A., Olson A.J., Whitten C.L., Arendshorst W.J. Superoxide mediates acute renal vasoconstriction produced by angiotensin II and catecholamines by a mechanism independent of nitric oxide. *Am J Physiol Heart Circ Physiol* ;**292**:H83-H92;2007
- [138] Liu J.Q., Folz R.J. Extracellular superoxide enhances 5-HT-induced murine pulmonary artery vasoconstriction. *Am J Physiol Lung Cell Mol Physiol* ;**287**:L111-L118;2004
- [139] Didion S.P., Hathaway C.A., Faraci F.M. Superoxide levels and function of cerebral blood vessels after inhibition of CuZn-SOD. *Am J Physiol Heart Circ Physiol* ;**281**:H1697-H1703;2001
- [140] Gupte S.A., Kaminski P.M., George S., Kouznestova L., Olson S.C., Mathew R. et al. Peroxide generation by p47phox-Src activation of Nox2 has a key role in protein kinase C-induced arterial smooth muscle contraction. *Am J Physiol Heart Circ Physiol* ;**296**:H1048-H1057;2009
- [141] Katsuyama M., Fan C., Yabe-Nishimura C. NADPH oxidase is involved in prostaglandin F2alpha-induced hypertrophy of vascular smooth muscle cells: induction of NOX1 by PGF2alpha. *J Biol Chem* ;**277**:13438-42;2002
- [142] Rice K.M., Uddemarri S., Desai D.H., Morrison R.G., Harris R., Wright G.L. et al. PGF2alpha-associated vascular smooth muscle hypertrophy is ROS dependent and involves the activation of mTOR, p70S6k, and PTEN. *Prostaglandins Other Lipid Mediat* ;**85**:49-57;2008
- [143] Zhang M., Dong Y., Xu J., Xie Z., Wu Y., Song P. et al. Thromboxane receptor activates the AMP-activated protein kinase in vascular smooth muscle cells via hydrogen peroxide. *Circ Res* ;**102**:328-37;2008
- [144] Medhora M., Chen Y., Gruenloh S., Harland D., Bodiga S., Zielonka J. et al. 20-HETE increases superoxide production and activates NADPH oxidase in pulmonary artery endothelial cells. *Am J Physiol Lung Cell Mol Physiol* ;**294**:L902-L911;2008
- [145] Singh N.K., Wang D., Kundumani-Sridharan V., Van Q.D., Niu J., Rao G.N. 15-Lipoxygenase-1-enhanced Src-Janus kinase 2-signal transducer and activator of transcription 3 stimulation and monocyte chemoattractant protein-1 expression require redox-sensitive activation of epidermal growth factor receptor in vascular wall remodeling. *J Biol Chem* ;**286**:22478-88;2011
- [146] Cogolludo A., Frazziano G., Cobeno L., Moreno L., Lodi F., Villamor E. et al. Role of reactive oxygen species in Kv channel inhibition and vasoconstriction induced by TP receptor activation in rat pulmonary arteries. *Ann N Y Acad Sci* ;**1091**:41-51;2006
- [147] Roztocil E., Nicholl S.M., Davies M.G. Sphingosine-1-phosphate-induced oxygen free radical generation in smooth muscle cell migration requires Galpha12/13 protein-mediated phospholipase C activation. *J Vasc Surg* ;**46**:1253-9;2007
- [148] Shaifta Y., Snetkov V.A., Prieto-Lloret J., Knock G.A., Smirnov S.V., Aaronson P.I. et al. Sphingosylphosphorylcholine potentiates vasoreactivity and voltage-gated Ca²⁺ entry via NOX1 and reactive oxygen species. *Cardiovasc Res* ;**106**:121-30;2015
- [149] Tanimoto T., Lungu A.O., Berk B.C. Sphingosine 1-phosphate transactivates the platelet-derived growth factor beta receptor and epidermal growth factor receptor in vascular smooth muscle cells. *Circ Res* ;**94**:1050-8;2004

- [150] Roztocil E., Nicholl S.M., Davies M.G. Mechanisms of sphingosine-1-phosphate-induced akt-dependent smooth muscle cell migration. *Surgery* ;**145**:34-41;2009
- [151] Warboys C.M., Toh H.B., Fraser P.A. Role of NADPH oxidase in retinal microvascular permeability increase by RAGE activation. *Invest Ophthalmol Vis Sci* ;**50**:1319-28;2009
- [152] Csanyi G., Yao M., Rodriguez A.I., Al G., I, Sharifi-Sanjani M., Frazziano G. et al. Thrombospondin-1 regulates blood flow via CD47 receptor-mediated activation of NADPH oxidase 1. *Arterioscler Thromb Vasc Biol* ;**32**:2966-73;2012
- [153] Kurrasch-Orbaugh D.M., Parrish J.C., Watts V.J., Nichols D.E. A complex signaling cascade links the serotonin_{2A} receptor to phospholipase A₂ activation: the involvement of MAP kinases. *J Neurochem* ;**86**:980-91;2003
- [154] Touyz R.M., Schiffrin E.L. Measurement of phospholipase d activation in vascular smooth muscle cells. *Methods Mol Med* ;**51**:355-62;2001
- [155] Chakraborti S., Sarkar J., Chowdhury A., Chakraborti T. Role of ADP ribosylation factor6-Cytohesin1-PhospholipaseD signaling axis in U46619 induced activation of NADPH oxidase in pulmonary artery smooth muscle cell membrane. *Arch Biochem Biophys* ;**633**:1-14;2017
- [156] Yamakawa T., Tanaka S., Yamakawa Y., Kamei J., Numaguchi K., Motley E.D. et al. Lysophosphatidylcholine activates extracellular signal-regulated kinases 1/2 through reactive oxygen species in rat vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol* ;**22**:752-8;2002
- [157] Zafari A.M., Ushio-Fukai M., Minieri C.A., Akers M., Lassegue B., Griendling K.K. Arachidonic acid metabolites mediate angiotensin II-induced NADH/NADPH oxidase activity and hypertrophy in vascular smooth muscle cells. *Antioxid Redox Signal* ;**1**:167-79;1999
- [158] Frey R.S., Rahman A., Kefer J.C., Minshall R.D., Malik A.B. PKC ζ regulates TNF- α -induced activation of NADPH oxidase in endothelial cells. *Circ Res* ;**90**:1012-9;2002
- [159] Heinloth A., Heermeier K., Raff U., Wanner C., Galle J. Stimulation of NADPH oxidase by oxidized low-density lipoprotein induces proliferation of human vascular endothelial cells. *J Am Soc Nephrol* ;**11**:1819-25;2000
- [160] Holland J.A., Meyer J.W., Schmitt M.E., Sauro M.D., Johnson D.K., Abdul-Karim R.W. et al. Low-density lipoprotein stimulated peroxide production and endocytosis in cultured human endothelial cells: mechanisms of action. *Endothelium* ;**5**:191-207;1997
- [161] Choy J.S., Lu X., Yang J., Zhang Z.D., Kassab G.S. Endothelial actin depolymerization mediates NADPH oxidase-superoxide production during flow reversal. *Am J Physiol Heart Circ Physiol* ;**306**:H69-H77;2014
- [162] Ungvari Z., Csiszar A., Huang A., Kaminski P.M., Wolin M.S., Koller A. High pressure induces superoxide production in isolated arteries via protein kinase C-dependent activation of NAD(P)H oxidase. *Circulation* ;**108**:1253-8;2003
- [163] Streeter J., Schickling B.M., Jiang S., Stanic B., Thiel W.H., Gakhar L. et al. Phosphorylation of Nox1 regulates association with NoxA1 activation domain. *Circ Res* ;**115**:911-8;2014

- [164] Inoguchi T., Li P., Umeda F., Yu H.Y., Kakimoto M., Imamura M. et al. High glucose level and free fatty acid stimulate reactive oxygen species production through protein kinase C--dependent activation of NAD(P)H oxidase in cultured vascular cells. *Diabetes* ;**49**:1939-45;2000
- [165] Rathore R., Zheng Y.M., Niu C.F., Liu Q.H., Korde A., Ho Y.S. et al. Hypoxia activates NADPH oxidase to increase [ROS]_i and [Ca²⁺]_i through the mitochondrial ROS-PKCepsilon signaling axis in pulmonary artery smooth muscle cells. *Free Radic Biol Med* ;**45**:1223-31;2008
- [166] Li J.M., Wheatcroft S., Fan L.M., Kearney M.T., Shah A.M. Opposing roles of p47phox in basal versus angiotensin II-stimulated alterations in vascular O₂- production, vascular tone, and mitogen-activated protein kinase activation. *Circulation* ;**109**:1307-13;2004
- [167] Grote K., Flach I., Luchtefeld M., Akin E., Holland S.M., Drexler H. et al. Mechanical stretch enhances mRNA expression and proenzyme release of matrix metalloproteinase-2 (MMP-2) via NAD(P)H oxidase-derived reactive oxygen species. *Circ Res* ;**92**:e80-e86;2003
- [168] Liu S., Ma X., Gong M., Shi L., Lincoln T., Wang S. Glucose down-regulation of cGMP-dependent protein kinase I expression in vascular smooth muscle cells involves NAD(P)H oxidase-derived reactive oxygen species. *Free Radic Biol Med* ;**42**:852-63;2007
- [169] Lv P., Miao S.B., Shu Y.N., Dong L.H., Liu G., Xie X.L. et al. Phosphorylation of smooth muscle 22alpha facilitates angiotensin II-induced ROS production via activation of the PKCdelta-P47phox axis through release of PKCdelta and actin dynamics and is associated with hypertrophy and hyperplasia of vascular smooth muscle cells in vitro and in vivo. *Circ Res* ;**111**:697-707;2012
- [170] Shaw S., Wang X., Redd H., Alexander G.D., Isales C.M., Marrero M.B. High glucose augments the angiotensin II-induced activation of JAK2 in vascular smooth muscle cells via the polyol pathway. *J Biol Chem* ;**278**:30634-41;2003
- [171] Zhu L.H., Wang L., Wang D., Jiang H., Tang Q.Z., Yan L. et al. Puerarin attenuates high-glucose-and diabetes-induced vascular smooth muscle cell proliferation by blocking PKCbeta2/Rac1-dependent signaling. *Free Radic Biol Med* ;**48**:471-82;2010
- [172] Forbes L.V., Moss S.J., Segal A.W. Phosphorylation of p67phox in the neutrophil occurs in the cytosol and is independent of p47phox. *FEBS Lett* ;**449**:225-9;1999
- [173] Saito S., Frank G.D., Mifune M., Ohba M., Utsunomiya H., Motley E.D. et al. Ligand-independent trans-activation of the platelet-derived growth factor receptor by reactive oxygen species requires protein kinase C-delta and c-Src. *J Biol Chem* ;**277**:44695-700;2002
- [174] MacKay C.E., Shaifta Y., Snetkov V.V., Francois A.A., Ward J.P.T., Knock G.A. ROS-dependent activation of RhoA/Rho-kinase in pulmonary artery: Role of Src-family kinases and ARHGEF1. *Free Radic Biol Med* ;**110**:316-31;2017
- [175] Griendling K.K., Sorescu D., Lassegue B., Ushio-Fukai M. Modulation of protein kinase activity and gene expression by reactive oxygen species and their role in vascular physiology and pathophysiology. *Arterioscler Thromb Vasc Biol* ;**20**:2175-83;2000
- [176] Knock G.A., Ward J.P. Redox regulation of protein kinases as a modulator of vascular function. *Antioxid Redox Signal* ;**15**:1531-47;2011

- [177] MacKay C.E., Knock G.A. Control of vascular smooth muscle function by Src-family kinases and reactive oxygen species in health and disease. *J Physiol* ;**593**:3815-28;2015
- [178] Duru E.A., Fu Y., Davies M.G. Urokinase requires NAD(P)H oxidase to transactivate the epidermal growth factor receptor. *Surgery* ;**152**:879-85;2012
- [179] Chowdhury A.K., Watkins T., Parinandi N.L., Saatian B., Kleinberg M.E., Usatyuk P.V. et al. Src-mediated tyrosine phosphorylation of p47phox in hyperoxia-induced activation of NADPH oxidase and generation of reactive oxygen species in lung endothelial cells. *J Biol Chem* ;**280**:20700-11;2005
- [180] Jackson E.K., Gao L., Zhu C. Mechanism of the vascular angiotensin II/alpha2-adrenoceptor interaction. *J Pharmacol Exp Ther* ;**314**:1109-16;2005
- [181] Moyers J.S., Bouton A.H., Parsons S.J. The sites of phosphorylation by protein kinase C and an intact SH2 domain are required for the enhanced response to beta-adrenergic agonists in cells overexpressing c-src. *Mol Cell Biol* ;**13**:2391-400;1993
- [182] Igishi T., Gutkind J.S. Tyrosine kinases of the Src family participate in signaling to MAP kinase from both Gq and Gi-coupled receptors. *Biochem Biophys Res Commun* ;**244**:5-10;1998
- [183] Nino G., Hu A., Grunstein J.S., McDonough J., Kreiger P.A., Josephson M.B. et al. G Protein betagamma-subunit signaling mediates airway hyperresponsiveness and inflammation in allergic asthma. *PLoS One* ;**7**:e32078;2012
- [184] Li Y., Liu J., Zhan X. Tyrosine phosphorylation of cortactin is required for H₂O₂-mediated injury of human endothelial cells. *J Biol Chem* ;**275**:37187-93;2000
- [185] Usatyuk P.V., Romer L.H., He D., Parinandi N.L., Kleinberg M.E., Zhan S. et al. Regulation of hyperoxia-induced NADPH oxidase activation in human lung endothelial cells by the actin cytoskeleton and cortactin. *J Biol Chem* ;**282**:23284-95;2007
- [186] Patil S., Bunderson M., Wilham J., Black S.M. Important role for Rac1 in regulating reactive oxygen species generation and pulmonary arterial smooth muscle cell growth. *Am J Physiol Lung Cell Mol Physiol* ;**287**:L1314-L1322;2004
- [187] Kong G., Lee S., Kim K.S. Inhibition of rac1 reduces PDGF-induced reactive oxygen species and proliferation in vascular smooth muscle cells. *J Korean Med Sci* ;**16**:712-8;2001
- [188] Tabet F., Schiffrin E.L., Touyz R.M. Mitogen-activated protein kinase activation by hydrogen peroxide is mediated through tyrosine kinase-dependent, protein kinase C-independent pathways in vascular smooth muscle cells: upregulation in spontaneously hypertensive rats. *J Hypertens* ;**23**:2005-12;2005
- [189] Chiarugi P., Pani G., Giannoni E., Taddei L., Colavitti R., Raugei G. et al. Reactive oxygen species as essential mediators of cell adhesion: the oxidative inhibition of a FAK tyrosine phosphatase is required for cell adhesion. *J Cell Biol* ;**161**:933-44;2003
- [190] Gadepalli R., Singh N.K., Kundumani-Sridharan V., Heckle M.R., Rao G.N. Novel role of proline-rich nonreceptor tyrosine kinase 2 in vascular wall remodeling after balloon injury. *Arterioscler Thromb Vasc Biol* ;**32**:2652-61;2012

- [191] Frank G.D., Mifune M., Inagami T., Ohba M., Sasaki T., Higashiyama S. et al. Distinct mechanisms of receptor and nonreceptor tyrosine kinase activation by reactive oxygen species in vascular smooth muscle cells: role of metalloprotease and protein kinase C-delta. *Mol Cell Biol* ;**23**:1581-9;2003
- [192] Kirabo A., Kearns P.N., Jarajapu Y.P., Sasser J.M., Oh S.P., Grant M.B. et al. Vascular smooth muscle Jak2 mediates angiotensin II-induced hypertension via increased levels of reactive oxygen species. *Cardiovasc Res* ;**91**:171-9;2011
- [193] Meng D., Shi X., Jiang B.H., Fang J. Insulin-like growth factor-I (IGF-I) induces epidermal growth factor receptor transactivation and cell proliferation through reactive oxygen species. *Free Radic Biol Med* ;**42**:1651-60;2007
- [194] Oeckler R.A., Kaminski P.M., Wolin M.S. Stretch enhances contraction of bovine coronary arteries via an NAD(P)H oxidase-mediated activation of the extracellular signal-regulated kinase mitogen-activated protein kinase cascade. *Circ Res* ;**92**:23-31;2003
- [195] Bokemeyer D., Schmitz U., Kramer H.J. Angiotensin II-induced growth of vascular smooth muscle cells requires an Src-dependent activation of the epidermal growth factor receptor. *Kidney Int* ;**58**:549-58;2000
- [196] Kwon J., Lee S.R., Yang K.S., Ahn Y., Kim Y.J., Stadtman E.R. et al. Reversible oxidation and inactivation of the tumor suppressor PTEN in cells stimulated with peptide growth factors. *Proc Natl Acad Sci U S A* ;**101**:16419-24;2004
- [197] Baumer A.T., ten F.H., Sauer H., Wartenberg M., Kappert K., Schnabel P. et al. Phosphatidylinositol 3-kinase-dependent membrane recruitment of Rac-1 and p47phox is critical for alpha-platelet-derived growth factor receptor-induced production of reactive oxygen species. *J Biol Chem* ;**283**:7864-76;2008
- [198] Fleming I.N., Gray A., Downes C.P. Regulation of the Rac1-specific exchange factor Tiam1 involves both phosphoinositide 3-kinase-dependent and -independent components. *Biochem J* ;**351**:173-82;2000
- [199] Han J., Luby-Phelps K., Das B., Shu X., Xia Y., Mosteller R.D. et al. Role of substrates and products of PI 3-kinase in regulating activation of Rac-related guanosine triphosphatases by Vav. *Science* ;**279**:558-60;1998
- [200] Paulsen C.E., Truong T.H., Garcia F.J., Homann A., Gupta V., Leonard S.E. et al. Peroxide-dependent sulfenylation of the EGFR catalytic site enhances kinase activity. *Nat Chem Biol* ;**8**:57-64;2011
- [201] Mifune M., Ohtsu H., Suzuki H., Nakashima H., Brailoiu E., Dun N.J. et al. G protein coupling and second messenger generation are indispensable for metalloprotease-dependent, heparin-binding epidermal growth factor shedding through angiotensin II type-1 receptor. *J Biol Chem* ;**280**:26592-9;2005
- [202] Banfi B., Clark R.A., Steger K., Krause K.H. Two novel proteins activate superoxide generation by the NADPH oxidase NOX1. *J Biol Chem* ;**278**:3510-3;2003
- [203] Ambasta R.K., Schreiber J.G., Janiszewski M., Busse R., Brandes R.P. Noxa1 is a central component of the smooth muscle NADPH oxidase in mice. *Free Radic Biol Med* ;**41**:193-201;2006

- [204] Kawano M., Miyamoto K., Kaito Y., Sumimoto H., Tamura M. Noxa1 as a moderate activator of Nox2-based NADPH oxidase. *Arch Biochem Biophys* ;**519**:1-7;2012
- [205] Chen F., Pandey D., Chadli A., Catravas J.D., Chen T., Fulton D.J. Hsp90 regulates NADPH oxidase activity and is necessary for superoxide but not hydrogen peroxide production. *Antioxid Redox Signal* ;**14**:2107-19;2011
- [206] Madrigal-Matute J., Fernandez-Garcia C.E., Gomez-Guerrero C., Lopez-Franco O., Munoz-Garcia B., Egido J. et al. HSP90 inhibition by 17-DMAG attenuates oxidative stress in experimental atherosclerosis. *Cardiovasc Res* ;**95**:116-23;2012
- [207] Janiszewski M., Lopes L.R., Carmo A.O., Pedro M.A., Brandes R.P., Santos C.X. et al. Regulation of NAD(P)H oxidase by associated protein disulfide isomerase in vascular smooth muscle cells. *J Biol Chem* ;**280**:40813-9;2005
- [208] Gimenez M., Verissimo-Filho S., Wittig I., Schickling B.M., Hahner F., Schurmann C. et al. Redox Activation of Nox1 (NADPH Oxidase 1) Involves an Intermolecular Disulfide Bond Between Protein Disulfide Isomerase and p47(phox) in Vascular Smooth Muscle Cells. *Arterioscler Thromb Vasc Biol* ;**39**:224-36;2019
- [209] Breton-Romero R., Gonzalez de O.C., Romero N., Sanchez-Gomez F.J., de A.C., Porras A. et al. Critical role of hydrogen peroxide signaling in the sequential activation of p38 MAPK and eNOS in laminar shear stress. *Free Radic Biol Med* ;**52**:1093-100;2012
- [210] Sturrock A., Cahill B., Norman K., Huecksteadt T.P., Hill K., Sanders K. et al. Transforming growth factor-beta1 induces Nox4 NAD(P)H oxidase and reactive oxygen species-dependent proliferation in human pulmonary artery smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* ;**290**:L661-L673;2006
- [211] Nisimoto Y., Jackson H.M., Ogawa H., Kawahara T., Lambeth J.D. Constitutive NADPH-dependent electron transferase activity of the Nox4 dehydrogenase domain. *Biochemistry* ;**49**:2433-42;2010
- [212] Serrander L., Cartier L., Bedard K., Banfi B., Lardy B., Plastre O. et al. NOX4 activity is determined by mRNA levels and reveals a unique pattern of ROS generation. *Biochem J* ;**406**:105-14;2007
- [213] Martyn K.D., Frederick L.M., von L.K., Dinauer M.C., Knaus U.G. Functional analysis of Nox4 reveals unique characteristics compared to other NADPH oxidases. *Cell Signal* ;**18**:69-82;2006
- [214] von L.K., Noack D., Wood M.R., Friedman J.S., Knaus U.G. Structural insights into Nox4 and Nox2: motifs involved in function and cellular localization. *Mol Cell Biol* ;**30**:961-75;2010
- [215] Ray R., Murdoch C.E., Wang M., Santos C.X., Zhang M., Alom-Ruiz S. et al. Endothelial Nox4 NADPH oxidase enhances vasodilatation and reduces blood pressure in vivo. *Arterioscler Thromb Vasc Biol* ;**31**:1368-76;2011
- [216] Kuroda J., Nakagawa K., Yamasaki T., Nakamura K., Takeya R., Kuribayashi F. et al. The superoxide-producing NAD(P)H oxidase Nox4 in the nucleus of human vascular endothelial cells. *Genes Cells* ;**10**:1139-51;2005

- [217] Somers M.J., Burchfield J.S., Harrison D.G. Evidence for a NADH/NADPH oxidase in human umbilical vein endothelial cells using electron spin resonance. *Antioxid Redox Signal* ;**2**:779-87;2000
- [218] Ambasta R.K., Kumar P., Griendling K.K., Schmidt H.H., Busse R., Brandes R.P. Direct interaction of the novel Nox proteins with p22phox is required for the formation of a functionally active NADPH oxidase. *J Biol Chem* ;**279**:45935-41;2004
- [219] Zana M., Peterfi Z., Kovacs H.A., Toth Z.E., Enyedi B., Morel F. et al. Interaction between p22(phox) and Nox4 in the endoplasmic reticulum suggests a unique mechanism of NADPH oxidase complex formation. *Free Radic Biol Med* ;**116**:41-9;2018
- [220] Brandes R.P., Schroder K. Composition and functions of vascular nicotinamide adenine dinucleotide phosphate oxidases. *Trends Cardiovasc Med* ;**18**:15-9;2008
- [221] Hu T., Ramachandrarao S.P., Siva S., Valancius C., Zhu Y., Mahadev K. et al. Reactive oxygen species production via NADPH oxidase mediates TGF-beta-induced cytoskeletal alterations in endothelial cells. *Am J Physiol Renal Physiol* ;**289**:F816-F825;2005
- [222] Meng D., Lv D.D., Fang J. Insulin-like growth factor-I induces reactive oxygen species production and cell migration through Nox4 and Rac1 in vascular smooth muscle cells. *Cardiovasc Res* ;**80**:299-308;2008
- [223] Zhuang J., Jiang T., Lu D., Luo Y., Zheng C., Feng J. et al. NADPH oxidase 4 mediates reactive oxygen species induction of CD146 dimerization in VEGF signal transduction. *Free Radic Biol Med* ;**49**:227-36;2010
- [224] Gorin Y., Ricono J.M., Kim N.H., Bhandari B., Choudhury G.G., Abboud H.E. Nox4 mediates angiotensin II-induced activation of Akt/protein kinase B in mesangial cells. *Am J Physiol Renal Physiol* ;**285**:F219-F229;2003
- [225] Zhang F., Jin S., Yi F., Xia M., Dewey W.L., Li P.L. Local production of O₂⁻ by NAD(P)H oxidase in the sarcoplasmic reticulum of coronary arterial myocytes: cADPR-mediated Ca²⁺ regulation. *Cell Signal* ;**20**:637-44;2008
- [226] Sutliff R.L., Hilenski L.L., Amanso A.M., Parastatidis I., Dikalova A.E., Hansen L. et al. Polymerase delta interacting protein 2 sustains vascular structure and function. *Arterioscler Thromb Vasc Biol* ;**33**:2154-61;2013
- [227] Lee S., Gharavi N.M., Honda H., Chang I., Kim B., Jen N. et al. A role for NADPH oxidase 4 in the activation of vascular endothelial cells by oxidized phospholipids. *Free Radic Biol Med* ;**47**:145-51;2009
- [228] Lyle A.N., Deshpande N.N., Taniyama Y., Seidel-Rogol B., Pounkova L., Du P. et al. Poldip2, a novel regulator of Nox4 and cytoskeletal integrity in vascular smooth muscle cells. *Circ Res* ;**105**:249-59;2009
- [229] Datla S.R., McGrail D.J., Vukelic S., Huff L.P., Lyle A.N., Pounkova L. et al. Poldip2 controls vascular smooth muscle cell migration by regulating focal adhesion turnover and force polarization. *Am J Physiol Heart Circ Physiol* ;**307**:H945-H957;2014

- [230] Li L., Lai E.Y., Luo Z., Solis G., Griendling K.K., Taylor W.R. et al. Superoxide and hydrogen peroxide counterregulate myogenic contractions in renal afferent arterioles from a mouse model of chronic kidney disease. *Kidney Int* ;**92**:625-33;2017
- [231] Banfi B., Tirone F., Durussel I., Knisz J., Moskwa P., Molnar G.Z. et al. Mechanism of Ca²⁺ activation of the NADPH oxidase 5 (NOX5). *J Biol Chem* ;**279**:18583-91;2004
- [232] BelAiba R.S., Djordjevic T., Petry A., Diemer K., Bonello S., Banfi B. et al. NOX5 variants are functionally active in endothelial cells. *Free Radic Biol Med* ;**42**:446-59;2007
- [233] Chen F., Haigh S., Yu Y., Benson T., Wang Y., Li X. et al. Nox5 stability and superoxide production is regulated by C-terminal binding of Hsp90 and CO-chaperones. *Free Radic Biol Med* ;**89**:793-805;2015
- [234] Chen F., Barman S., Yu Y., Haigh S., Wang Y., Black S.M. et al. Caveolin-1 is a negative regulator of NADPH oxidase-derived reactive oxygen species. *Free Radic Biol Med* ;**73**:201-13;2014
- [235] Tirone F., Cox J.A. NADPH oxidase 5 (NOX5) interacts with and is regulated by calmodulin. *FEBS Lett* ;**581**:1202-8;2007
- [236] Tirone F., Radu L., Craescu C.T., Cox J.A. Identification of the binding site for the regulatory calcium-binding domain in the catalytic domain of NOX5. *Biochemistry* ;**49**:761-71;2010
- [237] Sakurada R., Odagiri K., Hakamata A., Kamiya C., Wei J., Watanabe H. Calcium Release from Endoplasmic Reticulum Involves Calmodulin-Mediated NADPH Oxidase-Derived Reactive Oxygen Species Production in Endothelial Cells. *Int J Mol Sci* ;**20**;2019
- [238] Jagnandan D., Church J.E., Banfi B., Stuehr D.J., Marrero M.B., Fulton D.J. Novel mechanism of activation of NADPH oxidase 5. calcium sensitization via phosphorylation. *J Biol Chem* ;**282**:6494-507;2007
- [239] Chen F., Yu Y., Haigh S., Johnson J., Lucas R., Stepp D.W. et al. Regulation of NADPH oxidase 5 by protein kinase C isoforms. *PLoS One* ;**9**:e88405;2014
- [240] Pandey D., Gratton J.P., Rafikov R., Black S.M., Fulton D.J. Calcium/calmodulin-dependent kinase II mediates the phosphorylation and activation of NADPH oxidase 5. *Mol Pharmacol* ;**80**:407-15;2011
- [241] Pandey D., Fulton D.J. Molecular regulation of NADPH oxidase 5 via the MAPK pathway. *Am J Physiol Heart Circ Physiol* ;**300**:H1336-H1344;2011
- [242] El J.A., Valente A.J., Lechleiter J.D., Gamez M.J., Pearson D.W., Nauseef W.M. et al. Novel redox-dependent regulation of NOX5 by the tyrosine kinase c-Abl. *Free Radic Biol Med* ;**44**:868-81;2008
- [243] Davis Volk A.P., Moreland J.G. ROS-containing endosomal compartments: implications for signaling. *Methods Enzymol* ;**535**:201-24;2014
- [244] Janssen-Heininger Y.M., Mossman B.T., Heintz N.H., Forman H.J., Kalyanaraman B., Finkel T. et al. Redox-based regulation of signal transduction: principles, pitfalls, and promises. *Free Radic Biol Med* ;**45**:1-17;2008

- [245] Oakley F.D., Abbott D., Li Q., Engelhardt J.F. Signaling components of redox active endosomes: the redoxosomes. *Antioxid Redox Signal* ;**11**:1313-33;2009
- [246] Ushio-Fukai M. Compartmentalization of redox signaling through NADPH oxidase-derived ROS. *Antioxid Redox Signal* ;**11**:1289-99;2009
- [247] Huff L.P., Kikuchi D.S., Faidley E., Forrester S.J., Tsai M.Z., Lassegue B. et al. Polymerase-delta-interacting protein 2 activates the RhoGEF epithelial cell transforming sequence 2 in vascular smooth muscle cells. *Am J Physiol Cell Physiol* ;**316**:C621-C631;2019
- [248] Al G., I, Frazziano G., Rodriguez A.I., Csanyi G., Maniar S., St Croix C.M. et al. Aquaporin 1, Nox1, and Ask1 mediate oxidant-induced smooth muscle cell hypertrophy. *Cardiovasc Res* ;**97**:134-42;2013
- [249] Bienert G.P., Moller A.L., Kristiansen K.A., Schulz A., Moller I.M., Schjoerring J.K. et al. Specific aquaporins facilitate the diffusion of hydrogen peroxide across membranes. *J Biol Chem* ;**282**:1183-92;2007
- [250] Tie L., Wang D., Shi Y., Li X. Aquaporins in Cardiovascular System. *Adv Exp Med Biol* ;**969**:105-13;2017
- [251] Hawkins B.J., Madesh M., Kirkpatrick C.J., Fisher A.B. Superoxide flux in endothelial cells via the chloride channel-3 mediates intracellular signaling. *Mol Biol Cell* ;**18**:2002-12;2007
- [252] Choi H., Stark R.J., Raja B.S., Dikalova A., Lamb F.S. Apoptosis signal-regulating kinase 1 activation by Nox1-derived oxidants is required for TNFalpha receptor endocytosis. *Am J Physiol Heart Circ Physiol* ;**316**:H1528-H1537;2019
- [253] Rathaus M., Bernheim J. Oxygen species in the microvascular environment: regulation of vascular tone and the development of hypertension. *Nephrol Dial Transplant* ;**17**:216-21;2002
- [254] Chu X., Filali M., Stanic B., Takapoo M., Sheehan A., Bhalla R. et al. A critical role for chloride channel-3 (ClC-3) in smooth muscle cell activation and neointima formation. *Arterioscler Thromb Vasc Biol* ;**31**:345-51;2011
- [255] Miller F.J., Jr., Filali M., Huss G.J., Stanic B., Chamseddine A., Barna T.J. et al. Cytokine activation of nuclear factor kappa B in vascular smooth muscle cells requires signaling endosomes containing Nox1 and ClC-3. *Circ Res* ;**101**:663-71;2007
- [256] Prinz N., Clemens N., Canisius A., Lackner K.J. Endosomal NADPH-oxidase is critical for induction of the tissue factor gene in monocytes and endothelial cells. Lessons from the antiphospholipid syndrome. *Thromb Haemost* ;**109**:525-31;2013
- [257] Oakley F.D., Smith R.L., Engelhardt J.F. Lipid rafts and caveolin-1 coordinate interleukin-1beta (IL-1beta)-dependent activation of NFkappaB by controlling endocytosis of Nox2 and IL-1beta receptor 1 from the plasma membrane. *J Biol Chem* ;**284**:33255-64;2009
- [258] Tsutsumi R., Harizanova J., Stockert R., Schroder K., Bastiaens P.I.H., Neel B.G. Assay to visualize specific protein oxidation reveals spatio-temporal regulation of SHP2. *Nat Commun* ;**8**:466;2017

- [259] Li Q., Zhang Y., Marden J.J., Banfi B., Engelhardt J.F. Endosomal NADPH oxidase regulates c-Src activation following hypoxia/reoxygenation injury. *Biochem J* ;**411**:531-41;2008
- [260] Shahin W.S., Engelhardt J.F. Isolation of Redox-Active Endosomes (Redoxosomes) and Assessment of NOX Activity. *Methods Mol Biol* ;**1982**:461-72;2019
- [261] Mumbengegwi D.R., Li Q., Li C., Bear C.E., Engelhardt J.F. Evidence for a superoxide permeability pathway in endosomal membranes. *Mol Cell Biol* ;**28**:3700-12;2008
- [262] Anilkumar N., San J.G., Sawyer I., Santos C.X., Sand C., Brewer A.C. et al. A 28-kDa splice variant of NADPH oxidase-4 is nuclear-localized and involved in redox signaling in vascular cells. *Arterioscler Thromb Vasc Biol* ;**33**:e104-e112;2013
- [263] Yoshida L.S., Tsunawaki S. Expression of NADPH oxidases and enhanced H₂O₂-generating activity in human coronary artery endothelial cells upon induction with tumor necrosis factor- α . *Int Immunopharmacol* ;**8**:1377-85;2008
- [264] Kuroda J., Ago T., Matsushima S., Zhai P., Schneider M.D., Sadoshima J. NADPH oxidase 4 (Nox4) is a major source of oxidative stress in the failing heart. *Proc Natl Acad Sci U S A* ;**107**:15565-70;2010
- [265] Frazziano G., Al G., I, Baust J., Shiva S., Champion H.C., Pagano P.J. Nox-derived ROS are acutely activated in pressure overload pulmonary hypertension: indications for a seminal role for mitochondrial Nox4. *Am J Physiol Heart Circ Physiol* ;**306**:H197-H205;2014
- [266] Tang H., Hao Q., Rutherford S.A., Low B., Zhao Z.J. Inactivation of SRC family tyrosine kinases by reactive oxygen species in vivo. *J Biol Chem* ;**280**:23918-25;2005
- [267] Sundaresan M., Yu Z.X., Ferrans V.J., Irani K., Finkel T. Requirement for generation of H₂O₂ for platelet-derived growth factor signal transduction. *Science* ;**270**:296-9;1995
- [268] Lee S.R., Kwon K.S., Kim S.R., Rhee S.G. Reversible inactivation of protein-tyrosine phosphatase 1B in A431 cells stimulated with epidermal growth factor. *J Biol Chem* ;**273**:15366-72;1998
- [269] Tanabe Y., Saito M., Ueno A., Nakamura M., Takeishi K., Nakayama K. Mechanical stretch augments PDGF receptor beta expression and protein tyrosine phosphorylation in pulmonary artery tissue and smooth muscle cells. *Mol Cell Biochem* ;**215**:103-13;2000
- [270] Lim Y., Lim S.T., Tomar A., Gardel M., Bernard-Trifilo J.A., Chen X.L. et al. PyK2 and FAK connections to p190Rho guanine nucleotide exchange factor regulate RhoA activity, focal adhesion formation, and cell motility. *J Cell Biol* ;**180**:187-203;2008
- [271] Knock G.A., Snetkov V.A., Shafta Y., Drndarski S., Ward J.P., Aaronson P.I. Role of src-family kinases in hypoxic vasoconstriction of rat pulmonary artery. *Cardiovasc Res* ;**80**:453-62;2008
- [272] Knock G.A., Shafta Y., Snetkov V.A., Vowles B., Drndarski S., Ward J.P. et al. Interaction between src family kinases and rho-kinase in agonist-induced Ca²⁺-sensitization of rat pulmonary artery. *Cardiovasc Res* ;**77**:570-9;2008
- [273] Di S.J., Semenchuk L.A., Lauer J. Vanadate-induced contraction of smooth muscle and enhanced protein tyrosine phosphorylation. *Arch Biochem Biophys* ;**304**:386-91;1993

- [274] Frank G.D., Eguchi S., Inagami T., Motley E.D. N-acetylcysteine inhibits angiotensin II-mediated activation of extracellular signal-regulated kinase and epidermal growth factor receptor. *Biochem Biophys Res Commun* ;**280**:1116-9;2001
- [275] Weber D.S., Taniyama Y., Rocic P., Seshiah P.N., Dechert M.A., Gerthoffer W.T. et al. Phosphoinositide-dependent kinase 1 and p21-activated protein kinase mediate reactive oxygen species-dependent regulation of platelet-derived growth factor-induced smooth muscle cell migration. *Circ Res* ;**94**:1219-26;2004
- [276] Oshikawa J., Urao N., Kim H.W., Kaplan N., Razvi M., McKinney R. et al. Extracellular SOD-derived H₂O₂ promotes VEGF signaling in caveolae/lipid rafts and post-ischemic angiogenesis in mice. *PLoS One* ;**5**:e10189;2010
- [277] Yogi A., Mercure C., Touyz J., Callera G.E., Montezano A.C., Aranha A.B. et al. Renal redox-sensitive signaling, but not blood pressure, is attenuated by Nox1 knockout in angiotensin II-dependent chronic hypertension. *Hypertension* ;**51**:500-6;2008
- [278] Mahadev K., Motoshima H., Wu X., Ruddy J.M., Arnold R.S., Cheng G. et al. The NAD(P)H oxidase homolog Nox4 modulates insulin-stimulated generation of H₂O₂ and plays an integral role in insulin signal transduction. *Mol Cell Biol* ;**24**:1844-54;2004
- [279] Laniyonu A.A., Saifeddine M., Yang S.G., Hollenberg M.D. Tyrosine kinase inhibitors and the contractile action of G-protein-linked vascular agonists. *Can J Physiol Pharmacol* ;**72**:1075-85;1994
- [280] Gonzalez-Pacheco F.R., Caramelo C., Castilla M.A., Deudero J.J., Arias J., Yague S. et al. Mechanism of vascular smooth muscle cells activation by hydrogen peroxide: role of phospholipase C gamma. *Nephrol Dial Transplant* ;**17**:392-8;2002
- [281] Konishi H., Tanaka M., Takemura Y., Matsuzaki H., Ono Y., Kikkawa U. et al. Activation of protein kinase C by tyrosine phosphorylation in response to H₂O₂. *Proc Natl Acad Sci U S A* ;**94**:11233-7;1997
- [282] Mehdi M.Z., Pandey N.R., Pandey S.K., Srivastava A.K. H₂O₂-induced phosphorylation of ERK1/2 and PKB requires tyrosine kinase activity of insulin receptor and c-Src. *Antioxid Redox Signal* ;**7**:1014-20;2005
- [283] Yang Z.W., Zheng T., Wang J., Zhang A., Altura B.T., Altura B.M. Hydrogen peroxide induces contraction and raises [Ca²⁺]_i in canine cerebral arterial smooth muscle: participation of cellular signaling pathways. *Naunyn Schmiedebergs Arch Pharmacol* ;**360**:646-53;1999
- [284] Minetti M., Mallozzi C., Di Stasi A.M. Peroxynitrite activates kinases of the src family and upregulates tyrosine phosphorylation signaling. *Free Radic Biol Med* ;**33**:744-54;2002
- [285] Peters G.H., Frimurer T.M., Olsen O.H. Electrostatic evaluation of the signature motif (H/V)CX5R(S/T) in protein-tyrosine phosphatases. *Biochemistry* ;**37**:5383-93;1998
- [286] Barford D., Flint A.J., Tonks N.K. Crystal structure of human protein tyrosine phosphatase 1B. *Science* ;**263**:1397-404;1994
- [287] Meng T.C., Fukada T., Tonks N.K. Reversible oxidation and inactivation of protein tyrosine phosphatases in vivo. *Mol Cell* ;**9**:387-99;2002

- [288] Salmeen A., Barford D. Functions and mechanisms of redox regulation of cysteine-based phosphatases. *Antioxid Redox Signal* ;**7**:560-77;2005
- [289] Salsman S.J., Hensley K., Floyd R.A. Sensitivity of protein tyrosine phosphatase activity to the redox environment, cytochrome C, and microperoxidase. *Antioxid Redox Signal* ;**7**:1078-88;2005
- [290] Tonks N.K. Redox redux: revisiting PTPs and the control of cell signaling. *Cell* ;**121**:667-70;2005
- [291] Sanchez-Gomez F.J., Calvo E., Breton-Romero R., Fierro-Fernandez M., Anilkumar N., Shah A.M. et al. NOX4-dependent Hydrogen peroxide promotes shear stress-induced SHP2 sulfenylation and eNOS activation. *Free Radic Biol Med* ;**89**:419-30;2015
- [292] Shimizu H., Toyama O., Shiota M., Kim-Mitsuyama S., Miyazaki H. Protein tyrosine phosphatase LMW-PTP exhibits distinct roles between vascular endothelial and smooth muscle cells. *J Recept Signal Transduct Res* ;**25**:19-33;2005
- [293] Shimizu H., Nakagawa Y., Murakami C., Aoki N., Kim-Mitsuyama S., Miyazaki H. Protein tyrosine phosphatase PTPepsilon negatively regulates PDGF beta-receptor signaling induced by high glucose and PDGF in vascular smooth muscle cells. *Am J Physiol Cell Physiol* ;**299**:C1144-C1152;2010
- [294] Baas A.S., Berk B.C. Differential activation of mitogen-activated protein kinases by H₂O₂ and O₂⁻ in vascular smooth muscle cells. *Circ Res* ;**77**:29-36;1995
- [295] Guilluy C., Bregeon J., Toumaniantz G., Rolli-Derkinderen M., Retailleau K., Loufrani L. et al. The Rho exchange factor Arhgef1 mediates the effects of angiotensin II on vascular tone and blood pressure. *Nat Med* ;**16**:183-90;2010
- [296] Shaifta Y., Irechukwu N., Prieto-Lloret J., MacKay C.E., Marchon K.A., Ward J.P. et al. Divergent modulation of Rho-kinase and Ca(2+) influx pathways by Src family kinases and focal adhesion kinase in airway smooth muscle. *Br J Pharmacol* ;**172**:5265-80;2015
- [297] Sato H., Sato M., Kanai H., Uchiyama T., Iso T., Ohyama Y. et al. Mitochondrial reactive oxygen species and c-Src play a critical role in hypoxic response in vascular smooth muscle cells. *Cardiovasc Res* ;**67**:714-22;2005
- [298] Ben Mahdi M.H., Andrieu V., Pasquier C. Focal adhesion kinase regulation by oxidative stress in different cell types. *IUBMB Life* ;**50**:291-9;2000
- [299] Berk B.C. Vascular smooth muscle growth: autocrine growth mechanisms. *Physiol Rev* ;**81**:999-1030;2001
- [300] Mehdi M.Z., Azar Z.M., Srivastava A.K. Role of receptor and nonreceptor protein tyrosine kinases in H₂O₂-induced PKB and ERK1/2 signaling. *Cell Biochem Biophys* ;**47**:1-10;2007
- [301] Roskoski R., Jr. Src kinase regulation by phosphorylation and dephosphorylation. *Biochem Biophys Res Commun* ;**331**:1-14;2005
- [302] Sun G., Ramdas L., Wang W., Vinci J., McMurray J., Budde R.J. Effect of autophosphorylation on the catalytic and regulatory properties of protein tyrosine kinase Src. *Arch Biochem Biophys* ;**397**:11-7;2002

- [303] Giannoni E., Buricchi F., Rauei G., Ramponi G., Chiarugi P. Intracellular reactive oxygen species activate Src tyrosine kinase during cell adhesion and anchorage-dependent cell growth. *Mol Cell Biol* ;**25**:6391-403;2005
- [304] Akhand A.A., Pu M., Senga T., Kato M., Suzuki H., Miyata T. et al. Nitric oxide controls src kinase activity through a sulfhydryl group modification-mediated Tyr-527-independent and Tyr-416-linked mechanism. *J Biol Chem* ;**274**:25821-6;1999
- [305] Heppner D.E., Dustin C.M., Liao C., Hristova M., Veith C., Little A.C. et al. Direct cysteine sulfenylation drives activation of the Src kinase. *Nat Commun* ;**9**:4522;2018
- [306] Jo A., Park H., Lee S.H., Ahn S.H., Kim H.J., Park E.M. et al. SHP-2 binds to caveolin-1 and regulates Src activity via competitive inhibition of CSK in response to H₂O₂ in astrocytes. *PLoS One* ;**9**:e91582;2014
- [307] Mills J.E., Whitford P.C., Shaffer J., Onuchic J.N., Adams J.A., Jennings P.A. A novel disulfide bond in the SH2 Domain of the C-terminal Src kinase controls catalytic activity. *J Mol Biol* ;**365**:1460-8;2007
- [308] Singh D.K., Kumar D., Siddiqui Z., Basu S.K., Kumar V., Rao K.V. The strength of receptor signaling is centrally controlled through a cooperative loop between Ca²⁺ and an oxidant signal. *Cell* ;**121**:281-93;2005
- [309] Azar Z.M., Mehdi M.Z., Srivastava A.K. Activation of insulin-like growth factor type-1 receptor is required for H₂O₂-induced PKB phosphorylation in vascular smooth muscle cells. *Can J Physiol Pharmacol* ;**84**:777-86;2006
- [310] Griending K.K., Ushio-Fukai M. Reactive oxygen species as mediators of angiotensin II signaling. *Regul Pept* ;**91**:21-7;2000
- [311] Knock G.A., Snetkov V.A., Shaifta Y., Connolly M., Drndarski S., Noah A. et al. Superoxide constricts rat pulmonary arteries via Rho-kinase-mediated Ca(2+) sensitization. *Free Radic Biol Med* ;**46**:633-42;2009
- [312] Rosado J.A., Redondo P.C., Salido G.M., Gomez-Arteta E., Sage S.O., Pariente J.A. Hydrogen peroxide generation induces pp60src activation in human platelets: evidence for the involvement of this pathway in store-mediated calcium entry. *J Biol Chem* ;**279**:1665-75;2004
- [313] Andreev J., Galisteo M.L., Kranenburg O., Logan S.K., Chiu E.S., Okigaki M. et al. Src and Pyk2 mediate G-protein-coupled receptor activation of epidermal growth factor receptor (EGFR) but are not required for coupling to the mitogen-activated protein (MAP) kinase signaling cascade. *J Biol Chem* ;**276**:20130-5;2001
- [314] Calalb M.B., Polte T.R., Hanks S.K. Tyrosine phosphorylation of focal adhesion kinase at sites in the catalytic domain regulates kinase activity: a role for Src family kinases. *Mol Cell Biol* ;**15**:954-63;1995
- [315] Srinivasan R., Forman S., Quinlan R.A., Ohanian J., Ohanian V. Regulation of contractility by Hsp27 and Hic-5 in rat mesenteric small arteries. *Am J Physiol Heart Circ Physiol* ;**294**:H961-H969;2008

- [316] Frank G.D., Eguchi S. Activation of tyrosine kinases by reactive oxygen species in vascular smooth muscle cells: significance and involvement of EGF receptor transactivation by angiotensin II. *Antioxid Redox Signal* ;**5**:771-80;2003
- [317] Jay D.B., Papaharalambus C.A., Seidel-Rogol B., Dikalova A.E., Lassegue B., Griending K.K. Nox5 mediates PDGF-induced proliferation in human aortic smooth muscle cells. *Free Radic Biol Med* ;**45**:329-35;2008
- [318] Vepa S., Scribner W.M., Parinandi N.L., English D., Garcia J.G., Natarajan V. Hydrogen peroxide stimulates tyrosine phosphorylation of focal adhesion kinase in vascular endothelial cells. *Am J Physiol* ;**277**:L150-L158;1999
- [319] Gozin A., Franzini E., Andrieu V., Da C.L., Rollet-Labelle E., Pasquier C. Reactive oxygen species activate focal adhesion kinase, paxillin and p130cas tyrosine phosphorylation in endothelial cells. *Free Radic Biol Med* ;**25**:1021-32;1998
- [320] Littler C.M., Morris K.G., Jr., Fagan K.A., McMurtry I.F., Messing R.O., Dempsey E.C. Protein kinase C-epsilon-null mice have decreased hypoxic pulmonary vasoconstriction. *Am J Physiol Heart Circ Physiol* ;**284**:H1321-H1331;2003
- [321] Steinberg S.F. Structural basis of protein kinase C isoform function. *Physiol Rev* ;**88**:1341-78;2008
- [322] Gopalakrishna R., Jaken S. Protein kinase C signaling and oxidative stress. *Free Radic Biol Med* ;**28**:1349-61;2000
- [323] Gopalakrishna R., Gundimeda U., Schiffman J.E., McNeill T.H. A direct redox regulation of protein kinase C isoenzymes mediates oxidant-induced neuritogenesis in PC12 cells. *J Biol Chem* ;**283**:14430-44;2008
- [324] Chakraborti S., Chakraborti T. Down-regulation of protein kinase C attenuates the oxidant hydrogen peroxide-mediated activation of phospholipase A2 in pulmonary vascular smooth muscle cells. *Cell Signal* ;**7**:75-83;1995
- [325] Amberg G.C., Earley S., Glapa S.A. Local regulation of arterial L-type calcium channels by reactive oxygen species. *Circ Res* ;**107**:1002-10;2010
- [326] Pourmahram G.E., Snetkov V.A., Shaifta Y., Drndarski S., Knock G.A., Aaronson P.I. et al. Constriction of pulmonary artery by peroxide: role of Ca²⁺ release and PKC. *Free Radic Biol Med* ;**45**:1468-76;2008
- [327] Li L., Lai E.Y., Wellstein A., Welch W.J., Wilcox C.S. Differential effects of superoxide and hydrogen peroxide on myogenic signaling, membrane potential, and contractions of mouse renal afferent arterioles. *Am J Physiol Renal Physiol* ;**310**:F1197-F1205;2016
- [328] Zhu L.J., Klutho P.J., Scott J.A., Xie L., Luczak E.D., Dibbern M.E. et al. Oxidative activation of the Ca(2+)/calmodulin-dependent protein kinase II (CaMKII) regulates vascular smooth muscle migration and apoptosis. *Vascul Pharmacol* ;**60**:75-83;2014
- [329] Hasegawa T., Bando A., Tsuchiya K., Abe S., Okamoto M., Kirima K. et al. Enzymatic and nonenzymatic formation of reactive oxygen species from 6-anilino-5,8-quinolinequinone. *Biochim Biophys Acta* ;**1670**:19-27;2004

- [330] Li X.A., Bianchi C., Sellke F.W. Rat aortic smooth muscle cell density affects activation of MAP kinase and Akt by menadione and PDGF homodimer BB. *J Surg Res* ;**100**:197-204;2001
- [331] Jimenez-Altayo F., Briones A.M., Giraldo J., Planas A.M., Salaices M., Vila E. Increased superoxide anion production by interleukin-1 β impairs nitric oxide-mediated relaxation in resistance arteries. *J Pharmacol Exp Ther* ;**316**:42-52;2006
- [332] Wilcox C.S. Oxidative stress and nitric oxide deficiency in the kidney: a critical link to hypertension? *Am J Physiol Regul Integr Comp Physiol* ;**289**:R913-R935;2005
- [333] Bae O.N., Lim K.M., Han J.Y., Jung B.I., Lee J.Y., Noh J.Y. et al. U-shaped dose response in vasomotor tone: a mixed result of heterogenic response of multiple cells to xenobiotics. *Toxicol Sci* ;**103**:181-90;2008
- [334] Lai E.Y., Wellstein A., Welch W.J., Wilcox C.S. Superoxide modulates myogenic contractions of mouse afferent arterioles. *Hypertension* ;**58**:650-6;2011
- [335] Pfister S.L., Nithipatikom K., Campbell W.B. Role of superoxide and thromboxane receptors in acute angiotensin II-induced vasoconstriction of rabbit vessels. *Am J Physiol Heart Circ Physiol* ;**300**:H2064-H2071;2011
- [336] Rogers P.A., Dick G.M., Knudson J.D., Focardi M., Bratz I.N., Swafford A.N., Jr. et al. H₂O₂-induced redox-sensitive coronary vasodilation is mediated by 4-aminopyridine-sensitive K⁺ channels. *Am J Physiol Heart Circ Physiol* ;**291**:H2473-H2482;2006
- [337] Heumuller S., Wind S., Barbosa-Sicard E., Schmidt H.H., Busse R., Schroder K. et al. Apocynin is not an inhibitor of vascular NADPH oxidases but an antioxidant. *Hypertension* ;**51**:211-7;2008
- [338] Schluter T., Steinbach A.C., Steffen A., Rettig R., Grisk O. Apocynin-induced vasodilation involves Rho kinase inhibition but not NADPH oxidase inhibition. *Cardiovasc Res* ;**80**:271-9;2008
- [339] Jin L., Ying Z., Webb R.C. Activation of Rho/Rho kinase signaling pathway by reactive oxygen species in rat aorta. *Am J Physiol Heart Circ Physiol* ;**287**:H1495-H1500;2004
- [340] Rhoades R.A., Packer C.S., Roepke D.A., Jin N., Meiss R.A. Reactive oxygen species alter contractile properties of pulmonary arterial smooth muscle. *Can J Physiol Pharmacol* ;**68**:1581-9;1990
- [341] Ardanaz N., Beierwaltes W.H., Pagano P.J. Comparison of H₂O₂-induced vasoconstriction in the abdominal aorta and mesenteric artery of the mouse. *Vascul Pharmacol* ;**47**:288-94;2007
- [342] Ardanaz N., Beierwaltes W.H., Pagano P.J. Distinct hydrogen peroxide-induced constriction in multiple mouse arteries: potential influence of vascular polarization. *Pharmacol Rep* ;**60**:61-7;2008
- [343] Pelaez N.J., Braun T.R., Paul R.J., Meiss R.A., Packer C.S. H₂O₂ mediates Ca²⁺- and MLC(20) phosphorylation-independent contraction in intact and permeabilized vascular muscle. *Am J Physiol Heart Circ Physiol* ;**279**:H1185-H1193;2000

- [344] Sotnikova R. Investigation of the mechanisms underlying H₂O₂-evoked contraction in the isolated rat aorta. *Gen Pharmacol* ;**31**:115-9;1998
- [345] Pelaez N.J., Osterhaus S.L., Mak A.S., Zhao Y., Davis H.W., Packer C.S. MAPK and PKC activity are not required for H₂O₂-induced arterial muscle contraction. *Am J Physiol Heart Circ Physiol* ;**279**:H1194-H1200;2000
- [346] Jin N., Rhoades R.A. Activation of tyrosine kinases in H₂O₂-induced contraction in pulmonary artery. *Am J Physiol* ;**272**:H2686-H2692;1997
- [347] Sheehan D.W., Giese E.C., Gugino S.F., Russell J.A. Characterization and mechanisms of H₂O₂-induced contractions of pulmonary arteries. *Am J Physiol* ;**264**:H1542-H1547;1993
- [348] Tabet F., Savoia C., Schiffrin E.L., Touyz R.M. Differential calcium regulation by hydrogen peroxide and superoxide in vascular smooth muscle cells from spontaneously hypertensive rats. *J Cardiovasc Pharmacol* ;**44**:200-8;2004
- [349] Okatani Y., Watanabe K., Wakatsuki A., Sagara Y. Vasospastic action of hydrogen peroxide in human umbilical artery: relation to protein kinase C and calcium influx. *Acta Obstet Gynecol Scand* ;**76**:897-902;1997
- [350] Gao Y.J., Hirota S., Zhang D.W., Janssen L.J., Lee R.M. Mechanisms of hydrogen-peroxide-induced biphasic response in rat mesenteric artery. *Br J Pharmacol* ;**138**:1085-92;2003
- [351] Snetkov V.A., Smirnov S.V., Kua J., Aaronson P.I., Ward J.P., Knock G.A. Superoxide differentially controls pulmonary and systemic vascular tone through multiple signalling pathways. *Cardiovasc Res* ;**89**:214-24;2011
- [352] Thakali K., Davenport L., Fink G.D., Watts S.W. Cyclooxygenase, p38 mitogen-activated protein kinase (MAPK), extracellular signal-regulated kinase MAPK, Rho kinase, and Src mediate hydrogen peroxide-induced contraction of rat thoracic aorta and vena cava. *J Pharmacol Exp Ther* ;**320**:236-43;2007
- [353] Oeckler R.A., Arcuino E., Ahmad M., Olson S.C., Wolin M.S. Cytosolic NADH redox and thiol oxidation regulate pulmonary arterial force through ERK MAP kinase. *Am J Physiol Lung Cell Mol Physiol* ;**288**:L1017-L1025;2005
- [354] Lucchesi P.A., Belmadani S., Matrougui K. Hydrogen peroxide acts as both vasodilator and vasoconstrictor in the control of perfused mouse mesenteric resistance arteries. *J Hypertens* ;**23**:571-9;2005
- [355] Vogel P.A., Yang X., Moss N.G., Arendshorst W.J. Superoxide enhances Ca²⁺ entry through L-type channels in the renal afferent arteriole. *Hypertension* ;**66**:374-81;2015
- [356] Ilatovskaya D.V., Blass G., Palygin O., Levchenko V., Pavlov T.S., Grzybowski M.N. et al. A NOX4/TRPC6 Pathway in Podocyte Calcium Regulation and Renal Damage in Diabetic Kidney Disease. *J Am Soc Nephrol* ;**29**:1917-27;2018
- [357] Muralidharan P., Cserne S.H., Ingley E., Hool L. Evidence for redox sensing by a human cardiac calcium channel. *Sci Rep* ;**6**:19067;2016

- [358] Yang L., Xu J., Minobe E., Yu L., Feng R., Kameyama A. et al. Mechanisms underlying the modulation of L-type Ca^{2+} channel by hydrogen peroxide in guinea pig ventricular myocytes. *J Physiol Sci* ;**63**:419-26;2013
- [359] Chiamvimonvat N., O'Rourke B., Kamp T.J., Kallen R.G., Hofmann F., Flockerzi V. et al. Functional consequences of sulfhydryl modification in the pore-forming subunits of cardiovascular Ca^{2+} and Na^{+} channels. *Circ Res* ;**76**:325-34;1995
- [360] Scragg J.L., Dallas M.L., Wilkinson J.A., Varadi G., Peers C. Carbon monoxide inhibits L-type Ca^{2+} channels via redox modulation of key cysteine residues by mitochondrial reactive oxygen species. *J Biol Chem* ;**283**:24412-9;2008
- [361] Iesaki T., Wolin M.S. Thiol oxidation activates a novel redox-regulated coronary vasodilator mechanism involving inhibition of Ca^{2+} influx. *Arterioscler Thromb Vasc Biol* ;**20**:2359-65;2000
- [362] Poteser M., Romanin C., Schreibmayer W., Mayer B., Groschner K. S-nitrosation controls gating and conductance of the $\alpha 1$ subunit of class C L-type Ca^{2+} channels. *J Biol Chem* ;**276**:14797-803;2001
- [363] Gui P., Wu X., Ling S., Stotz S.C., Winkfein R.J., Wilson E. et al. Integrin receptor activation triggers converging regulation of Cav1.2 calcium channels by c-Src and protein kinase A pathways. *J Biol Chem* ;**281**:14015-25;2006
- [364] Ren C., Zhang J., Philipson K.D., Kotlikoff M.I., Blaustein M.P., Matteson D.R. Activation of L-type Ca^{2+} channels by protein kinase C is reduced in smooth muscle-specific $\text{Na}^{+}/\text{Ca}^{2+}$ exchanger knockout mice. *Am J Physiol Heart Circ Physiol* ;**298**:H1484-H1491;2010
- [365] Wu X., Davis G.E., Meininger G.A., Wilson E., Davis M.J. Regulation of the L-type calcium channel by $\alpha 5\beta 1$ integrin requires signaling between focal adhesion proteins. *J Biol Chem* ;**276**:30285-92;2001
- [366] Rogers P.A., Chilian W.M., Bratz I.N., Bryan R.M., Jr., Dick G.M. H_2O_2 activates redox- and 4-aminopyridine-sensitive Kv channels in coronary vascular smooth muscle. *Am J Physiol Heart Circ Physiol* ;**292**:H1404-H1411;2007
- [367] Graham S., Ding M., Ding Y., Sours-Brothers S., Luchowski R., Gryczynski Z. et al. Canonical transient receptor potential 6 (TRPC6), a redox-regulated cation channel. *J Biol Chem* ;**285**:23466-76;2010
- [368] Chen T.X., Xu X.Y., Zhao Z., Zhao F.Y., Gao Y.M., Yan X.H. et al. Hydrogen peroxide is a critical regulator of the hypoxia-induced alterations of store-operated Ca^{2+} entry into rat pulmonary arterial smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* ;**312**:L477-L487;2017
- [369] Zhang J., Zhou J., Cai L., Lu Y., Wang T., Zhu L. et al. Extracellular calcium-sensing receptor is critical in hypoxic pulmonary vasoconstriction. *Antioxid Redox Signal* ;**17**:471-84;2012
- [370] Chow J.Y., Estrema C., Orneles T., Dong X., Barrett K.E., Dong H. Calcium-sensing receptor modulates extracellular Ca^{2+} entry via TRPC-encoded receptor-operated channels in human aortic smooth muscle cells. *Am J Physiol Cell Physiol* ;**301**:C461-C468;2011

- [371] Becker S., Moir L.M., Snetkov V.A., Aaronson P.I. Hypoxic pulmonary vasoconstriction in intact rat intrapulmonary arteries is not initiated by inhibition of Na⁺-Ca²⁺ exchange. *Am J Physiol Lung Cell Mol Physiol* ;**293**:L982-L990;2007
- [372] Hu Q., Zheng G., Zweier J.L., Deshpande S., Irani K., Ziegelstein R.C. NADPH oxidase activation increases the sensitivity of intracellular Ca²⁺ stores to inositol 1,4,5-trisphosphate in human endothelial cells. *J Biol Chem* ;**275**:15749-57;2000
- [373] Suzuki Y.J., Ford G.D. Superoxide stimulates IP₃-induced Ca²⁺ release from vascular smooth muscle sarcoplasmic reticulum. *Am J Physiol* ;**262**:H114-H116;1992
- [374] Lopez R.J., Byrne S., Vukcevic M., Sekulic-Jablanovic M., Xu L., Brink M. et al. An RYR1 mutation associated with malignant hyperthermia is also associated with bleeding abnormalities. *Sci Signal* ;**9**:ra68;2016
- [375] Liao B., Zheng Y.M., Yadav V.R., Korde A.S., Wang Y.X. Hypoxia induces intracellular Ca²⁺ release by causing reactive oxygen species-mediated dissociation of FK506-binding protein 12.6 from ryanodine receptor 2 in pulmonary artery myocytes. *Antioxid Redox Signal* ;**14**:37-47;2011
- [376] Yan Y., Wei C.L., Zhang W.R., Cheng H.P., Liu J. Cross-talk between calcium and reactive oxygen species signaling. *Acta Pharmacol Sin* ;**27**:821-6;2006
- [377] Fellner S.K., Arendshorst W.J. Angiotensin II, reactive oxygen species, and Ca²⁺ signaling in afferent arterioles. *Am J Physiol Renal Physiol* ;**289**:F1012-F1019;2005
- [378] Lee S., Paudel O., Jiang Y., Yang X.R., Sham J.S. CD38 mediates angiotensin II-induced intracellular Ca(2+) release in rat pulmonary arterial smooth muscle cells. *Am J Respir Cell Mol Biol* ;**52**:332-41;2015
- [379] Park D.R., Nam T.S., Kim Y.W., Bae Y.S., Kim U.H. Oxidative activation of type III CD38 by NADPH oxidase-derived hydrogen peroxide in Ca(2+) signaling. *FASEB J* ;**33**:3404-19;2019
- [380] Adachi T., Weisbrod R.M., Pimentel D.R., Ying J., Sharov V.S., Schoneich C. et al. S-Glutathiolation by peroxynitrite activates SERCA during arterial relaxation by nitric oxide. *Nat Med* ;**10**:1200-7;2004
- [381] Bidwell P.A., Liu G.S., Nagarajan N., Lam C.K., Haghighi K., Gardner G. et al. HAX-1 regulates SERCA2a oxidation and degradation. *J Mol Cell Cardiol* ;**114**:220-33;2018
- [382] Somlyo A.P., Somlyo A.V. Ca²⁺ sensitivity of smooth muscle and nonmuscle myosin II: modulated by G proteins, kinases, and myosin phosphatase. *Physiol Rev* ;**83**:1325-58;2003
- [383] Wilson D.P., Susnjar M., Kiss E., Sutherland C., Walsh M.P. Thromboxane A₂-induced contraction of rat caudal arterial smooth muscle involves activation of Ca²⁺ entry and Ca²⁺ sensitization: Rho-associated kinase-mediated phosphorylation of MYPT1 at Thr-855, but not Thr-697. *Biochem J* ;**389**:763-74;2005
- [384] Amano M., Fukata Y., Kaibuchi K. Regulation and functions of Rho-associated kinase. *Exp Cell Res* ;**261**:44-51;2000
- [385] Heo J., Campbell S.L. Mechanism of redox-mediated guanine nucleotide exchange on redox-active Rho GTPases. *J Biol Chem* ;**280**:31003-10;2005

- [386] Aghajanian A., Wittchen E.S., Campbell S.L., Burridge K. Direct activation of RhoA by reactive oxygen species requires a redox-sensitive motif. *PLoS One* ;**4**:e8045;2009
- [387] Maekawa M., Ishizaki T., Boku S., Watanabe N., Fujita A., Iwamatsu A. et al. Signaling from Rho to the actin cytoskeleton through protein kinases ROCK and LIM-kinase. *Science* ;**285**:895-8;1999
- [388] de Rezende F.F., Martins L.A., Niland S., Wittig I., Heide H., Schroder K. et al. Integrin $\alpha 7 \beta 1$ is a redox-regulated target of hydrogen peroxide in vascular smooth muscle cell adhesion. *Free Radic Biol Med* ;**53**:521-31;2012
- [389] Gunst S.J., Zhang W. Actin cytoskeletal dynamics in smooth muscle: a new paradigm for the regulation of smooth muscle contraction. *Am J Physiol Cell Physiol* ;**295**:C576-C587;2008
- [390] Zemlickova E., Johannes F.J., Aitken A., Dubois T. Association of CPI-17 with protein kinase C and casein kinase I. *Biochem Biophys Res Commun* ;**316**:39-47;2004
- [391] Dessy C., Kim I., Sougnéz C.L., Laporte R., Morgan K.G. A role for MAP kinase in differentiated smooth muscle contraction evoked by α -adrenoceptor stimulation. *Am J Physiol* ;**275**:C1081-C1086;1998
- [392] Trappanese D.M., Sivilich S., Ets H.K., Kako F., Autieri M.V., Moreland R.S. Regulation of mitogen-activated protein kinase by protein kinase C and mitogen-activated protein kinase phosphatase-1 in vascular smooth muscle. *Am J Physiol Cell Physiol* ;**310**:C921-C930;2016
- [393] Adachi T., Pimentel D.R., Heibeck T., Hou X., Lee Y.J., Jiang B. et al. S-glutathiolation of Ras mediates redox-sensitive signaling by angiotensin II in vascular smooth muscle cells. *J Biol Chem* ;**279**:29857-62;2004
- [394] Gao Q., Zhao X., Ahmad M., Wolin M.S. Mitochondrial-derived hydrogen peroxide inhibits relaxation of bovine coronary arterial smooth muscle to hypoxia through stimulation of ERK MAP kinase. *Am J Physiol Heart Circ Physiol* ;**297**:H2262-H2269;2009
- [395] Meloche S., Landry J., Huot J., Houle F., Marceau F., Giasson E. p38 MAP kinase pathway regulates angiotensin II-induced contraction of rat vascular smooth muscle. *Am J Physiol Heart Circ Physiol* ;**279**:H741-H751;2000
- [396] Zagorac D., Yamaura K., Zhang C., Roman R.J., Harder D.R. The effect of superoxide anion on autoregulation of cerebral blood flow. *Stroke* ;**36**:2589-94;2005
- [397] Bidani A.K., Griffin K.A., Williamson G., Wang X., Loutzenhiser R. Protective importance of the myogenic response in the renal circulation. *Hypertension* ;**54**:393-8;2009
- [398] Petrone A., Sap J. Emerging issues in receptor protein tyrosine phosphatase function: lifting fog or simply shifting? *J Cell Sci* ;**113 (Pt 13)**:2345-54;2000
- [399] Carlstrom M., Wilcox C.S., Arendshorst W.J. Renal autoregulation in health and disease. *Physiol Rev* ;**95**:405-511;2015
- [400] Prysyazhna O., Rudyk O., Eaton P. Single atom substitution in mouse protein kinase G eliminates oxidant sensing to cause hypertension. *Nat Med* ;**18**:286-90;2012

- [401] Bautista-Nino P.K., van der Stel M., Batenburg W.W., de V.R., Roks A.J.M., Danser A.H.J. Endothelium-derived hyperpolarizing factor and protein kinase G α activation: H₂O₂ versus S-nitrosothiols. *Eur J Pharmacol* ;**827**:112-6;2018
- [402] Matoba T., Shimokawa H., Nakashima M., Hirakawa Y., Mukai Y., Hirano K. et al. Hydrogen peroxide is an endothelium-derived hyperpolarizing factor in mice. *J Clin Invest* ;**106**:1521-30;2000
- [403] Zhang D.X., Borbouse L., Gebremedhin D., Mendoza S.A., Zinkevich N.S., Li R. et al. H₂O₂-induced dilation in human coronary arterioles: role of protein kinase G dimerization and large-conductance Ca²⁺-activated K⁺ channel activation. *Circ Res* ;**110**:471-80;2012
- [404] Morikawa K., Fujiki T., Matoba T., Kubota H., Hatanaka M., Takahashi S. et al. Important role of superoxide dismutase in EDHF-mediated responses of human mesenteric arteries. *J Cardiovasc Pharmacol* ;**44**:552-6;2004
- [405] Leung H.S., Leung F.P., Yao X., Ko W.H., Chen Z.Y., Vanhoutte P.M. et al. Endothelial mediators of the acetylcholine-induced relaxation of the rat femoral artery. *Vascul Pharmacol* ;**44**:299-308;2006
- [406] Edwards D.H., Li Y., Griffith T.M. Hydrogen peroxide potentiates the EDHF phenomenon by promoting endothelial Ca²⁺ mobilization. *Arterioscler Thromb Vasc Biol* ;**28**:1774-81;2008
- [407] Park S.W., Noh H.J., Sung D.J., Kim J.G., Kim J.M., Ryu S.Y. et al. Hydrogen peroxide induces vasorelaxation by enhancing 4-aminopyridine-sensitive K_v currents through S-glutathionylation. *Pflugers Arch* ;**467**:285-97;2015
- [408] Miura H., Bosnjak J.J., Ning G., Saito T., Miura M., Gutterman D.D. Role for hydrogen peroxide in flow-induced dilation of human coronary arterioles. *Circ Res* ;**92**:e31-e40;2003
- [409] Saitoh S., Kiyooka T., Rocic P., Rogers P.A., Zhang C., Swafford A. et al. Redox-dependent coronary metabolic dilation. *Am J Physiol Heart Circ Physiol* ;**293**:H3720-H3725;2007
- [410] Liu Y., Bubolz A.H., Mendoza S., Zhang D.X., Gutterman D.D. H₂O₂ is the transferrable factor mediating flow-induced dilation in human coronary arterioles. *Circ Res* ;**108**:566-73;2011
- [411] Liu C., Ngai C.Y., Huang Y., Ko W.H., Wu M., He G.W. et al. Depletion of intracellular Ca²⁺ stores enhances flow-induced vascular dilatation in rat small mesenteric artery. *Br J Pharmacol* ;**147**:506-15;2006
- [412] Juncos L.A., Garvin J., Carretero O.A., Ito S. Flow modulates myogenic responses in isolated microperfused rabbit afferent arterioles via endothelium-derived nitric oxide. *J Clin Invest* ;**95**:2741-8;1995
- [413] Montezano A.C., Touyz R.M. Reactive oxygen species and endothelial function--role of nitric oxide synthase uncoupling and Nox family nicotinamide adenine dinucleotide phosphate oxidases. *Basic Clin Pharmacol Toxicol* ;**110**:87-94;2012
- [414] Siu K.L., Gao L., Cai H. Differential Roles of Protein Complexes NOX1-NOXO1 and NOX2-p47phox in Mediating Endothelial Redox Responses to Oscillatory and Unidirectional Laminar Shear Stress. *J Biol Chem* ;**291**:8653-62;2016

- [415] Wind S., Beuerlein K., Armitage M.E., Taye A., Kumar A.H., Janowitz D. et al. Oxidative stress and endothelial dysfunction in aortas of aged spontaneously hypertensive rats by NOX1/2 is reversed by NADPH oxidase inhibition. *Hypertension* ;**56**:490-7;2010
- [416] Avdonin P.V., Nadeev A.D., Mironova G.Y., Zharkikh I.L., Avdonin P.P., Goncharov N.V. Enhancement by Hydrogen Peroxide of Calcium Signals in Endothelial Cells Induced by 5-HT1B and 5-HT2B Receptor Agonists. *Oxid Med Cell Longev* ;**2019**:1701478;2019
- [417] Hu Q., Yu Z.X., Ferrans V.J., Takeda K., Irani K., Ziegelstein R.C. Critical role of NADPH oxidase-derived reactive oxygen species in generating Ca²⁺ oscillations in human aortic endothelial cells stimulated by histamine. *J Biol Chem* ;**277**:32546-51;2002
- [418] Raignault A., Bolduc V., Lesage F., Thorin E. Pulse pressure-dependent cerebrovascular eNOS regulation in mice. *J Cereb Blood Flow Metab* ;**37**:413-24;2017
- [419] De Keulenaer G.W., Chappell D.C., Ishizaka N., Nerem R.M., Alexander R.W., Griending K.K. Oscillatory and steady laminar shear stress differentially affect human endothelial redox state: role of a superoxide-producing NADH oxidase. *Circ Res* ;**82**:1094-101;1998
- [420] Duerschmidt N., Stielow C., Muller G., Pagano P.J., Morawietz H. NO-mediated regulation of NAD(P)H oxidase by laminar shear stress in human endothelial cells. *J Physiol* ;**576**:557-67;2006
- [421] Paravicini T.M., Miller A.A., Drummond G.R., Sobey C.G. Flow-induced cerebral vasodilatation in vivo involves activation of phosphatidylinositol-3 kinase, NADPH-oxidase, and nitric oxide synthase. *J Cereb Blood Flow Metab* ;**26**:836-45;2006
- [422] Lu X., Guo X., Wassall C.D., Kemple M.D., Unthank J.L., Kassab G.S. Reactive oxygen species cause endothelial dysfunction in chronic flow overload. *J Appl Physiol (1985)* ;**110**:520-7;2011
- [423] Matsuno K., Yamada H., Iwata K., Jin D., Katsuyama M., Matsuki M. et al. Nox1 is involved in angiotensin II-mediated hypertension: a study in Nox1-deficient mice. *Circulation* ;**112**:2677-85;2005
- [424] Thompson J.A., Larion S., Mintz J.D., Belin de Chantemele E.J., Fulton D.J., Stepp D.W. Genetic Deletion of NADPH Oxidase 1 Rescues Microvascular Function in Mice With Metabolic Disease. *Circ Res* ;**121**:502-11;2017
- [425] Jung O., Schreiber J.G., Geiger H., Pedrazzini T., Busse R., Brandes R.P. gp91phox-containing NADPH oxidase mediates endothelial dysfunction in renovascular hypertension. *Circulation* ;**109**:1795-801;2004
- [426] Tzima E., Irani-Tehrani M., Kiosses W.B., Dejana E., Schultz D.A., Engelhardt B. et al. A mechanosensory complex that mediates the endothelial cell response to fluid shear stress. *Nature* ;**437**:426-31;2005
- [427] Xu Z., Yu Y., Duh E.J. Vascular endothelial growth factor upregulates expression of ADAMTS1 in endothelial cells through protein kinase C signaling. *Invest Ophthalmol Vis Sci* ;**47**:4059-66;2006

- [428] Chou M.T., Wang J., Fujita D.J. Src kinase becomes preferentially associated with the VEGFR, KDR/Flk-1, following VEGF stimulation of vascular endothelial cells. *BMC Biochem* ;**3**:32;2002
- [429] Garrett T.A., van Buul J.D., Burridge K. VEGF-induced Rac1 activation in endothelial cells is regulated by the guanine nucleotide exchange factor Vav2. *Exp Cell Res* ;**313**:3285-97;2007
- [430] Milovanova T., Chatterjee S., Hawkins B.J., Hong N., Sorokina E.M., Debolt K. et al. Caveolae are an essential component of the pathway for endothelial cell signaling associated with abrupt reduction of shear stress. *Biochim Biophys Acta* ;**1783**:1866-75;2008
- [431] Noel J., Wang H., Hong N., Tao J.Q., Yu K., Sorokina E.M. et al. PECAM-1 and caveolae form the mechanosensing complex necessary for NOX2 activation and angiogenic signaling with stopped flow in pulmonary endothelium. *Am J Physiol Lung Cell Mol Physiol* ;**305**:L805-L818;2013
- [432] Chatterjee S., Browning E.A., Hong N., Debolt K., Sorokina E.M., Liu W. et al. Membrane depolarization is the trigger for PI3K/Akt activation and leads to the generation of ROS. *Am J Physiol Heart Circ Physiol* ;**302**:H105-H114;2012
- [433] Chatterjee S., Nieman G.F., Christie J.D., Fisher A.B. Shear stress-related mechanosignaling with lung ischemia: lessons from basic research can inform lung transplantation. *Am J Physiol Lung Cell Mol Physiol* ;**307**:L668-L680;2014
- [434] Gratton J.P., Fontana J., O'Connor D.S., Garcia-Cardena G., McCabe T.J., Sessa W.C. Reconstitution of an endothelial nitric-oxide synthase (eNOS), hsp90, and caveolin-1 complex in vitro. Evidence that hsp90 facilitates calmodulin stimulated displacement of eNOS from caveolin-1. *J Biol Chem* ;**275**:22268-72;2000
- [435] Landmesser U., Dikalov S., Price S.R., McCann L., Fukai T., Holland S.M. et al. Oxidation of tetrahydrobiopterin leads to uncoupling of endothelial cell nitric oxide synthase in hypertension. *J Clin Invest* ;**111**:1201-9;2003
- [436] Chabot F., Mitchell J.A., Quinlan G.J., Evans T.W. Characterization of the vasodilator properties of peroxynitrite on rat pulmonary artery: role of poly (adenosine 5'-diphosphoribose) synthase. *Br J Pharmacol* ;**121**:485-90;1997
- [437] Fraile M.L., Conde M.V., Sanz L., Moreno M.J., Marco E.J., Lopez de Pablo A.L. Different influence of superoxide anions and hydrogen peroxide on endothelial function of isolated cat cerebral and pulmonary arteries. *Gen Pharmacol* ;**25**:1197-205;1994
- [438] Morikawa K., Shimokawa H., Matoba T., Kubota H., Akaike T., Talukder M.A. et al. Pivotal role of Cu,Zn-superoxide dismutase in endothelium-dependent hyperpolarization. *J Clin Invest* ;**112**:1871-9;2003
- [439] Selemidis S., Dusting G.J., Peshavariya H., Kemp-Harper B.K., Drummond G.R. Nitric oxide suppresses NADPH oxidase-dependent superoxide production by S-nitrosylation in human endothelial cells. *Cardiovasc Res* ;**75**:349-58;2007
- [440] Qian J., Chen F., Kovalenkov Y., Pandey D., Moseley M.A., Foster M.W. et al. Nitric oxide reduces NADPH oxidase 5 (Nox5) activity by reversible S-nitrosylation. *Free Radic Biol Med* ;**52**:1806-19;2012

- [441] Craige S.M., Chen K., Pei Y., Li C., Huang X., Chen C. et al. NADPH oxidase 4 promotes endothelial angiogenesis through endothelial nitric oxide synthase activation. *Circulation* ;**124**:731-40;2011
- [442] Cai H., Davis M.E., Drummond G.R., Harrison D.G. Induction of endothelial NO synthase by hydrogen peroxide via a Ca(2+)/calmodulin-dependent protein kinase II/janus kinase 2-dependent pathway. *Arterioscler Thromb Vasc Biol* ;**21**:1571-6;2001
- [443] Hwang J., Saha A., Boo Y.C., Sorescu G.P., McNally J.S., Holland S.M. et al. Oscillatory shear stress stimulates endothelial production of O₂⁻ from p47phox-dependent NAD(P)H oxidases, leading to monocyte adhesion. *J Biol Chem* ;**278**:47291-8;2003
- [444] Hwang J., Ing M.H., Salazar A., Lassegue B., Griendling K., Navab M. et al. Pulsatile versus oscillatory shear stress regulates NADPH oxidase subunit expression: implication for native LDL oxidation. *Circ Res* ;**93**:1225-32;2003
- [445] Brandes R.P., Schmitz-Winnenthal F.H., Feletou M., Godecke A., Huang P.L., Vanhoutte P.M. et al. An endothelium-derived hyperpolarizing factor distinct from NO and prostacyclin is a major endothelium-dependent vasodilator in resistance vessels of wild-type and endothelial NO synthase knockout mice. *Proc Natl Acad Sci U S A* ;**97**:9747-52;2000
- [446] Garland C.J., Dora K.A. EDH: endothelium-dependent hyperpolarization and microvascular signalling. *Acta Physiol (Oxf)* ;**219**:152-61;2017
- [447] McNeish A.J., Dora K.A., Garland C.J. Possible role for K⁺ in endothelium-derived hyperpolarizing factor-linked dilatation in rat middle cerebral artery. *Stroke* ;**36**:1526-32;2005
- [448] Knot H.J., Standen N.B., Nelson M.T. Ryanodine receptors regulate arterial diameter and wall [Ca²⁺] in cerebral arteries of rat via Ca²⁺-dependent K⁺ channels. *J Physiol* ;**508 (Pt 1)**:211-21;1998
- [449] Yuan Q., Yang J., Santulli G., Reiken S.R., Wronska A., Kim M.M. et al. Maintenance of normal blood pressure is dependent on IP3R1-mediated regulation of eNOS. *Proc Natl Acad Sci U S A* ;**113**:8532-7;2016
- [450] Wilson C., Munoz-Palma E., Henriquez D.R., Palmisano I., Nunez M.T., Di G.S. et al. A Feed-Forward Mechanism Involving the NOX Complex and RyR-Mediated Ca²⁺ Release During Axonal Specification. *J Neurosci* ;**36**:11107-19;2016
- [451] Andersson D.A., Gentry C., Moss S., Bevan S. Transient receptor potential A1 is a sensory receptor for multiple products of oxidative stress. *J Neurosci* ;**28**:2485-94;2008
- [452] Sawada Y., Hosokawa H., Matsumura K., Kobayashi S. Activation of transient receptor potential ankyrin 1 by hydrogen peroxide. *Eur J Neurosci* ;**27**:1131-42;2008
- [453] Earley S., Gonzales A.L., Crnich R. Endothelium-dependent cerebral artery dilation mediated by TRPA1 and Ca²⁺-Activated K⁺ channels. *Circ Res* ;**104**:987-94;2009
- [454] Earley S. Endothelium-dependent cerebral artery dilation mediated by transient receptor potential and Ca²⁺-activated K⁺ channels. *J Cardiovasc Pharmacol* ;**57**:148-53;2011

- [455] Qian X., Francis M., Solodushko V., Earley S., Taylor M.S. Recruitment of dynamic endothelial Ca²⁺ signals by the TRPA1 channel activator AITC in rat cerebral arteries. *Microcirculation* ;**20**:138-48;2013
- [456] Sullivan M.N., Gonzales A.L., Pires P.W., Bruhl A., Leo M.D., Li W. et al. Localized TRPA1 channel Ca²⁺ signals stimulated by reactive oxygen species promote cerebral artery dilation. *Sci Signal* ;**8**:ra2;2015
- [457] Feletou M., Vanhoutte P.M. Endothelium-dependent hyperpolarization of canine coronary smooth muscle. *Br J Pharmacol* ;**93**:515-24;1988
- [458] Edwards G., Dora K.A., Gardener M.J., Garland C.J., Weston A.H. K⁺ is an endothelium-derived hyperpolarizing factor in rat arteries. *Nature* ;**396**:269-72;1998
- [459] Matoba T., Shimokawa H. Hydrogen peroxide is an endothelium-derived hyperpolarizing factor in animals and humans. *J Pharmacol Sci* ;**92**:1-6;2003
- [460] Eichhorn B., Muller G., Leuner A., Sawamura T., Ravens U., Morawietz H. Impaired vascular function in small resistance arteries of LOX-1 overexpressing mice on high-fat diet. *Cardiovasc Res* ;**82**:493-502;2009
- [461] Takaki A., Morikawa K., Murayama Y., Yamagishi H., Hosoya M., Ohashi J. et al. Roles of endothelial oxidases in endothelium-derived hyperpolarizing factor responses in mice. *J Cardiovasc Pharmacol* ;**52**:510-7;2008
- [462] Yada T., Shimokawa H., Hiramatsu O., Kajita T., Shigeto F., Goto M. et al. Hydrogen peroxide, an endogenous endothelium-derived hyperpolarizing factor, plays an important role in coronary autoregulation in vivo. *Circulation* ;**107**:1040-5;2003
- [463] Matoba T., Shimokawa H., Morikawa K., Kubota H., Kunihiro I., Urakami-Harasawa L. et al. Electron spin resonance detection of hydrogen peroxide as an endothelium-derived hyperpolarizing factor in porcine coronary microvessels. *Arterioscler Thromb Vasc Biol* ;**23**:1224-30;2003
- [464] Saitoh S., Zhang C., Tune J.D., Potter B., Kiyooka T., Rogers P.A. et al. Hydrogen peroxide: a feed-forward dilator that couples myocardial metabolism to coronary blood flow. *Arterioscler Thromb Vasc Biol* ;**26**:2614-21;2006
- [465] Yi L., Morgan J.T., Ragsdale S.W. Identification of a thiol/disulfide redox switch in the human BK channel that controls its affinity for heme and CO. *J Biol Chem* ;**285**:20117-27;2010
- [466] Burgoyne J.R., Madhani M., Cuello F., Charles R.L., Brennan J.P., Schroder E. et al. Cysteine redox sensor in PKG α enables oxidant-induced activation. *Science* ;**317**:1393-7;2007
- [467] Burke-Wolin T., Abate C.J., Wolin M.S., Gurtner G.H. Hydrogen peroxide-induced pulmonary vasodilation: role of guanosine 3',5'-cyclic monophosphate. *Am J Physiol* ;**261**:L393-L398;1991
- [468] Wolin M.S. Reactive oxygen species and the control of vascular function. *Am J Physiol Heart Circ Physiol* ;**296**:H539-H549;2009

- [469] Egea J., Fabregat I., Frapart Y.M., Ghezzi P., Gorlach A., Kietzmann T. et al. European contribution to the study of ROS: A summary of the findings and prospects for the future from the COST action BM1203 (EU-ROS). *Redox Biol* ;**13**:94-162;2017
- [470] Cuevas S., Villar V.A.M., Jose P.A. Genetic polymorphisms associated with reactive oxygen species and blood pressure regulation. *Pharmacogenomics J* ;2019
- [471] Masi S., Uliana M., Virdis A. Angiotensin II and vascular damage in hypertension: Role of oxidative stress and sympathetic activation. *Vascul Pharmacol* ;**115**:13-7;2019
- [472] Massaro M., Scoditti E., Carluccio M.A., De C.R. Oxidative stress and vascular stiffness in hypertension: A renewed interest for antioxidant therapies? *Vascul Pharmacol* ;**116**:45-50;2019
- [473] Chabrashvili T., Kitiyakara C., Blau J., Karber A., Aslam S., Welch W.J. et al. Effects of ANG II type 1 and 2 receptors on oxidative stress, renal NADPH oxidase, and SOD expression. *Am J Physiol Regul Integr Comp Physiol* ;**285**:R117-R124;2003
- [474] Montezano A.C., Dulak-Lis M., Tsiropoulou S., Harvey A., Briones A.M., Touyz R.M. Oxidative stress and human hypertension: vascular mechanisms, biomarkers, and novel therapies. *Can J Cardiol* ;**31**:631-41;2015
- [475] do Vale G.T., Tirapelli C.R. Are reactive oxygen species important mediators of vascular dysfunction? *Curr Hypertens Rev* ;2019
- [476] Pinheiro L.C., Oliveira-Paula G.H. Sources and effects of oxidative stress in hypertension. *Curr Hypertens Rev* ;2019
- [477] Sorriento D., De L.N., Trimarco B., Iaccarino G. The Antioxidant Therapy: New Insights in the Treatment of Hypertension. *Front Physiol* ;**9**:258;2018
- [478] Al-Kuraishy H.M., Al-Gareeb A.I., Al-Maiah T.J. Concept and connotation of oxidative stress in preeclampsia. *J Lab Physicians* ;**10**:276-82;2018
- [479] Pechanova O., Simko F. Chronic antioxidant therapy fails to ameliorate hypertension: potential mechanisms behind. *J Hypertens Suppl* ;**27**:S32-S36;2009
- [480] Daiber A., Munzel T. Organic Nitrate Therapy, Nitrate Tolerance, and Nitrate-Induced Endothelial Dysfunction: Emphasis on Redox Biology and Oxidative Stress. *Antioxid Redox Signal* ;**23**:899-942;2015
- [481] Ahmad K.A., Yuan Y.D., Nawaz W., Ze H., Zhuo C.X., Talal B. et al. Antioxidant therapy for management of oxidative stress induced hypertension. *Free Radic Res* ;**51**:428-38;2017
- [482] Taysi S., Tascan A.S., Ugur M.G., Demir M. Radicals, Oxidative/Nitrosative Stress and Preeclampsia. *Mini Rev Med Chem* ;**19**:178-93;2019
- [483] Briones A.M., Tabet F., Callera G.E., Montezano A.C., Yogi A., He Y. et al. Differential regulation of Nox1, Nox2 and Nox4 in vascular smooth muscle cells from WKY and SHR. *J Am Soc Hypertens* ;**5**:137-53;2011

- [484] Morawietz H., Weber M., Rueckschloss U., Lauer N., Hacker A., Kojda G. Upregulation of vascular NAD(P)H oxidase subunit gp91phox and impairment of the nitric oxide signal transduction pathway in hypertension. *Biochem Biophys Res Commun* ;**285**:1130-5;2001
- [485] Rahali S., Li Y., Anand-Srivastava M.B. Contribution of oxidative stress and growth factor receptor transactivation in natriuretic peptide receptor C-mediated attenuation of hyperproliferation of vascular smooth muscle cells from SHR. *PLoS One* ;**13**:e0191743;2018
- [486] Madamanchi N.R., Runge M.S. Redox signaling in cardiovascular health and disease. *Free Radic Biol Med* ;**61**:473-501;2013
- [487] Cifuentes M.E., Rey F.E., Carretero O.A., Pagano P.J. Upregulation of p67(phox) and gp91(phox) in aortas from angiotensin II-infused mice. *Am J Physiol Heart Circ Physiol* ;**279**:H2234-H2240;2000
- [488] Fukui T., Ishizaka N., Rajagopalan S., Laursen J.B., Capers Q., Taylor W.R. et al. p22phox mRNA expression and NADPH oxidase activity are increased in aortas from hypertensive rats. *Circ Res* ;**80**:45-51;1997
- [489] De Batista P.R., Palacios R., Martin A., Hernanz R., Medici C.T., Silva M.A. et al. Toll-like receptor 4 upregulation by angiotensin II contributes to hypertension and vascular dysfunction through reactive oxygen species production. *PLoS One* ;**9**:e104020;2014
- [490] Gavazzi G., Banfi B., Deffert C., Fiette L., Schappi M., Herrmann F. et al. Decreased blood pressure in NOX1-deficient mice. *FEBS Lett* ;**580**:497-504;2006
- [491] Rueckschloss U., Quinn M.T., Holtz J., Morawietz H. Dose-dependent regulation of NAD(P)H oxidase expression by angiotensin II in human endothelial cells: protective effect of angiotensin II type 1 receptor blockade in patients with coronary artery disease. *Arterioscler Thromb Vasc Biol* ;**22**:1845-51;2002
- [492] Rodriguez A.I., Csanyi G., Ranayhossaini D.J., Feck D.M., Blose K.J., Assatourian L. et al. MEF2B-Nox1 signaling is critical for stretch-induced phenotypic modulation of vascular smooth muscle cells. *Arterioscler Thromb Vasc Biol* ;**35**:430-8;2015
- [493] Paravicini T.M., Chrissobolis S., Drummond G.R., Sobey C.G. Increased NADPH-oxidase activity and Nox4 expression during chronic hypertension is associated with enhanced cerebral vasodilatation to NADPH in vivo. *Stroke* ;**35**:584-9;2004
- [494] Chen D., Zang Y.H., Qiu Y., Zhang F., Chen A.D., Wang J.J. et al. BCL6 Attenuates Proliferation and Oxidative Stress of Vascular Smooth Muscle Cells in Hypertension. *Oxid Med Cell Longev* ;**2019**:5018410;2019
- [495] Carrillo-Sepulveda M.A., Keen H.L., Davis D.R., Grobe J.L., Sigmund C.D. Role of vascular smooth muscle PPARgamma in regulating AT1 receptor signaling and angiotensin II-dependent hypertension. *PLoS One* ;**9**:e103786;2014
- [496] Touyz R.M., Anagnostopoulou A., Camargo L.L., Rios F.J., Montezano A.C. Vascular Biology of Superoxide-Generating NADPH Oxidase 5-Implications in Hypertension and Cardiovascular Disease. *Antioxid Redox Signal* ;**30**:1027-40;2019
- [497] Yu P., Han W., Villar V.A., Yang Y., Lu Q., Lee H. et al. Unique role of NADPH oxidase 5 in oxidative stress in human renal proximal tubule cells. *Redox Biol* ;**2**:570-9;2014

- [498] Kumar R., Kohli S., Ali Z., Duhan K., Ram R., Gupta M. et al. CYBA (p22phox) variants associate with blood pressure and oxidative stress markers in hypertension: a replication study in populations of diverse altitudes. *Hypertens Res* ;**38**:498-506;2015
- [499] Wang Y., Chen F., Le B., Stepp D.W., Fulton D.J. Impact of Nox5 polymorphisms on basal and stimulus-dependent ROS generation. *PLoS One* ;**9**:e100102;2014
- [500] Kontos H.A., Wei E.P. Hydroxyl radical-dependent inactivation of guanylate cyclase in cerebral arterioles by methylene blue and by LY83583. *Stroke* ;**24**:427-34;1993
- [501] Violi F., Sanguigni V., Carnevale R., Plebani A., Rossi P., Finocchi A. et al. Hereditary deficiency of gp91(phox) is associated with enhanced arterial dilatation: results of a multicenter study. *Circulation* ;**120**:1616-22;2009
- [502] Fardella C., Rodriguez-Portales J.A. Intracellular calcium and blood pressure: comparison between primary hyperparathyroidism and essential hypertension. *J Endocrinol Invest* ;**18**:827-32;1995
- [503] Fleckenstein-Grun G., Frey M., Thimm F., Hofgartner W., Fleckenstein A. Calcium overload-an important cellular mechanism in hypertension and arteriosclerosis. *Drugs* ;**44 Suppl 1**:23-30;1992
- [504] Loukotova J., Kunes J., Zicha J. Cytosolic free calcium response to angiotensin II in aortic VSMC isolated from male and female SHR. *Physiol Res* ;**47**:507-10;1998
- [505] Arai T., Ohyanagi M., Shibuya J., Iwasaki T. Increased function of the voltage-dependent calcium channels, without increase of Ca²⁺ release from the sarcoplasmic reticulum in the arterioles of spontaneous hypertensive rats. *Am J Hypertens* ;**12**:1236-42;1999
- [506] Touyz R.M., Schiffrin E.L. Role of calcium influx and intracellular calcium stores in angiotensin II-mediated calcium hyper-responsiveness in smooth muscle from spontaneously hypertensive rats. *J Hypertens* ;**15**:1431-9;1997
- [507] Pesic A., Madden J.A., Pesic M., Rusch N.J. High blood pressure upregulates arterial L-type Ca²⁺ channels: is membrane depolarization the signal? *Circ Res* ;**94**:e97-104;2004
- [508] Erne P., Hermsmeyer K. Intracellular vascular muscle Ca²⁺ modulation in genetic hypertension. *Hypertension* ;**14**:145-51;1989
- [509] Neusser M., Tepel M., Golinski P., Holthues J., Spieker C., Zhu Z. et al. Different calcium storage pools in vascular smooth muscle cells from spontaneously hypertensive and normotensive Wistar-Kyoto rats. *J Hypertens* ;**12**:533-8;1994
- [510] Seko T., Ito M., Kureishi Y., Okamoto R., Moriki N., Onishi K. et al. Activation of RhoA and inhibition of myosin phosphatase as important components in hypertension in vascular smooth muscle. *Circ Res* ;**92**:411-8;2003
- [511] Lee D.L., Webb R.C., Jin L. Hypertension and RhoA/Rho-kinase signaling in the vasculature: highlights from the recent literature. *Hypertension* ;**44**:796-9;2004
- [512] Amin F., Ahmad A., Feroz A., Khaqi S.S., Khan M.S., Tabrez S. et al. An update on the association of protein kinases with cardiovascular diseases. *Curr Pharm Des* ;2019

- [513] Touyz R.M., Alves-Lopes R., Rios F.J., Camargo L.L., Anagnostopoulou A., Arner A. et al. Vascular smooth muscle contraction in hypertension. *Cardiovasc Res* ;**114**:529-39;2018
- [514] Yuan T.Y., Chen D., Chen Y.C., Zhang H.F., Niu Z.R., Jiao X.Z. et al. A novel hypertensive crisis rat model established by excessive norepinephrine infusion and the potential therapeutic effects of Rho-kinase inhibitors on it. *Biomed Pharmacother* ;**109**:1867-75;2019
- [515] Denniss S.G., Jeffery A.J., Rush J.W. RhoA-Rho kinase signaling mediates endothelium- and endoperoxide-dependent contractile activities characteristic of hypertensive vascular dysfunction. *Am J Physiol Heart Circ Physiol* ;**298**:H1391-H1405;2010
- [516] Chen X., Touyz R.M., Park J.B., Schiffrin E.L. Antioxidant effects of vitamins C and E are associated with altered activation of vascular NADPH oxidase and superoxide dismutase in stroke-prone SHR. *Hypertension* ;**38**:606-11;2001
- [517] Kawada N., Imai E., Karber A., Welch W.J., Wilcox C.S. A mouse model of angiotensin II slow pressor response: role of oxidative stress. *J Am Soc Nephrol* ;**13**:2860-8;2002
- [518] Chen X., Patel K., Connors S.G., Mendonca M., Welch W.J., Wilcox C.S. Acute antihypertensive action of Tempol in the spontaneously hypertensive rat. *Am J Physiol Heart Circ Physiol* ;**293**:H3246-H3253;2007
- [519] Wilcox C.S., Pearlman A. Chemistry and antihypertensive effects of tempol and other nitroxides. *Pharmacol Rev* ;**60**:418-69;2008
- [520] Dikalova A., Clempus R., Lassegue B., Cheng G., McCoy J., Dikalov S. et al. Nox1 overexpression potentiates angiotensin II-induced hypertension and vascular smooth muscle hypertrophy in transgenic mice. *Circulation* ;**112**:2668-76;2005
- [521] Qi M., Elion E.A. MAP kinase pathways. *J Cell Sci* ;**118**:3569-72;2005
- [522] Illario M., Cavallo A.L., Bayer K.U., Di M.T., Fenzi G., Rossi G. et al. Calcium/calmodulin-dependent protein kinase II binds to Raf-1 and modulates integrin-stimulated ERK activation. *J Biol Chem* ;**278**:45101-8;2003
- [523] Li N., Wang C., Wu Y., Liu X., Cao X. Ca(2+)/calmodulin-dependent protein kinase II promotes cell cycle progression by directly activating MEK1 and subsequently modulating p27 phosphorylation. *J Biol Chem* ;**284**:3021-7;2009
- [524] Frank G.D., Saito S., Motley E.D., Sasaki T., Ohba M., Kuroki T. et al. Requirement of Ca(2+) and PKCdelta for Janus kinase 2 activation by angiotensin II: involvement of PYK2. *Mol Endocrinol* ;**16**:367-77;2002
- [525] Liu Y., Suzuki Y.J., Day R.M., Fanburg B.L. Rho kinase-induced nuclear translocation of ERK1/ERK2 in smooth muscle cell mitogenesis caused by serotonin. *Circ Res* ;**95**:579-86;2004
- [526] Banes A.K., Shaw S.M., Tawfik A., Patel B.P., Ogbi S., Fulton D. et al. Activation of the JAK/STAT pathway in vascular smooth muscle by serotonin. *Am J Physiol Cell Physiol* ;**288**:C805-C812;2005
- [527] Abe J., Berk B.C. Fyn and JAK2 mediate Ras activation by reactive oxygen species. *J Biol Chem* ;**274**:21003-10;1999

- [528] Lee K.M., Kang H.A., Park M., Lee H.Y., Song M.J., Ko K. et al. Interleukin-24 suppresses the growth of vascular smooth muscle cells by inhibiting H₂O₂-induced reactive oxygen species production. *Pharmacology* ;**90**:332-41;2012
- [529] Li G., Li Y.Y., Sun J.E., Lin W.H., Zhou R.X. ILK-PI3K/AKT pathway participates in cutaneous wound contraction by regulating fibroblast migration and differentiation to myofibroblast. *Lab Invest* ;**96**:741-51;2016
- [530] Scott J.A., Xie L., Li H., Li W., He J.B., Sanders P.N. et al. The multifunctional Ca²⁺/calmodulin-dependent kinase II regulates vascular smooth muscle migration through matrix metalloproteinase 9. *Am J Physiol Heart Circ Physiol* ;**302**:H1953-H1964;2012
- [531] Maheswaranathan M., Gole H.K., Fernandez I., Lassegue B., Griendling K.K., San M.A. Platelet-derived growth factor (PDGF) regulates Slingshot phosphatase activity via Nox1-dependent auto-dephosphorylation of serine 834 in vascular smooth muscle cells. *J Biol Chem* ;**286**:35430-7;2011
- [532] Montenegro M.F., Valdivia A., Smolensky A., Verma K., Taylor W.R., San M.A. Nox4-dependent activation of cofilin mediates VSMC reorientation in response to cyclic stretching. *Free Radic Biol Med* ;**85**:288-94;2015
- [533] Mercure M.Z., Ginnan R., Singer H.A. CaM kinase II delta2-dependent regulation of vascular smooth muscle cell polarization and migration. *Am J Physiol Cell Physiol* ;**294**:C1465-C1475;2008
- [534] Forrester S.J., Kikuchi D.S., Hernandez M.S., Xu Q., Griendling K.K. Reactive Oxygen Species in Metabolic and Inflammatory Signaling. *Circ Res* ;**122**:877-902;2018
- [535] Wu J., Saleh M.A., Kirabo A., Itani H.A., Montaniel K.R., Xiao L. et al. Immune activation caused by vascular oxidation promotes fibrosis and hypertension. *J Clin Invest* ;**126**:50-67;2016
- [536] Nikolopoulos S.N., Turner C.E. Integrin-linked kinase (ILK) binding to paxillin LD1 motif regulates ILK localization to focal adhesions. *J Biol Chem* ;**276**:23499-505;2001
- [537] Jiang M., Bujo H., Ohwaki K., Unoki H., Yamazaki H., Kanaki T. et al. Ang II-stimulated migration of vascular smooth muscle cells is dependent on LR11 in mice. *J Clin Invest* ;**118**:2733-46;2008
- [538] Tamborindéguy M.T., Matte B.F., Ramos G.O., Alves A.M., Bernardi L., Lamers M.L. NADPH-oxidase-derived ROS alters cell migration by modulating adhesions dynamics. *Biol Cell* ;**110**:225-36;2018
- [539] Lehoux S., Esposito B., Merval R., Tedgui A. Differential regulation of vascular focal adhesion kinase by steady stretch and pulsatility. *Circulation* ;**111**:643-9;2005
- [540] Xing Z., Chen H.C., Nowlen J.K., Taylor S.J., Shalloway D., Guan J.L. Direct interaction of v-Src with the focal adhesion kinase mediated by the Src SH2 domain. *Mol Biol Cell* ;**5**:413-21;1994
- [541] Hellstrand P., Albinsson S. Stretch-dependent growth and differentiation in vascular smooth muscle: role of the actin cytoskeleton. *Can J Physiol Pharmacol* ;**83**:869-75;2005

- [542] Rigacci S., Rovida E., Dello S.P., Berti A. Low Mr phosphotyrosine protein phosphatase associates and dephosphorylates p125 focal adhesion kinase, interfering with cell motility and spreading. *J Biol Chem* ;**277**:41631-6;2002
- [543] Burridge K., Sastry S.K., Sallee J.L. Regulation of cell adhesion by protein-tyrosine phosphatases. I. Cell-matrix adhesion. *J Biol Chem* ;**281**:15593-6;2006
- [544] Touyz R.M., Wu X.H., He G., Salomon S., Schiffrin E.L. Increased angiotensin II-mediated Src signaling via epidermal growth factor receptor transactivation is associated with decreased C-terminal Src kinase activity in vascular smooth muscle cells from spontaneously hypertensive rats. *Hypertension* ;**39**:479-85;2002
- [545] Rao G.N. Hydrogen peroxide induces complex formation of SHC-Grb2-SOS with receptor tyrosine kinase and activates Ras and extracellular signal-regulated protein kinases group of mitogen-activated protein kinases. *Oncogene* ;**13**:713-9;1996
- [546] Frank G.D., Motley E.D., Inagami T., Eguchi S. PYK2/CAKbeta represents a redox-sensitive tyrosine kinase in vascular smooth muscle cells. *Biochem Biophys Res Commun* ;**270**:761-5;2000
- [547] Madamanchi N.R., Li S., Patterson C., Runge M.S. Thrombin regulates vascular smooth muscle cell growth and heat shock proteins via the JAK-STAT pathway. *J Biol Chem* ;**276**:18915-24;2001
- [548] Gajos-Draus A., Duda M., Beresewicz A. Cardiac and renal upregulation of Nox2 and NF-kappaB and repression of Nox4 and Nrf2 in season- and diabetes-mediated models of vascular oxidative stress in guinea-pig and rat. *Physiol Rep* ;**5**;2017
- [549] Zhang J., Chen J., Yang J., Xu C.W., Pu P., Ding J.W. et al. Resveratrol attenuates oxidative stress induced by balloon injury in the rat carotid artery through actions on the ERK1/2 and NF-kappa B pathway. *Cell Physiol Biochem* ;**31**:230-41;2013
- [550] Cabal-Hierro L., Lazo P.S. Signal transduction by tumor necrosis factor receptors. *Cell Signal* ;**24**:1297-305;2012
- [551] Yamamoto T., Suzuki T., Kobayashi A., Wakabayashi J., Maher J., Motohashi H. et al. Physiological significance of reactive cysteine residues of Keap1 in determining Nrf2 activity. *Mol Cell Biol* ;**28**:2758-70;2008
- [552] Weber D.S., Rocic P., Mellis A.M., Laude K., Lyle A.N., Harrison D.G. et al. Angiotensin II-induced hypertrophy is potentiated in mice overexpressing p22phox in vascular smooth muscle. *Am J Physiol Heart Circ Physiol* ;**288**:H37-H42;2005
- [553] Bhatia K., Elmarakby A.A., El-Remessy A.B., Sullivan J.C. Oxidative stress contributes to sex differences in angiotensin II-mediated hypertension in spontaneously hypertensive rats. *Am J Physiol Regul Integr Comp Physiol* ;**302**:R274-R282;2012
- [554] Graton M.E., Potje S.R., Troiano J.A., Vale G.T., Perassa L.A., Nakamune A.C.M.S. et al. Apocynin alters redox signaling in conductance and resistance vessels of spontaneously hypertensive rats. *Free Radic Biol Med* ;**134**:53-63;2018

- [555] Beswick R.A., Dorrance A.M., Leite R., Webb R.C. NADH/NADPH oxidase and enhanced superoxide production in the mineralocorticoid hypertensive rat. *Hypertension* ;**38**:1107-11;2001
- [556] Armando I., Wang X., Villar V.A., Jones J.E., Asico L.D., Escano C. et al. Reactive oxygen species-dependent hypertension in dopamine D2 receptor-deficient mice. *Hypertension* ;**49**:672-8;2007
- [557] Lu W., Kang J., Hu K., Tang S., Zhou X., Xu L. et al. The role of the Nox4-derived ROS-mediated RhoA/Rho kinase pathway in rat hypertension induced by chronic intermittent hypoxia. *Sleep Breath* ;**21**:667-77;2017
- [558] Zeng S.Y., Yang L., Yan Q.J., Gao L., Lu H.Q., Yan P.K. Nox1/4 dual inhibitor GKT137831 attenuates hypertensive cardiac remodelling associating with the inhibition of ADAM17-dependent proinflammatory cytokines-induced signalling pathways in the rats with abdominal artery constriction. *Biomed Pharmacother* ;**109**:1907-14;2019
- [559] Teixeira G., Szyndralewicz C., Molango S., Carnesecchi S., Heitz F., Wiesel P. et al. Therapeutic potential of NADPH oxidase 1/4 inhibitors. *Br J Pharmacol* ;**174**:1647-69;2017
- [560] Tang Y., Huang Q., Liu C., Ou H., Huang D., Peng F. et al. p22phox promotes Ang-II-induced vascular smooth muscle cell phenotypic switch by regulating KLF4 expression. *Biochem Biophys Res Commun* ;**514**:280-6;2019
- [561] Ngarashi D., Fujikawa K., Ferdaus M.Z., Zahid H.M., Ohara H., Nabika T. Dual inhibition of NADPH oxidases and xanthine oxidase potently prevents salt-induced stroke in stroke-prone spontaneously hypertensive rats. *Hypertens Res* ;2019
- [562] Touyz R.M., Mercure C., He Y., Javeshghani D., Yao G., Callera G.E. et al. Angiotensin II-dependent chronic hypertension and cardiac hypertrophy are unaffected by gp91phox-containing NADPH oxidase. *Hypertension* ;**45**:530-7;2005
- [563] Wang H.D., Xu S., Johns D.G., Du Y., Quinn M.T., Cayatte A.J. et al. Role of NADPH oxidase in the vascular hypertrophic and oxidative stress response to angiotensin II in mice. *Circ Res* ;**88**:947-53;2001
- [564] Ren Y., D'Ambrosio M.A., Liu R., Pagano P.J., Garvin J.L., Carretero O.A. Enhanced myogenic response in the afferent arteriole of spontaneously hypertensive rats. *Am J Physiol Heart Circ Physiol* ;**298**:H1769-H1775;2010
- [565] Bendall J.K., Rinze R., Adlam D., Tatham A.L., de B.J., Wilson N. et al. Endothelial Nox2 overexpression potentiates vascular oxidative stress and hemodynamic response to angiotensin II: studies in endothelial-targeted Nox2 transgenic mice. *Circ Res* ;**100**:1016-25;2007
- [566] Bouabout G., Ayme-Dietrich E., Jacob H., Champy M.F., Birling M.C., Pavlovic G. et al. Nox4 genetic inhibition in experimental hypertension and metabolic syndrome. *Arch Cardiovasc Dis* ;**111**:41-52;2018
- [567] Di M.E., Gray S.P., Kennedy K., Szyndralewicz C., Lyle A.N., Lassegue B. et al. NOX4-derived reactive oxygen species limit fibrosis and inhibit proliferation of vascular smooth muscle cells in diabetic atherosclerosis. *Free Radic Biol Med* ;**97**:556-67;2016

- [568] Smyrniak I., Zhang X., Zhang M., Murray T.V., Brandes R.P., Schroder K. et al. Nicotinamide adenine dinucleotide phosphate oxidase-4-dependent upregulation of nuclear factor erythroid-derived 2-like 2 protects the heart during chronic pressure overload. *Hypertension* ;**65**:547-53;2015
- [569] Holterman C.E., Thibodeau J.F., Towaj C., Gutsol A., Montezano A.C., Parks R.J. et al. Nephropathy and elevated BP in mice with podocyte-specific NADPH oxidase 5 expression. *J Am Soc Nephrol* ;**25**:784-97;2014
- [570] Jha J.C., Banal C., Okabe J., Gray S.P., Hettige T., Chow B.S.M. et al. NADPH Oxidase Nox5 Accelerates Renal Injury in Diabetic Nephropathy. *Diabetes* ;**66**:2691-703;2017
- [571] Morio Y., Carter E.P., Oka M., McMurtry I.F. EDHF-mediated vasodilation involves different mechanisms in normotensive and hypertensive rat lungs. *Am J Physiol Heart Circ Physiol* ;**284**:H1762-H1770;2003
- [572] Torok J. Histamine-induced relaxation in pulmonary artery of normotensive and hypertensive rats: relative contribution of prostanoids, nitric oxide and hyperpolarization. *Physiol Res* ;**49**:107-14;2000
- [573] Tanaka S., Shiroto T., Godo S., Saito H., Ikumi Y., Ito A. et al. Important role of endothelium-dependent hyperpolarization in the pulmonary microcirculation in male mice: implications for hypoxia-induced pulmonary hypertension. *Am J Physiol Heart Circ Physiol* ;**314**:H940-H953;2018
- [574] Kuwahira I., Gonzalez N.C., Heisler N., Piiper J. Changes in regional blood flow distribution and oxygen supply during hypoxia in conscious rats. *J Appl Physiol (1985)* ;**74**:211-4;1993
- [575] Marshall C., Mamary A.J., Verhoeven A.J., Marshall B.E. Pulmonary artery NADPH-oxidase is activated in hypoxic pulmonary vasoconstriction. *Am J Respir Cell Mol Biol* ;**15**:633-44;1996
- [576] Weissmann N., Tadic A., Hanze J., Rose F., Winterhalder S., Nollen M. et al. Hypoxic vasoconstriction in intact lungs: a role for NADPH oxidase-derived H₂O₂? *Am J Physiol Lung Cell Mol Physiol* ;**279**:L683-L690;2000
- [577] Archer S.L., Reeve H.L., Michelakis E., Puttagunta L., Waite R., Nelson D.P. et al. O₂ sensing is preserved in mice lacking the gp91 phox subunit of NADPH oxidase. *Proc Natl Acad Sci U S A* ;**96**:7944-9;1999
- [578] Weissmann N., Zeller S., Schafer R.U., Turowski C., Ay M., Quanz K. et al. Impact of mitochondria and NADPH oxidases on acute and sustained hypoxic pulmonary vasoconstriction. *Am J Respir Cell Mol Biol* ;**34**:505-13;2006
- [579] Ahmad M., Kelly M.R., Zhao X., Kandhi S., Wolin M.S. Roles for Nox4 in the contractile response of bovine pulmonary arteries to hypoxia. *Am J Physiol Heart Circ Physiol* ;**298**:H1879-H1888;2010
- [580] Gupte S.A., Okada T., McMurtry I.F., Oka M. Role of pentose phosphate pathway-derived NADPH in hypoxic pulmonary vasoconstriction. *Pulm Pharmacol Ther* ;**19**:303-9;2006
- [581] Archer S.L., Souil E., Dinh-Xuan A.T., Schremmer B., Mercier J.C., El Y.A. et al. Molecular identification of the role of voltage-gated K⁺ channels, Kv1.5 and Kv2.1, in hypoxic

- pulmonary vasoconstriction and control of resting membrane potential in rat pulmonary artery myocytes. *J Clin Invest* ;**101**:2319-30;1998
- [582] Robertson T.P., Dipp M., Ward J.P., Aaronson P.I., Evans A.M. Inhibition of sustained hypoxic vasoconstriction by Y-27632 in isolated intrapulmonary arteries and perfused lung of the rat. *Br J Pharmacol* ;**131**:5-9;2000
 - [583] Wang Z., Lanner M.C., Jin N., Swartz D., Li L., Rhoades R.A. Hypoxia inhibits myosin phosphatase in pulmonary arterial smooth muscle cells: role of Rho-kinase. *Am J Respir Cell Mol Biol* ;**29**:465-71;2003
 - [584] Zhou J., Zhang J., Lu Y., Huang S., Xiao R., Zeng X. et al. Mitochondrial transplantation attenuates hypoxic pulmonary vasoconstriction. *Oncotarget* ;**7**:31284-98;2016
 - [585] Song T., Zheng Y.M., Wang Y.X. Cross Talk Between Mitochondrial Reactive Oxygen Species and Sarcoplasmic Reticulum Calcium in Pulmonary Arterial Smooth Muscle Cells. *Adv Exp Med Biol* ;**967**:289-98;2017
 - [586] Robertson T.P., Hague D., Aaronson P.I., Ward J.P. Voltage-independent calcium entry in hypoxic pulmonary vasoconstriction of intrapulmonary arteries of the rat. *J Physiol* ;**525 Pt 3**:669-80;2000
 - [587] Weissmann N., Voswinckel R., Hardebusch T., Rosseau S., Ghofrani H.A., Schermuly R. et al. Evidence for a role of protein kinase C in hypoxic pulmonary vasoconstriction. *Am J Physiol* ;**276**:L90-L95;1999
 - [588] Rathore R., Zheng Y.M., Li X.Q., Wang Q.S., Liu Q.H., Ginnan R. et al. Mitochondrial ROS-PKCepsilon signaling axis is uniquely involved in hypoxic increase in [Ca²⁺]_i in pulmonary artery smooth muscle cells. *Biochem Biophys Res Commun* ;**351**:784-90;2006
 - [589] Mittal M., Gu X.Q., Pak O., Pamenter M.E., Haag D., Fuchs D.B. et al. Hypoxia induces Kv channel current inhibition by increased NADPH oxidase-derived reactive oxygen species. *Free Radic Biol Med* ;**52**:1033-42;2012
 - [590] Svoboda L.K., Reddie K.G., Zhang L., Vesely E.D., Williams E.S., Schumacher S.M. et al. Redox-sensitive sulfenic acid modification regulates surface expression of the cardiovascular voltage-gated potassium channel Kv1.5. *Circ Res* ;**111**:842-53;2012
 - [591] Simonneau G., Montani D., Celermajer D.S., Denton C.P., Gatzoulis M.A., Krowka M. et al. Haemodynamic definitions and updated clinical classification of pulmonary hypertension. *Eur Respir J* ;**53**;2019
 - [592] Ismail S., Sturrock A., Wu P., Cahill B., Norman K., Huecksteadt T. et al. NOX4 mediates hypoxia-induced proliferation of human pulmonary artery smooth muscle cells: the role of autocrine production of transforming growth factor- β 1 and insulin-like growth factor binding protein-3. *Am J Physiol Lung Cell Mol Physiol* ;**296**:L489-L499;2009
 - [593] Veit F., Pak O., Egemnazarov B., Roth M., Kosanovic D., Seimetz M. et al. Function of NADPH oxidase 1 in pulmonary arterial smooth muscle cells after monocrotaline-induced pulmonary vascular remodeling. *Antioxid Redox Signal* ;**19**:2213-31;2013
 - [594] Dennis K.E., Aschner J.L., Milatovic D., Schmidt J.W., Aschner M., Kaplowitz M.R. et al. NADPH oxidases and reactive oxygen species at different stages of chronic hypoxia-induced

- pulmonary hypertension in newborn piglets. *Am J Physiol Lung Cell Mol Physiol* ;**297**:L596-L607;2009
- [595] Barman S.A., Chen F., Su Y., Dimitropoulou C., Wang Y., Catravas J.D. et al. NADPH oxidase 4 is expressed in pulmonary artery adventitia and contributes to hypertensive vascular remodeling. *Arterioscler Thromb Vasc Biol* ;**34**:1704-15;2014
- [596] Norton C.E., Broughton B.R., Jernigan N.L., Walker B.R., Resta T.C. Enhanced depolarization-induced pulmonary vasoconstriction following chronic hypoxia requires EGFR-dependent activation of NAD(P)H oxidase 2. *Antioxid Redox Signal* ;**18**:1777-88;2013
- [597] Hood K.Y., Mair K.M., Harvey A.P., Montezano A.C., Touyz R.M., MacLean M.R. Serotonin Signaling Through the 5-HT1B Receptor and NADPH Oxidase 1 in Pulmonary Arterial Hypertension. *Arterioscler Thromb Vasc Biol* ;**37**:1361-70;2017
- [598] Wedgwood S., Lakshminrusimha S., Czech L., Schumacker P.T., Steinhorn R.H. Increased p22(phox)/Nox4 expression is involved in remodeling through hydrogen peroxide signaling in experimental persistent pulmonary hypertension of the newborn. *Antioxid Redox Signal* ;**18**:1765-76;2013
- [599] Liu J.Q., Zelko I.N., Erbynn E.M., Sham J.S., Folz R.J. Hypoxic pulmonary hypertension: role of superoxide and NADPH oxidase (gp91phox). *Am J Physiol Lung Cell Mol Physiol* ;**290**:L2-10;2006
- [600] Hoshikawa Y., Ono S., Suzuki S., Tanita T., Chida M., Song C. et al. Generation of oxidative stress contributes to the development of pulmonary hypertension induced by hypoxia. *J Appl Physiol (1985)* ;**90**:1299-306;2001
- [601] Redout E.M., van der Toorn A., Zuidwijk M.J., van de Kolk C.W., van Echteld C.J., Musters R.J. et al. Antioxidant treatment attenuates pulmonary arterial hypertension-induced heart failure. *Am J Physiol Heart Circ Physiol* ;**298**:H1038-H1047;2010
- [602] Freund-Michel V., Guibert C., Dubois M., Courtois A., Marthan R., Savineau J.P. et al. Reactive oxygen species as therapeutic targets in pulmonary hypertension. *Ther Adv Respir Dis* ;**7**:175-200;2013
- [603] Wong C.M., Bansal G., Pavlickova L., Marcocci L., Suzuki Y.J. Reactive oxygen species and antioxidants in pulmonary hypertension. *Antioxid Redox Signal* ;**18**:1789-96;2013
- [604] Aggarwal S., Gross C.M., Sharma S., Fineman J.R., Black S.M. Reactive oxygen species in pulmonary vascular remodeling. *Compr Physiol* ;**3**:1011-34;2013
- [605] Nagaoka T., Gebb S.A., Karoor V., Homma N., Morris K.G., McMurtry I.F. et al. Involvement of RhoA/Rho kinase signaling in pulmonary hypertension of the fawn-hooded rat. *J Appl Physiol (1985)* ;**100**:996-1002;2006
- [606] Truong L., Zheng Y.M., Wang Y.X. Mitochondrial Rieske iron-sulfur protein in pulmonary artery smooth muscle: A key primary signaling molecule in pulmonary hypertension. *Arch Biochem Biophys* ;**664**:68-75;2019
- [607] Ward J.P., McMurtry I.F. Mechanisms of hypoxic pulmonary vasoconstriction and their roles in pulmonary hypertension: new findings for an old problem. *Curr Opin Pharmacol* ;**9**:287-96;2009

- [608] Xing X.Q., Gan Y., Wu S.J., Chen P., Zhou R., Xiang X.D. Rho-kinase as a potential therapeutic target for the treatment of pulmonary hypertension. *Drug News Perspect* ;**19**:517-22;2006
- [609] Weise-Cross L., Resta T.C., Jernigan N.L. Redox Regulation of Ion Channels and Receptors in Pulmonary Hypertension. *Antioxid Redox Signal* ;2019
- [610] Wan J., Yamamura A., Zimnicka A.M., Voiriot G., Smith K.A., Tang H. et al. Chronic hypoxia selectively enhances L- and T-type voltage-dependent Ca²⁺ channel activity in pulmonary artery by upregulating Cav1.2 and Cav3.2. *Am J Physiol Lung Cell Mol Physiol* ;**305**:L154-L164;2013
- [611] Dahan D., Ducret T., Quignard J.F., Marthan R., Savineau J.P., Esteve E. Implication of the ryanodine receptor in TRPV4-induced calcium response in pulmonary arterial smooth muscle cells from normoxic and chronically hypoxic rats. *Am J Physiol Lung Cell Mol Physiol* ;**303**:L824-L833;2012
- [612] Wang J., Weigand L., Lu W., Sylvester J.T., Semenza G.L., Shimoda L.A. Hypoxia inducible factor 1 mediates hypoxia-induced TRPC expression and elevated intracellular Ca²⁺ in pulmonary arterial smooth muscle cells. *Circ Res* ;**98**:1528-37;2006
- [613] Forrest A.S., Joyce T.C., Huebner M.L., Ayon R.J., Wiwchar M., Joyce J. et al. Increased TMEM16A-encoded calcium-activated chloride channel activity is associated with pulmonary hypertension. *Am J Physiol Cell Physiol* ;**303**:C1229-C1243;2012
- [614] Smith K.A., Ayon R.J., Tang H., Makino A., Yuan J.X. Calcium-Sensing Receptor Regulates Cytosolic [Ca (2+)] and Plays a Major Role in the Development of Pulmonary Hypertension. *Front Physiol* ;**7**:517;2016
- [615] Broughton B.R., Jernigan N.L., Norton C.E., Walker B.R., Resta T.C. Chronic hypoxia augments depolarization-induced Ca²⁺-sensitization in pulmonary vascular smooth muscle through superoxide-dependent stimulation of RhoA. *Am J Physiol Lung Cell Mol Physiol* ;2009
- [616] Cantoni S., Cavalli S., Pastore F., Accetta A., Pala D., Vaccaro F. et al. Pharmacological characterization of a highly selective Rho kinase (ROCK) inhibitor and its therapeutic effects in experimental pulmonary hypertension. *Eur J Pharmacol* ;**850**:126-34;2019
- [617] Fike C.D., Dikalova A., Slaughter J.C., Kaplowitz M.R., Zhang Y., Aschner J.L. Reactive oxygen species-reducing strategies improve pulmonary arterial responses to nitric oxide in piglets with chronic hypoxia-induced pulmonary hypertension. *Antioxid Redox Signal* ;**18**:1727-38;2013
- [618] Antigny F., Hautefort A., Meloche J., Belacel-Ouari M., Manoury B., Rucker-Martin C. et al. Potassium Channel Subfamily K Member 3 (KCNK3) Contributes to the Development of Pulmonary Arterial Hypertension. *Circulation* ;**133**:1371-85;2016
- [619] Fu L.C., Lv Y., Zhong Y., He Q., Liu X., Du L.Z. Tyrosine phosphorylation of Kv1.5 is upregulated in intrauterine growth retardation rats with exaggerated pulmonary hypertension. *Braz J Med Biol Res* ;**50**:e6237;2017
- [620] Mondejar-Parreno G., Callejo M., Barreira B., Morales-Cano D., Esquivel-Ruiz S., Moreno L. et al. miR-1 is increased in pulmonary hypertension and downregulates Kv1.5 channels in rat pulmonary arteries. *J Physiol* ;**597**:1185-97;2019

- [621] Hautefort A., Mendes-Ferreira P., Sabourin J., Manaud G., Bertero T., Rucker-Martin C. et al. Bmpr2 Mutant Rats Develop Pulmonary and Cardiac Characteristics of Pulmonary Arterial Hypertension. *Circulation* ;**139**:932-48;2019
- [622] Liu Y., Li M., Warburton R.R., Hill N.S., Fanburg B.L. The 5-HT transporter transactivates the PDGFbeta receptor in pulmonary artery smooth muscle cells. *FASEB J* ;**21**:2725-34;2007
- [623] Pullamsetti S.S., Berghausen E.M., Dabral S., Tretyn A., Butrous E., Savai R. et al. Role of Src tyrosine kinases in experimental pulmonary hypertension. *Arterioscler Thromb Vasc Biol* ;**32**:1354-65;2012
- [624] Fukai K., Nakamura A., Hoshino A., Nakanishi N., Okawa Y., Ariyoshi M. et al. Pyk2 aggravates hypoxia-induced pulmonary hypertension by activating HIF-1alpha. *Am J Physiol Heart Circ Physiol* ;**308**:H951-H959;2015
- [625] Bijli K.M., Kang B.Y., Sutliff R.L., Hart C.M. Proline-rich tyrosine kinase 2 downregulates peroxisome proliferator-activated receptor gamma to promote hypoxia-induced pulmonary artery smooth muscle cell proliferation. *Pulm Circ* ;**6**:202-10;2016
- [626] Zhang X., Shao R., Gao W., Sun G., Liu Y., Fa X. Inhibition of miR-361-5p suppressed pulmonary artery smooth muscle cell survival and migration by targeting ABCA1 and inhibiting the JAK2/STAT3 pathway. *Exp Cell Res* ;**363**:255-61;2018
- [627] Lu X., Bijli K.M., Ramirez A., Murphy T.C., Kleinhenz J., Hart C.M. Hypoxia downregulates PPARgamma via an ERK1/2-NF-kappaB-Nox4-dependent mechanism in human pulmonary artery smooth muscle cells. *Free Radic Biol Med* ;**63**:151-60;2013
- [628] Fan Z., Li C., Qin C., Xie L., Wang X., Gao Z. et al. Role of the PI3K/AKT pathway in modulating cytoskeleton rearrangements and phenotype switching in rat pulmonary arterial vascular smooth muscle cells. *DNA Cell Biol* ;**33**:12-9;2014
- [629] Goncharov D.A., Kudryashova T.V., Ziai H., Ihida-Stansbury K., Delisser H., Krymskaya V.P. et al. Mammalian target of rapamycin complex 2 (mTORC2) coordinates pulmonary artery smooth muscle cell metabolism, proliferation, and survival in pulmonary arterial hypertension. *Circulation* ;**129**:864-74;2014
- [630] Guzy R.D., Hoyos B., Robin E., Chen H., Liu L., Mansfield K.D. et al. Mitochondrial complex III is required for hypoxia-induced ROS production and cellular oxygen sensing. *Cell Metab* ;**1**:401-8;2005
- [631] Adesina S.E., Kang B.Y., Bijli K.M., Ma J., Cheng J., Murphy T.C. et al. Targeting mitochondrial reactive oxygen species to modulate hypoxia-induced pulmonary hypertension. *Free Radic Biol Med* ;**87**:36-47;2015
- [632] Liu P., Gu Y., Luo J., Ye P., Zheng Y., Yu W. et al. Inhibition of Src activation reverses pulmonary vascular remodeling in experimental pulmonary arterial hypertension via Akt/mTOR/HIF-1<alpha> signaling pathway. *Exp Cell Res* ;**380**:36-46;2019
- [633] Green D.E., Kang B.Y., Murphy T.C., Hart C.M. Peroxisome proliferator-activated receptor gamma (PPARgamma) regulates thrombospondin-1 and Nox4 expression in hypoxia-induced human pulmonary artery smooth muscle cell proliferation. *Pulm Circ* ;**2**:483-91;2012

- [634] Sweeney L., Voelkel N.F. Estrogen exposure, obesity and thyroid disease in women with severe pulmonary hypertension. *Eur J Med Res* ;**14**:433-42;2009
- [635] Li S., Tabar S.S., Malec V., Eul B.G., Klepetko W., Weissmann N. et al. NOX4 regulates ROS levels under normoxic and hypoxic conditions, triggers proliferation, and inhibits apoptosis in pulmonary artery adventitial fibroblasts. *Antioxid Redox Signal* ;**10**:1687-98;2008
- [636] Mittal M., Roth M., Konig P., Hofmann S., Dony E., Goyal P. et al. Hypoxia-dependent regulation of nonphagocytic NADPH oxidase subunit NOX4 in the pulmonary vasculature. *Circ Res* ;**101**:258-67;2007
- [637] Iwata K., Ikami K., Matsuno K., Yamashita T., Shiba D., Ibi M. et al. Deficiency of NOX1/nicotinamide adenine dinucleotide phosphate, reduced form oxidase leads to pulmonary vascular remodeling. *Arterioscler Thromb Vasc Biol* ;**34**:110-9;2014
- [638] Seta F., Rahmani M., Turner P.V., Funk C.D. Pulmonary oxidative stress is increased in cyclooxygenase-2 knockdown mice with mild pulmonary hypertension induced by monocrotaline. *PLoS One* ;**6**:e23439;2011
- [639] Nisbet R.E., Graves A.S., Kleinhenz D.J., Rupnow H.L., Reed A.L., Fan T.H. et al. The role of NADPH oxidase in chronic intermittent hypoxia-induced pulmonary hypertension in mice. *Am J Respir Cell Mol Biol* ;**40**:601-9;2009
- [640] Wedgwood S., Lakshminrusimha S., Schumacker P.T., Steinhorn R.H. Cyclic stretch stimulates mitochondrial reactive oxygen species and Nox4 signaling in pulmonary artery smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* ;**309**:L196-L203;2015
- [641] Liu Z., Duan Y.L., Ge S.L., Zhang C.X., Gong W.H., Xu J.J. Effect of estrogen on right ventricular remodeling of monocrotaline-induced pulmonary arterial hypertension in rats and its mechanism. *Eur Rev Med Pharmacol Sci* ;**23**:1742-50;2019
- [642] Jiang Q., Fu X., Tian L., Chen Y., Yang K., Chen X. et al. NOX4 mediates BMP4-induced upregulation of TRPC1 and 6 protein expressions in distal pulmonary arterial smooth muscle cells. *PLoS One* ;**9**:e107135;2014
- [643] Diebold I., Petry A., Hess J., Gorchach A. The NADPH oxidase subunit NOX4 is a new target gene of the hypoxia-inducible factor-1. *Mol Biol Cell* ;**21**:2087-96;2010
- [644] Green D.E., Murphy T.C., Kang B.Y., Kleinhenz J.M., Szyndralewicz C., Page P. et al. The Nox4 inhibitor GKT137831 attenuates hypoxia-induced pulmonary vascular cell proliferation. *Am J Respir Cell Mol Biol* ;**47**:718-26;2012
- [645] Diebold I., Flugel D., Becht S., BelAiba R.S., Bonello S., Hess J. et al. The hypoxia-inducible factor-2alpha is stabilized by oxidative stress involving NOX4. *Antioxid Redox Signal* ;**13**:425-36;2010
- [646] Barman S.A., Fulton D. Adventitial Fibroblast Nox4 Expression and ROS Signaling in Pulmonary Arterial Hypertension. *Adv Exp Med Biol* ;**967**:1-11;2017
- [647] Veith C., Kraut S., Wilhelm J., Sommer N., Quanz K., Seeger W. et al. NADPH oxidase 4 is not involved in hypoxia-induced pulmonary hypertension. *Pulm Circ* ;**6**:397-400;2016

- [648] Rudyk O., Rowan A., Prysyazhna O., Krasemann S., Hartmann K., Zhang M. et al. Oxidation of PKGI α mediates an endogenous adaptation to pulmonary hypertension. *Proc Natl Acad Sci U S A* ;2019
- [649] Zielonka J., Kalyanaraman B. Small-molecule luminescent probes for the detection of cellular oxidizing and nitrating species. *Free Radic Biol Med* ;**128**:3-22;2018
- [650] Liu J.Q., Sham J.S., Shimoda L.A., Kuppusamy P., Sylvester J.T. Hypoxic constriction and reactive oxygen species in porcine distal pulmonary arteries. *Am J Physiol Lung Cell Mol Physiol* ;**285**:L322-L333;2003
- [651] Lee S.J., Seo K.W., Yun M.R., Bae S.S., Lee W.S., Hong K.W. et al. 4-Hydroxynonenal enhances MMP-2 production in vascular smooth muscle cells via mitochondrial ROS-mediated activation of the Akt/NF-kappaB signaling pathways. *Free Radic Biol Med* ;**45**:1487-92;2008
- [652] McNally J.S., Davis M.E., Giddens D.P., Saha A., Hwang J., Dikalov S. et al. Role of xanthine oxidoreductase and NAD(P)H oxidase in endothelial superoxide production in response to oscillatory shear stress. *Am J Physiol Heart Circ Physiol* ;**285**:H2290-H2297;2003
- [653] Abdalla S., Will J.A. Potentiation of the hypoxic contraction of guinea-pig isolated pulmonary arteries by two inhibitors of superoxide dismutase. *Gen Pharmacol* ;**26**:785-92;1995