King's Research Portal

DOI:

[10.1038/s41390-019-0717-9](https://doi.org/10.1038/s41390-019-0717-9)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Story, L., Zhang, T., Steinweg, J. K., Hutter, J., Matthew, J., Dassios, T., Seed, P. T., Pasupathy, D., Allsop, J., Hajnal, J. V., Greenough, A., Shennan, A. H., & Rutherford, M. (2020). Foetal lung volumes in pregnant women who deliver very preterm: a pilot study. *Pediatric Research*, 87, 1066–1071. <https://doi.org/10.1038/s41390-019-0717-9>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Fetal Lung Volumes in Pregnancies that Deliver Very Preterm: a Pilot Study

Cite this article as: Lisa Story, Tong Zhang, Johannes K Steinweg, Jana Hutter, Jacqueline Matthew, Dassios Theodore, Paul T Seed, Dharmintra Pasupathy, Joanna Allsop, Joseph V Hajnal, Anne Greenough, Andrew H Shennan and Mary Rutherford, Fetal Lung Volumes in Pregnancies that Deliver Very Preterm: a Pilot Study, *Pediatric Research* doi:10.1038/s41390-019-0717-9

This Author Accepted Manuscript is a PDF file of an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. <https://www.nature.com/authors/policies/license.html#AAMtermsV1>

Author accepted manuscript

Fetal Lung Volumes in Pregnancies that Deliver Very Preterm: a Pilot Study

Lisa Story^{1*}, Tong Zhang², Johannes K Steinweg², Jana Hutter², Jacqueline Matthew², Dassios Theodore¹, Mr Paul T Seed¹, Dharmindra Pasupathy¹, Joanna Allsop², Joseph V Hajnal², Anne Greenough¹, Andrew H Shennan¹, Mary Rutherford².

1 Department of Women and Children's Health, King's College London, UK

2 Department of Perinatal Health and Imaging, King's College London, UK

Corresponding Author:

Dr Lisa Story

Department of Women and Children's Health

St Thomas' Hospital

London

SE1 7EH

E-mail: lisa.story@kcl.ac.uk

Telephone: +44 (0) 20 7188 7118 x53618

Statement of Financial Support:

This work was supported by: the Wellcome Trust IEH Award (102431), the iFIND project, Engineering and Physical Sciences Research Council, the National Institute for Health Research, Medical Research Council, Tommy's

and the Biomedical Research Centre at Guy's and St Thomas' NHS Foundation Trust.

Disclosures Statement:

AS is chief investigator on a number of trials funded by NIHR and charity sources related to preterm birth prediction and prevention. Hologic, Biomedical and Qiagen have provided samples for these studies. Hologic have provided funding (paid to the institution) to evaluate technical performance of their samples. No other disclosures.

Category of Study:

Clinical Research Article.

Contributions

Dr LS and MR devised the study. LS recruited patients, analysed the data with the assistance PS, and wrote the manuscript. JA was the research radiographer who acquired the MRI images of the preterm cohort. TZ developed the techniques used for lung segmentation and images and protocols were optimised by Dr JH and Prof JH. JM assisted with the calculation of body volumes. DP, TD, AG and AS assisted with data interpretation and writing of the manuscript. JS assisted in the collection of neonatal outcome data.

Abstract

Background

Infants born preterm are at increased risk of pulmonary morbidity. The contribution of antenatal factors to impairments in lung structure/function has not been fully elucidated. This study aimed to compare standardized lung volumes from fetuses that delivered <32 weeks gestation with fetuses that delivered >37 weeks.

Methods

Fourteen women who delivered <32 weeks gestation and 56 women who delivered >37 underwent a fetal MRI. Slice-volume reconstruction was then used and the fetal lungs were then segmented using multi-atlas approaches. Body volumes were calculated by manual segmentation and lung:body volume ratios generated.

Results

Mean gestation at MRI of the preterm group was 27^{+2} weeks (SD 2.9, range 20^{+6} - 31^{+3}) and control group 25^{+3} weeks (SD 4.7 range 20^{+5} - 31^{+6}). Mean gestation at delivery of the preterm group was 29^{+2} (SD 2.6, range 22^{+0} - 32^{+0}). Lung:body volume ratios and fetal lung volumes were smaller in fetuses that delivered preterm both with and without preterm premature rupture of membranes compared to those born at term ($p < 0.001$ in all cases).

Conclusion

Fetuses that deliver very preterm had reduced lung volumes when standardized for fetal size, irrespective of ruptured membranes. These are

novel findings and suggest an antenatal aetiology of insult and possible focus for future preventative therapies.

Author accepted manuscript

Introduction

Preterm birth (PTB) is the most important single determinant of adverse infant outcome with regards to survival and quality of life.(1) Morbidity is inversely correlated to gestational age (GA), and the most significant adverse outcomes are associated with delivery before 32 weeks.

Although the advent of antenatal steroids has significantly improved respiratory morbidity of infants born preterm, specifically respiratory distress syndrome (RDS), it remains a significant complication associated with PTB.(2) RDS results from failure of adequate lung expansion post delivery due to surfactant deficiency. Many of these infants progress to prolonged supplementary oxygen requirement with additional effects from ventilator and hyperoxic induced acute lung injury.(3) This chronic evolving condition is referred to as bronchopulmonary dysplasia (BPD) and complications can persist into adulthood.(4) The aetiology is complex and multifactorial: genetic predisposition and infection have been implicated in its development.(2) Various factors are thought to culminate in inflammatory cytokine and chemokine release, ultimately disrupting alveolar and microvascular development of the peripheral lung.(5) The understanding of BPD associated with PTB, is hampered by a lack of antenatally derived assessments that can be utilized to predict long-term outcomes and mechanisms underlying the disease pathogenesis.(2)

Infection/inflammation has been implicated in the aetiology of PTB and approximately 80% of cases of PTB prior to 28 weeks, have evidence of significant microbial colonization within placental parenchyma.(6) It is therefore hypothesized that infection/inflammation associated with events triggering PTB may additionally affect antenatal lung development. No studies have explored lung volumes standardized for fetal weight that deliver very preterm or standardized the lung volume for the overall size of the fetus. This study therefore has two aims: 1) to establish MRI derived normative ranges for fetal lung volumes standardising for fetal size and 2) to compare standardised lung volumes in fetuses that subsequently delivered <32 weeks gestation with a group of control fetuses that delivered at term.

Methods

Mothers of fetuses at high-risk of PTB were prospectively recruited from the antenatal ward and the Preterm Surveillance Clinic at St Thomas' Hospital London between December 2015 and October 2017.

Inclusion criteria were: GA 20-32 weeks, high-risk of PTB encompassing either asymptomatic women with either a history of previous PTB, late miscarriage >16 weeks or cervical surgery with a >50% risk of PTB in the next two weeks (based on an algorithm derived from quantitative cervico-vaginal fibronectin and cervical length(7)) or PPROM. Exclusion criteria were: fetuses known to have structural or chromosomal abnormalities, multiple pregnancies, active labour, inability to give informed consent, pregnancy complications such as pre-eclampsia or gestational diabetes, contraindications to MRI such as claustrophobia or a recently sited metallic implant.

Following assessment of eligibility, women were invited to participate and written consent obtained. Fetal MRI was performed using a 1.5T Philips Ingenia MRI System (Philips Medical systems, Best, the Netherlands) with a 32-channel cardiac coil placed around the mother's abdomen. Imaging of the fetal trunk was performed using T2 weighted single shot turbo spin echo (ssTSE), acquired in three orthogonal planes using the following scanning parameters: TR=25991 ms, TE=80ms, slice thickness 2.5mm, slice overlap 1.25 mm and flip angle=90°. In addition T2 weighted ssTSE images were acquired (using a larger slice thickness/lower resolution) to provide coverage

of the whole uterus, again in three orthogonal planes. Two subjects were imaged at 3T on a Philips Achieva system using otherwise similar procedures.

To obtain volumetric data of the lungs, for each subject, all available T2 weighted ssTSE stacks covering the fetal thorax, which were typically corrupted by fetal movement between individual slices, were motion corrected and combined to produce a single 3D volumetric image, using slice to volume reconstruction (SVR).(8, 9).

Normal anatomical atlases were derived from the volumetric reconstructions of 25 previously imaged normal subjects (20^{+1} - 31^{+6} weeks GA). A mask was initially placed over the fetal thorax and then lung volumes were manually segmented for each of these normal subjects from the SVR image using ITK SNAP (version 2.2.0)(10).

Lung segmentation was then performed automatically on the study fetuses using the age appropriate atlases as follows: eight age matched atlas subjects were selected and each was first affine registered to the study fetus segmented target image to achieve a global pose alignment(10). Free form deformation (FFD) based non-linear registration with control points spaced at 3 mm intervals was used to estimate the local deformations. After registration the atlas segments were transformed to the target subject using linear interpolation and a final segmentation was determined by 'majority vote'.(10).

Segmentation of the fetal body and amniotic fluid was performed using the uterine ssFSE images using ITK-SNAP (version 2.2.0) in a two-step process. In each case a rough automatic segmentation of the body was obtained based on image contrast while utilizing user-defined thresholds. Editing was then performed manually. Intra and inter-observer variability had previously been confirmed (intraclass correlation co-efficient (ICC) intra-observer variability 0.85 and inter-observer ICC 0.971). Lung: body volumes were then calculated.

A control group was identified from the intelligent fetal imaging and diagnosis project (www.iFINDproject.com). This is a study conducted at King's College London which aims to improve fetal ultrasound imaging through automated image processing. Part of this study combines conventional ultrasound imaging with more detailed MRI scans to build a comprehensive map of fetal anatomy to use for computer assisted diagnosis of fetal anomalies. Cases were selected from low risk pregnancies where the MRI was performed between 20⁺⁰ and 32⁺⁰ weeks gestation and delivery occurred >37 weeks. All fetuses underwent the same imaging sequences and reconstruction as described above.

Details of maternal demographics, timing of steroid administration, delivery and neonatal parameters were collected including: gestation at delivery, sex of infant, birthweight, birthweight centile, neonatal unit admission, number of days of invasive ventilation, Continuous Positive Airway Pressure (CPAP), and need for supplemental oxygen.

Data were assessed for normality using distributional plots of standardized residuals. Demographic and neonatal outcome data were analysed using a student t-test where data were continuous and Chi-squared when categorical. For the low risk control cases delivering at term, obtained from the iFIND study, the normal growth trajectory of lung volume and lung:body volume ratios between 20 and 32 weeks gestation was estimated by maximum likelihood regression, using the Stata command `xrml`.⁽¹¹⁾ Gestation-adjusted centiles were then calculated for all lung volumes and lung:body volume ratios. ROC curves were generated for low lung:body volume ratio and low gestation adjusted lung volume centiles as predictors of PTB. Adjusting for the effects of gestation, multiple regression analysis was used to produce estimates between fetuses that delivered preterm and those that delivered at term for lung and body volumes and lung:body volume ratios. Multiple regression analysis was also used to compare amniotic fluid volumes between fetuses that delivered preterm with and without ruptured membranes and the control group, accounting for the effects of gestation.

For neonatal outcome data (not normally distributed) the Spearman Rank Correlation was used. Statistical analysis was performed using the SPSS software package (version 25, SPSS IBM) and Stata version 14.0 (StataCorp, College Station, Texas, USA).

This study was conducted under the ethics: 07/H0707/105 and 14/LO/1806

Results

Thirty-eight women were identified as eligible to participate in the study group: 35 agreed to undergo MRI imaging: 17 delivered prior to MRI (median two days from agreeing to participate, range 0-11 days). Eighteen patients had an MRI: three delivered >32 weeks. Lung reconstruction was not possible in one case leaving 14 suitable for analysis. Six women had ruptured and eight intact membranes at the time of imaging. Twelve were performed on a 1.5T and two on a 3T scanner. Eleven of the 14 cases had antenatal steroids prior to imaging. Fifty-six controls were identified from the iFIND study (all healthy volunteers) as a control group.

Clinical characteristics of the cohorts can be seen in Table 1 and lung volume, body volume and lung:body volume ratios in Table 2. Distribution plots for lung volumes and lung:body volume ratios from 20-32 weeks gestation derived from the control group of fetuses are shown in Figure 1. In fetuses that delivered at term, the lung volume comprised 4.2% of the total body volume (standard deviation 0.6%) compared to 3.4% in the preterm group (standard deviation 0.6%).

The gestation adjusted lung volume centiles and the lung:body volume ratio were tested as predictors of PTB (Figure 2). Fetuses that delivered preterm had significantly lower lung volumes, body volumes and lung:body volume ratios at the time of scan than those that delivered at term, allowing for gestational change (Table 2 and Figure 3). Excluding the two fetuses scanned at 3T did not affect this finding (accounting for different magnetic

field strengths potentially causing contrast differences and measurement discrepancies). These results were observed in both groups: fetuses with ruptured (mean 33900 mm³ SD 13500) and intact membranes (mean 27200mm³ SD 13200 p<0.001 in both cases). Body volumes were also significantly smaller in fetuses that subsequently delivered preterm (both with (mean 981000 mm³ SD 397000, p=0.007) and without (mean 861000mm³ SD 421000 p=0.009) ruptured membranes) when adjusted for GA at MRI. When standardised for fetal size, fetuses that delivered preterm had significantly lower lung:body volume ratios. These effects were again observed in both fetuses that had intact membranes (mean 0.035 SD 0.006) and those with ruptured membranes (mean 0.032 SD 0.007) at the time of MRI (p<0.001 in both cases).

Low values of both lung volume and lung:body volume ratio appeared predictive of preterm delivery with the gestation adjusted centile being superior.

There was no significant difference in the amniotic fluid volume between the control group and fetuses that delivered <32 weeks with intact membranes.

There was one intrapartum death at 22⁺⁰ weeks gestation. The mean number of days from MRI to delivery in the preterm group was 10.5 (range 0-48). The median number of days of rupture of membranes to MRI in the group that had PPRM was 4 (range 3-18). Two fetuses that delivered >37 weeks were

admitted to the neonatal unit, one for a few hours with suspected RDS and the second for two days with hypoglycaemia not requiring respiratory support.

There was no correlation between respiratory outcomes (length of ventilation and supplemental oxygen use) and antenatal lung:body volume ratios (supplementary table S1). Placental histology was available for 12 of the cases: ten had confirmed chorioamnionitis.

Discussion

We have reported normal ranges for MRI derived fetal lung volumes and lung:body volume ratios of fetuses 20-32 weeks gestation. We found a significant reduction in lung volumes in both fetuses with and without ruptured membranes that delivered <32 weeks when standardized for body volume compared to fetuses that delivered >37 weeks. Fetuses that delivered preterm were globally smaller, when antenatal MRI derived body volumes were compared with a control population that delivered at term.

Absolute Lung Volumes

Previous studies have reported a reduction in absolute MRI derived lung volumes in fetuses with PPROM in comparison with control cohorts(12, 13). This finding may partly be explained by the fact that lung volume in utero also comprises of some amniotic fluid. However, our finding that a reduction in lung volume also occurs in fetuses with intact membranes is a novel one. Unlike the previous studies in fetuses with PPROM, this study is also the first to standardize the lung volume for overall fetal size.

Body Volumes

Our study has demonstrated that fetuses that delivered preterm were globally smaller, on antenatal MRI imaging, however there was no statistical difference in the birthweight centile between fetuses that delivered preterm and those that delivered at term.

There has been discussion as to whether reference centiles for preterm babies should be based on 'typical' preterm infants, or on healthy growing fetuses that will be delivered at term. The World Health Organisation birthweight centiles(14, 15) (used to evaluate birthweight centiles in this study) and the INTERGROWTH centiles(16) both use information from 'typical' preterm deliveries. Gardosi's customized birthweight centiles, as well as adjusting for maternal and neonatal features, use healthy term deliveries for their reference sample and then extrapolate backwards for earlier gestations.(17)

This study, and other studies using MRI scanning for fetal volume, allows direct measurement of body volume in fetuses who later delivered both term and preterm. Using volume as a proxy for weight, our results contribute to the discussion on fetal reference standards noted above. Unfortunately, due to the limitations of data capture in this study it was not possible to calculate customized birthweight centiles, however, if our results are confirmed, in a larger more ethnically representative sample, they would imply that the factors that drive spontaneous PTB may have a detrimental effect on

fetal growth antenatally. This is supported by the findings of Partap et al, who undertook a prospective cohort study of nulliparous women with a singleton pregnancy assessing 3892 women at 20 and 28 weeks gestation using ultrasound. 2.5% of women underwent a spontaneous PTB. It was noted that slow growth velocity of the femur was associated with an increased risk of spontaneous PTB.(18)

Lung:body volume ratios

Lung:body volumes were created in order to standardize for fetal size. The lung:body volume ratio was consistent across gestation in the control group. In fetuses that delivered <32 weeks, lung:body volume ratios were significantly lower than in fetuses that delivered >37 weeks gestation. This finding was consistent in fetuses with both ruptured and intact membranes. Low values of both lung volume and lung:body volume ratio appeared predictive of preterm delivery with the gestation adjusted centile being superior.

Lung:body volume ratios have also been derived using MRI to investigate fetuses with congenital diaphragmatic hernia,(19). Although 44 normal fetuses were assessed the normative data was not explicitly reported in their paper. No studies have previously used lung:body volume ratios to investigate fetuses that deliver preterm.

Although a major determinant of lung volume will be amniotic fluid volume, we have also demonstrated that fetuses that have intact membranes that deliver

prior to 32 weeks of gestation also have a reduction in lung:body volume ratios. We hypothesize that this may be due to cytokines implicated in the aetiology of PTB also affecting pulmonary development. Infection/inflammation is reported to be associated in pathways mediating PTB(20), common even with intact membranes. Fetal membranes express immune sensors and mount a response to microbial products with inflammatory activation(21), which can include cytokines that promote neutrophil and monocyte infiltration.(22) Elevated levels of interleukin (IL)1- β and IL-8 have been reported in umbilical cord blood of neonates delivered with evidence of the fetal inflammatory response.(23) 83% percent of cases that delivered preterm in this study had histological evidence of chorioamnionitis, where histology was available, post delivery.

Although a number of studies have demonstrated that exposure to intra-amniotic infection is protective for the development of RDS,(24) the relationship between BPD and antenatal infection/inflammation is less well defined. In PTB normal pulmonary development is disrupted.(25) Pathology specimens of infants with BPD demonstrate a decreased number of alveoli which are larger and more simplified in structure and blunted pulmonary microvascular growth.(26) Intrauterine infection has been shown to spread to the lung causing injury and remodeling in a sheep model(27) resulting in persistent changes in lung morphometry(28) and vascular development.(29, 30). It is therefore possible that these pathological processes are affecting fetal lung development.

Numbers in this study are currently small, however no strong correlations were observed between antenatal lung:body volume ratios and short-term neonatal respiratory outcomes. Further studies will therefore be warranted to explore this association, increasing the sample size to confirm the findings and taking into account the multiple confounding factors perinatally well as the varying times between the antenatal MRI and delivery. The association between markers of infection/inflammation both in maternal samples, umbilical cord blood and neonatal samples with imaging findings will also need further exploration.

Conclusions

Although it should be noted that numbers in this study are small to date it does appear that antenatal lung and body volumes are smaller in fetuses that subsequently deliver preterm. In the future, more work is warranted to see if these findings are reproducible in a larger group of fetuses. If this is the case the findings are novel and may suggest an antenatal aetiology of insult. This knowledge may facilitate individualization of postnatal treatment plans and provide a possible focus for future preventive therapies.

Acknowledgements

This work was undertaken with the support of patients and staff at St Thomas's Hospital London and was supported by: the Wellcome Trust IEH Award (102431), the iFIND project, Engineering and Physical Sciences Research Council, the National Institute for Health Research, Medical Research Council, Tommy's and the Biomedical Research Centre at Guy's

and St Thomas' NHS Foundation Trust. Miss Amelie Spangenberg (medical student at King's College London) assisted with analysis of the amniotic fluid volume data.

Author accepted manuscript

Author accepted manuscript

References

1. Saigal S, Doyle LW 2008 An overview of mortality and sequelae of preterm birth from infancy to adulthood. *Lancet* 371:261-269.
2. Maitre NL, Ballard RA, Ellenberg JH, Davis SD, Greenberg JM, Hamvas A, Pryhuber GS, Prematurity, Respiratory Outcomes P 2015 Respiratory consequences of prematurity: evolution of a diagnosis and development of a comprehensive approach. *J Perinatol* 35:313-321.
3. Bolton CE, Bush A, Hurst JR, Kotecha S, McGarvey L 2015 Lung consequences in adults born prematurely. *Thorax* 70:574-580.
4. Hilgendorff A, O'Reilly MA 2015 Bronchopulmonary dysplasia early changes leading to long-term consequences. *Front Med (Lausanne)* 2:2.
5. Jobe AJ 1999 The new BPD: an arrest of lung development. *Pediatr Res* 46:641-643.
6. Onderdonk AB, Hecht JL, McElrath TF, Delaney ML, Allred EN, Leviton A, Investigators ES 2008 Colonization of second-trimester placenta parenchyma. *Am J Obstet Gynecol* 199:52 e51-52 e10.
7. Kuhrt K, Hezelgrave N, Foster C, Seed PT, Shennan AH 2015 Development and validation of a predictive tool for spontaneous preterm birth, incorporating quantitative fetal fibronectin, in symptomatic women. *Ultrasound Obstet Gynecol*.
8. Kuklisova-Murgasova M, Quaghebeur G, Rutherford MA, Hajnal JV, Schnabel JA 2012 Reconstruction of fetal brain MRI with intensity matching and complete outlier removal. *Med Image Anal* 16:1550-1564.
9. Kainz B, Steinberger M, Wein W, et al. 2015 Fast Volume Reconstruction From Motion Corrupted Stacks of 2D Slices. *IEEE Trans Med Imaging* 34:1901-1913.
10. Rueckert D, Sonoda LI, Hayes C, Hill DL, Leach MO, Hawkes DJ 1999 Nonrigid registration using free-form deformations: application to breast MR images. *IEEE Trans Med Imaging* 18:712-721.
11. Royston P, Wright EM 1998 How to construct 'normal ranges' for fetal variables. *Ultrasound Obstet Gynecol* 11:30-38.
12. Kaspran G, Brugger PC, Helmer H, Langer M, Balassy C, Prayer D 2006 [Fetal lung development on MRT. Normal course and impairment due to premature rupture of membranes]. *Radiologe* 46:120-127.
13. Messerschmidt A, Patariaia A, Helmer H, et al. 2011 Fetal MRI for prediction of neonatal mortality following preterm premature rupture of the fetal membranes. *Pediatr Radiol* 41:1416-1420.
14. Freeman JV, Cole TJ, Chinn S, Jones PR, White EM, Preece MA 1995 Cross sectional stature and weight reference curves for the UK, 1990. *Arch Dis Child* 73:17-24.
15. Norris T, Seaton SE, Manktelow BN, et al. 2018 Updated birth weight centiles for England and Wales. *Arch Dis Child Fetal Neonatal Ed* 103:F577-F582.
16. Villar J, Papageorghiou AT, Pang R, et al. International F, Newborn Growth Consortium for the 21st C 2014 The likeness of fetal growth and newborn size across non-isolated populations in the INTERGROWTH-21st Project: the Fetal Growth Longitudinal Study and Newborn Cross-Sectional Study. *Lancet Diabetes Endocrinol* 2:781-792.

17. Gardosi J, Francis A, Turner S, Williams M 2018 Customized growth charts: rationale, validation and clinical benefits. *Am J Obstet Gynecol* 218:S609-S618.
18. Partap U, Sovio U, Smith GC 2016 Fetal Growth and the Risk of Spontaneous Preterm Birth in a Prospective Cohort Study of Nulliparous Women. *Am J Epidemiol* 184:110-119.
19. Weidner M, Hagelstein C, Debus A, et al. 2014 MRI-based ratio of fetal lung volume to fetal body volume as a new prognostic marker in congenital diaphragmatic hernia. *AJR Am J Roentgenol* 202:1330-1336.
20. Keelan JA 2018 Intrauterine inflammatory activation, functional progesterone withdrawal, and the timing of term and preterm birth. *J Reprod Immunol* 125:89-99.
21. Hoang M, Potter JA, Gysler SM, et al. 2014 Human fetal membranes generate distinct cytokine profiles in response to bacterial Toll-like receptor and nod-like receptor agonists. *Biol Reprod* 90:39.
22. Kim CJ, Romero R, Chaemsathong P, Chaiyasit N, Yoon BH, Kim YM 2015 Acute chorioamnionitis and funisitis: definition, pathologic features, and clinical significance. *Am J Obstet Gynecol* 213:S29-52.
23. Mestan K, Yu Y, Thorsen P, et al. 2009 Cord blood biomarkers of the fetal inflammatory response. *J Matern Fetal Neonatal Med* 22:379-387.
24. Kunzmann S, Collins JJ, Kuypers E, Kramer BW 2013 Thrown off balance: the effect of antenatal inflammation on the developing lung and immune system. *Am J Obstet Gynecol* 208:429-437.
25. Balany J, Bhandari V 2015 Understanding the Impact of Infection, Inflammation, and Their Persistence in the Pathogenesis of Bronchopulmonary Dysplasia. *Front Med (Lausanne)* 2:90.
26. Husain AN, Siddiqui NH, Stocker JT 1998 Pathology of arrested acinar development in postsurfactant bronchopulmonary dysplasia. *Hum Pathol* 29:710-717.
27. Kramer BW, Ladenburger A, Kunzmann S, et al. 2009 Intravenous lipopolysaccharide-induced pulmonary maturation and structural changes in fetal sheep. *Am J Obstet Gynecol* 200:195 e191-110.
28. Willet KE, Jobe AH, Ikegami M, Newnham J, Brennan S, Sly PD 2000 Antenatal endotoxin and glucocorticoid effects on lung morphometry in preterm lambs. *Pediatr Res* 48:782-788.
29. Kallapur SG, Bachurski CJ, Le Cras TD, Joshi SN, Ikegami M, Jobe AH 2004 Vascular changes after intra-amniotic endotoxin in preterm lamb lungs. *Am J Physiol Lung Cell Mol Physiol* 287:L1178-1185.
30. Thomas W, Seidenspinner S, Kramer BW, et al. 2011 Airway angiopoietin-2 in ventilated very preterm infants: association with prenatal factors and neonatal outcome. *Pediatr Pulmonol* 46:777-784.

Figure 1: Normal ranges and 3, 10, 50, 90 and 97th centiles for MRI derived (a) lung volumes and (b) lung:body volume ratios between 20 and 32 weeks gestation obtained from a 1.5T MRI scanner.

The expected mean lung volume (cm³), Z-score and gestation-adjusted centile at a given gestational age (GA) were estimated:

$$m = -447.4 \times ga + 233.5 \times \log_e(ga) + 576$$

$$SD = 0.1821 \times \text{expected mean}$$

$$Z \text{ score for any individual fetus} = (\text{measured lung volume} - m)/SD$$

The expected lung: body volume ratio was estimated, with no detectable variation of healthy lung:body volume ratios with gestational age:

$$m = 0.0419881, SD = 0.0062148$$

$$Z \text{ score for any individual fetus} = (\text{measured lung volume} - m)/SD$$

The expected lung: body volume ratio was estimated, with no detectable variation of healthy lung:body volume ratios with gestational age:

$$m = 0.0419881, SD = 0.0062148$$

$$Z \text{ score for any individual fetus} = (\text{measured lung volume} - m)/SD$$

Figure 2: ROC curve of prediction of prematurity from antenatal MRI scans: lung volume, gestation adjusted centile and lung:body volume ratio.

Figure 3: Graphs illustrating the relationship between gestation at MRI and (a) lung volume (b) body volume and (c) lung: body volume ratio in

fetuses that delivered <32 weeks gestation and those who delivered >37 weeks gestation. The line of best fit and 95% confidence interval is given for the control group.

Author accepted manuscript

Characteristic	Preterm cohort (n=14)	Term cohort (n=56)	P
Maternal age, y Mean (SD) Range	35 (5.7) 28-45	33 (4.7) 20-42	0.26
BMI, kg/m ² Mean $\pm$ SD Range	23.5 (2.9) 21-29	23.3 (4.7) 20-42	0.67
Ethnicity n (%) White Black South Asian East Asian Other	6 (43) 3 (21) 3 (21) 1 (7) 1 (7)	48 (85) 1 (2) 1 (2) 1 (2) 5 (9)	<0.001
Parity n (%) 0 1 2 3 4	8 (57) 3 (23) 1 (8) 1 (8) 1 (8)	37 (66) 16 (29) 3 (5) 0 0	0.01
GA at MRI, wk Mean (SD) Range	27.2 (3.3) 20.6-31.4	25.4 (2.5) 20.7-31.9	0.09
GA at birth, wk Mean (SD) Range	29.2 (2.6) 22-31.9	40 (1.2) 37.3-42.1	<0.001
Birthweight, g Mean (SD) Range	1353 (330) 770-1875	3429 (489) 2200-4560	<0.001
Birthweight Centile of Livebirths			

0-3	0	3 (5)	0.49
3-10	0	4 (7)	
10-25	2 (15)	8 (14)	
25-50	4 (31)	10 (18)	
50-75	5 (38)	19 (34)	
75-90	0	7 (13)	
90-97	2 (15)	4 (7)	
97-100	0	1 (2)	
Sex of Infant n (%)			0.25
Female	7 (50)	35 (63)	
Male	6 (43)	21 (38)	
Undetermined	1 (7)	0	
Outcome n (%)			0.2
Live to discharge	13 (93)	56 (100)	
Neonatal/intrapartum death	1 (7)		

Table 1: Clinical Characteristics of the Cohort (t-test used for analysis where data was continuous and Chi-squared where data categorical)

Variable	Term Cohort Mean (SD) (n=56)	Preterm Cohort Mean (SD) (n=14)	Difference in Preterm Cohort (95% CI)	P
Lung Volume, mm ³	34700 (17000)	31000 (13300)	-13300 (-18500 to -8100)	<0.001
Body Volume, mm ³	8235300 (725000)	929500 (396300)	-136300 (-208300 to -64400)	<0.001
Lung:Body Volume Ratio	0.042 (0.006)	0.034 (0.006)	-0.008 (-0.12 to -0.004)	<0.001

Table 2: of MRI parameters of fetal MRIs controlling for the effect of gestational age at MRI scan by linear regression.

(a)

(b)