



King's Research Portal

DOI:

[10.1113/JP279518](https://doi.org/10.1113/JP279518)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Baburamani, A. A., & Arichi, T. (2020). Complementing cooling: the ongoing search for an effective adjunct to therapeutic hypothermia. *The Journal of physiology*, 598(5), 905-906. <https://doi.org/10.1113/JP279518>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Complementing cooling: the ongoing search for an effective adjunct to therapeutic hypothermiaAna A. Baburamani¹, Tomoki Arichi^{1,2,3}

¹Centre for the Developing Brain, School of Biomedical Engineering & Imaging Sciences, King's College London, King's Health Partners, St Thomas' Hospital, London, SE1 7EH; ² Department of Bioengineering, Imperial College London, South Kensington, London SW7 2AZ; ³ Children's Neurosciences, Evelina London Children's Hospital, London, UK.

Running Title: Adjunct Therapies to Hypothermia

Hypoxic-ischaemic encephalopathy (HIE) remains a leading cause of neonatal death and adverse neurodevelopmental outcome, affecting 2-4 in 1000 live births. In cases of moderate to severe HIE, the current standard of care in near-term or term infants is therapeutic hypothermia initiated within 6 hours of the asphyxic insult. Preclinical and clinical studies have been fundamental in establishing the optimal temperature reduction (3°-5°C) and treatment duration (up to 72h); and importantly have shown that longer or deeper cooling protocols do not improve neuroprotection and in some instances have even demonstrated decreased efficacy. Furthermore, as 8-9 infants need to be treated with optimised therapeutic hypothermia for one infant to survive and/or reduce long term neurodisability

This is an Accepted Article that has been peer-reviewed and approved for publication in the The Journal of Physiology, but has yet to undergo copy-editing and proof correction. Please cite this article as an 'Accepted Article'; [doi: 10.1113/JP279518](https://doi.org/10.1113/JP279518).

This article is protected by copyright. All rights reserved.

(Edwards *et al.*, 2010), there remains a clear need to identify adjunct therapies that can further improve outcomes for this vulnerable population.

One such adjunct therapy is erythropoietin (Epo), which has been demonstrated in preclinical studies and a small number of clinical studies to reduce cellular signs of brain injury, macroscopic evidence of injury as assessed by MRI, and result in trends towards improved clinical outcomes in HIE. Moreover, it is an attractive neuroprotective agent as it has an essential role for normal neurodevelopment and has been implicated in injury repair mechanisms through eliciting acute (anti-apoptotic, anti-inflammatory, anti-oxidant and anti-excitotoxic) and regenerative (angiogenesis, neurogenesis, oligodendrogenesis) mechanisms. In addition, the pharmacokinetics of high dose recombinant (rEpo) are now well defined and importantly, it has also been shown to be safe when given in combination with hypothermia in neonates with HIE (Wu *et al.*, 2012).

In this issue of *The Journal of Physiology*, Wassink *et al.*, (2020) investigate whether high-dose rEpo combined with moderate hypothermia in a fetal sheep model of severe global cerebral ischaemia provides added neuroprotection over hypothermia alone. To ensure clinical relevance, their protocol included delayed onset of both hypothermia and treatment, with the infusion of high-dose rEpo initiated 3 hours after the ischaemic insult. Importantly, whilst independently neuroprotective in comparison to the ischaemia-vehicle, Wassink *et al.*, (2020) found there was no enhanced benefit from combining rEpo with therapeutic hypothermia on assessment of both physiological parameters (EEG, heart rate, carotid artery blood flow) and cellular histology. Of added concern, ischaemia-rEpo-hypothermia was also associated with increased cortical caspase-3 positive cells, suggesting increased ongoing cell death even after 7 days of recovery in the combined therapy group.

Although the results of Phase 3 clinical trials of rEpo are still awaited, this study highlights existing concerns about its efficacy as an adjunct clinical therapy for HIE. Along with rEpo, several other agents have also been identified through preclinical studies as promising post-natal adjunct therapies to compliment hypothermia (Robertson *et al.*, 2012; Wassink *et al.*, 2019). However, to date

unfortunately none have been able to demonstrate significant effects as adjunct agents with therapeutic hypothermia in a large clinical trial. Xenon, a monoatomic gas which readily crosses the blood-brain barrier, was found to be of significant benefit in preclinical studies but when taken to a randomised control trial, there was no added benefit when inhaled Xenon was combined with therapeutic hypothermia (Azzopardi *et al.*, 2016). Melatonin similarly is known to cross the blood brain barrier and is an endogenous hormone which is thought have a key role in neural repair through reducing oxidative stress. Although a single small clinical study (Aly *et al.*, 2015) described significant improvements in MRI white matter appearances and short term developmental outcome, a larger scale clinical trial has not been completed. Moreover, the promising results of pre-clinical piglet studies should be considered with caution as effectiveness with delayed initiation of treatment and the potentially adverse effects of the ethanol used as a diluent to deliver melatonin have not been studied in detail (Robertson *et al.*, 2013; Davidson *et al.*, 2015). There are also several clinical trials in progress of autologous stem cells derived from cord blood or the placenta, although their precise mechanism of action for this application is still unclear and thus defining their optimal use remains challenging (Bennet *et al.*, 2012).

So for now, as the search continues, where to next? It is important to consider that the success of therapeutic hypothermia was achieved through a bench to bedside translational pathway which included first establishing the cellular nature and delayed natural history of the underlying brain injury, the mechanisms of potential neuroprotection and then the optimal delivery of the therapy. A vital step in this pathway is therefore to also have preclinical models and protocols which can recapitulate the conditions and standards of clinical care to maintain relevance. Through this, it is likely that the key to identifying the next adjunct therapy will be to understand how the mechanism of action of any potential agent interacts with and hopefully directly complements hypothermia; rather than acting through the same biological route and potentially competing/interfering with hypothermic neuroprotection as appears to be the case with rEpo. Any novel therapy must also be feasible to deliver in the clinical environment, have a favourable risk/benefit ratio, all within the context of significantly improving outcome above and beyond a proven successful therapy.

Words: 843

Additional Information

Conflicts of Interests

None declared.

Author Contributions

Both authors have equally contributed to conception, drafting and revising the manuscript, having read and approved the final version. They agree to be accountable for all aspects of the work.

Funding

We gratefully acknowledge financial support from Sparks and Great Ormond Street Hospital Children's Charity [V5318]; the Wellcome/EPSRC Centre for Medical Engineering [WT 203148/Z/16/Z]; Dr Arichi is supported by an MRC Clinician Scientist fellowship [MR/P008712/1]. We also gratefully acknowledge support from National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) based at Guy's and St Thomas' NHS Foundation Trust and King's College London and supported by the NIHR Clinical Research Facility (CRF) at Guy's and St Thomas'. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

References

Aly H, Elmahdy H, El-Dib M, Rowisha M, Awany M, El-Gohary T, Elbatch M, Hamisa M & El-Mashad AR. (2015). Melatonin use for neuroprotection in perinatal asphyxia: a randomized controlled pilot study. *J Perinatol* **35**, 186-191.

This article is protected by copyright. All rights reserved.

Azzopardi D, Robertson NJ, Bainbridge A, Cady E, Charles-Edwards G, Deierl A, Fagiolo G, Franks NP, Griffiths J, Hajnal J, Juszczak E, Kapetanakis B, Linsell L, Maze M, Omar O, Strohm B, Tusor N & Edwards AD. (2016). Moderate hypothermia within 6 h of birth plus inhaled xenon versus moderate hypothermia alone after birth asphyxia (TOBY-Xe): a proof-of-concept, open-label, randomised controlled trial. *Lancet neurology* **15**, 145-153.

Bennet L, Tan S, Van den Heuvel L, Derrick M, Groenendaal F, van Bel F, Juul S, Back SA, Northington F, Robertson NJ, Mallard C & Gunn AJ. (2012). Cell therapy for neonatal hypoxia-ischemia and cerebral palsy. *Ann Neurol* **71**, 589-600.

Davidson JO, Wassink G, van den Heuvel LG, Bennet L & Gunn AJ. (2015). Therapeutic Hypothermia for Neonatal Hypoxic-Ischemic Encephalopathy - Where to from Here? *Front Neurol* **6**, 198.

Edwards AD, Brocklehurst P, Gunn AJ, Halliday H, Juszczak E, Levene M, Strohm B, Thoresen M, Whitelaw A & Azzopardi D. (2010). Neurological outcomes at 18 months of age after moderate hypothermia for perinatal hypoxic ischaemic encephalopathy: synthesis and meta-analysis of trial data. *BMJ* **340**, c363.

Robertson NJ, Faulkner S, Fleiss B, Bainbridge A, Andorka C, Price D, Powell E, Lecky-Thompson L, Thei L, Chandrasekaran M, Hristova M, Cady EB, Gressens P, Golay X & Raivich G. (2013). Melatonin augments hypothermic neuroprotection in a perinatal asphyxia model. *Brain* **136**, 90-105.

Robertson NJ, Tan S, Groenendaal F, van Bel F, Juul SE, Bennet L, Derrick M, Back SA, Valdez RC, Northington F, Gunn AJ & Mallard C. (2012). Which neuroprotective agents are ready for bench to bedside translation in the newborn infant? *J Pediatr* **160**, 544-552 e544.

Wassink G, Davidson JO, Dhillon SK, Zhou K, Bennet L, Thoresen M & Gunn AJ. (2019). Therapeutic Hypothermia in Neonatal Hypoxic-Ischemic Encephalopathy. *Curr Neurol Neurosci Rep* **19**, 2.

Wassink G, Davidson JO, Fraser M, Yuill CA, Bennet L & Gunn AJ. (2020). Non-additive effects of adjunct erythropoietin therapy with therapeutic hypothermia after global cerebral ischaemia in near-term fetal sheep. *J Physiol*.

Wu YW, Bauer LA, Ballard RA, Ferriero DM, Glidden DV, Mayock DE, Chang T, Durand DJ, Song D, Bonifacio SL, Gonzalez FF, Glass HC & Juul SE. (2012). Erythropoietin for neuroprotection in neonatal encephalopathy: safety and pharmacokinetics. *Pediatrics* **130**, 683-691.