King's Research Portal

DOI:

[10.1176/appi.ps.201100521](https://doi.org/10.1176/appi.ps.201100521)

Document Version

Early version, also known as pre-print

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Slade, M. (2012). Everyday Solutions for Everyday Problems: How Mental Health Systems Can Support Recovery. *Psychiatric Services*, 63(7), 702-704. <https://doi.org/10.1176/appi.ps.201100521>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Everyday solutions for everyday problems

Acknowledgements

The author is grateful to Miles Rinaldi for help in developing these ideas.

Disclosures

No interests to disclose.

Abstract

People who experience mental illness can be viewed as either fundamentally different from, or fundamentally like, everyone else in society. Recovery-oriented mental health systems focus on commonality. In practice, this involves an orientation towards supporting everyday solutions for everyday problems, rather than on providing specialist treatments for mental illness-related problems. This change is illustrated in relation to help offered with housing, employment, relationships and spirituality. Interventions may contribute to the process of striving for a life worth living, but are a means not an end. Mental health systems which offer treatments in support of the individual's life goals are very different to those which treat patients in their best interests. The strongest contribution of mental health services to recovery is to support everyday solutions to everyday problems.

Mainstream and exceptionalist public policy

Is it more helpful to view people who experience mental illness as fundamentally different from, or fundamentally similar to, other members of their community? This question underpins a debate about the best balance between mainstream public policy and exceptionalist mental health-specific policy in the USA (1). Summarising a series of articles arguing for a shift towards mainstream policy, the authors identify two conclusions. First, understanding the contribution of mental health research to mainstream public policy is in the interests of all. Second, this shift changes the metrics for evaluating success, away from traditional clinical imperatives such as symptomatology and relapse rates, and towards valued social roles such as residential stability and labor force participation.

This shift towards emphasising commonality over difference is mirrored in the UK. Current mental health policy identifies six outcomes to improve mental health for the population: better physical health, supporting recovery, improving experience of services, reducing avoidable harm, decreasing stigma, and improving population well-being (2). This changes the balance from previous policy arrangements, away from special policy for dealing with mental health problems and towards integration of mental health policy into mainstream social policy – reflected in its title “*No health without mental health*”.

Although less explored, the same question arises at the level of the individual. Two classes of problem and two classes of solution can be distinguished.

People using mental health services often have both everyday problems (of the sort experienced by others in their community) and more specifically mental illness-related problems (e.g. symptoms, cognitive difficulties). Service provision needs to balance work across these two classes of problems. Similarly, everyday solutions as used by others in their community can be contrasted with specialist solutions (*i.e.* treatments). Again, the challenge for services is achieving the right balance. The traditional approach has perhaps been to prioritise specialist solutions for mental illness-related problems - treat the person so they can subsequently get on with their life. However, in a recovery-oriented mental health system, the balance changes towards supporting everyday solutions for everyday problems.

An illness or a person?

A recovery orientation is central to mental health policy throughout the English-speaking world. The scientific evidence underpinning this re-orientation has been collated (3, 4). What are the practical implications?

Development and consolidation of identity is central to recovery (5). The construct of identity has emerged from three academic disciplines (4). A sociological understanding of identity emphasises **commonality**, the ways in which we are like other people. A psychological understanding emphasises **difference**, the ways in which we are unlike others. A philosophical understanding emphasises **permanence**, persistence of identity over time and space. Mental illness creates a sense of difference - being alone or helpless or tainted or hopeless - in the individual. Clinical processes which

further emphasise difference, such as assessment processes focussed on intrapsychic deficits, can inadvertently contribute to the development and maintenance of a stigmatised identity as a mental patient. Indeed, the focus on intrapsychic deficits to the exclusion of intrapsychic strengths (e.g. artistic skills) and environmental deficits (e.g. poverty) and strengths (e.g. cultural resilience) is a long-standing criticism of the mental health system (6). Supporting recovery involves amplifying commonality and permanence rather than difference and transience. This can be summarised as “*Recovery begins when you find someone or something to connect with*”. The job of the system is to support this connection with self (permanence) and others (commonality).

The central shift involved in a recovery-oriented system is therefore from seeing a patient - someone who is fundamentally different and so needs treatment before they can get on with their life - to seeing a person who is fundamentally similar to people without mental illness in their efforts to “*live the most fulfilling lives that they can live*” (7). What does this mean for clinicians?

Supporting recovery

International best practice in supporting recovery identifies four domains of action: Supporting personally defined recovery (what you do), Working relationship (how you do it), Organizational commitment, and Promoting citizenship (8). This last domain has been the least investigated, yet plausibly is the most influential. Improving social inclusion and community integration

may involve a much wider role for clinicians than providing treatments (9), with more attention paid to supporting the person to make connections in their life, and to the creation of inclusive communities: “*Social inclusion must come down to somewhere to live, something to do, someone to love. It’s as simple - and as complicated - as that*” (10). How can clinicians support people to meet housing, employment and relationship goals?

For housing, the requirement for mental health or substance abuse treatment as a precondition for obtaining housing differentiates two approaches. Emerging evidence suggests the everyday solution (no precondition – ‘housing first’ model) leads to greater consumer satisfaction than the specialist solution (‘supported housing’ model) (11). The assumption that being a good tenant should be a requirement for obtaining tenancy has proved unhelpful, e.g. the impact of sobriety at program entry on outcome is minimal (12)

For employment, the empirical data are clear – the everyday solution of supporting people to obtain and maintain community-based competitive employment through Individual Placement and Support schemes is more effective than sheltered workshops providing non-competitive employment (13). This points to the benefits of the principle of everyday solutions for everyday problems, with compensatory supports provided as needed in order to access those everyday solutions. The idea of being clinically fixed – or ‘work-readiness’ as it is sometimes expressed – has proven to be unhelpful.

Does this principle extend to intimate relationships? Quite possibly. If the concept of 'readiness' has proved unhelpful in relation to housing and employment, then perhaps it is equally unhelpful when social skills training is provided for someone who wants a relationship. The alternative, everyday solution is to support the person to have access to a pool of potential partners with whom they can learn relationship-building skills. Examples include going with the person to join a community-based social or sports club, do voluntary work, use an internet dating service, or go on a speed dating event. These proactive approaches are stretching, in different ways, for the individual and the clinician. The individual may need support to take on these challenges, and it may be more helpful to frame them as learning opportunities rather than expecting initial success. Ongoing involvement and debriefing may well be required as the person learns to cope with the ups-and-downs these experiences will involve. Similarly, the clinician may need support through supervision to move beyond constraining clinical beliefs, such as the importance of being 'better' before doing normal things like dating.

If the principle of everyday solutions for everyday problems is accepted, then other existing clinical approaches also become open for debate. For example, the search for meaning in life is perhaps universal. Mental illness can profoundly change perceptions of self, world and others. The experience of mental ill-health therefore involves identity challenge like many physical disorders. Yet people with mental illness are just as likely to be seeking meaning in their life as any other citizen. If everyday solutions are to be the instinctive orientation of mental health workers, then explanatory models of

mental disorder need to be used judiciously – to provide an account of the experience of mental illness, but also to avoid imposing a specialist solution by creating a uni-dimensional engulfing identity, such as a schizophrenic patient. Allowing space for diverse explanations across different dimensions becomes important, including clinical, functional, physical, social and – most challengingly – existential (14). Whether due to a legacy of Freudian views of religion as regressive and pathological, or because professionals have (statistically) abnormal spiritual views – 90% of the US population believe in a personal God, compared with 24% of clinical or counselling psychologists (15) - evidence indicates that discussing spirituality is challenging for workers (16). Providing opportunities to access spiritual experiences (e.g. reading scripture, prayer, attending places of worship, on-line religious resources), uplifting secular experiences (e.g. through exposure to art, literature, poetry, dance, music, science or nature) and everyday spiritual methods for coping with adversity (e.g. marking boundaries, spiritual purification, spiritual reframing) are all potential areas of future clinical effort.

New ways of working

Overall, the argument is to reverse some priorities. People with mental illness don't need treatment - they need a life. Treatment may contribute to the process of striving for a life worth living, but is a means not an end.

There are of course caveats. Some of the above suggestions will be premature for people in the early stages of recovery, but resilience is developed by engagement rather than avoidance in life. Some people have

mental illness-related problems for which there are no obvious everyday solutions, although these may not be the person's highest priority, and many mental illness-related problems dissipate when everyday problems are resolved. Finally, some people find that mental illness-related problems such as derogatory voices, memory problems or paranoia impede their efforts to find everyday solutions. Offering specialist treatments is often warranted and can be a helpful building block in constructing an identity as a person in recovery, although clinicians need to retain the modesty consistent with the reality that many people find non-clinical ways of self-managing their mental health problems.

But the overarching principle is to ensure that the impoverished expectations, clinical preoccupations and stigmatising beliefs sometimes held by mental health workers do not preclude everyday ways of addressing common (in both senses) human problems. Mental health systems which offer treatments in support of the individual's life goals may look different to those which treat patients in their best interests. Arguably, the strongest contribution of mental health services to recovery is to support everyday solutions to everyday problems.

References

1. Goldman HH, Glied SA, Alegria M: Conclusion: Mental Health in the Mainstream of Public Policy. *American Journal of Psychiatry* 166:1215-6, 2009.
2. HM Government: No health without mental health. Delivering better mental health outcomes for people of all ages. London: Department of Health, 2011.
3. Warner R: Recovery from schizophrenia and the recovery model. *Current Opinion in Psychiatry* 22:374-80, 2009.

4. Slade M: Personal recovery and mental illness. A guide for mental health professionals. Cambridge, Cambridge University Press, 2009.
5. Leamy M, Bird V, Le Boutillier C, et al: A conceptual framework for personal recovery in mental health: systematic review and narrative synthesis. *British Journal of Psychiatry* 199:445-52, 2011.
6. Wright BA, Lopez SJ: Widening the Diagnostic Focus. A Case for Including Human Strengths and Environmental Resources. In: Snyder CR, Lopez SJ, editors. *Handbook of Positive Psychology*. New York, Oxford University Press, 2002. p. 26-44.
7. Silverstein SM, Bellack A: A scientific agenda for the concept of recovery as it applies to schizophrenia. *Clinical Psychology Review* 28:1108-1124, 2008.
8. Le Boutillier C, Leamy M, Bird VJ, et al: What does recovery mean in practice? A qualitative analysis of international recovery-oriented practice guidance. *Psychiatric Services* 62:1470-1476, 2011.
9. Slade M: Mental illness and well-being: the central importance of positive psychology and recovery approaches. *BMC Health Services Research*. 10:26, 2010.
10. Dunn S: Creating accepting communities: report of the Mind inquiry into social exclusion and mental health problems. London, Mind, 1999.
11. Robbins PC, Callahan L, Monahan J: Perceived coercion to treatment and housing satisfaction in housing-first and supportive housing programs. *Psychiatric Services* 60:1251-1253, 2009.
12. Schinka JA, Casey RJ, KasproW W, et al: Requiring Sobriety at Program Entry: Impact on Outcomes in Supported Transitional Housing for Homeless Veterans. *Psychiatric Services* 62:1325-1330, 2011.
13. Pogoda TK, Cramer IE, Rosenheck RA, et al: Qualitative Analysis of Barriers to Implementation of Supported Employment in the Department of Veterans Affairs. *Psychiatric Services* 62:1289-1295, 2011.
14. Whitley R, Drake R: Recovery: A Dimensional Approach. *Psychiatric Services* 61:1248-1250, 2010.
15. Pargament KI: *Spiritually Integrated Psychotherapy*. New York, Guilford, 2007.
16. Huguelet P, Mohr S, Betrisey C, et al: A Randomized Trial of Spiritual Assessment of Outpatients With Schizophrenia: Patients' and Clinicians' Experience. *Psychiatric Services* 62:79-86, 2011.