King's Research Portal

DOI:

[10.1002/eat.23246](https://doi.org/10.1002/eat.23246)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Ambwani, S., Cardi, V., Albano, G., Cao, L., Crosby, R. D., Macdonald, P., Schmidt, U., & Treasure, J. (2020). A multicenter audit of outpatient care for adult anorexia nervosa: Symptom trajectory, service use, and evidence in support of "early stage" versus "severe and enduring" classification. *The International journal of eating disorders*, 53(8), 1337-1348. <https://doi.org/10.1002/eat.23246>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

A multicentre audit of outpatient care for adult anorexia nervosa: Symptom trajectory, service use, and evidence in support “early stage” versus “severe and enduring” classification.

Suman Ambwani ^{a*+}, Valentina Cardi^{b+}, Gaia Albano^{b,c}, Li Cao^{d,e}, Ross D. Crosby^{d,e}, Pamela Macdonald^b, Ulrike Schmidt^b, and Janet Treasure^b

⁺SA and VC should be considered joint first author

^aDepartment of Psychology, Dickinson College, Carlisle, PA

^bSection of Eating Disorders, Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King’s College London, London, UK

^cDepartment of Psychology, Educational Sciences and Human Movement, Palermo, Italy

^dSanford Center for Biobehavioral Research, Fargo, ND

^eDepartment of Psychiatry and Behavioral Sciences, University of North Dakota School of Medicine and Health Sciences, Fargo, ND

Abstract: 259 words

Manuscript: 4592 words

Acknowledgments

This manuscript presents independent research funded by the National Institute for Health Research (NIHR) under its Research for Patient Benefit (RfPB) Programme (Grant Reference Number PB-PG-0712-28041). The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. JT and US acknowledge financial support from the National Institute for Health Research (NIHR) Specialist Biomedical Research Centre for Mental Health award to the South London and Maudsley NHS Foundation Trust and the Institute of Psychiatry, King's College London. US is supported by an NIHR Senior Investigator Award. SA thanks the Research and Development Committee at Dickinson College for providing sabbatical research funds in support of this project.

We also acknowledge the support of the National Institute for Health Research Clinical Research Network (NIHR CRN) and our participating centres: Avon and Wiltshire Mental Health NHS Trust, Barnet Enfield and Haringey Mental Health NHS Trust, Berkshire Healthcare NHS Foundation Trust, Cambridgeshire and Peterborough NHS Foundation Trust, Central and North West London NHS Foundation Trust, Coventry and Warwickshire Partnership NHS Trust, Cumbria Partnership NHS Foundation Trust, Derbyshire Healthcare Foundation Trust, Dorset Healthcare University NHS Foundation Trust, Kent and Medway NHS, Leicestershire Partnership NHS Trust, Lincolnshire Partnership NHS Foundation Trust, Northamptonshire Healthcare NHS Foundation Trust, North Essex Partnership NHS Foundation Trust, Nottinghamshire Healthcare NHS Foundation Trust, South London and Maudsley NHS Foundation Trust, South Staffordshire and Shropshire healthcare NHS Foundation Trust, South Essex Partnership University NHS Foundation Trust, South West London and St George's Mental Health NHS Trust, Surrey and Borders Partnership NHS Foundation Trust, Sussex

1 Partnership NHS Foundation Trust, Worcestershire Health and Care NHS Trust. We are grateful
2 to all our study participants for their patience and perseverance.

3 **Data availability**

4 The authors had full access to the trial data and continue to do so. The deidentified study
5 data will be made publicly available by December 2020.

6

7

8

Abstract

Background: We explored the clinical utility of “staging” anorexia nervosa (AN) based on duration of illness and psychological wellbeing. We also investigated 12-month symptom trajectories and service usage in a large cohort of patients with AN assessed for outpatient treatment. **Method:** We conducted secondary analyses on data from a multisite clinical trial of adults with AN ($N=187$) recruited from 22 NHS England specialist eating disorder outpatient services into a digital treatment augmentation study. Clinical outcomes and service use were measured at post-intervention (six weeks), 6 and 12 months. We grouped patients into two categories: “early stage” (illness duration <3 years; $n=60$) and “severe and enduring” stage (SE-AN; $n=41$) indicated by distress (Depression Anxiety and Stress Scales, DASS ≥ 60) and illness duration (≥ 7 years). **Results:** Overall, patients reported large to moderate improvements from baseline to 12 months in body mass index, eating disorder symptoms, and moderate improvements in mood and work/social adjustment, and 23.6% met criteria for recovery. However, patients classified as SE-AN reported higher rates of accessing intensive services, higher eating disorder symptomatology, and poorer work/social adjustment at baseline, and lower rates of improvement in work/social adjustment at 12 months compared to “early stage” respondents. **Discussion:** Although present findings suggest overall symptomatic improvements, exploratory results highlight marked differences in course and service use between people at different stages of AN and suggest a need to consider staging for clinical decision-making. Treatment research differentiating between clinical subtypes of AN and adoption of a more personalized approach may ensure that services and care pathways better fit patient needs.

Keywords. Anorexia nervosa, classification, staging, outpatient care, treatment outcomes, illness duration, social adjustment.

1 **Trial Registration number:** ISRCTN ClinicalTrials.gov NCT02336841

2

1 A multicentre audit of outpatient care for adult anorexia nervosa: Symptom trajectory, service
2 use, and evidence in support “early stage” versus “severe and enduring” classification.

3 Anorexia nervosa is a debilitating illness that often runs a chronic course and is
4 associated with increased mortality rates, substantial physical and psychological comorbidities,
5 and adverse social consequences (Arcelus, Mitchell, Wales, & Nielsen, 2011). Many patients
6 gradually recover, but longitudinal studies indicate that 61% of people with anorexia nervosa
7 continue to exhibit illness symptoms at 5-year follow-up (Støving, Andries, Brixen, Bilenberg, &
8 Hørder, 2011), and nearly 40% of remain ill at 22-year follow-up (Eddy et al., 2017). Recent
9 findings suggest that although interventions generally yield small improvements in body weight
10 (with greater effects for specialized treatments over comparator treatments), these effects are not
11 maintained at follow-up, and treatment effects on psychological functioning are nonsignificant
12 (Murray, Quintana, Loeb, Griffiths, & Le Grange, 2018); thus, clinical outcomes seem to vary by
13 the nature of the assessment variable and timing of assessment. Indeed, treatment outcome
14 research for AN is frequently challenged by relatively short follow-up periods, small sample
15 sizes, and limited ecological validity given the highly controlled nature of most clinical trials.
16 Other difficulties include differences in inclusion criteria (e.g., whether or not a minimum BMI
17 cut-off is used), differences in reporting and access to rescue treatment (e.g., the numbers who
18 are admitted to higher levels of care), differences in protocols for those who receive higher levels
19 of care, as well as differences in outcome measures (e.g., BMI versus psychosocial functioning).
20 Moreover, there is disagreement among experts regarding the salience (and characteristics) of
21 illness severity and chronicity for treatment outcomes: whereas some highlight the clinical
22 impacts of neuroprogression (Treasure, Stein, & Maguire, 2015), others posit that “severe and
23 enduring” classification schemes have little clinical utility (Raykos, Erceg-Hurn, McEvoy,

1 Fursland, & Waller, 2018). These controversies underscore the need to better understand the
2 trajectory of illness, use of services, and roles of illness duration and severity among individuals
3 receiving treatment for adult anorexia nervosa.

4 Although patients advocate for more research to understand the processes of symptom
5 duration and recovery (van Furth, van der Meer, & Cowan, 2016), there are many uncertainties
6 related to planning appropriate care pathways for adult anorexia nervosa (NHS England, 2019;
7 NICE, 2017). Research suggests that early interventions produce good outcomes (McClelland et
8 al., 2018) whereas people with an enduring form of illness respond less well (Russell, Szmukler,
9 Dare, & Eisler, 1987) and longer duration predicts poorer eating disorder outcomes (Fichter,
10 Quadflieg, & Hedlund, 2006). These findings suggest that a staging model may be applicable to
11 anorexia nervosa (Treasure et al., 2015). However, there is currently no accepted marker of
12 disease progression or clinical features that might be used to stage the illness (Wonderlich et al.,
13 2012). Following Russell's lead it has been common to define the "early stage" of illness as less
14 than 3 years in duration (Treasure et al., 2015), a classification scheme that has demonstrated
15 some clinical utility. For instance, research suggests that in adolescence those in the early stage
16 responded well to family based therapy (Russell et al., 1987). More recently it has been found
17 that a tailored intervention for those in the early stage of illness improves outcome (Fukutomi et
18 al., 2019; McClelland et al., 2018) and this has been cited as an example of good practice in a
19 recent UK Commissioning report (NHS England, 2019).

20 There is less consensus in the field regarding the definition of later stages of illness in
21 anorexia nervosa. Terms such as "severe and enduring" anorexia nervosa (SE-AN), "chronic,"
22 "persistent," and "treatment-resistant" have been used to describe those who have been ill for a
23 longer period of time (Broomfield, Stedal, Touyz, & Rhodes, 2017). Notably, severity and

1 duration of illness appear to represent distinct dimensions, suggesting that reliance solely on the
2 “enduringness” component would be insufficient (Coutinho & Brandão, 2019; Wildes et al.,
3 2017). A modal cut-off of 7 years is commonly employed to describe “long” illness duration and
4 markers of severity including body mass index and number of treatment attempts have been
5 proposed but not received consensus (see (Broomfield et al., 2017) for a review). For instance,
6 body mass index is used as a marker of severity in DSM-5 but several studies have questioned
7 this proposal (Machado, Grilo, & Crosby, 2017; Smith et al., 2017). Instead, patients identified
8 as having severe and persistent forms of illness have highlighted negative mood and poorer
9 social functioning as core illness markers (Arkell & Robinson, 2008), suggesting that these may
10 be valuable indicators of chronicity in anorexia nervosa. Indeed, patients and carers prioritise
11 social function and mood as relevant treatment outcomes (Himmerich et al., 2018; Linville,
12 Brown, Sturm, & McDougal, 2012; Pettersen & Rosenvinge, 2002). Moreover, emotional
13 distress is well-established as an underlying maintenance factor for eating disorders (Schmidt &
14 Treasure, 2006; Treasure & Schmidt, 2013). For example, a network analysis found that
15 depression, anxiety and interpersonal sensitivity were nodes with the highest centrality in the
16 network of symptoms in AN (Solmi, Collantoni, Meneguzzo, Tenconi, & Favaro, 2019). Another
17 network analysis from a large population of adolescents with anorexia nervosa found that
18 depressive symptoms and personal alienation were central features that played a key bridging
19 role (Monteleone et al., 2019). Treatment studies also support the centrality of mood in AN: for
20 instance, proof of concept studies testing neuromodulation interventions used for treatment
21 resistance in depression and obsessive-compulsive disorder have shown some promise with an
22 improvement in mood as the main initial effect, albeit with mixed impact on AN
23 symptomatology (Dalton et al., 2018; Lipsman et al., 2017; McClelland, Kekic, Campbell, &

Schmidt, 2016). Mood-related symptoms may also offer prognostic utility in AN: one longitudinal study of eating disorders reported that patients with a comorbid major depressive disorder diagnosis at baseline were six times more likely to remain unrecovered from restrictive anorexia nervosa 22 years later than those without depression (Franko et al., 2018). A systematic review of prevention and early intervention programs similarly highlighted the need to direct greater attention toward comorbid depressive symptoms in eating disorders: although most programs reduced eating disorder symptoms, over half were unsuccessful in reducing concurrent depressive symptoms and those that specifically targeted negative mood were no more likely to reduce depressive symptoms (Rodgers & Paxton, 2014). Thus, characterizing SE-AN through criteria that attend to severity of distress as well as duration of illness may bridge the gap across conflicting areas of evidence and thereby inform therapeutic targets.

There is also disagreement among experts about the salience of illness chronicity for treatment outcomes, with some arguing that the “severe and enduring” classification has little clinical utility. For example one recent study indicated that duration of illness, BMI and failed inpatient treatment do not predict outcome at the end of inpatient treatment (Wildes et al., 2017) and a lower powered case series (n=66) from an Italian inpatient centre failed to find differences in 12 month outcomes among those with SE-AN as defined as long duration (> 7 years) and failed previous treatment (Calugi, El Ghoch, & Dalle Grave, 2017). Another study, a case series involving outpatient cognitive behavioral therapy (n=134, with n=69 completers) concluded that duration of illness, eating disorder symptoms, and body mass index did not predict treatment outcomes, but did not indicate how many respondents were classified in the 3-, 7-, and 11-year illness duration categories (Raykos et al., 2018). Thus, it is likely that controversies regarding the

clinical utility of SE-AN classification arise in part from the variable design and quality of the studies and thus underscore the need for more research.

The present study explored the clinical utility of a definition of “severe and enduring” anorexia nervosa that considered illness duration as well as severity of psychological distress in one of the largest cohorts of patients with anorexia nervosa in the UK (Cardi et al., 2015; Cardi et al., 2019). To provide a comparison with previously reported outcomes for recipients of outpatient care, we first described changes from baseline to 12 months for individuals receiving NHS outpatient services for AN on key clinical variables (BMI, eating disorder symptoms, depression, anxiety, and stress, and work and social adjustment) and reported rates of recovery at 12 months and service usage for inpatient, daypatient, and outpatient services over the past year. To more specifically explore the clinical utility of staging classifications (such as “early stage” versus “severe and enduring” forms of illness), we assessed differences in symptomatology at baseline, changes from baseline to 12 months, rates of recovery, and service usage for individuals classified according to these schemes.

Method

Full details of the study protocol for the SHARED trial (Cardi et al., 2015) and the primary trial outcomes (Cardi et al., 2019) are published. As intervention groups did not exhibit significant differences in eating disorder symptoms, BMI, psychological well-being, and work/social adjustment at baseline or at any of the follow-up periods, here we report on secondary data analyses from the combined sample.

Setting and recruitment

Participants were recruited from 22 NHS adult ED outpatient services across the UK (see acknowledgements for details) between April 2015 and December 2016. The authors assert that

all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008. All procedures involving human subjects/patients were approved by a National Research Ethics Service Committee (London Brent, approval number 14/LO/1347) and from the local research and development offices at the participating centres. Written informed consent was obtained from all patients.

Participants

Participants were recruited by clinicians and clinical study officers from consecutive assessments if 1) they were aged 16 or over, 2) had a Diagnostic and Statistical Manual of Mental Disorders, 5th Edition (DSM-5) or ICD-10 diagnosis of AN or other specified feeding or eating disorder (OSFED) with features of AN and a body mass index (BMI) of 18.5 kg/m² or below. Participants were considered ineligible for the following reasons 1) insufficient knowledge of English and/or 2) severe mental or physical illness needing treatment (i.e. psychosis or diabetes mellitus). Participants completed study assessments through IESO Digital Health (<http://www.iesohealth.com>) at baseline, six weeks, six months and 12 months. Clinical teams provided additional details regarding BMI and outpatient service use at six weeks, six months and 12 months.

Measures

Demographic information and patient history. Participants self-reported age, gender, ethnicity, years of education, employment and social status, duration of illness, time of illness onset, diagnosis and first treatment received, previous hospital admissions, psychiatric comorbidity and medication, and height and weight (which were used to calculate Body Mass

Index BMI = weight (kg) / [height (m)]²). Patient height and weight was also measured by the clinical sites and reported to the research team.

Eating Disorder Examination Questionnaire (Fairburn & Beglin, 1994). The EDE-Q is a 36-item self-report measure of eating disorder symptomatology. Items are scored on a six-point Likert scale, and higher scores indicate more severe eating disorder symptomatology. In the current study, we used the EDE-Q global score to assess eating disorder symptomatology (EDE-Q Cronbach's $\alpha = 0.91$).

Depression, Anxiety and Stress Scales (Lovibond & Lovibond, 1995). The DASS-21 is a 21-item self-report measure of psychological distress over the past seven days. Items are scored on a four-point Likert scale, with higher scores indicating greater distress. The DASS-21 includes three subscales to assess symptoms of anxiety, depression, and stress, as well as an overall score to reflect general distress (DASS-total Cronbach's $\alpha = 0.92$).

Work and Social Adjustment Scale (Marks, 1986). The WSAS is a five-item self-report measure of perceived impairment in everyday functioning due to an identified problem. The scale evaluates functioning in the following domains: work, home management, social and private leisure activities, and close relationships. Scores range from 0 to 8 for each item, with higher overall scores indicating more severe functional impairment (WSAS Cronbach's $\alpha: 0.72$).

Participant service usage. Participants self-reported their usage of various clinical services (outpatient, daypatient, and inpatient services) related to the eating disorder over the previous 6 and 12 months. If patients did not report data on service use, we asked clinical centers to supplement this information.

Intervention

Each centre delivered their usual treatment. Treatment in NHS clinics follows the NICE guidelines and quality standards (NICE, 2017), which recommend outpatient care in the first instance with transfer to day/inpatient if and when the level of risk is high. The form of treatment is determined only by clinical decision of level of need and is free at the point of delivery. The group randomised to receive the adjunctive intervention received early phase 6-week online chat guidance delivered via a digital platform along with psychoeducational material supplemented with recovery narratives (Cardi et al., 2015). The intervention was designed to enhance motivation and decrease engagement with the psychopathology of anorexia nervosa and facilitate the development of a recovery identity.

Data Analyses

All analyses were conducted in SPSS Version 26.0. We used multiple imputation for missing data at the 6-month and 12-month assessments, and conducted paired t-tests to assess changes on key clinical indicators (BMI, EDE-Q, DASS, WSAS) from baseline to the follow-up assessments for the full sample. We conducted frequency analyses to assess service usage related to the eating disorder (past hospitalization assessed at baseline, and use of inpatient, day patient, and outpatient services over the past 6 months and 12 months) and recovery. We defined “recovery” as a combination of BMI (≥ 18.5) and eating disorder symptoms (EDE-Q 1 standard deviation above the nonclinical mean, or EDE-Q < 2.77 ; (Mond, Hay, Rodgers, Owen, & Beumont, 2004)) following past research (Couturier & Lock, 2006).

For our exploratory analyses comparing baseline functioning and outcomes for respondents in the “early stage” vs. “severe and enduring” illness, we defined “early stage” as illness duration less than 3 years (Brown et al., 2018; Russell et al., 1987). Past studies on SE-AN typically rely on duration, but not severity, criteria in defining a “severe and enduring”

group, and although related, severity is not equivalent to illness duration. For instance, among our respondents, duration of illness was correlated with one indicator of illness severity, BMI ($r = -.25, p = .001$), but correlations with other clinical indicators at baseline (such as EDE-Q, DASS, and WSAS) were small and non-significant. Moreover, the revised cognitive interpersonal model posits that secondary neuroprogressive changes associated with high stress, anhedonia, and elevated fears are relevant for clinical functioning and outcomes in AN (Treasure et al., under review). Thus, a composite variable comprising severity and duration indicators likely captures unique variance above and beyond duration of illness alone. We evaluated several options for defining SE-AN through a composite of duration and BMI, duration and EDE-Q, duration and WSAS, and duration and DASS, and these comparisons are reported in the supplementary materials. However, we chose to define SE-AN in the current investigation as a combination of severity of psychological distress (DASS-total ≥ 60 ; (Beaufort, De Weert-Van Oene, Buwalda, De Leeuw, & Goudriaan, 2017) and duration of illness (≥ 7 years; (Robinson, 2009; Touyz et al., 2013) due to the centrality of psychological distress for the phenomenology of disordered eating.

We then conducted a chi-square analysis comparing groups on lifetime and past year usage of intensive services (day/inpatient care), followed by independent samples t-tests comparing “early stage” vs. “severe and enduring” groups on baseline functioning (EDE-Q, BMI, WSAS). Finally, we conducted mixed repeated measures ANOVAs to compare changes from baseline to 12 months across “early stage” and “severe and enduring” classification groups on BMI, EDE-Q, and WSAS.

Results

Participant characteristics

One-hundred eighty-seven participants completed the baseline assessment (see Table 1 for demographic and descriptive information). There was large variation in the clinical characteristics. Many had a long duration of illness ($M = 7.76$ years; $SD = 8.91$ years) with a 5-year gap between self-recognized eating disorder symptoms and the start of treatment and their BMI had fallen by 6.3 kg/m^2 . Over a quarter had previously received inpatient care and reported psychiatric co-morbidity, and 44.3% were receiving psychiatric medication. Both eating disorder psychopathology (EDE-Q) and levels of depression, anxiety, and stress (DASS) were elevated and psychosocial function was impaired (WSAS).

Adherence to the follow up protocol

For questionnaire measures (EDE-Q, DASS, WSAS) there was 75.4% completion at 6 months and 62.6% at 12 months. For service usage, 74.9% reported on their service usage at 6 months and 61.5% at 12 months. We obtained some missing data (weight and service use) from clinical sites, resulting in BMI data for 70% of participants at 6 and 12 months.

Clinical outcomes, service use, and rates of “recovery” at follow-up

Paired t-tests using multiple imputation data indicated small improvements from baseline to 6 months in distress (DASS Hedge’s $g = 0.28$) and work and social adjustment (WSAS Hedge’s $g = 0.30$), and moderate improvements in BMI (Hedge’s $g = 0.57$) and eating disorder symptoms (EDE-Q Hedge’s $g = 0.54$) (see Table 2). Small to moderate improvements on these outcomes were maintained at 12 months, with the only exception of BMI, which was associated with a larger effect size at the 12 month assessment (Hedge’s $g = 0.85$).

Among those who participated in the 12-month assessment, frequency analyses indicated that a large proportion remained in outpatient care (86/115; 74.8%). Data from the 6-month and 12-month assessment periods indicated that over a third of respondents (44/123; 35.8%) accessed

intensive care (i.e., inpatient or daypatient services) at some point over the past year. Among those who continued with follow-up assessments at 12 months ($n = 123$), 23.6% of respondents met BMI and EDE-Q criteria for recovery¹.

Exploratory analyses comparing clinical outcomes for “early stage” vs. “severe and enduring” respondents

Service Usage. A chi-square analysis of independence compared “early stage” and “severe and enduring” respondents in their usage of intensive services (inpatient/daypatient) either at baseline or at follow-up assessments. Results indicate that SE-AN respondents were more likely to have received (29/35; 82.9%) intensive care at some point as compared to “early stage” respondents (18/39 = 46.2%), $\chi^2(1) = 10.72, p = .001$.

Baseline differences. A series of independent samples t-tests comparing the “early stage” and SE-AN groups on key clinical indicators at baseline indicated moderate sized, significant group differences for EDE-Q and WSAS, and small and nonsignificant differences for BMI (see Table 3).

12 month outcomes. Mixed repeated measures ANOVAs compared changes from baseline to 12-months for BMI, EDE-Q, and WSAS by “early stage” versus “severe and enduring” classification. Results indicated main effects for time and group, with improvements over time on all three markers, but poorer functioning on all three markers for the SE-AN group (see Figures 1-3). Moreover, there was a significant time x illness group interaction for WSAS, indicating relatively less improvement on this indicator over time for the SE-AN group (see Table 3). There was a trend toward significance for the time x illness group interactions for BMI

¹ Given different published EDE-Q norms for community women, we also tested an alternate criterion for “recovery” ($EDE-Q < 1.79$ per Aardoom et al., 2012). Based on these criteria ($EDE-Q < 1.79$ and $BMI \geq 18.5$), 13.8% of respondents met the threshold for recovery at 12 months.

and EDE-Q ($p = .06$) and examination of the box plots suggested a similar pattern with lower rates of improvement over time for the SE-AN group.

Exploratory analyses evaluating clinical outcomes based on varying SE-AN classification schemes

A series of exploratory independent samples t-tests comparing non-SE-AN and SE-AN respondents by varying classification schemes (i.e., duration combined with either high DASS, low BMI, high EDE-Q, or high WSAS) suggested several differences in clinical functioning at baseline across groups with SE-AN respondents exhibiting poorer clinical functioning (medium to large effect sizes) than non-SE-AN respondents at baseline. However, results varied as a function of the classification scheme and the nature of the outcome variable (see Table S1). We then conducted a series of exploratory mixed repeated measures ANOVAs comparing changes from baseline to 12 months on clinical indicators for the SE-AN vs. non-SE-AN respondents. In general, results indicated improvements over time, and differences in outcomes based on classification group, but differences in the trajectory of change over time by classification group were nonsignificant when SE-AN was defined by WSAS, EDE-Q, and in most cases, BMI. However, results indicated significant interaction effects when SE-AN was defined by DASS, such that the rates of improvement over time for BMI, EDE-Q, and WSAS were smaller for the SE-AN than non-SE-AN respondents.

Discussion

The current study reported clinical outcomes, service use, and differences by staging classification for people receiving outpatient care for anorexia nervosa in the UK. Overall, we found large to moderate improvements in BMI, eating disorder symptoms and moderate changes in mood symptoms, and work and social adjustment at 12 months, which is consistent with

1 previously reported improvements in BMI among recipients of adult outpatient treatment (Zeeck
2 et al., 2018). However, previously published rates of accessing intensive services are more
3 mixed, with estimates ranging from 11.9%-15.6% in a UK trial (Schmidt et al., 2015) to 23%-
4 41% in a German trial (Zipfel et al., 2014), and our finding that 35.8% reported transitioning to a
5 higher intensity of treatment (in/day patient care) during the year fall within the higher end of the
6 spectrum. Among those who continued with follow-up assessments at 12 months ($n = 123$), less
7 than a quarter (23.6%) of respondents met criteria for recovery as defined by body mass index
8 and eating disorder symptoms. These lower than desirable levels of improvement are consistent
9 with results from a recent systematic review and meta-analysis, which reported high levels of
10 residual symptoms even among patients with AN who were considered remitted or recovered
11 (Tomba, Tecuta, Crocetti, Squarcio, & Tomei, 2019).

12 Our major objective was to explore the clinical utility of classification schemes for
13 anorexia nervosa based on markers of psychological distress and illness duration. Approximately
14 36% (60/166) of our respondents were classified as “early stage” and 25% (41/166) were
15 classified as “severe and enduring.” Patients in the SE-AN group reported more lifetime
16 hospitalizations and worse eating disorder symptoms, psychological functioning and work and
17 social wellbeing compared to patients with shorter illness duration. Patients in the SE-AN group
18 also reported less symptomatic changes over time compared to the “early phase” group,
19 especially in relation to work and social adjustment. Exploratory results also suggest less
20 improvement in eating disorder symptomatology, work and social adjustment, and BMI over
21 time for those classified as SE-AN as compared to those identified as non-SE-AN. These data
22 support the clinical utility of SE-AN classification based on the previously established threshold
23 of 7 years of illness with the addition of emotional distress as an index of severity.

1 There is very little evidence on the specific treatment of individuals classified as having
2 “severe and enduring” anorexia nervosa apart from a few investigations (Andries, Frystyk,
3 Flyvbjerg, & Støving, 2014; Calugi et al., 2017; Dalton et al., 2018; Lipsman et al., 2017; Touyz
4 et al., 2013). Nevertheless, many studies in adults include a large number of patients that fall into
5 this subgroup and suggest lower response to treatment: for instance, studies testing olanzapine
6 (Attia et al., 2019) and dronabinol (Andries et al., 2014) report small improvements in weight
7 gain but little change in other domains. Thus, there is an urgent need for research on
8 interventions targeting individuals with a chronic course of illness (Wonderlich et al., 2012).

9 **Strengths and limitations**

10 The present study reported on service use and clinical outcomes for the largest, most
11 representative sample of patients receiving outpatient services for AN in the UK over a 12-month
12 follow-up period. Attrition from protocol follow-up (30%) was within the range reported in a
13 systematic review of previous studies in anorexia nervosa (DeJong, Broadbent, & Schmidt,
14 2012) and therefore we used multiple imputation for missing data where possible. A strength of
15 the study is that it fulfils proposed benchmark criteria for AN intervention research including
16 reporting on separate physiological (BMI), cognitive/affective (EDE-Q, DASS) and
17 social/functional (WSAS) outcome indicators, in addition to providing details about service use
18 over the course of a follow-up period (Murray, Loeb, & Le Grange, 2018). Moreover, given
19 recent scrutiny of bias in anorexia nervosa treatment outcome research (Murray et al., 2019), we
20 are pleased to report that we participated in pre-registration of the protocol (Cardi et al., 2015),
21 completed the trial as proposed, published the primary outcomes (Cardi et al., 2019), and will be
22 making study materials and data publicly available to facilitate replication and extension.
23 However, our analyses comparing “early stage” and “SE-AN” responses involved modest

1 samples, given the stringent criteria used to define these groups, and we did not statistically
2 control for our multiple comparisons due to the exploratory nature of our analyses. Thus, present
3 findings are limited and require further replication and extension in larger samples of participants
4 receiving outpatient care for anorexia nervosa.

5 **Clinical implications**

6 The NICE guidelines (NICE, 2004, 2017) recommend that the care pathway for anorexia
7 nervosa should begin with outpatient treatment, however, there is likely a great deal of service
8 variability across treatment centers and patients may have been assigned to waiting lists
9 following the assessment period. In our study, there were small to large improvements across
10 clinical indicators in this naturalistic setting, but less than a quarter had reached the criteria for
11 “recovery” at 12 months. Over a third of patients were transferred to day/inpatient care because
12 of medical and or psychiatric risk, and over three quarters of patients remained in treatment at 12
13 months. These data offer a point of comparison for outpatient care outcomes, whilst highlighting
14 the need to more closely scrutinize the nature of treatment as usual across care settings, including
15 an assessment of their ability to provide care without delay.

16 There remains uncertainty in the literature as to whether categorization by “early stage”
17 vs. “severe and enduring” has implications for treatment and services. Our findings lend support
18 for a model where longer duration of illness and poorer psychological wellbeing might require
19 more specific treatment efforts. For instance, individuals with SE-AN may benefit from
20 neuromodulation interventions targeting treatment-resistant depression, such as deep brain
21 stimulation and repetitive transcranial magnetic stimulation, or from pharmacological approaches
22 such as ketamine and psilocybin (Treasure et al., under review). Indeed, our investigation
23 highlights the salience of distress as an indicator of illness severity, thus, directing greater

1 attention toward the role of such distress could clarify useful mechanisms for additional
2 intervention and suggest possibilities for a combined approach targeting both mood and eating
3 disorder symptomatology. However, we acknowledge that staging efforts should consider
4 associated risks for increased stigmatization and pursue strategies to mitigate the potential for
5 such harm.

6 Although some clinical practice guidelines have been developed to address the needs of
7 patients with SE-AN (e.g. guidelines from the Royal Australian and New Zealand College of
8 Psychiatrists (Hay et al., 2014), the Royal College of Psychiatrists' MARSIPAN (Robinson &
9 Rhys Jones, 2018), and The British Columbia Ministry of Health (Geller, Goodrich, Chan,
10 Cockell, & Srikaneswaran, n.d.)), we believe that further work is needed to characterize the
11 clinical needs of this patient group and to implement a combined approach that targets both
12 mood and eating disorder symptoms.

13 **Conclusion**

14 We provide evidence of low “recovery” rates (23%) and relatively high rates of
15 admission to more intense forms of treatment (36%) among a large UK-based population of
16 patients with anorexia nervosa assessed for outpatient treatment. This is despite overall large
17 improvements in BMI and moderate improvements in eating disorder psychopathology.
18 Moreover, treatment gains were more modest for patients with a severe and enduring form of
19 illness characterized by longer duration and worse psychological wellbeing. We suggest that
20 using treatment strategies to provide rapid relief of symptoms of low mood and greater support
21 with psychosocial functioning may be useful to enhance standard care for adult anorexia nervosa.

References

- Andries, A., Frystyk, J., Flyvbjerg, A., & Støving, R. K. (2014). Dronabinol in severe, enduring anorexia nervosa: A randomized controlled trial. *International Journal of Eating Disorders*, 47(1), 18–23. <https://doi.org/10.1002/eat.22173>
- Arcelus, J., Mitchell, A. J., Wales, J., & Nielsen, S. (2011). Mortality Rates in Patients With Anorexia Nervosa and Other Eating Disorders. *Archives of General Psychiatry*, 68(7), 724. <https://doi.org/10.1001/archgenpsychiatry.2011.74>
- Arkell, J., & Robinson, P. (2008). A pilot case series using qualitative and quantitative methods: Biological, psychological and social outcome in severe and enduring eating disorder (anorexia nervosa). *International Journal of Eating Disorders*, 41(7), 650–656. <https://doi.org/10.1002/eat.20546>
- Attia, E., Steinglass, J. E., Timothy Walsh, B., Wang, Y., Wu, P., Schreyer, C., ... Marcus, M. D. (2019). Olanzapine versus placebo in adult outpatients with anorexia nervosa: A randomized clinical trial. *American Journal of Psychiatry*, 176(6), 449–456. <https://doi.org/10.1176/appi.ajp.2018.18101125>
- Beaufort, I. N., De Weert-Van Oene, G. H., Buwalda, V. A. J., De Leeuw, J. R. J., & Goudriaan, A. E. (2017). The Depression, Anxiety and Stress Scale (DASS-21) as a Screener for Depression in Substance Use Disorder Inpatients: A Pilot Study. *European Addiction Research*, 23(5), 260–268. <https://doi.org/10.1159/000485182>
- Broomfield, C., Stedal, K., Touyz, S., & Rhodes, P. (2017). Labeling and defining severe and enduring anorexia nervosa: A systematic review and critical analysis. *International Journal of Eating Disorders*, 50(6), 611–623. <https://doi.org/10.1002/eat.22715>
- Brown, A., McClelland, J., Boysen, E., Mountford, V., Glennon, D., & Schmidt, U. (2018). The

- 1 FREED Project (first episode and rapid early intervention in eating disorders): service
2 model, feasibility and acceptability. *Early Intervention in Psychiatry*, 12(2), 250–257.
3 <https://doi.org/10.1111/eip.12382>
- 4 Calugi, S., El Ghoch, M., & Dalle Grave, R. (2017). Intensive enhanced cognitive behavioural
5 therapy for severe and enduring anorexia nervosa: A longitudinal outcome study. *Behaviour*
6 *Research and Therapy*, 89, 41–48. <https://doi.org/10.1016/j.brat.2016.11.006>
- 7 Cardi, V., Ambwani, S., Crosby, R., Macdonald, P., Todd, G., Park, J., ... Treasure, J. (2015).
8 Self-Help And Recovery guide for Eating Disorders (SHARED): Study protocol for a
9 randomized controlled trial. *Trials*, 16(1). <https://doi.org/10.1186/s13063-015-0701-6>
- 10 Cardi, Valentina, Albano, G., Ambwani, S., Cao, L., Crosby, R. D., Macdonald, P., ... Treasure,
11 J. (2019). A randomised clinical trial to evaluate the acceptability and efficacy of an early
12 phase, online, guided augmentation of outpatient care for adults with anorexia nervosa.
13 *Psychological Medicine*, 1–12. <https://doi.org/10.1017/S0033291719002824>
- 14 Coutinho, F., & Brandão, I. (2019). Severe and enduring Anorexia Nervosa: a brief narrative
15 review about the concept and its therapeutic options LETTER TO THE EDITOR Citation:
16 Coutinho et al. Severe and enduring Anorexia Nervosa: a brief nar-rative review about the
17 concept and its therapeutic options INTERN ATIONAL JOURNAL OF AND CLINICAL
18 NEUROSCIENCES MENTAL HEALTH Severe and enduring Anorexia Nervosa: concepts
19 and therapeutic options. *International Journal of Clinical Neurosciences and Mental*
20 *Health*, 6, 3. <https://doi.org/10.21035/ijcnmh.2019.6.3>
- 21 Couturier, J., & Lock, J. (2006). What is recovery in adolescent anorexia nervosa? *International*
22 *Journal of Eating Disorders*, 39(7), 550–555. <https://doi.org/10.1002/eat.20309>
- 23 Dalton, B., Bartholdy, S., McClelland, J., Kekic, M., Rennalls, S. J., Werthmann, J., ... Schmidt,

- U. (2018). Randomised controlled feasibility trial of real versus sham repetitive transcranial magnetic stimulation treatment in adults with severe and enduring anorexia nervosa: the TIARA study. *BMJ Open*, 8(7), e021531. <https://doi.org/10.1136/bmjopen-2018-021531>
- DeJong, H., Broadbent, H., & Schmidt, U. (2012). A systematic review of dropout from treatment in outpatients with anorexia nervosa. *International Journal of Eating Disorders*, 45(5), 635–647. <https://doi.org/10.1002/eat.20956>
- Eddy, K. T., Tabri, N., Thomas, J. J., Murray, H. B., Keshaviah, A., Hastings, E., ... Franko, D. L. (2017). Recovery From Anorexia Nervosa and Bulimia Nervosa at 22-Year Follow-Up. *The Journal of Clinical Psychiatry*, 78(02), 184–189. <https://doi.org/10.4088/JCP.15m10393>
- Fairburn, C. G., & Beglin, S. J. (1994). Assessment of eating disorder psychopathology: Interview or self-report questionnaire? *International Journal of Eating Disorders* 2J, 16, 363–370.
- Fichter, M. M., Quadflieg, N., & Hedlund, S. (2006). Twelve-year course and outcome predictors of anorexia nervosa. *International Journal of Eating Disorders*, 39(2), 87–100. <https://doi.org/10.1002/eat.20215>
- Franko, D. L., Tabri, N., Keshaviah, A., Murray, H. B., Herzog, D. B., Thomas, J. J., ... Eddy, K. T. (2018). Predictors of long-term recovery in anorexia nervosa and bulimia nervosa: Data from a 22-year longitudinal study. *Journal of Psychiatric Research*, 96, 183–188. <https://doi.org/10.1016/j.jpsychires.2017.10.008>
- Fukutomi, A., Austin, A., McClelland, J., Brown, A., Glennon, D., Mountford, V., ... Schmidt, U. (2019). First episode rapid early intervention for eating disorders: A two-year follow-up. *Early Intervention in Psychiatry*. <https://doi.org/10.1111/eip.12881>

- 1 Geller, J., Goodrich, S., Chan, K., Cockell, S., & Srikameswaran, S. (n.d.). *Clinical practice*
2 *guidelines for the BC eating disorders continuum of services*. Retrieved from
3 [http://mh.providencehealthcare.org/sites/default/files/BC Eating Disorders Clinical Practice](http://mh.providencehealthcare.org/sites/default/files/BC%20Eating%20Disorders%20Clinical%20Practice%20Guidelines.pdf)
4 [Guidelines.pdf](http://mh.providencehealthcare.org/sites/default/files/BC Eating Disorders Clinical Practice Guidelines.pdf)
- 5 Hay, P., Chinn, D., Forbes, D., Madden, S., Newton, R., Sugenor, L., ... Ward, W. (2014). Royal
6 Australian and New Zealand College of Psychiatrists clinical practice guidelines for the
7 treatment of eating disorders. *Australian & New Zealand Journal of Psychiatry*, 48(11),
8 977–1008. <https://doi.org/10.1177/0004867414555814>
- 9 Himmerich, H., Joaquim, M., Bentley, J., Kan, C., Dornik, J., Treasure, J., & Schmidt, U. (2018,
10 August 1). Psychopharmacological options for adult patients with anorexia nervosa: The
11 patients' and carers' perspectives. *CNS Spectrums*. Cambridge University Press.
12 <https://doi.org/10.1017/S1092852917000529>
- 13 Linville, D., Brown, T., Sturm, K., & McDougal, T. (2012). Eating Disorders and Social
14 Support: Perspectives of Recovered Individuals. *Eating Disorders*, 20(3), 216–231.
15 <https://doi.org/10.1080/10640266.2012.668480>
- 16 Lipsman, N., Lam, E., Volpini, M., Sutandar, K., Twose, R., Giacobbe, P., ... Lozano, A. M.
17 (2017). Deep brain stimulation of the subcallosal cingulate for treatment-refractory anorexia
18 nervosa: 1 year follow-up of an open-label trial. *The Lancet Psychiatry*, 4(4), 285–294.
19 [https://doi.org/10.1016/S2215-0366\(17\)30076-7](https://doi.org/10.1016/S2215-0366(17)30076-7)
- 20 Lovibond, S., & Lovibond, P. (1995). *Manual for the Depression Anxiety and Stress Scales* (2nd
21 ed.). Sydney: Psychology Foundation.
- 22 Machado, P. P. P., Grilo, C. M., & Crosby, R. D. (2017). Evaluation of the DSM-5 Severity
23 Indicator for Anorexia Nervosa. *European Eating Disorders Review*, 25(3), 221–223.

1 <https://doi.org/10.1002/erv.2508>

2 Marks, I. (1986). *Behavioral Psychotherapy*. Bristol, UK: John Wright, now published by

3 I.Marks, Institute of Psychiatry, London.

4 McClelland, J., Hodsoll, J., Brown, A., Lang, K., Boysen, E., Flynn, M., ... Schmidt, U. (2018).

5 A pilot evaluation of a novel First Episode and Rapid Early Intervention service for Eating
6 Disorders (FREED). *European Eating Disorders Review*, 26(2), 129–140.

7 <https://doi.org/10.1002/erv.2579>

8 McClelland, J., Kekic, M., Campbell, I. C., & Schmidt, U. (2016). Repetitive Transcranial

9 Magnetic Stimulation (rTMS) Treatment in Enduring Anorexia Nervosa: A Case Series.

10 *European Eating Disorders Review*, 24(2), 157–163. <https://doi.org/10.1002/erv.2414>

11 Mond, J. M., Hay, P. J., Rodgers, B., Owen, C., & Beumont, P. J. V. (2004). Validity of the

12 Eating Disorder Examination Questionnaire (EDE-Q) in screening for eating disorders in
13 community samples. *Behaviour Research and Therapy*, 42(5), 551–567.

14 [https://doi.org/https://doi.org/10.1016/S0005-7967\(03\)00161-X](https://doi.org/https://doi.org/10.1016/S0005-7967(03)00161-X)

15 Monteleone, A. M., Mereu, A., Cascino, G., Criscuolo, M., Castiglioni, M. C., Pellegrino, F., ...

16 Zanna, V. (2019). Re-conceptualization of anorexia nervosa psychopathology: A network

17 analysis study in adolescents with short duration of the illness. *The International Journal of*

18 *Eating Disorders*. <https://doi.org/10.1002/eat.23137>

19 Murray, S. B., Compte, E. J., Quintana, D. S., Mitchison, D., Griffiths, S., & Nagata, J. M.

20 (2019). Registration, reporting, and replication in clinical trials: The case of anorexia
21 nervosa. *International Journal of Eating Disorders*, eat.23187.

22 <https://doi.org/10.1002/eat.23187>

23 Murray, S. B., Loeb, K. L., & Le Grange, D. (2018). Treatment outcome reporting in anorexia

- nervosa: time for a paradigm shift? *Journal of Eating Disorders*, 6(1), 10.
<https://doi.org/10.1186/s40337-018-0195-1>
- Murray, S. B., Quintana, D. S., Loeb, K. L., Griffiths, S., & Le Grange, D. (2018). Treatment outcomes for anorexia nervosa: a systematic review and meta-analysis of randomized controlled trials. *Psychological Medicine*, 1–10.
<https://doi.org/10.1017/S0033291718002088>
- NHS England. (2019). Adult Eating Disorders: Community, Inpatient and Intensive Day Patient Care – Guidance for commissioners and providers. Retrieved October 27, 2019, from <https://www.england.nhs.uk/publication/adult-eating-disorders-community-inpatient-and-intensive-day-patient-care-guidance-for-commissioners-and-providers/>
- NICE. (2004). *Eating disorders - core interventions in the treatment and management of anorexia nervosa, bulimia nervosa and related eating disorders*. NICE Clinical Guideline No 9. London, UK. Retrieved from <http://www.nice.org.uk>
- NICE. (2017). *Eating disorders: Recognition and treatment*. (NC69). London, UK. Retrieved from <https://www.nice.org.uk/guidance/ng69>
- Pettersen, G., & Rosenvinge, J. H. (2002). Improvement and recovery from eating disorders: a patient perspective. *Eating Disorders*, 10(1), 61–71. <https://doi.org/10.1002/erv.425>
- Raykos, B. C., Erceg-Hurn, D. M., McEvoy, P. M., Fursland, A., & Waller, G. (2018). Severe and enduring anorexia nervosa? Illness severity and duration are unrelated to outcomes from cognitive behaviour therapy. *Journal of Consulting and Clinical Psychology*, 86(8), 702–709. <https://doi.org/10.1037/ccp0000319>
- Robinson, P. (2009). *Severe and Enduring Eating Disorder (SEED) : Management of Complex Presentations of Anorexia and Bulimia Nervosa*. John Wiley & Sons.

- 1 Robinson, P., & Rhys Jones, W. (2018). MARSIPAN: management of really sick patients with
2 anorexia nervosa. *BJPsych Advances*, 24(1), 20–32. <https://doi.org/10.1192/bja.2017.2>
- 3 Rodgers, R. F., & Paxton, S. J. (2014). The impact of indicated prevention and early intervention
4 on co-morbid eating disorder and depressive symptoms: a systematic review. *Journal of*
5 *Eating Disorders*, 2(1), 30. <https://doi.org/10.1186/s40337-014-0030-2>
- 6 Russell, G. F. M., Szmukler, G. I., Dare, C., & Eisler, I. (1987). An Evaluation of Family
7 Therapy in Anorexia Nervosa and Bulimia Nervosa. *Archives of General Psychiatry*,
8 44(12), 1047–1056. <https://doi.org/10.1001/archpsyc.1987.01800240021004>
- 9 Schmidt, U., Magill, N., Renwick, B., Keyes, A., Kenyon, M., Dejong, H., ... Landau, S. (2015).
10 The Maudsley Outpatient Study of Treatments for Anorexia Nervosa and Related
11 Conditions (MOSAIC): Comparison of the Maudsley Model of Anorexia Nervosa
12 Treatment for Adults (MANTRA) with specialist supportive clinical management (SSCM)
13 in outpatients with broadly defined anorexia nervosa: A randomized controlled trial.
14 *Journal of Consulting and Clinical Psychology*, 83(4), 796–807.
15 <https://doi.org/10.1037/ccp0000019>
- 16 Schmidt, U., & Treasure, J. (2006). Anorexia nervosa: Valued and visible. A cognitive-
17 interpersonal maintenance model and its implications for research and practice. *British*
18 *Journal of Clinical Psychology*, 45(3), 343–366. <https://doi.org/10.1348/014466505X53902>
- 19 Smith, K. E., Ellison, J. M., Crosby, R. D., Engel, S. G., Mitchell, J. E., Crow, S. J., ...
20 Wonderlich, S. A. (2017). The validity of DSM-5 severity specifiers for anorexia nervosa,
21 bulimia nervosa, and binge-eating disorder. *International Journal of Eating Disorders*,
22 50(9), 1109–1113. <https://doi.org/10.1002/eat.22739>
- 23 Solmi, M., Collantoni, E., Meneguzzo, P., Tenconi, E., & Favaro, A. (2019). Network analysis of

- 1 specific psychopathology and psychiatric symptoms in patients with anorexia nervosa.
2 *European Eating Disorders Review*, 27(1), 24–33. <https://doi.org/10.1002/erv.2633>
- 3 Støving, R. K., Andries, A., Brixen, K., Bilenberg, N., & Hørder, K. (2011). Gender differences
4 in outcome of eating disorders: A retrospective cohort study. *Psychiatry Research*, 186(2–
5 3), 362–366. <https://doi.org/10.1016/J.PSYCHRES.2010.08.005>
- 6 Tomba, E., Tecuta, L., Crocetti, E., Squarcio, F., & Tomei, G. (2019). Residual eating disorder
7 symptoms and clinical features in remitted and recovered eating disorder patients: A
8 systematic review with meta-analysis. *International Journal of Eating Disorders*. John
9 Wiley and Sons Inc. <https://doi.org/10.1002/eat.23095>
- 10 Touyz, S., Le Grange, D., Lacey, H., Hay, P., Smith, R., Maguire, S., ... Crosby, R. D. (2013).
11 Treating severe and enduring anorexia nervosa: a randomized controlled trial.
12 *Psychological Medicine*, 43(12), 2501–2511. <https://doi.org/10.1017/S0033291713000949>
- 13 Treasure, J., & Schmidt, U. (2013). The cognitive-interpersonal maintenance model of anorexia
14 nervosa revisited: a summary of the evidence for cognitive, socio-emotional and
15 interpersonal predisposing and perpetuating factors. *Journal of Eating Disorders*, 1(1), 13.
16 <https://doi.org/10.1186/2050-2974-1-13>
- 17 Treasure, J., Stein, D., & Maguire, S. (2015). Has the time come for a staging model to map the
18 course of eating disorders from high risk to severe enduring illness? An examination of the
19 evidence. *Early Intervention in Psychiatry*, 9(3), 173–184.
20 <https://doi.org/10.1111/eip.12170>
- 21 Treasure, J., Willmott, D., Ambwani, S., Cardi, V., Clark Bryan, D., Rowlands, K., & Schmidt,
22 U. (under review). Cognitive interpersonal model for anorexia nervosa revisited: the
23 perpetuating factors that contribute to the development of the severe and enduring illness.

- 1 van Furth, E. F., van der Meer, A., & Cowan, K. (2016, August 1). Top 10 research priorities for
2 eating disorders. *The Lancet Psychiatry*. Elsevier Ltd. [https://doi.org/10.1016/S2215-](https://doi.org/10.1016/S2215-0366(16)30147-X)
3 [0366\(16\)30147-X](https://doi.org/10.1016/S2215-0366(16)30147-X)
- 4 Wildes, J. E., Forbush, K. T., Hagan, K. E., Marcus, M. D., Attia, E., Gianini, L. M., & Wu, W.
5 (2017). Characterizing severe and enduring anorexia nervosa: An empirical approach.
6 *International Journal of Eating Disorders*, 50(4), 389–397.
7 <https://doi.org/10.1002/eat.22651>
- 8 Wonderlich, S., Mitchell, J. E., Crosby, R. D., Myers, T. C., Kadlec, K., Lahaise, K., ...
9 Schander, L. (2012, May). Minimizing and treating chronicity in the eating disorders: A
10 clinical overview. *International Journal of Eating Disorders*.
11 <https://doi.org/10.1002/eat.20978>
- 12 Zeeck, A., Herpertz-Dahlmann, B., Friederich, H.-C., Brockmeyer, T., Resmark, G., Hagenah,
13 U., ... Hartmann, A. (2018). Psychotherapeutic Treatment for Anorexia Nervosa: A
14 Systematic Review and Network Meta-Analysis. *Frontiers in Psychiatry*, 9, 158.
15 <https://doi.org/10.3389/fpsyt.2018.00158>
- 16 Zipfel, S., Wild, B., Groß, G., Friederich, H.-C., Teufel, M., Schellberg, D., ... Herzog, W.
17 (2014). Focal psychodynamic therapy, cognitive behaviour therapy, and optimised
18 treatment as usual in outpatients with anorexia nervosa (ANTOP study): randomised
19 controlled trial. *The Lancet*, 383(9912), 127–137. [https://doi.org/10.1016/S0140-](https://doi.org/10.1016/S0140-6736(13)61746-8)
20 [6736\(13\)61746-8](https://doi.org/10.1016/S0140-6736(13)61746-8)