



## King's Research Portal

DOI:

[10.1530/JOE-19-0568](https://doi.org/10.1530/JOE-19-0568)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Simpson, S. J. S., Smith, L. I. F., Jones, P. M., & Bowe, J. E. (2020). UCN2: A new candidate influencing pancreatic  $\beta$ -cell adaptations in pregnancy. *Journal of Endocrinology*, 245(2), 247–257 .  
<https://doi.org/10.1530/JOE-19-0568>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

1 **UCN2: A new candidate influencing pancreatic  $\beta$ -cell adaptations to pregnancy**

2 **Sian.J.S.Simpson\*, Lorna.I.F.Smith, Peter.M.Jones, James.E.Bowe**

3

4 **Department of Diabetes, School of Life Course Sciences, Faculty of Life Science and**  
5 **Medicine, King's College London, London, United Kingdom**

6

7 **Corresponding author:**

8 Miss Sian.J.S.Simpson

9 Hodgkin Building

10 King's College London

11 Guy's Campus

12 London, SE1 1UL

13 [sian.simpson@kcl.ac.uk](mailto:sian.simpson@kcl.ac.uk)

14

15 **Short title**

16 UCN2 and  $\beta$ -cell adaptations to pregnancy

17

18 **Keywords**

19 Insulin, islet,  $\beta$ -cell adaptation, pregnancy, corticotropin-releasing hormone, urocortin

20

21 **Word count:**

22

23

24

25

**26 Abstract**

27 The corticotropin releasing hormone (CRH) family of peptides, including urocortin (UCN) 1,  
28 2 and 3, are established hypothalamic neuroendocrine peptides, regulating the physiological  
29 and behaviour responses to stress indirectly, via the hypothalamic-pituitary-adrenal (HPA)  
30 axis. More recently, these peptides have been implicated in diverse roles in peripheral organs  
31 through direct signalling, including in placental and pancreatic islet physiology. CRH has  
32 been shown to stimulate insulin release through activation of its cognate receptors, CRH  
33 receptor 1 (CRHR1) and 2. However, the physiological significance of this is unknown. We  
34 have previously reported that during mouse pregnancy, expression of CRH peptides increase  
35 in mouse placenta suggesting that these peptides may play a role in various biological  
36 functions associated with pregnancy, particularly the pancreatic islet adaptations that occur in  
37 the pregnant state to compensate for the physiological increase in maternal insulin resistance.  
38 In the current study, we show that mouse pregnancy is associated with increased circulating  
39 levels of UCN2 and that when we pharmacologically block endogenous CRHR signalling in  
40 pregnant mice, impairment of glucose tolerance is observed. This effect on glucose tolerance  
41 was comparable to that displayed with specific CRHR2 blockade and not with specific  
42 CRHR1 blockade. No effects on insulin sensitivity or the proliferative capacity of  $\beta$ -cells  
43 were detected. Thus, CRHR2 signalling appears to be involved in  $\beta$ -cell adaptive responses to  
44 pregnancy in the mouse, with endogenous placental UCN2 being the likely signal mediating  
45 this.

46

47

48

49

50

51

52 **Introduction**

53 The corticotropin releasing hormone (CRH) peptide family comprises CRH and the  
54 structurally related urocortin peptides (UCN1, UCN2 and UCN3). These neuroendocrine  
55 peptides are best known for their involvement in regulating the physiological and behavioural  
56 responses to stress, through the cognate G-protein coupled receptors (GPCRs), CRH receptor  
57 1 (CRHR1) and CRH receptor 2 (CRHR2) (Bakshi et al., 2002; Chen et al., 1993; Lovenberg  
58 et al., 1995; Weninger et al., 1999), as part of the hypothalamic-pituitary-adrenal (HPA) axis.  
59 More recent evidence suggests additional, diverse, extra-hypothalamic roles for these  
60 peptides in peripheral organs (Chatoo et al., 2018; Chatzaki et al., 2019; Paschos et al., 2013).  
61 Thus, CRH expression has been reported in the adrenal gland and the gastrointestinal tract  
62 (Suda et al., 1984); UCN1 is expressed in heart, skin and adipose tissue (Kimura et al., 2002;  
63 Seres et al., 2004; Wierzbicka et al., 2017); and UCN2 and UCN3 have been detected in  
64 peripheral blood cells, skeletal muscle, pancreas and gestational tissues such as fetal  
65 membranes and placental villi (Petraglia et al., 2010). CRHR1 and CRHR2 are also  
66 expressed in a wide range of tissues, including cardiac myocytes, the adrenal gland, adipose  
67 tissue, skeletal muscle and skin (Hillhouse and Grammatopoulos, 2001), also suggesting  
68 physiological roles for the CRH peptide family unrelated to the HPA axis. However, under  
69 normal circumstances, levels of the peptides in the peripheral circulation are low (Ng et al.,  
70 2004; Sasaki et al., 1987), suggesting that the peptides may be produced locally to function as  
71 autocrine or paracrine agents in tissues where the respective receptors are also expressed (Li  
72 et al., 2013; van der Meulen et al., 2015; Zouboulis et al., 2002).

73

74 There is increasing evidence that the CRH peptide family may be involved in peripheral  
75 metabolic control via direct actions on insulin-secreting  $\beta$ -cells in pancreatic islets of

76 Langerhans (Li et al., 2007; Schmid et al., 2011). Both CRHR1 and CRHR2 are expressed in  
77 rodent (Kanno et al., 1999; Schmid et al., 2011) and human islets (Amisten et al., 2013),  
78 whilst *in vitro* administration of exogenous CRH stimulates insulin secretion from mouse and  
79 human islets as well as enhancing proliferation in neonatal rat  $\beta$ -cells (Huising et al., 2010).  
80 Similarly,  $\beta$ -cell derived UCN3 has been implicated in the local regulation of both insulin and  
81 glucagon release (Li et al., 2007). Despite the evidence demonstrating direct effects of  
82 exogenous CRH on islet function, the physiological relevance of this interaction is unclear,  
83 given the islets would not normally be exposed to significant levels of peptides of the CRH  
84 family. There is some evidence that placentally-derived CRH and urocortins are involved in  
85 various biological functions associated with pregnancy (Thomson, 2013; You et al., 2014).  
86 Thus, pregnancy represents one possible physiological state in which the effects of the CRH  
87 family on islet function may play a role.

88

89 During pregnancy, maternal insulin resistance increases and this is compensated for by  
90 increases in  $\beta$ -cell mass and enhanced insulin secretory responses to elevations in plasma  
91 glucose (Baeyens et al., 2016; Pasek and Gannon, 2013; Xue et al., 2010). We have recently  
92 reported an upregulation of CRH, UCN2 and UCN3 mRNA expression in mouse placenta on  
93 gestational day 12 (Drynda et al., 2018), which correlates to the initiation of  $\beta$ -cell  
94 adaptations in rodent pregnancy (Rieck and Kaestner, 2010). Similarly, in human pregnancy,  
95 levels of CRH in the peripheral circulation increase as gestation progresses (Campbell et al.,  
96 1987; Sasaki et al., 1987) and CRH immunoreactivity has been reported in human placenta  
97 (Grino et al., 1987), consistent with a placental source for the circulating CRH. In the current  
98 study, we have therefore investigated a potential role for the CRH peptide family in the  
99 regulation of glucose homeostasis during pregnancy.

100

101

102 **Materials and methods**103 ***Animals***

104 Female Institute of Cancer Research (ICR) mice (8-12 weeks of age, Envigo, Bicester, UK)  
105 were used for *in vivo* studies. This is a commonly used outbred mouse strain with very good  
106 reproductive and maternal characteristics. All animals were housed under controlled,  
107 pathogen free conditions (12-hour light/ dark cycle (0700 – 1900 hr lights on), temperature  
108 22±2°C) and provided with standard chow diet and water ad libitum. For timed pregnancy  
109 studies, female mice were mated with male ICR mice and the presence of vaginal plug  
110 assessed daily and denoted day 1 of pregnancy if present. Age-matched female mice were  
111 used for non-pregnant studies, with procedures carried out at the same time intervals as  
112 described for pregnancy studies. All procedures were conducted under approval by King's  
113 College London Animal Welfare and Ethical Review Board and were undertaken in  
114 accordance with United Kingdom Home Office Regulations.

115

116 ***Islet isolation and insulin secretion in vitro***

117 For *in vitro* insulin secretion studies, pancreatic islets were isolated from female ICR mice  
118 via collagenase digestion of the exocrine pancreas, as described previously (Rackham et al.,  
119 2016). Isolated islets were subsequently maintained at 37°C in RPMI (Sigma, UK)  
120 supplemented with 10% [vol./vol.] fetal bovine serum, 2 mmol/l glutamine and 100 U/ml  
121 penicillin/0.1 mg/ml streptomycin for 24 hours before use. Islets were loaded into a multi-  
122 channel, temperature controlled perfusion system, as described previously (Liu et al., 2013),  
123 and pre-perfused for 1 hour with physiological salt buffer (Bowe et al., 2019) containing 2  
124 mmol/l glucose before being exposed to 20 mmol/l glucose in the presence or absence of the  
125 CRHR agonists, CRH (50 nmol/l, Sigma, UK), Stressin I (100 nmol/l, Tocris, UK) or UCN2

126 (100 nmol/l, Sigma, UK) at 37°C. Perifusate samples were collected every 2 minutes and  
127 insulin secretion was quantified using an in-house insulin radioimmunoassay (Jones et al.,  
128 1988).

129

### 130 *In vivo osmotic minipump studies*

131 Osmotic minipumps (ALZET®, Model 1002, Charles River, UK) were implanted  
132 subcutaneously into pregnant or non-pregnant mice to chronically administer test agents.  
133 Surgical implantation of osmotic minipumps was carried out on day 7 of pregnancy (or  
134 equivalent time interval for non-pregnant mice) under isoflurane anaesthesia (Isothesia®,  
135 Henry Schein®). Minipumps were loaded with physiological saline, non-specific CRHR  
136 antagonist ( $\alpha$ -helical CRF<sub>9-41</sub>, 1 mg/ml, Tocris, UK) or receptor specific CRHR antagonists,  
137 Antalarmin hydrochloride (1 mg/ml, Tocris, UK) or Antisauvagine-30 (3 mg/ml, Tocris, UK)  
138 for CRHR1 and R2, respectively. Test agents were delivered at a rate of 0.25  $\mu$ l /hour for a  
139 total period of 11 days. Assessment of glucose tolerance and insulin tolerance were  
140 conducted on gestational days 16 and 18, respectively.

141

### 142 *Assessment of glucose homeostasis*

143 Intraperitoneal glucose tolerance tests (IPGTT) were conducted on day 16 of gestation. Mice  
144 were fasted from 0900 for 6 hours and then administered with glucose (2 g/kg, Sigma, UK).  
145 Blood sampling was performed by small tail prick at time points 0, 15, 30, 60, 90 and 120  
146 min following glucose administration to determine blood glucose levels using an Accu-Chek  
147 glucose meter (Roche Diagnostics, UK). Intraperitoneal insulin tolerance tests (IPITT) were  
148 conducted on day 18 of gestation. Mice were again fasted from 0900 for 6 hours prior to  
149 metabolic testing and were subsequently administered with insulin (0.75 IU/kg, Sigma, UK).

150 Blood sampling was performed by small tail prick at time points 0, 15, 30, 45 and 60 min  
151 following insulin injection to determine blood glucose levels.

### 152 ***Measurements of circulating CRH-related peptides***

153 On day 18, animals were euthanised by intraperitoneal injection of terminal anaesthesia  
154 (Euthatal®, Merial Animal Health Ltd, UK) and terminal blood samples were collected via  
155 cardiac puncture into sterile heparin-coated tubes. Samples were also collected from control  
156 pregnant mice on day 16. Samples were centrifuged (1,800 rcf, 20 minutes, 4°C) and the  
157 subsequent plasma stored at -20°C for later assay of circulating peptide levels using  
158 commercially available ELISA kits (CRH: CEA835Mu, Cloud-Clone Corp, USA; UCN1:  
159 CEA231Mu, Cloud-Clone Corp, USA; UCN2: MOFI00425, ELISAGenie, UK; UCN3:  
160 CED140Mu, Cloud-Clone Corp, USA) following the manufacturers' instructions.

161

### 162 ***Quantification of mRNA expression***

163 Isolated female islets from non-pregnant and pregnant (day 16) mice were immediately snap  
164 frozen in liquid nitrogen following purification from the exocrine pancreas for subsequent  
165 RNA extraction using RNeasy Mini Kit (Qiagen, UK) and High Capacity cDNA Reverse  
166 Transcription Kit (Applied Biosystems™, UK) for cDNA synthesis, as described previously  
167 (Drynda et al., 2018). Placenta samples were also collected after termination at day 18 of  
168 pregnancy and snap frozen. RNA extraction and cDNA conversion were conducted as  
169 described above. Islet CRH receptor and placental CRH ligand mRNA expression was  
170 subsequently quantified by quantitative reverse transcription polymerase chain reaction  
171 (qRT-PCR) using SYBR Green PCR Kit (QuantiTect, Qiagen, UK) and a LC96 Light Cycler  
172 (Roche Diagnostics, UK). QuantiTect primer assays were used for expression analysis of  
173 genes of interest using Glyceraldehyde 3-phosphate dehydrogenase (*Gapdh*) as the  
174 housekeeping gene (Mouse *Crh*- QT01055789, *Ucn1*- QT00326879, *Ucn2*-QT01556534,



175 *Ucn3*-QT00302267, *Crhr1*-QT00106232, *Crhr2*-QT00151543, *Gapdh*-QT01658692, Qiagen,  
176 UK).

177

### 178 ***Assessment of $\beta$ -cell mass***

179 For osmotic minipump studies, bromo-deoxy-uridine (BrdU, 1 mg/ml, Sigma, UK) was  
180 administered in the drinking water from day 14 to day 18 of pregnancy with fresh BrdU  
181 drinking water being replaced every two days. After termination at day 18, pancreata were  
182 dissected, fixed in 4% paraformaldehyde (Sigma, UK) and embedded in paraffin wax before  
183 being cut into 5 $\mu$ m thick sections using Leica microtome (RM2255). Representative sections  
184 (3-4 sections per animal), approximately 150  $\mu$ m apart, were co-stained with guinea pig anti-  
185 insulin antibody (1:200, Dako, UK) to visualise islet  $\beta$ -cells and monoclonal mouse anti-  
186 BrdU antibody (1:100, Sigma, UK) to identify proliferating cells as previously described  
187 (Bowe et al., 2019). Images were taken on Nikon Eclipse TE2000-U fluorescent microscope  
188 and quantification of BrdU positive  $\beta$ -cells and  $\beta$ -cell area was performed using ImageJ 1.49c  
189 software.

190

### 191 ***Statistical analysis***

192 Statistical analysis was performed using GraphPad Prism 8.0 software. For comparison  
193 between two groups, unpaired, two-tailed students t-test was used. For *in vivo* glucose and  
194 insulin tolerance tests, two-way repeated measures ANOVA was used, followed by Tukey's  
195 multiple comparison test to identify significance between multiple groups.

196

197

## 198 **Results**

### 199 ***CRH receptor gene expression profile in pregnancy***

200 Islets isolated from non-pregnant and pregnant (d.16) female mice expressed both *Crhr1* and  
201 *Crhr2* mRNAs, as shown in Figure 1. As expected, *Crhr1* expression in islets was higher  
202 than *Crhr2* expression, displaying an analogous expression pattern for the receptors to that in  
203 the pituitary, a classical target for CRH. Islet *Crhr1* mRNA expression was significantly  
204 reduced during pregnancy compared to non-pregnant levels (Fig. 1A), whereas islet *Crhr2*  
205 mRNA levels were unchanged between non-pregnant and pregnant animals (Fig. 2B). Thus,  
206 islets express receptors for the entire CRH family of peptides.

207

### 208 ***Effects of CRH receptor stimulation on insulin secretion***

209 Activating either CRHR1 or CRHR2 enhanced glucose-induced insulin secretion from  
210 isolated mouse islets in a dynamic perfusion system, as shown in Figure 2. Exposure to 20  
211 mmol/l glucose initiated a rapid increase in insulin secretion, which was further potentiated  
212 by the addition of CRH (acting as a non-specific CRHR1 and CRHR2 agonist, Fig. 2A);  
213 Stressin I (a CRHR1 specific agonist, Fig. 2C); or of UCN2 (a CRHR2 specific agonist, Fig.  
214 2E). Area under the curve quantification of glucose stimulated insulin secretion (30-50 min)  
215 confirms the significant potentiation of insulin secretion in the presence of stimulatory  
216 concentrations of glucose, induced by all CRH receptor agonists tested (Fig. 2B, D, F).  
217 CRHR agonists had no significant effect on insulin secretion at a sub-stimulatory  
218 concentration of glucose (Data not shown, 2 mmol/l glucose: Control:  $0.056 \pm 0.010$   
219 ng/islet/hr vs + 50 nmol/l CRH:  $0.045 \pm 0.009$  vs + 100 nmol/l Stressin I:  $0.034 \pm 0.007$  vs +  
220 100 nmol/l Ucn2:  $0.053 \pm 0.008$ , mean  $\pm$  SEM, n=9 observations  $p > 0.999$ ). Thus, activation  
221 of CRHR1 or CRHR2 potentiates glucose-stimulated insulin secretion from islet  $\beta$ -cells.

222

### 223 ***Circulating CRH and urocortin profile during pregnancy***

224 qRT-PCR measurements demonstrated that mRNAs for *Crh*, *Ucn1*, *Ucn2* and *Ucn3* were all  
225 expressed by mouse placenta at day 18 at similar levels (Fig. 3A), confirming our previous  
226 observations (Drynda et al., 2018). Furthermore, all four peptides were detected in the  
227 peripheral circulation, with UCN2 being the most abundant circulating CRHR agonist (Fig.  
228 3B). The circulating levels of CRH, UCN1 and UCN3 were unchanged between non-  
229 pregnant and pregnant female mice. However, circulating levels of UCN2 were elevated  
230 almost 2-fold by day 16 of pregnancy when compared to age-matched virgin female controls  
231 (Fig. 3B). Thus, the pancreatic islets are likely to be exposed to elevated levels of UCN2  
232 during pregnancy, with the placenta being the most likely source for the increased levels.  
233 Therefore, the candidate ligand of the CRH family to play a physiological role in the islet  
234 adaptation to pregnancy appears to be UCN2.

235

### 236 *Effect of pharmacologically blocking endogenous CRH receptor signalling during* 237 *pregnancy*

238 The consequences of pharmacological blockade of CRH receptor signalling *in vivo* was  
239 assessed in both non-pregnant and pregnant mice, revealing a pregnancy- and receptor-  
240 specific phenotype, as shown in Figure 4. As expected, intraperitoneal administration of  
241 glucose, elevated blood glucose levels within 15 minutes in both pregnant and non-pregnant  
242 mice (Fig. 4A, E). Chronic pharmacological blockade of total CRHR signalling during  
243 pregnancy with a non-selective antagonist,  $\alpha$ -helical CRF<sub>9-41</sub>, resulted in a mild impairment to  
244 glucose tolerance, with significantly higher blood glucose concentrations at 15 minutes after  
245 glucose administration, compared to saline controls (Fig. 4A). Chronic administration of the  
246 CRHR2 antagonist, Antisauvagine-30, resulted in a similar impairment to glucose tolerance  
247 in pregnant mice, but not in animals treated with the specific CRHR1 antagonist, Antalarmin  
248 hydrochloride (Fig. 4A, B). These data are consistent with an endogenous ligand, acting via

249 CRHR2, playing a physiological role in maintaining normal glucose tolerance during  
250 pregnancy. All pregnant mice were insulin resistant by day 18 of pregnancy as indicated by  
251 the failure to respond to exogenous insulin administration and lowering of blood glucose,  
252 however, none of the CRHR antagonists had any detectable effects on insulin sensitivity (Fig.  
253 4C, D). Chronic treatment of non-pregnant female mice with  $\alpha$ -helical CRF<sub>9-41</sub> to block total  
254 CRHR signalling had no significant effect on glucose tolerance or insulin sensitivity (Fig.  
255 4E-H). Given the lack of effect of  $\alpha$ -helical CRF<sub>9-41</sub>, receptor specific antagonists were not  
256 tested outside of pregnancy. Thus, CRHR2 activation by an endogenous ligand is involved in  
257 maintaining glucose homeostasis specifically during pregnancy.

258

259 In addition to effects on whole body glucose homeostasis, pregnancy in mice is also  
260 associated with an increased rate of  $\beta$ -cell proliferation to increase the functional  $\beta$ -cell mass  
261 (Rieck and Kaestner, 2010). This was evaluated by BrdU<sup>+</sup>  $\beta$ -cell staining (Fig. 5A, B).  
262 Chronic blockade of total CRHR signalling during pregnancy using  $\alpha$ -helical CRH<sub>9-41</sub>, had no  
263 significant effects on  $\beta$ -cell proliferation,  $\beta$ -cell size or the average insulin<sup>+</sup>  $\beta$ -cell area, as  
264 shown in Figure 5C-E. The effects of CRHR activation on glucose homeostasis during  
265 pregnancy are therefore most likely direct effects on the  $\beta$ -cell to enhance insulin secretion  
266 rather than to increase the  $\beta$ -cell mass.

267

268

## 269 **Discussion**

270 During pregnancy, the metabolic profile of the mother adapts to ensure a sufficient supply of  
271 energy for the developing fetus. A progressive increase in maternal insulin resistance across  
272 pregnancy represents a key mechanism for increasing fuel availability to the fetus (Freemark,  
273 2006; Newbern and Freemark, 2011). This insulin resistance is compensated for by an

274 increase in the maternal functional  $\beta$ -cell mass and enhanced insulin secretory responses  
275 (Baeyens et al., 2016). Failure of the  $\beta$ -cell to adapt to the maternal metabolic load can lead  
276 to maternal glucose intolerance and, eventually, to overt gestational diabetes (Plows et al.,  
277 2018; Zhang et al., 2010). In rodent models, the early  $\beta$ -cell adaptations to pregnancy involve  
278 non-placental signals (Drynda et al., 2015), but as placentation is established and pregnancy  
279 progresses, the placenta becomes an important endocrine organ, secreting numerous  
280 hormonal signals, which influence maternal and fetal physiology (Jansson, 2016). The  
281 lactogenic hormones, prolactin and placental lactogen, are important pregnancy associated  
282 signals, well-established to act via  $\beta$ -cell prolactin receptors to induce  $\beta$ -cell mass  
283 expansion and enhance insulin secretion (Brelje et al., 1993; Huang et al., 2009; Sorenson et  
284 al., 1993; Vasavada et al., 2000). These effects may be mediated, at least in part, by an  
285 upregulation of intra-islet serotonin (Kim et al., 2010; Ohara-Imaizumi et al., 2013).  
286 However, the mouse placenta expresses approximately 80 different ligands for which  $\beta$ -cells  
287 express the cognate GPCRs (Drynda et al., 2018) and it is unlikely that the lactogenic  
288 hormones are the only signals involved in regulating islet adaptations. These placental  
289 ligands include a number of peptides more usually associated with hypothalamic  
290 neuroendocrine functions. We have recently identified kisspeptin as an important placental  
291 signal regulating  $\beta$ -cell function during pregnancy (Bowe et al., 2019). The current study  
292 extends these observations to implicate another classical hypothalamic neuroendocrine  
293 system, the CRH peptide family, in placental control of  $\beta$ -cell function.

294

295 The expression profile of CRH receptors in mouse islets is consistent with previous reports  
296 confirming the expression of both *Crhr1* and *Crhr2* using mouse (Huising et al., 2011) or  
297 human (Amisten et al., 2013) islets. These observations suggest that islet cells have an innate  
298 capacity to recognise and respond to circulating CRH and the urocortin peptides. The

299 decreased expression levels of *Crhr1* during pregnancy is also suggestive of a shift in the  
300 receptor ratio to potentially direct *Crhr2* signalling under the influence of placental signals.  
301 Accordingly, our *in vitro* measurements of insulin secretion from isolated islets,  
302 demonstrated that activation of either CRHR1 or CRHR2 significantly potentiates glucose-  
303 stimulated insulin secretion (GSIS). Similar to other  $\beta$ -cell GPCRs, activation of CRHR1 and  
304 CRHR2 only enhanced insulin secretion in the presence of a stimulatory concentration of  
305 glucose, suggesting that the physiological function of receptor activation is to modulate the  
306 extent of the insulin secretory response to elevated glucose concentrations, rather than to  
307 initiate secretion. Our dynamic measurements of insulin secretion from isolated islets  
308 correspond with studies using mouse or human islets in static incubations (Huisling et al.,  
309 2010; O'Carroll et al., 2008) and imply that increased levels of CRHR agonists will result in  
310 an enhanced glucose-induced insulin secretory response. However, whilst previous studies  
311 have suggested a role for the CRH family in regulating islet function, the physiological  
312 purpose of this effect was unclear.

313

314 Placental expression and secretion of CRHR agonists is contentious. Earlier studies detected  
315 CRH mRNA and immunoreactivity in placentae from humans and non-human primates (Frim  
316 et al., 1988; Robinson et al., 1989; Sasaki et al., 1987), but failed to detect it in non-primate  
317 species including lemur, guinea pig and rat (Robinson et al., 1989). In human pregnancy,  
318 levels of CRH in the peripheral circulation increase as gestation progresses (Campbell et al.,  
319 1987; Sasaki et al., 1987). It has thus been suggested that the physiological purpose of this  
320 increase is in regulating parturition through modulation of signals controlling myometrium  
321 contractility and inflammation (McLean et al., 1995; Thomson, 2013; You et al., 2014).  
322 Contrary to human pregnancy, placental CRH in rodents is not thought to have a significant  
323 role in initiating parturition, with evidence of a more influential role in facilitating

324 implantation particularly during murine pregnancy (Athanasakis et al., 1999). Increased  
325 expression of UCN2 mRNA and protein has been reported in both human and mouse  
326 gestational tissues (including fetal membranes, myometrium and placenta) (Voltolini et al.,  
327 2015), although conflicting reports suggest no significant change in circulating levels of  
328 UCN1, UCN2 or UCN3 during human pregnancy (Pepels et al., 2010). In the current study  
329 we detected the expression of mRNAs for all members of the CRH family in mouse placenta.  
330 Circulating levels of CRH, UCN1 and UCN3 were unchanged in pregnant and non-pregnant  
331 mice, suggesting that these ligands are not released by the mouse placenta at significant  
332 levels, however circulating levels of UCN2 were significantly increased during gestation. The  
333 circulating concentrations of UCN2 which we detected during pregnancy are close to the  
334 reported EC50 values for CRHR2 (Dautzenberg et al., 2004; Hauger et al., 2003; Patel et al.,  
335 2012) and are consistent with  $\beta$ -cell CRHR2 activation in response to pregnancy signals.  
336 These observations are also consistent with the placenta being the source of the increased  
337 circulating UCN2 during mouse pregnancy, analogous to the increases in placentally derived  
338 kisspeptin in the circulation during mouse and human pregnancy (Bowe et al., 2019; Dhillon et  
339 al., 2006; Mark et al., 2013), and suggest that it may potentially play a physiological role  
340 during pregnancy. However, it cannot be ruled out that the pregnancy-associated UCN2  
341 derives from an alternative peripheral source, such as skin or skeletal muscle where it is also  
342 highly expressed (Chen et al., 2004).

343

344 Irrespective of its source, our *in vivo* studies suggest a role for circulating UCN2 in the  
345 regulation of  $\beta$ -cell insulin secretory responses during mouse pregnancy. Thus,  
346 pharmacological blockade of CRHR2 impaired glucose tolerance in pregnant mice, but a  
347 similar impairment was not observed with CRHR1 blockade, nor in non-pregnant females.  
348 The lack of effect of *in vivo* CRHR blockade on insulin resistance during pregnancy suggests

349 that the impaired glucose tolerance reflects a  $\beta$ -cell targeted effect, consistent with our *in*  
350 *vitro* observations of enhanced insulin secretion in response to CRHR2 activation. Most  
351 placental hormones involved in  $\beta$ -cell adaptations to pregnancy exert dual effects to acutely  
352 increase the rate of insulin secretion from individual  $\beta$ -cells, and chronically to induce  
353 expansion of the functional  $\beta$ -cell pool. These compensatory mechanisms ensure that the  
354 mother can sustain a robust insulin secretory response to elevated plasma glucose, especially  
355 in the prevailing insulin resistant environment. Under normal circumstances the rate of  $\beta$ -cell  
356 proliferation is very low, but chronic exposure to lactogenic hormones (Baeyens et al., 2016;  
357 Brelje et al., 1993; Huang et al., 2009) or to kisspeptin (Bowe et al., 2019) during gestation  
358 increases the rate of  $\beta$ -cell proliferation, and so increases the functional  $\beta$ -cell mass both *in*  
359 *vitro* and *in vivo*. In the current study, chronic blockade of total CRH receptors during  
360 pregnancy had no significant effects on  $\beta$ -cell size or proliferation, nor on the overall  $\beta$ -cell  
361 mass. This provides further evidence that the impairment to glucose tolerance *in vivo* during  
362 pregnancy, is due to an endogenous ligand, specifically targeting CRHR2, enhancing  $\beta$ -cell  
363 insulin secretion. The physiological significance of these differences in modes of action of  
364 placental factors is uncertain, but there may be therapeutic advantages in the ability of UCN2  
365 to enhance glucose-induced insulin secretion without targeting the clinical challenges of  
366 manipulating  $\beta$ -cell proliferation.

367

368 The variability of maternal glycaemia throughout pregnancy can range from normal/mild  
369 glucose intolerance, to severe in the case of gestational diabetes. The pharmacological  
370 blockade of CRHR2 signalling during pregnancy appears to reveal a transient and mild  
371 glucose intolerance in comparison to the more profound defect in glucose tolerance displayed  
372 by mutant PRLR mice (Huang et al., 2009). Given the importance of maintaining appropriate  
373 maternal glycaemic control during pregnancy, it is perhaps not surprising that there are



374 multiple control mechanisms that ensure an integrated  $\beta$ -cell insulin secretory response.  
375 Therefore, the mild phenotype displayed may have been compensated by complementary  
376 signals to prevent major disruptions to glucose homeostasis.

377

378 In summary, we have demonstrated that CRHR2 signalling is involved in  $\beta$ -cell adaptive  
379 responses to pregnancy in the mouse, with endogenous placental UCN2 being the likely  
380 signal mediating this adaptation. Unlike other identified placental signals, the effects of  
381 UCN2 appear to be confined to amplifying glucose-induced insulin secretion without  
382 concomitant alterations in the  $\beta$ -cell mass. Blocking the endogenous CRHR2 agonist during  
383 gestation induces a mild glucose intolerance rather than overt gestational diabetes suggesting  
384 that UCN2 may act in concert with other placental signals to fine-tune the compensatory  $\beta$ -  
385 cell adaptations to maternal insulin resistance during pregnancy. Deciphering the interplay  
386 between these different signals will lead to a more comprehensive understanding of the  
387 pathophysiology of gestational diabetes and may offer novel diagnostic or therapeutic  
388 strategies.

389

### 390 **Declaration of interest, Funding and Acknowledgements**

391 The authors declare that there is no conflict of interest that could be perceived as prejudicing  
392 the impartiality of the research reported. This work was supported by grant funding from both  
393 the Diabetes Research and Wellness Foundation (DRWF) (SCA/OF/12/18) and the Medical  
394 Research Council (MRC) Doctoral Training Program studentship (Sian JS Simpson).

395

396

397

398

399  
400  
401  
402  
403  
404  
405  
406

407 **Bibliography**

408

- 409 Amisten, S., Salehi, A., Rorsman, P., Jones, P.M., Persaud, S.J., 2013. An atlas and  
410 functional analysis of G-protein coupled receptors in human islets of Langerhans. *Pharmacol.*  
411 *Ther.* 139, 359–391. <https://doi.org/10.1016/j.pharmthera.2013.05.004>
- 412 Athanassakis, I., Farmakiotis, V., Aifantis, I., Gravanis, A., Vassiliadis, S., 1999. Expression  
413 of corticotrophin-releasing hormone in the mouse uterus: participation in embryo  
414 implantation. *J. Endocrinol.* 163, 221–227. <https://doi.org/10.1677/joe.0.1630221>
- 415 Baeyens, L., Hindi, S., Sorenson, R.L., German, M.S., 2016.  $\beta$ -Cell Adaptation in Pregnancy.  
416 *Diabetes Obes Metab* 18, 63–70. <https://doi.org/10.1111/dom.12716>
- 417 Bakshi, V.P., Smith-Roe, S., Newman, S.M., Grigoriadis, D.E., Kalin, N.H., 2002. Reduction  
418 of Stress-Induced Behavior by Antagonism of Corticotropin-Releasing Hormone 2 (CRH2)  
419 Receptors in Lateral Septum or CRH1 Receptors in Amygdala. *J. Neurosci.* 22, 2926–2935.  
420 <https://doi.org/10.1523/JNEUROSCI.22-07-02926.2002>
- 421 Bowe, J.E., Hill, T.G., Hunt, K.F., Smith, L.I.F., Simpson, S.J.S., Amiel, S.A., Jones, P.M.,  
422 2019. A role for placental kisspeptin in  $\beta$  cell adaptation to pregnancy. *JCI Insight* 4.  
423 <https://doi.org/10.1172/jci.insight.124540>

424 Brelje, T.C., Scharp, D.W., Lacy, P.E., Ogren, L., Talamantes, F., Robertson, M., Friesen,  
425 H.G., Sorenson, R.L., 1993. Effect of homologous placental lactogens, prolactins, and growth  
426 hormones on islet B-cell division and insulin secretion in rat, mouse, and human islets:  
427 implication for placental lactogen regulation of islet function during pregnancy.  
428 *Endocrinology* 132, 879–887. <https://doi.org/10.1210/endo.132.2.8425500>

429 Campbell, E.A., Linton, E.A., Wolfe, C.D.A., Scraggs, P.R., Jones, M.T., Lowry, P.J., 1987.  
430 Plasma Corticotropin-Releasing Hormone Concentrations During Pregnancy and Parturition.  
431 *None* 64, 1054–1059. <https://doi.org/10.1210/jcem-64-5-1054>

432 Chato, M., Li, Y., Ma, Z., Coote, J., Du, J., Chen, X., 2018. Involvement of Corticotropin-  
433 Releasing Factor and Receptors in Immune Cells in Irritable Bowel Syndrome. *Front.*  
434 *Endocrinol.* 9. <https://doi.org/10.3389/fendo.2018.00021>

435 Chatzaki, E., Kefala, N., Drosos, I., Lalidou, F., Baritaki, S., 2019. Do urocortins have a role  
436 in treating cardiovascular disease? *Drug Discovery Today* 24, 279–284.  
437 <https://doi.org/10.1016/j.drudis.2018.09.004>

438 Chen, A., Blount, A., Vaughan, J., Brar, B., Vale, W., 2004. Urocortin II Gene Is Highly  
439 Expressed in Mouse Skin and Skeletal Muscle Tissues: Localization, Basal Expression in  
440 Corticotropin-Releasing Factor Receptor (CRFR) 1- and CRFR2-Null Mice, and Regulation  
441 by Glucocorticoids. *Endocrinology* 145, 2445–2457. <https://doi.org/10.1210/en.2003-1570>

442 Chen, R., Lewis, K.A., Perrin, M.H., Vale, W.W., 1993. Expression cloning of a human  
443 corticotropin-releasing-factor receptor. *Proc. Natl. Acad. Sci. U.S.A.* 90, 8967–8971.

444 Dautzenberg, F.M., Gutknecht, E., Linden, I.V. der, Olivares-Reyes, J.A., Dürrenberger, F.,  
445 Hauger, R.L., 2004. Cell-type specific calcium signaling by corticotropin-releasing factor  
446 type 1 (CRF1) and 2a (CRF2(a)) receptors: phospholipase C-mediated responses in human  
447 embryonic kidney 293 but not SK-N-MC neuroblastoma cells. *Biochemical Pharmacology*  
448 68, 1833–1844. <https://doi.org/10.1016/j.bcp.2004.07.013>

- 449 Dhillon, W.S., et al., 2006. Plasma kisspeptin is raised in patients with gestational  
450 trophoblastic neoplasia and falls during treatment. *Am. J. Physiol. Endocrinol. Metab.* 291,  
451 E878-884. <https://doi.org/10.1152/ajpendo.00555.2005>
- 452 Drynda, R., Persaud, S.J., Bowe, J.E., Jones, P.M., 2018. The Placental Secretome:  
453 Identifying Potential Cross-Talk Between Placenta and Islet  $\beta$ -Cells. *Cell. Physiol. Biochem.*  
454 45, 1165–1171. <https://doi.org/10.1159/000487357>
- 455 Drynda, R., Peters, C.J., Jones, P.M., Bowe, J.E., 2015. The role of non-placental signals in  
456 the adaptation of islets to pregnancy. *Horm. Metab. Res.* 47, 64–71. [https://doi.org/10.1055/s-](https://doi.org/10.1055/s-0034-1395691)  
457 0034-1395691
- 458 Freemark, M., 2006. Regulation of Maternal Metabolism by Pituitary and Placental  
459 Hormones: Roles in Fetal Development and Metabolic Programming. *HRP* 65, 41–49.  
460 <https://doi.org/10.1159/000091505>
- 461 Frim, D.M., Emanuel, R.L., Robinson, B.G., Smas, C.M., Adler, G.K., Majzoub, J.A., 1988.  
462 Characterization and gestational regulation of corticotropin-releasing hormone messenger  
463 RNA in human placenta. *J Clin Invest* 82, 287–292.
- 464 Grino, M., Chrousos, G.P., Margioris, A.N., 1987. The corticotropin releasing hormone gene  
465 is expressed in human placenta. *Biochem. Biophys. Res. Commun.* 148, 1208–1214.  
466 [https://doi.org/10.1016/s0006-291x\(87\)80261-9](https://doi.org/10.1016/s0006-291x(87)80261-9)
- 467 Hauger, R.L., Grigoriadis, D.E., Dallman, M.F., Plotsky, P.M., Vale, W.W., Dautzenberg,  
468 F.M., 2003. International Union of Pharmacology. XXXVI. Current Status of the  
469 Nomenclature for Receptors for Corticotropin-Releasing Factor and Their Ligands.  
470 *Pharmacol Rev* 55, 21–26. <https://doi.org/10.1124/pr.55.1.3>
- 471 Hillhouse, E.W., Grammatopoulos, D.K., 2001. Characterising the Corticotropin-releasing  
472 Hormone (CRH) Receptors Mediating CRH and Urocortin Actions During Human  
473 Pregnancy and Labour. *Stress* 4, 235–246. <https://doi.org/10.3109/10253890109014748>

- 474 Huang, C., Snider, F., Cross, J.C., 2009. Prolactin receptor is required for normal glucose  
475 homeostasis and modulation of beta-cell mass during pregnancy. *Endocrinology* 150, 1618–  
476 1626. <https://doi.org/10.1210/en.2008-1003>
- 477 Huising, M.O., Pilbrow, A.P., Matsumoto, M., van der Meulen, T., Park, H., Vaughan, J.M.,  
478 Lee, S., Vale, W.W., 2011. Glucocorticoids Differentially Regulate the Expression of CRFR1  
479 and CRFR2 $\alpha$  in MIN6 Insulinoma Cells and Rodent Islets. *Endocrinology* 152, 138–150.  
480 <https://doi.org/10.1210/en.2010-0791>
- 481 Huising, M.O., van der Meulen, T., Vaughan, J.M., Matsumoto, M., Donaldson, C.J., Park,  
482 H., Billestrup, N., Vale, W.W., 2010. CRFR1 is expressed on pancreatic beta cells, promotes  
483 beta cell proliferation, and potentiates insulin secretion in a glucose-dependent manner. *Proc.*  
484 *Natl. Acad. Sci. U.S.A.* 107, 912–917. <https://doi.org/10.1073/pnas.0913610107>
- 485 Jansson, T., 2016. Placenta plays a critical role in maternal–fetal resource allocation.  
486 *Proceedings of the National Academy of Sciences* 113, 11066–11068.  
487 <https://doi.org/10.1073/pnas.1613437113>
- 488 Jones, P.M., SALMONT, D.M.W., Howell, S.L., 1988. Protein phosphorylation in electrically  
489 permeabilized islets of Langerhans 254, 7.
- 490 Kanno, T., Suga, S., Nakano, K., Kamimura, N., Wakui, M., 1999. Corticotropin-releasing  
491 factor modulation of Ca<sup>2+</sup> influx in rat pancreatic beta-cells. *Diabetes* 48, 1741–1746.
- 492 Kim, H., et al., 2010. Serotonin regulates pancreatic beta cell mass during pregnancy. *Nat.*  
493 *Med.* 16, 804–808. <https://doi.org/10.1038/nm.2173>
- 494 Kimura, Y., Takahashi, K., Totsune, K., Muramatsu, Y., Kaneko, C., Darnel, A.D., Suzuki,  
495 T., Ebina, M., Nukiwa, T., Sasano, H., 2002. Expression of urocortin and corticotropin-  
496 releasing factor receptor subtypes in the human heart. *J. Clin. Endocrinol. Metab.* 87, 340–  
497 346. <https://doi.org/10.1210/jcem.87.1.8160>

- 498 Li, C., Chen, P., Vaughan, J., Lee, K.-F., Vale, W., 2007. Urocortin 3 regulates glucose-  
499 stimulated insulin secretion and energy homeostasis. *Proc. Natl. Acad. Sci. U.S.A.* 104,  
500 4206–4211. <https://doi.org/10.1073/pnas.0611641104>
- 501 Li, J., et al., 2013. Urocortin 2 autocrine/paracrine and pharmacologic effects to activate  
502 AMP-activated protein kinase in the heart. *PNAS* 110, 16133–16138.  
503 <https://doi.org/10.1073/pnas.1312775110>
- 504 Liu, B., Hassan, Z., Amisten, S., King, A.J., Bowe, J.E., Huang, G.C., Jones, P.M., Persaud,  
505 S.J., 2013. The novel chemokine receptor, G-protein-coupled receptor 75, is expressed by  
506 islets and is coupled to stimulation of insulin secretion and improved glucose homeostasis.  
507 *Diabetologia* 56, 2467–2476. <https://doi.org/10.1007/s00125-013-3022-x>
- 508 Lovenberg, T.W., Liaw, C.W., Grigoriadis, D.E., Clevenger, W., Chalmers, D.T., De Souza,  
509 E.B., Oltersdorf, T., 1995. Cloning and characterization of a functionally distinct  
510 corticotropin-releasing factor receptor subtype from rat brain. *Proc. Natl. Acad. Sci. U.S.A.*  
511 92, 836–840.
- 512 Mark, P.J., Jones, M.L., Lewis, J.L., Waddell, B.J., Smith, J.T., 2013. Kiss1 and Kiss1r  
513 mRNA expression in the rat placenta: changes with gestational age and regulation by  
514 glucocorticoids. *Placenta* 34, 657–662. <https://doi.org/10.1016/j.placenta.2013.04.012>
- 515 McLean, M., Bisits, A., Davies, J., Woods, R., Lowry, P., Smith, R., 1995. A placental clock  
516 controlling the length of human pregnancy. *Nature Medicine* 1, 460–463.  
517 <https://doi.org/10.1038/nm0595-460>
- 518 Newbern, D., Freemark, M., 2011. Placental hormones and the control of maternal  
519 metabolism and fetal growth. *Curr Opin Endocrinol Diabetes Obes* 18, 409–416.  
520 <https://doi.org/10.1097/MED.0b013e32834c800d>

- 521 O'Carroll, A.-M., Howell, G.M., Roberts, E.M., Lolait, S.J., 2008. Vasopressin potentiates  
522 corticotropin-releasing hormone-induced insulin release from mouse pancreatic  $\beta$ -cells. *J*  
523 *Endocrinol* 197, 231–239. <https://doi.org/10.1677/JOE-07-0645>
- 524 Ohara-Imaizumi, M., et al., 2013. Serotonin regulates glucose-stimulated insulin secretion  
525 from pancreatic  $\beta$  cells during pregnancy. *Proc. Natl. Acad. Sci. U.S.A.* 110, 19420–19425.  
526 <https://doi.org/10.1073/pnas.1310953110>
- 527 Paschos, K., Chouridou, E., Koureta, M., Lambropoulou, M., Kolios, G., Chatzaki, E., 2013.  
528 The corticotropin releasing factor system in the liver: expression, actions and possible  
529 implications in hepatic physiology and pathology. *HORMONES* 12, 236–245.  
530 <https://doi.org/10.14310/horm.2002.1407>
- 531 Pasek, R.C., Gannon, M., 2013. Advancements and challenges in generating accurate animal  
532 models of gestational diabetes mellitus. *Am. J. Physiol. Endocrinol. Metab.* 305, E1327-  
533 1338. <https://doi.org/10.1152/ajpendo.00425.2013>
- 534 Patel, K., Rademaker, M., Kirkpatrick, C., Charles, C., Fisher, S., Yandle, T., Richards, A.,  
535 2012. Comparative pharmacokinetics and pharmacodynamics of urocortins 1, 2 and 3 in  
536 healthy sheep. *British Journal of Pharmacology* 166, 1916–1925.  
537 <https://doi.org/10.1111/j.1476-5381.2012.01904.x>
- 538 Pepels, P.P.L.M., Spaanderman, M.E.A., Hermus, A.R.M.M., Lotgering, F.K., Sweep,  
539 C.G.J., 2010. Placental urocortin-2 and -3: endocrine or paracrine functioning during healthy  
540 pregnancy? *Placenta* 31, 475–481. <https://doi.org/10.1016/j.placenta.2010.03.012>
- 541 Petraglia, F., Imperatore, A., Challis, J.R.G., 2010. Neuroendocrine Mechanisms in  
542 Pregnancy and Parturition. *Endocrine Reviews* 31, 783–816. [https://doi.org/10.1210/er.2009-](https://doi.org/10.1210/er.2009-0019)  
543 0019

- 544 Plows, J.F., Stanley, J.L., Baker, P.N., Reynolds, C.M., Vickers, M.H., 2018. The  
545 Pathophysiology of Gestational Diabetes Mellitus. *Int J Mol Sci* 19.  
546 <https://doi.org/10.3390/ijms19113342>
- 547 Rackham, C.L., Vargas, A.E., Hawkes, R.G., Amisten, S., Persaud, S.J., Austin, A.L.F.,  
548 King, A.J.F., Jones, P.M., 2016. Annexin A1 Is a Key Modulator of Mesenchymal Stromal  
549 Cell-Mediated Improvements in Islet Function. *Diabetes* 65, 129–139.  
550 <https://doi.org/10.2337/db15-0990>
- 551 Rieck, S., Kaestner, K.H., 2010. Expansion of beta-cell mass in response to pregnancy.  
552 *Trends Endocrinol. Metab.* 21, 151–158. <https://doi.org/10.1016/j.tem.2009.11.001>
- 553 Robinson, B.G., Arbiser, J.L., Emanuel, R.L., Majzoub, J.A., 1989. Species-specific placental  
554 corticotropin releasing hormone messenger RNA and peptide expression. *Molecular and*  
555 *Cellular Endocrinology* 62, 337–341. [https://doi.org/10.1016/0303-7207\(89\)90022-1](https://doi.org/10.1016/0303-7207(89)90022-1)
- 556 Sasaki, A., Shinkawa, O., Margioris, A.N., Liotta, A.S., Sato, S., Murakami, O., Go, M.,  
557 Shimizu, Y., Hanew, K., Yoshinaga, K., 1987. Immunoreactive corticotropin-releasing  
558 hormone in human plasma during pregnancy, labor, and delivery. *J. Clin. Endocrinol. Metab.*  
559 64, 224–229. <https://doi.org/10.1210/jcem-64-2-224>
- 560 Schmid, J., et al., 2011. Modulation of pancreatic islets-stress axis by hypothalamic releasing  
561 hormones and 11 $\beta$ -hydroxysteroid dehydrogenase. *PNAS* 108, 13722–13727.  
562 <https://doi.org/10.1073/pnas.1110965108>
- 563 Seres, J., Bornstein, S.R., Seres, P., Willenberg, H.S., Schulte, K.M., Scherbaum, W.A.,  
564 Ehrhart-Bornstein, M., 2004. Corticotropin-Releasing Hormone System in Human Adipose  
565 Tissue. *None* 89, 965–970. <https://doi.org/10.1210/jc.2003-031299>
- 566 Sorenson, R.L., Brelje, T.C., Roth, C., 1993. Effects of steroid and lactogenic hormones on  
567 islets of Langerhans: a new hypothesis for the role of pregnancy steroids in the adaptation of



568 islets to pregnancy. *Endocrinology* 133, 2227–2234.  
569 <https://doi.org/10.1210/endo.133.5.8404674>

570 Suda, T., Tomori, N., Tozawa, F., Mouri, T., Demura, H., Shizume, K., 1984. Distribution  
571 and Characterization of Immunoreactive Corticotropin-Releasing Factor in Human Tissues.  
572 *None* 59, 861–866. <https://doi.org/10.1210/jcem-59-5-861>

573 Thomson, M., 2013. The physiological roles of placental corticotropin releasing hormone in  
574 pregnancy and childbirth. *J. Physiol. Biochem.* 69, 559–573. [https://doi.org/10.1007/s13105-](https://doi.org/10.1007/s13105-012-0227-2)  
575 [012-0227-2](https://doi.org/10.1007/s13105-012-0227-2)

576 van der Meulen, T., Donaldson, C.J., Cáceres, E., Hunter, A.E., Cowing-Zitron, C., Pound,  
577 L.D., Adams, M.W., Zembrzycki, A., Grove, K.L., Huising, M.O., 2015. Urocortin3  
578 mediates somatostatin-dependent negative feedback control of insulin secretion. *Nat Med* 21,  
579 769–776. <https://doi.org/10.1038/nm.3872>

580 Vasavada, R.C., Garcia-Ocaña, A., Zawalich, W.S., Sorenson, R.L., Dann, P., Syed, M.,  
581 Ogren, L., Talamantes, F., Stewart, A.F., 2000. Targeted expression of placental lactogen in  
582 the beta cells of transgenic mice results in beta cell proliferation, islet mass augmentation,  
583 and hypoglycemia. *J. Biol. Chem.* 275, 15399–15406.

584 Voltolini, C., Battersby, S., Novembri, R., Torricelli, M., Severi, F.M., Petraglia, F., Norman,  
585 J.E., 2015. Urocortin 2 Role in Placental and Myometrial Inflammatory Mechanisms at  
586 Parturition. *Endocrinology* 156, 670–679. <https://doi.org/10.1210/en.2014-1432>

587 Weninger, S.C., Dunn, A.J., Muglia, L.J., Dikkes, P., Miczek, K.A., Swiergiel, A.H.,  
588 Berridge, C.W., Majzoub, J.A., 1999. Stress-induced behaviors require the corticotropin-  
589 releasing hormone (CRH) receptor, but not CRH. *PNAS* 96, 8283–8288.  
590 <https://doi.org/10.1073/pnas.96.14.8283>

- 591 Wierzbicka, J.M., Żmijewski, M.A., Antoniewicz, J., Sobjanek, M., Slominski, A.T., 2017.  
592 Differentiation of Keratinocytes Modulates Skin HPA Analog. *Journal of Cellular Physiology*  
593 232, 154–166. <https://doi.org/10.1002/jcp.25400>
- 594 Xue, Y., Liu, C., Xu, Y., Yuan, Q., Xu, K., Mao, X., Chen, G., Wu, X., Brendel, M.D., Liu,  
595 Chao, 2010. Study on pancreatic islet adaptation and gene expression during pregnancy in  
596 rats. *Endocrine* 37, 83–97. <https://doi.org/10.1007/s12020-009-9273-0>
- 597 You, X., Liu, J., Xu, C., Liu, W., Zhu, X., Li, Y., Sun, Q., Gu, H., Ni, X., 2014.  
598 Corticotropin-releasing hormone (CRH) promotes inflammation in human pregnant  
599 myometrium: the evidence of CRH initiating parturition? *J. Clin. Endocrinol. Metab.* 99,  
600 E199-208. <https://doi.org/10.1210/jc.2013-3366>
- 601 Zhang, H., Zhang, J., Pope, C.F., Crawford, L.A., Vasavada, R.C., Jagasia, S.M., Gannon,  
602 M., 2010. Gestational Diabetes Mellitus Resulting From Impaired  $\beta$ -Cell Compensation in  
603 the Absence of FoxM1, a Novel Downstream Effector of Placental Lactogen. *Diabetes* 59,  
604 143–152. <https://doi.org/10.2337/db09-0050>
- 605 Zouboulis, C.C., Seltmann, H., Hiroi, N., Chen, W., Young, M., Oeff, M., Scherbaum, W.A.,  
606 Orfanos, C.E., McCann, S.M., Bornstein, S.R., 2002. Corticotropin-releasing hormone: an  
607 autocrine hormone that promotes lipogenesis in human sebocytes. *Proc. Natl. Acad. Sci.*  
608 U.S.A. 99, 7148–7153. <https://doi.org/10.1073/pnas.102180999>

609

### 610 **Figure legends**

611

612 **Figure. 1:** Expression of *Crhr1* (A) and *Crhr2* (B) mRNAs by isolated female islets in non-  
613 pregnancy (white bar) and pregnancy day 16 (PD.16; black bar). Anterior pituitary was used  
614 as a positive control (grey bar) and mRNA expression levels were quantified to the relative  
615 expression of housekeeping gene, *Gapdh*. *Crhr1* mRNA expression levels decreased

616 significantly during pregnancy (~60%) whereas levels of *Crhr2* expression were unchanged.  
617 Data are presented as mean + S.E.M., n=5, \*p<0.05; Students T-test non-pregnant vs PD.16.  
618 **Figure. 2:** Effect of exogenous CRH (A), CRHR1 specific agonist Stressin 1 (C) and  
619 CRHR2 specific agonist UCN2 (E) on dynamic insulin secretion from isolated, perfused  
620 female mouse islets. Islets were exposed to physiological buffer containing 20 mmol/l  
621 glucose only or supplemented with agonists between 30-50 mins. All CRHR agonists  
622 potentiated glucose-stimulated insulin secretion over that seen from control islets, as  
623 demonstrated by the rate of insulin secretion (A,C,E) and area under curve data (B,D,F).  
624 Data are presented as mean ± S.E.M., n= 3-4 per treatment group, AUC 20 mmol/l glucose +  
625 agonist, 30-50 min, \*p<0.05, \*\*p<0.01, \*\*\*p<0.001; Students T-test control vs agonist  
626 treatment.

627

628 **Figure. 3:** Expression of CRH and urocortins mRNAs in mouse placenta on day 18 of  
629 pregnancy (PD.18) (A) and circulating concentrations of CRH peptides during mouse  
630 pregnancy (PD.16) (B). Expression levels were quantified to the relative expression of  
631 housekeeping gene *Gapdh*. *Crh*, *Ucn1*, *Ucn2* and *Ucn3* mRNAs were all expressed by mouse  
632 placenta. Plasma levels of CRH, UCN1 and UCN3 were similar in pregnant and non-  
633 pregnant mice. However, plasma UCN2 was significantly elevated during pregnancy. Data  
634 presented as mean + S.E.M., n=6, \*\*\*p<0.001; Two-way ANOVA followed by Tukey's  
635 multiple comparisons test.

636


637 **Figure. 4:** Effects of chronic administration of CRHR antagonists on glucose homeostasis  
638 during pregnancy (A-D) and non-pregnancy (E-H). Pregnant mice (PD.16) treated with either  
639  $\alpha$ -helical CRF<sub>9-41</sub> or AS-30 (Antisauvagine-30) displayed a significant impairment in glucose  
640 tolerance 15 mins after glucose loading (2g/kg) when comparison to control mice

641 administered saline (solid black line circle icon). No difference in glucose tolerance was seen  
642 in mice administered AH (Antalarmin hydrochloride). AUC from 0-30 min for each  
643 treatment group is displayed in panel B. No change in overall insulin sensitivity was observed  
644 between all treatment groups (C). AUC from 0-60min for each treatment group is displayed  
645 in panel D, (n=7-19). In non-pregnant mice chronic administration of  $\alpha$ -helical CRF<sub>9-41</sub> had  
646 no significant effects on glucose tolerance (E) or insulin sensitivity (G). AUC for glucose  
647 tolerance 0-30 min and insulin sensitivity 0-60 min are displayed in panel F and H  
648 respectively, (n=5-6). Data are presented as mean  $\pm$  S.E.M., # (Control vs  $\alpha$ -helical CRF<sub>9-41</sub>) /  
649 \* (Control vs AS-30): 15 mins p<0.05; Two-way repeated measures ANOVA followed by  
650 Tukey's multiple comparisons test.

651


652 **Figure. 5:** Effect of chronic administration of a non-selective CRHR antagonist ( $\alpha$ -helical  
653 CRF<sub>9-41</sub>) on  $\beta$ -cell morphology during pregnancy. Representative images of immunostaining  
654 for the measurement of  $\beta$ -cell proliferation in control (A) and  $\alpha$ -helical CRF<sub>9-41</sub> (B) islets  
655 showing insulin staining (red) and BrdU staining (green). Mice administered BrdU from days  
656 14 – 18 of pregnancy displayed no significant differences in the percentage of BrdU labelled  
657  $\beta$ -cells between control and  $\alpha$ -helical CRF<sub>9-41</sub> treated mice (C). Average  $\beta$ -cell size (D) and  
658 average  $\beta$ -cell islet area (E) were also unchanged between control and antagonist treatments.  
659 Data presented showing quantification (3-4 sections/animal analysed) for individual animals  
660 with bar showing mean, n=8-9 animals per treatment group. Scale bar- 50  $\mu$ m

Figure 1




209x296mm (300 x 300 DPI)

Figure 2




209x296mm (300 x 300 DPI)

Figure 3



209x296mm (300 x 300 DPI)

Figure 4



209x296mm (300 x 300 DPI)



