



King's Research Portal

DOI:

[10.1017/S1368980019003914](https://doi.org/10.1017/S1368980019003914)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Dikariyanto, V., Berry, S. E., Pot, G. K., Francis, L., Smith, L., & Hall, W. L. (2020). Tree nut snack consumption is associated with better diet quality and CVD risk in the UK adult population: National Diet and Nutrition Survey (NDNS) 2008-2014. *Public Health Nutrition*, 23(17), 3160-3169. <https://doi.org/10.1017/S1368980019003914>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Tree nut snack consumption is associated with better diet quality and CVD risk in the UK adult population: National Diet and Nutrition Survey (NDNS) 2008–2014

Vita Dikariyanto*, Sarah E Berry, Gerda K Pot, Lucy Francis, Leanne Smith and Wendy L Hall*

Department of Nutritional Sciences, School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London, London SE1 9NH, UK

Submitted 13 July 2018: Final revision received 15 August 2019: Accepted 9 September 2019

Abstract

Objectives: To examine associations of tree nut snack (TNS) consumption with diet quality and cardiovascular disease (CVD) risk in UK adults from National Diet and Nutrition Survey (NDNS) 2008–2014.

Design: Cross-sectional analysis using data from 4-d food diaries, blood samples and physical measurements for CVD risk markers. To estimate diet quality, modified Mediterranean Diet Score (MDS) and modified Healthy Diet Score (HDS) were applied. Associations of TNS consumption with diet quality and markers of CVD risk were investigated using survey-adjusted multivariable linear regression adjusted for sex, age, ethnicity, socio-economic and smoking status, region of residency and total energy and alcohol intake.

Setting: UK free-living population.

Subjects: 4738 adults (≥ 19 years).

Results: TNS consumers had higher modified MDS and HDS relative to non-consumers. TNS consumers also had lower BMI, WC, SBP and DBP and higher HDL compared to non-consumers, although a dose-related fully adjusted significant association between increasing nut intake (g per 4184 kJ/1000 kcal energy intake) and lower marker of CVD risk was only observed for SBP. TNS consumption was also associated with higher intake of total fat, mono-, *n*-3 and *n*-6 polyunsaturated fatty acids, fibre, vitamin A, thiamin, folate, vitamin C, vitamin E, potassium, magnesium, phosphorus, selenium and iron; and lower intake of saturated fatty acids, *trans* fatty acids, total carbohydrate, starch, free sugar, sodium and chloride.

Conclusions: TNS consumers report better dietary quality and consumption was associated with lower CVD risk factors. Encouraging replacement of less healthy snacks with TNS should be encouraged as part of general dietary guidelines.

Keywords

Nuts
Cross-sectional analysis
Diet quality
CVD
Nutrients

An average of 2.55 snacks per day are consumed in the UK and Ireland, with over a third of these snacks being confectionary or crisps/popcorn/nuts⁽¹⁾. Nuts are a popular snack as shown by the growing trend for consuming tree nuts over the past 10 years⁽²⁾. North America was the region with the highest production; however, it was Europe that was the largest consumer in the world. Almonds (*Prunusdulcis*), walnuts (*Juglansregia*), pecans (*Caryaillinoensis*), pine nuts (*Pinuspinea*), cashews (*Anacardiumoccidentale*), macadamia nuts (*Macadamia*), hazelnuts (*Corylusavelana*), pistachios (*Pistaciavera*), Brazil nuts (*Bertholletiaexcelsa*) and chestnuts (*Castanea*) are examples of edible tree nuts that are produced commercially⁽²⁾.

Almonds, walnuts, pecans, pine nuts, cashews, macadamia nuts, hazelnuts, pistachios and Brazil nuts differ to some extent in their nutrient profiles. However, tree nuts are generally energy-dense, with a high proportion of fat made up of unsaturated fatty acids; low in sodium; and rich in plant-based protein, dietary fibre, and micronutrients, including niacin, vitamin B₆, vitamin E, vitamin K, folic acid, calcium, magnesium, potassium, selenium, phosphorus and zinc. Tree nuts are also rich in phytosterols and (poly)phenols, which promote antioxidant and anti-inflammatory pathways^(3–5). Because of these properties, tree nuts and health outcomes have been the focus of many human clinical trials and observational studies.

*Corresponding authors: Email vita.dikariyanto@kcl.ac.uk, wendy.hall@kcl.ac.uk



Previous dietary intervention studies showed that tree nut consumption resulted in lowered type-2 diabetes and cardiovascular disease (CVD) risk factors. Walnut, almond, pistachio, macadamia nut, cashew and hazelnut consumption favourably modified blood lipid profile^(6–15), mixed nuts improved insulin sensitivity⁽¹⁴⁾, and walnuts lowered inflammatory markers⁽¹⁶⁾ and improved endothelium-dependent vasodilation⁽¹⁷⁾, all of which would be predicted to reduce the risk of CVD. Furthermore, contrary to popular perception, nut-enriched diets are not linked with increased risk of weight gain⁽¹⁸⁾ and tree nut consumption has been shown to assist weight loss as part of an energy-restricted diet in obese or overweight subjects⁽¹⁹⁾.

Cross-sectional analysis of tree nut consumption and indicators of diet quality and cardiovascular health have also been undertaken. In the US adult population (≥ 19 years), using the National Health and Nutrition Examination Survey (NHANES) 2005–2010 database (n 14 386) based on 24-h dietary recalls, it was reported that tree nut consumption was linked to lower BMI, waist circumference (WC), systolic blood pressure (SBP), and insulin resistance index (HOMA-IR) and higher high-density lipoprotein (HDL-C) adjusted for age, sex, ethnicity, poverty index ratio, physical activity level, smoking status and alcohol intake⁽²⁰⁾. O'Neil *et al.* (2015) also showed that tree nut consumers, compared to non-consumers, had significantly higher diet quality scores (HEI-2005, a diet quality score widely used in the USA) and greater nutrient adequacy for dietary fibre, vitamin A, vitamin E, vitamin C, folate, calcium, iron, magnesium, zinc and potassium⁽²¹⁾.

The purpose of the current study was to examine associations between tree nut snack (TNS) consumption and diet quality, and CVD risk markers, in a nationally representative UK adult population, using data from the UK National Diet and Nutrition Survey (NDNS) rolling programme 2008–2014. Dietary data were derived from estimated 4-d food diaries in a population of 4738 adults (≥ 19 years)^(22,23), which differs from the NHANES analysis 2005–2010, which was based on two multiple pass 24-h dietary recalls in a larger population of 14 386 adults⁽²¹⁾. The hypothesis of the current study was that greater TNS consumption would be associated with higher diet quality, greater nutrient adequacy, and lower prevalence of CVD risk markers in UK adults.

Materials and Methods

The National Diet and Nutrition Survey Rolling Programme (NDNS-RP) and study population

The NDNS-RP is a long-running government-funded scheme to assess diet, nutrient intake and nutritional status of the general population (>1.5 years) living in private households in the UK (England, Scotland, Wales and North Ireland)^(22,23). Random sampling was carried out

on addresses throughout the UK. A single address could have multiple households and a household in an address was selected randomly. An adult in the household was also randomly selected. Selected participants were requested to complete a 4-d estimated food diary, interviewed to collect information, such as dietary habits, socio-demographic background and lifestyle as well as anthropometrically measured and blood sample taken^(22,23).

The survey involves two stages: (i) interview visits to collect information on socio-demography, administer the 4-d food diaries, and carry out anthropometric measurements, and (ii) a nurse visit to do further physical measurements and collect blood and 24-h urine samples^(22,23). Following venepuncture, an EDTA and a serum gel monovette tube from each participant's sample set were sent by post, to the Immunology and Biochemistry Laboratory at Addenbrooke's Hospital in Cambridge for prompt analysis. The remaining samples (lithium heparin, serum or fluoride blood monovette tubes) were processed and stored below -40°C (or at a maximum of -20°C where -40°C facilities were not available), before being transported on dry ice to the Human Nutrition Research (HNR) facility for analysis. The cross-sectional analysis reported here included data from adult participants (≥ 19 years, n 4738), who completed a 4-d estimated food diary in the NDNS-RP 2008–2014 (years 1–6)^(22,23).

CVD risk markers

Body mass index (BMI; kg/m^2), waist circumference (WC; cm), systolic blood pressure (SBP; mmHg), diastolic blood pressure (DBP; mmHg), total cholesterol (TC; mmol/l), triglycerides (TAG; mmol/l), HDL-cholesterol (HDL-C; mmol/l), LDL-cholesterol (LDL-C; mmol/l), TC:HDL-C (the ratio of TC and HDL-C) and C-reactive protein (CRP; mg/l) were CVD risk markers included in the analysis. Interviewer measurement protocols and procedures for blood sample collection, processing, analysis and quality controls are detailed elsewhere^(22,23). Body height and weight were measured using a portable stadiometer and a weight scale, and BMI was calculated by fieldworkers. WC measurement was taken using a tape measure. The discrepancy tolerances of repeat measurement readings were not detailed in the NDNS method protocols. Omron HEM907, an automated validated monitor, was used to measure blood pressure in a sitting position after a five-minute rest. Trained fieldworkers took blood pressure measurements three times and results were presented based on the mean value of second and third readings with one-minute intervals^(22,23).

Diet quality indices

To estimate diet quality, two existing diet scores were used: the Mediterranean Diet Score (MDS)⁽²⁴⁾ and Healthy Diet Score (HDS)⁽²⁵⁾. Maynard *et al.* (2004) developed HDS based on Healthy Diet Indicator (HDI) and the UK



guidelines at that point in time, as recommended by the Committee on Medical Aspects of Food Policy (COMA)⁽²⁵⁾. Modifications were applied to HDS for the current study to reflect UK current recommendations^(22,26–30), and nuts were removed from the MDS scoring system as appropriate for the current study on diet and health associations with nut consumption. The potential top score of the modified MDS remained the same: 9, but the modified HDS had a potential top score of 14 while the original HDS scoring range was 0–12 (see Table A2 in Appendices). Tables A1 and A2 in Appendices show original and modified items of MDS and HDS items, respectively.

Statistical analysis

Prior to statistical analysis, TNS intake was defined and determined. TNS consumption was defined as: (i) any amount of consumption or (ii) ≥ 7.08 g ($\frac{1}{4}$ oz) of TNS. The ≥ 7.08 g ($\frac{1}{4}$ oz) cut-off was adopted to facilitate comparisons with previous cross-sectional analysis of associations between tree nut consumption and dietary scores/nutrient adequacy in a US adult population⁽²¹⁾. Data on TNS consumption were isolated from the database prior to statistical analysis and total TNS intake was calculated. Tree nuts included were almonds, walnuts, pecans, pine nuts, cashews, macadamia nuts, hazelnuts, pistachios, Brazil nuts and chestnuts. Although the US Food and Drug Administration recognises coconuts as a tree nut, they were excluded since they are fruits of palm trees and not commonly consumed whole as a snack food. Peanuts were also excluded since they are classified as legumes.

Statistical analysis was carried out using SPSS IBM 23, and a two-sided *P*-value of 0.05 was considered statistically significant. Data are presented as adjusted means (95 % CI) for individual nutrient intakes, total diet quality scores as well as levels of CVD risk markers, and as medians (with IQRs) for the amount of TNS consumed and age. To examine whether there was a statistically significant association between tree nut consumption and alcohol and total energy intakes as well as demographic variables, that is, age, sex, ethnicity, socio-economic and smoking status and region of residency, survey-adjusted generalised linear model (GLM) with a binary logistic link function was used. Survey-adjusted GLM with a linear link function (predictors: age, sex, ethnicity, socio-economic and smoking status, region of residency, total energy and alcohol intake) was used to examine whether there were significant differences between TNS consumers and non-consumers in their diet quality scores, nutrient intakes and CVD risk markers. To investigate dose–response associations between TNS consumption (g/4184 kJ energy intake) and diet quality and CVD risk markers, survey-adjusted multivariable linear regression models were used adjusting for the same covariates mentioned above. Normal residual distributions were checked by visual inspection of histogram and Q–Q plots; data with non-normally distributed residuals were log

transformed using \log_{10} for analysis of survey-adjusted GLM and multivariable linear regression. The results of analysis were back transformed into the geometric mean values. Homoscedasticity was checked by plotting the standardised residuals of dependent variables and predictors.

During the analysis, the weight factor provided by the NDNS database resource was applied to adjust for non-response and known socio-economic differences in the survey to ensure that the data were nationally representative for the UK population and reducing selection bias and non-response bias^(31,32). The weight factor used is *wti_Y14* (weight for individual and diary-all ages, combined Y1–4) and *wti_Y56* (weight for individual and diary-all ages, combined Y5–6) for investigating differences in diet quality scores and nutrient intakes between TNS consumers and non-consumers, associations between tree nut consumption and demographic variables and multivariable linear regression including diet quality scores. Weight factors *wtn_Y14* (weight for nurse-all ages, combined Y1–4) and *wtn_Y56* (weight for nurse-all ages, combined Y5–6) were used for GLM and multivariable linear regression including variables BMI, WC and blood pressure; and *wtb_Y14* (weight for blood-all ages, combined Y1–4) and *wtb_Y56* (weight for blood-all ages, combined Y5–6) were used for GLM and multivariable linear regression for blood analyte variables including CRP and lipids^(31,32).

Results

Demographic information

Table 1 shows background characteristics of TNS consumers and non-consumers. Median TNS-A (any amount of TNS intake) consumption (*n* 484) contributed 0.8 % of total energy intake while median consumption in the TNS-B group (including individuals who consumed ≥ 7.08 g TNS per day, equivalent to $\frac{1}{4}$ oz, *n* 224) was 2.3 % of total energy intake. On average, TNS consumers were significantly older than non-consumers and were more likely to be female and non-smokers. TNS-A consumption was significantly associated with the demographic factors included, such as sex, ethnicity, socio-economic status, smoking status and region of residency. TNS-B consumption was also significantly associated with these demographic variables, except region of residency.

Diet quality scores

Geometric estimated marginal mean total scores of modified MDS were significantly higher in TNS-A consumers (5.9; 95 % CI 5.2, 6.6) compared with non-consumers (4.9; 95 % CI 4.4, 5.4; *P* < 0.001). Similarly, geometric estimated marginal mean total scores for the modified HDS were significantly higher in TNS-A consumers (6.1; 95 % CI 5.5, 6.8) compared with non-consumers (5.4; 95 % CI 4.9, 6.0; *P* < 0.001). Results for TNS-B consumers were

Table 1 Background characteristics of tree nut snack (TNS) consumers compared to non-consumers in the UK adult population (≥ 19 years) based on NDNS 2008–2014, *n* 4738

		TNS-A			TNS-B		
		Consumers, <i>n</i> 484	Non-consumers, <i>n</i> 4254	<i>P</i> -value	Consumers, <i>n</i> 224	Non-consumers, <i>n</i> 4514	<i>P</i> -value
Amount of tree nuts consumed	Gram						
	Median	6.5			14.0		
	IQR	10.8			10.6		
	% Total energy intake						
	Median	0.8			2.3		
	IQR	2.2			5.1		
Age				<0.001*			<0.001*
	Median	51	48		53	48	
	IQR	24	27		24	27	
Sex (%)	Male	31.1	41.6	<0.001*	32.8	40.8	<0.001*
	Female	68.9	58.4		67.2	59.2	
Ethnicity (%)	White	87.8	93.8	0.003*	88.9	93.4	0.016*
	Mixed ethnic group	1.7	0.9		0.9	1.0	
	Black or Black British	1.6	2.0		1.7	1.9	
	Asian or Asian British	5.9	2.2		5.4	2.4	
	Any other group	3.1	1.2		3.0	1.3	
Region (%)	England	68.1	54.8	0.003*	64.8	55.8	0.131
	Scotland	11.9	17.9		11.9	17.5	
	Wales	12.5	14.8		16.7	14.4	
	Northern Ireland	7.5	12.6		6.6	12.3	
Socio-economic status (%)	Higher managerial and professional occupations	27.2	13.7	<0.001*	25.9	14.6	<0.001*
	Lower managerial and professional occupations	31.2	23.1		26.4	23.9	
	Intermediate occupations	8.2	10.5		9.7	10.3	
	Small employers and own account workers	11.0	10.4		11.1	10.4	
	Lower supervisory and technical occupations	6.5	9.5		7.0	9.3	
	Semi-routine occupations	9.4	15.1		13.2	14.6	
	Routine occupations	3.4	12.9		4.5	12.2	
	Never worked	1.3	3.1		0.7	3.0	
	Others	1.9	1.7		1.5	1.8	
Smoking status (%)	Current smoker	11.7	25.0	<0.001*	12.5	24.2	<0.001*
	Ex-Regular smoker	25.7	23.6		27.3	23.7	
	Never regular smoker	62.6	51.4		60.2	52.2	
Alcohol intake (g/d)				0.012*			0.002*
	Median	6.0	0.7		4.4	1.8	
	IQR	18.4	16.7		17.0	16.9	
Energy intake (kJ/d)				<0.001*			<0.001*
	Unadjusted mean	7950.9	7325.3		8168.8	7350.5	
	SD	2023.8	2364.4		2087.0	2344.7	

This is a descriptive table. Survey-adjusted generalised linear model with a linear binary logistic function was used to investigate the association between TNS consumption and demographic variables.

**P* was <0.05 indicating a significant association.

almost identical (data not shown). To investigate dose–response associations between every gram increase in TNS consumption per 4184 kJ of adult’s energy intake and diet quality scores, the survey-adjusted regression model was adjusted for age, sex, ethnicity, socio-economic and smoking status, alcohol and energy intakes. There was no dose response observed in the scores of modified MDS and modified HDS ($P = 0.726$ and $P = 0.971$, respectively).

Nutrient intake

TNS consumers had significantly higher total energy, food energy, fat, *cis*-monounsaturated fatty acids, *cis* *n*-6 fatty acids, *cis* *n*-3 fatty acids (TNS-A only), intrinsic milk sugars and fibre intakes, as shown in Table 2. Saturated fatty

acids, *trans*-fatty acids, total carbohydrate, starch, non-milk extrinsic sugars, intrinsic milk sugar and starch and alcohol (TNS-B only) intakes were significantly lower in TNS consumers. For micronutrients, as shown in Table 2, fully adjusted analysis revealed that TNS consumers, relative to non-consumers, had significantly higher intakes of vitamin A (TNS-A only), vitamin E, thiamin, riboflavin (TNS-B only), folate, pantothenic acid, biotin, vitamin C, potassium, magnesium, phosphorus, iron, copper, zinc, manganese and selenium and lower intakes of sodium and chloride. However, there were no differences between groups for vitamins D, riboflavin (TNS-A only), niacin equivalents, vitamin B₆, vitamin B₁₂, calcium and iodine.

Table 2 Energy, macro- and micronutrient intake of tree nut snack consumers defined by any amount of daily consumption or ≥ 7.08 gram consumption per day, in the UK adult population (≥ 19 years) based on National Diet and Nutrition Survey 2008–2014, *n* 4738

Nutrients	TNS-A					TNS-B				
	Consumers		Non-consumers			Consumers		Non-consumers		
	<i>n</i> 484		<i>n</i> 4254			<i>n</i> 224		<i>n</i> 4514		
	Estimated marginal mean	95 % CI	Estimated marginal mean	95 % CI	<i>P</i> -value	Estimated marginal mean	95 % CI	Estimated marginal mean	95 % CI	<i>P</i> -value
Macronutrients (diet only, % food energy)*										
Total energy (kJ)	7365.1†	6335.8, 8393.9	6548.0	5532.5, 7563.0	<0.001	7628.3†	6581.4, 8675.5	6782.7	5771.8, 7793.5	<0.001
Food energy (kJ)	7127.0†	6155.9, 8098.6	6373.9	5415.4, 7332.0	<0.001	7417.8†	6430.0, 8405.2	6580.2	5626.6, 7533.3	<0.001
Protein	17.4	15.5, 19.2	17.3	15.5, 19.2	0.827	17.5	15.7, 19.4	17.3	15.5, 19.1	0.384
Fat	37.1†	34.2, 40.0	35.1	32.2, 38.0	<0.001	37.6†	34.7, 40.5	35.2	32.3, 38.0	<0.001
Saturated fatty acids	12.2†	10.6, 13.7	12.5	11.0, 14.0	0.035	11.8†	10.2, 13.3	12.3	10.8, 13.9	0.008
<i>cis</i> -Monounsaturated fatty acids	14.3†	13.0, 15.6	13.0	11.8, 14.3	<0.001	15.0†	13.6, 16.3	13.1	11.8, 14.4	<0.001
<i>cis</i> <i>n</i> -6 fatty acids	6.2†	5.5, 6.9	5.3	4.6, 6.0	<0.001	6.6 †	5.8, 7.3	5.4	4.7, 6.1	<0.001
<i>cis</i> <i>n</i> -3 fatty acids‡	1.1†	0.9, 1.3	1.0	0.8, 1.2	<0.001	1.0	0.8, 1.1	1.0	0.9, 1.1	0.469
<i>Trans</i> fatty acids	0.5†	0.4, 0.7	0.6	0.5, 0.7	<0.001	0.5†	0.3, 0.6	0.6	0.4, 0.7	<0.001
Carbohydrate	45.6†	42.4, 48.8	47.6	44.4, 50.8	<0.001	44.9†	41.7, 48.2	47.6	44.4, 50.7	<0.001
Total sugars	17.7	14.5, 20.9	17.2	14.1, 20.4	0.139	18.1	14.9, 21.4	17.8	14.6, 20.9	0.399
Starch	27.9†	25.1, 30.6	30.3	27.6, 33.0	<0.001	26.7†	24.0, 29.5	29.8	27.1, 32.4	<0.001
Non-milk extrinsic sugars	7.7†	4.7, 10.7	8.7	5.8, 11.7	0.001	7.5†	4.4, 10.5	9.0	6.1, 11.9	<0.001
Intrinsic milk sugars and starch	37.8†	34.8, 40.9	38.8	35.8, 41.9	0.001	34.3†	32.5, 36.2	35.9	34.4, 37.6	0.003
Intrinsic milk sugars‡	10.0†	8.2, 11.8	8.5	6.8, 10.3	<0.001	9.9†	8.1, 12.1	7.9	6.5, 9.6	<0.001
Non-starch polysaccharides (Englyst Fibre, g)	15.1†	13.2, 16.9	13.3	11.4, 15.1	<0.001	15.8†	13.9, 17.7	13.7	11.9, 15.5	<0.001
Alcohol (g)‡§	12.1	8.8, 16.5	12.9	9.6, 17.5	0.290	10.0†	7.1, 14.0	12.7	9.6, 16.9	0.013
Micronutrients**										
Vitamin A (retinol equivalents) (µg)‡§	904.9†	753.4, 1086.9	828.9	696.5, 986.5	0.024	966.1	794.0, 1175.2	885.5	750.9, 1044.2	0.128
Vitamin D (µg)‡§	2.5	2.1, 3.0	2.4	2.0, 2.8	0.213	2.4	2.0, 2.9	2.4	2.0, 2.8	0.912
Vitamin E (mg)‡	11.6†	10.1, 13.0	9.8	8.4, 11.2	<0.001	11.6†	10.6, 12.7	9.3	8.6, 10.1	<0.001
Thiamin (mg)	1.4†	1.2, 1.6	1.3	1.1, 1.5	0.001	1.5†	1.3, 1.7	1.3	1.1, 1.5	<0.001
Riboflavin (mg)	1.5	1.2, 1.7	1.4	1.2, 1.7	0.196	1.5†	1.3, 1.8	1.4	1.2, 1.7	0.003
Niacin equivalent (mg)	33.5	28.7, 38.3	33.8	29.1, 38.6	0.486	34.4	29.5, 39.3	33.8	29.1, 38.5	0.382
Vitamin B ₆ (mg)‡	1.9	1.5, 2.3	1.9	1.5, 2.3	0.992	1.9	1.7, 2.1	1.9	1.8, 2.1	0.530
Vitamin B ₁₂ (µg)‡§	4.6	3.6, 6.0	4.7	3.7, 6.0	0.777	4.5	3.9, 5.2	4.8	4.3, 5.5	0.154

Table 2 Continued

Nutrients	TNS-A					TNS-B				
	Consumers		Non-consumers		P-value	Consumers		Non-consumers		P-value
	n 484		n 4254			n 224		n 4514		
	Estimated marginal mean	95 % CI	Estimated marginal mean	95 % CI		Estimated marginal mean	95 % CI	Estimated marginal mean	95 % CI	
Folate (µg)	242.3†	201.9, 282.6	229.4	189.5, 269.2	0.001	255.7†	214.7, 296.6	237.0	197.5, 276.6	0.001
Pantothenic acid (mg)	5.6	4.6, 6.5	5.3	4.4, 6.3	0.007	5.9†	4.9, 6.8	5.3	4.4, 6.2	<0.001
Biotin (µg)	36.4†	30.1, 42.7	30.0	23.8, 36.3	<0.001	41.3†	34.9, 47.6	31.7	25.5, 37.9	<0.001
Vitamin C (mg)‡§	79.1†	58.4, 107.3	62.1	46.0, 83.8	<0.001	90.4†	75.3, 108.6	79.0	67.8, 92.3	0.012
Sodium (mg)	1854.8†	1596.0, 2113.6	2053.9	1798.5, 2309.3	<0.001	1732.2†	1469.6, 1994.8	2006.2	1752.6, 2259.8	<0.001
Potassium (mg)	2866.7†	2595.1, 3138.2	2645.3	2377.3, 2913.3	<0.001	3021.3†	2746.0, 3296.6	2694.8	2428.8, 2960.5	<0.001
Calcium (mg)	696.6	591.3, 801.8	702.8	599.0, 806.7	0.541	717.4	610.8, 824.0	709.6	606.7, 812.6	0.599
Magnesium (mg)	276.9†	251.0, 302.8	237.8	212.3, 263.4	<0.001	301.5†	275.3, 327.8	245.3	219.9, 270.6	<0.001
Phosphorus (mg)	1165.8†	1056.1, 1275.6	1125.4	1017.1, 1233.7	<0.001	1191.5†	1080.3, 1302.6	1126.2	1018.9, 1233.5	<0.001
Iron (mg)	11.1†	9.9, 12.4	10.4	9.1, 11.6	<0.001	11.2†	9.9, 12.4	10.5	9.3, 11.7	<0.001
Copper (mg)‡§	1.2†	1.0, 1.4	1.0	0.9, 1.2	<0.001	1.4†	1.3, 1.6	1.2	1.1, 1.3	<0.001
Zinc (mg)	9.0†	7.9, 10.0	8.8	7.7, 9.8	0.044	9.3†	8.2, 10.3	8.8	7.8, 9.8	0.002
Chloride (mg)	3017.1†	2631.3, 3402.9	3286.7	2906.0, 3667.5	<0.001	2877.8†	2486.4, 3269.1	3242.2	2864.3, 3620.1	<0.001
Manganese (mg)	3.3†	2.8, 3.8	2.8	2.3, 3.3	<0.001	3.5†	3.0, 4.0	2.9	2.4, 3.4	<0.001
Iodine (µg)	161.9	131.1, 192.7	158.7	128.3, 189.1	0.282	158.2	127.0, 189.4	158.6	128.5, 188.8	0.914
Selenium (µg)‡	56.9†	48.3, 65.4	51.9	43.4, 60.3	<0.001	54.9†	49.8, 60.5	50.5	46.6, 54.8	0.004

The actual sample size in the computation for vitamin A and vitamin D, for TNS-A consumers was 314 and for TNS-A non-consumers was 2172, whereas for TNS-B consumers was 138 and for TNS-B non-consumers was 2348. The actual sample size in the computation for alcohol, *cis-n3* fatty acids, intrinsic milk sugars and starch, vitamin E, vitamin B₆, vitamin B₁₂, vitamin C, copper and selenium for TNS-B consumers was 138 and for TNS-B non-consumers was 2348. There were no missing values in the computation for other nutrients as outcomes.

*Survey-adjusted generalised linear model (GLM) with a linear link function and predictors such as age, sex, ethnicity, region of residency, socio-economic and smoking status was used for energy intake as an outcome for TNS-A; survey-adjusted GLM with a linear link function and predictors such as age, sex, ethnicity, region of residency, socio-economic and smoking status, alcohol and energy intakes was used for other macronutrient intake outcomes for TNS-A; survey-adjusted GLM with a linear link function and predictors such as age, sex, ethnicity, region of residency, socio-economic and smoking status, and energy intake was used for alcohol intake as an outcome for TNS-A. The same statistical analysis was conducted for TNS-B, but region of residency was excluded from predictors.

†P < 0.05 showed a significant difference.

‡Geometric marginal means were presented due to non-normally distributed residual data in TNS-B population.

§Geometric marginal means were presented due to non-normally distributed residual data in TNS-A population.

**Survey-adjusted GLM with a linear link function and predictors: age, sex, ethnicity, region of residency, socio-economic and smoking status, alcohol and energy intakes was used for TNS-A. The same statistical analysis was conducted for TNS-B but region of residency was excluded from predictors.



CVD risk markers

Blood samples were not available from all participants, and anthropometric and blood pressure data were also missing. Associations between TNS consumption and CVD risk markers were analysed for the remaining participants. The estimated marginal mean (95 % CI) values of CVD risk markers are shown in Table 3. For TNS-A consumers, BMI, WC, SBP and DBP were significantly lower, and HDL was significantly higher compared to non-consumers. For those consuming >7.08 g TNS/d (TNS-B), only WC, SBP and DBP were significantly lower compared to non-consumers (data shown in online supplementary material). Survey-adjusted regression analysis showed that for every gram increase in TNS consumption per 4184 kJ of adults' energy intake (Table 3), SBP was significantly lower demonstrating a dose-response relationship ($P = 0.028$).

Discussion

Interventional and observational evidence suggests that replacing refined carbohydrate-based snacks with tree nut snacks may improve blood lipid profiles, management of body weight^(33,34) and nutrient intakes. However, TNS intakes in the general UK population have not been fully investigated. Previous studies have been conducted in the NHANES US adult population^(20,21) using multiple 24-h dietary recalls to collect food intake data. This cross-sectional analysis using a representative UK adult population revealed that just 10 % of respondents reported consuming any amount of TNS during their 4-d food intake recording period, just less than 5 % reported consuming more than 7.08 g (¼ oz) per day on average (around a handful over the 4-d period), and only 0.34 % reported consuming the US Food and Drug Administration recommendation of 42.5 g per day⁽³⁵⁾. The relatively small sub-population of TNS consumers was more likely to be female, white, older and living in England and less likely to be current smokers relative to non-consumers.

Increments in TNS consumption (g per 4184 kJ of energy intake) were not associated with significantly greater modified MDS and HDS in consumers. This lack of dose-response relationship could be due to the low consumption of TNS in the population (for TNS-A consumers, median 0.8 % of total energy intake and 6.5 g/d in terms of total weight intake; for TNS-B consumers, 2.3 % of total energy intake, and 14.0 g/d in terms of total weight intake). TNS consumption status may be an indicator of improved overall diet quality, but the actual amount consumed has very little practical impact.

Since TNS consumption status appears to act as a marker of healthy dietary patterns, it is not surprising that the overall nutrient intake profile of TNS consumers was more favourable compared to non-consumers. The contribution of non-milk extrinsic sugar intakes to energy was only marginally lower in TNS consumers (a difference of

Table 3 CVD risk marker values in UK adults (≥ 19 years) based on National Diet and Nutrition Survey 2008–2014, in consumers of any amount of tree nut snack (TNS-A) and non-consumers, and the association of tree nut snack consumption and risk markers

CVD risk marker	Consumers n 484			Non-consumers n 4254			Associations between tree nut consumption and CVD risk markers†			
	Estimated marginal mean	95 % CI	Value‡	Estimated marginal mean	95 % CI	Value‡	β	95 % CI	P-value	R ²
BMI (kg/m ²)§,	25.4	24.0, 26.8	26.3	25.0, 27.8	0.002*	0.002*	1.035	0.991, 1.081	0.128	0.121
WC (cm)¶	91.5	88.5, 94.6	94.2	91.5, 97.0	<0.001*	<0.001*	0.094	-0.080, 0.268	0.289	0.293
SBP (mmHg)††	119.7	116.2, 123.2	124.0	120.8, 127.1	<0.001*	<0.001*	-0.242	-0.458, -0.026	0.028*	0.286
DBP (mmHg)††	69.2	66.8, 71.7	72.0	69.7, 74.2	<0.001*	<0.001*	-0.034	-0.196, 0.127	0.677	0.033
TC (mmol/l)‡‡	4.9	4.5, 5.3	4.9	4.5, 5.3	0.627	0.627	0.011	-0.007, 0.029	0.218	0.109
TAG (mmol/l)§,§§	1.1	0.9, 1.3	1.1	0.9, 1.4	0.220	0.220	0.972	0.813, 1.164	0.757	0.084
HDL-C (mmol/l)‡‡	1.5	1.4, 1.7	1.4	1.3, 1.6	0.008*	0.008*	-0.001	-0.009, 0.007	0.754	0.277
LDL-C (mmol/l)¶¶	2.9	2.6, 3.3	2.9	2.6, 3.2	0.980	0.980	0.011	-0.006, 0.028	0.204	0.046
TC:HDL-C‡‡	3.5	3.1, 4.0	3.6	3.2, 4.0	0.412	0.412	0.016	-0.005, 0.037	0.143	0.163
CRP (mg/l)§,¶¶	1.9	1.3, 2.6	2.1	1.5, 2.9	0.062	0.062	1.194	0.933, 1.528	0.157	0.095

BMI, body mass index; WC, waist circumference; SBP, systolic blood pressure; DBP, diastolic blood pressure; TC, total cholesterol; TAG, triglycerides; HDL-C, high-density lipoprotein cholesterol; LDL-C, low-density lipoprotein cholesterol; CRP, C-reactive protein

†Survey-adjusted generalised linear model with a linear link function and predictors, such as age, sex, ethnicity, region of residency, socio-economic and smoking status, alcohol and energy intakes was used.

‡Survey-adjusted multivariable linear regression was used, adjusted for age, sex, ethnicity, region of residency, socio-economic and smoking status and alcohol intake.

§Geometric marginal means and geometric β values were presented due to non-normally distributed residual data. Geometric β values were interpreted as ratios of geometric means.

¶Due to missing data, sample sizes were as follows: tree nut snack-A consumers || 241, ¶ 384, †† 326, ‡‡ 274, §§¶¶ 176 and ¶¶¶ 176; non-consumers || 176, ¶ 310, †† 273, ‡‡ 213, §§ 161, |||| 2,096 and ¶¶¶ 1,164.

* $P < 0.05$ showed a significant difference.



1.0% of energy). Fibre (non-starch polysaccharides) intakes were 1.8 g higher in TNS consumers compared with non-consumers, but TNS intake is unlikely to contribute more than a third of this difference, with the remainder due to greater intakes of other fibre-rich foods.

The observations reported here and in the US population imply that TNS are usually eaten as a part of an overall healthier dietary pattern in industrialised countries⁽²¹⁾, which would be predicted to translate to better cardiovascular health outcomes. UK TNS consumers had significantly reduced BMI, WC, SBP and DBP and significantly higher HDL-C, but the slightly lower mean CRP in TNS consumers did not reach statistical significance compared with non-consumers. O'Neil *et al.* (2015) previously reported that ≥ 7.08 g tree nut consumption was associated with lower BMI and WC, as well as SBP and higher HDL-C, in the US adult population adjusted for the same covariates as used in the present analysis, plus physical activity level⁽²⁰⁾. Overall feeding trials have reported that higher tree nut consumption did not result in weight gain^(18,19), which may be related to their satiating/satiety-inducing properties⁽³⁶⁾, as well as limited lipid bioaccessibility⁽³⁷⁾. Since TNS consumers' median intake was low in the UK, observed differences in BMI and WC could be related to confounding factors such as physical activity levels, which was not considered in the present analysis due to lack of available data. Mean SBP was 4.3 mmHg lower and mean DBP was 2.8 mmHg lower in TNS consumers compared to non-consumers, a clinically meaningful difference that would be predicted to reduce risk of CVD. The SUN prospective cohort study reported that there was no association between tree nut consumption and blood pressure; the potential reasons could be an underestimated amount of nut consumption, no assessment on the change in nut consumption during follow-up, and no specific information on preparation method, for example, salted, roasted or raw⁽³⁸⁾. The Physician's Health Study observed blood pressure reduction only in lean volunteers⁽³⁹⁾. A recent meta-analysis of 21 randomised control trials reported that total nut consumption lowered SBP in participants without type 2 diabetes, and mixed nuts also lowered DBP⁽⁴⁰⁾. Although plasma CRP concentrations were not significantly different in the NDNS cohort, a cross-sectional study using data from the Nurse's Health Study and Health Professional Follow-Up Study revealed that consumers eating tree nuts ≥ 5 times weekly based on FFQ had significantly lower CRP⁽⁴¹⁾, suggesting larger differences in intake may be required to impact on systemic inflammatory markers. However, a meta-analysis of 20 randomised controlled trials suggested that tree nut consumption did not reduce CRP⁽⁴²⁾. In the current study, the amount of nuts consumed by consumers in the current UK cohort was low, and therefore, the SBP and DBP differences observed are likely to be the sum effect of an overall healthier dietary pattern including TNS⁽⁴³⁻⁴⁶⁾.

A significant difference was observed in HDL-C between TNS consumers and non-consumers. Cross-sectional analysis

in the US adult population also reported higher HDL-C in TNS consumers⁽²⁰⁾. There were no significant differences observed in other blood lipids. A recent meta-analysis of 61 interventional clinical trials revealed that tree nut intake reduced TC, TAG and LDL-C, and it was reported that the dosage of tree nut intake determined cholesterol lowering capacity rather than the nut types⁽⁴⁷⁾. A pooled analysis of 25 feeding trials conducted in seven countries demonstrated the reduction of TC, LDL-C and the ratio of TC to HDL-C but failed to report the increase of HDL-C in response to tree nut intake⁽⁴⁸⁾. These inconsistent associations of tree nut consumption and blood pressure, CRP and blood lipids between cross-sectional analysis and clinical trials could be due to different dosage and duration of consumption (duration of the study), residual confounding effects, characteristics such as baseline lipid profile, as well as study sample size relating to statistical power⁽⁴¹⁾.

Strengths of the current study include using a relatively large, nationally representative UK population, and the close agreement with results reported in a nationally representative US population suggests that findings may be generalisable to other industrialised countries with similar dietary profiles. The availability of estimated portion size food diary data over a 4-d period is considered to be one of the more accurate dietary assessment methods in large populations, although underreporting of energy intake is a well-known problem with this methodology that limits the conclusions that can be drawn. Furthermore, the use of 4-d estimated food diaries means that significant nut intakes on other days may have been missed and a significant proportion of TNS consumers may have been wrongly classified as non-consumers; analysis based on frequency of tree nut consumption was not possible. Available information on physical activity was incomplete so statistical analysis models could not be adjusted for this potentially confounding factor. Different types of tree nuts have differing nutrient profiles and potentially nutrient bioaccessibility, and therefore it may be misleading to group them altogether in terms of associations with CVD risk factors. In addition to that, missing data for CVD risk factors resulted in lower sample sizes.

In conclusion, the prevalence of TNS consumers in the UK adult population is estimated to be approximately 10%, and median intakes were low in the group classified as TNS consumers. Tree nut snack consumption was associated with higher diet quality scores and a more favourable nutrient intake profile. Tree nut snack consumption may be a marker of a healthy dietary pattern and is associated with lower adiposity and blood pressure. It is recommended that tree nuts should replace high refined carbohydrate-based snacks as part of a healthy diet. To determine the relative contribution of tree nuts to the sum impact of a healthier dietary pattern on risk of CVD, future randomised controlled trials should investigate the effect of replacing usual refined carbohydrate snacks with tree nuts on markers of cardiometabolic disease risk.





Acknowledgements

Acknowledgements: We thank the two anonymous reviewers for their valuable insights. **Financial support:** This work was supported in part by the Almond Board of California (grant number 15-BERRYS-01) and the Indonesia Endowment Fund for Education. The funders had no role in the design, analysis or writing of this article. **Conflict of interest:** None. **Authorship:** The authors' responsibilities were as follows: V.D., W.L.H., S.E.B. and G.K.P. designed the research; V.D. analysed the data; L.F. advised on analysis of the data; V.D. and W.L.H. wrote the manuscript; W.L.H. had primary responsibility for final content. All authors read, commented and approved the final manuscript. **Ethics of human subject participation:** The current study was conducted according to the guidelines laid down in the Declaration of Helsinki, and all procedures involving human subjects were approved by the National Health Service Local Research Ethics Committee covering each of the postal sectors. Written informed consent was obtained from all participants.

Supplementary material

For supplementary material accompanying this paper visit <https://doi.org/10.1017/S1368980019003914>

References

- Bord Bia – The Irish Food Board (2018) Healthy snacking UK and Ireland. <https://www.bordbia.ie/globalassets/bordbia.ie/industry/marketing-reports/consumer-reports/healthy-snacking-uk-and-ireland-january-2018.pdf> (accessed June 2019).
- International Nut & Dried Fruit Council (2016) Nuts & dried fruits: Statistical yearbook 2016/2017. <https://www.nutfruit.org/consumers/news/detail/statistical-yearbook-2016-17> (accessed September 2017).
- U.S. Department of Agriculture. Agriculture Research Service. Food Data Central (2019) Nuts, almonds. <https://fdc.nal.usda.gov/fdc-app.html#/food-details/170567/nutrients> (accessed June 2019).
- Chen CY & Blumberg JB (2008) Phytochemical composition of nuts. *Asia Pac J Clin Nutr* **17**, Suppl 1, 329–332.
- Maguire LS, O'Sullivan SM, Galvin K *et al.* (2004) Fatty acid profile, tocopherol, squalene and phytosterol content of walnuts, almonds, peanuts, hazelnuts and the macadamia nut. *Int J Food Sci Nutr* **55**, 171–178.
- Rajaram S, Haddad EH, Mejia A *et al.* (2009) Walnuts and fatty fish influence different serum lipid fractions in normal to mildly hyperlipidemic individuals: a randomized controlled study. *Am J Clin Nutr* **89**, 1657s–1663s.
- Torabian S, Haddad E, Cordero-MacIntyre Z *et al.* (2010) Long-term walnut supplementation without dietary advice induces favorable serum lipid changes in free-living individuals. *Eur J Clin Nutr* **64**, 274–279.
- Tapsell LC, Batterham MJ, Teuss G *et al.* (2009) Long-term effects of increased dietary polyunsaturated fat from walnuts on metabolic parameters in type II diabetes. *Eur J Clin Nutr* **63**, 1008–1015.
- Jenkins DJ, Kendall CW, Marchie A *et al.* (2008) Almonds reduce biomarkers of lipid peroxidation in older hyperlipidemic subjects. *J Nutr* **138**, 908–913.
- Gebauer SK, West SG, Kay CD *et al.* (2008) Effects of pistachios on cardiovascular disease risk factors and potential mechanisms of action: a dose-response study. *Am J Clin Nutr* **88**, 651–659.
- Griel AE, Cao Y, Bagshaw DD *et al.* (2008) A macadamia nut-rich diet reduces total and LDL-cholesterol in mildly hypercholesterolemic men and women. *J Nutr* **138**, 761–767.
- Mercanligil SM, Arslan P, Alasalvar C *et al.* (2007) Effects of hazelnut-enriched diet on plasma cholesterol and lipoprotein profiles in hypercholesterolemic adult men. *Eur J Clin Nutr* **61**, 212–220.
- Mukuddem-Petersen J, Stonehouse-Oosthuizen W, Jerling JC *et al.* (2007) Effects of a high walnut and high cashew nut diet on selected markers of the metabolic syndrome: a controlled feeding trial. *Br J Nutr* **97**, 1144–1153.
- Casas-Agustench P, Lopez-Uriarte P, Bullo M *et al.* (2011) Effects of one serving of mixed nuts on serum lipids, insulin resistance and inflammatory markers in patients with the metabolic syndrome. *Nutr Metab Cardiovasc Dis* **21**, 126–135.
- Ros E (2010) Health benefits of nut consumption. *Nutrients* **2**, 652–682.
- Jimenez-Gomez Y, Lopez-Miranda J, Blanco-Colio LM *et al.* (2009) Olive oil and walnut breakfasts reduce the postprandial inflammatory response in mononuclear cells compared with a butter breakfast in healthy men. *Atherosclerosis* **204**, e70–e76.
- Ros E, Nunez I, Perez-Heras A *et al.* (2004) A walnut diet improves endothelial function in hypercholesterolemic subjects: A randomized crossover trial. *Circulation* **109**, 1609–1614.
- Flores-Mateo G, Rojas-Rueda D, Basora J *et al.* (2013) Nut intake and adiposity: Meta-analysis of clinical trials. *Am J Clin Nutr* **97**, 1346–1355.
- Wien MA, Sabate JM, Ikle DN *et al.* (2003) Almonds vs complex carbohydrates in a weight reduction program. *Int J Obes Relat Metab Disord* **27**, 1365–1372.
- O'Neil CE, Fulgoni VL 3rd & Nicklas TA (2015) Tree Nut consumption is associated with better adiposity measures and cardiovascular and metabolic syndrome health risk factors in U.S. Adults: NHANES 2005–2010. *Nutr J* **14**, 64.
- O'Neil CE, Nicklas TA & Fulgoni VL 3rd (2015) Tree nut consumption is associated with better nutrient adequacy and diet quality in adults: National Health and Nutrition Examination Survey 2005–2010. *Nutrients* **7**, 595–607.
- Bates B, Lennox A, Prentice A *et al.* (2014) *National Diet and Nutrition Survey Results from Years 1, 2, 3 and 4 (combined) of the Rolling Programme (2008/2009–2011/2012)*. London: Public Health England.
- Bates B, Cox L, Nicholson S *et al.* (2016) *National Diet and Nutrition Survey Results from Years 5 and 6 (combined) of the Rolling Programme (2012/2013–2013/2014)*. London: Public Health England.
- Couto E, Boffetta P, Lagiou P *et al.* (2011) Mediterranean dietary pattern and cancer risk in the EPIC cohort. *Br J Cancer* **104**, 1493–1499.
- Maynard M, Ness AR, Abraham L *et al.* (2005) Selecting a healthy diet score: Lessons from a study of diet and health in early old age (the Boyd Orr cohort). *Public Health Nutr* **8**, 321–326.
- Nutrition Science Team. Public Health England (2016) *Government Recommendations for Energy and Nutrients for Males and Females Aged 1–18 years and 19+ years*. London: Department of Health.
- DH Cardiovascular Disease Team (2013) *Cardiovascular Disease Outcomes Strategy: Improving Outcomes for People with or at Risk of Cardiovascular Disease*. London: Department of Health.



28. Public Health England (2016) *The Eatwell Guide*. London, UK: Department of Health.
29. Scientific Advisory Committee on Nutrition (2018) *Draft Report: Saturated Fats and Health*. London: Department of Health.
30. Scientific Advisory Committee on Nutrition (2015) *Carbohydrates and Health*. London: The Stationery Office.
31. NatCen Social Research, The Medical Research Council Elsie Widdowson Laboratory (MRC EWL) & University College London (UCL) (2008) *National Diet and Nutrition Survey Years 1-4 2008/09-2011/12: User Guide for UK Data (core & country boost data)*. London: Public Health England.
32. NatCen Social Research & The Medical Research Council Elsie Widdowson Laboratory (MRC EWL) (2012) *National Diet and Nutrition Survey Years 5-6 2012/13-2013/14: User Guide for UK Data*. London: Public Health England.
33. Li Z, Song R, Nguyen C *et al.* (2010) Pistachio nuts reduce triglycerides and body weight by comparison to refined carbohydrate snack in obese subjects on a 12-week weight loss program. *J Am Coll Nutr* **29**, 198–203.
34. Rehm CD & Drewnowski A (2017) Replacing American snacks with tree nuts increases consumption of key nutrients among US children and adults: Results of an NHANES modeling study. *Nutr J* **16**, 17.
35. U. S. Food & Drug Administration (2003) Qualified health claims: Letter of enforcement discretion – nuts and coronary heart disease (Docket No 02P-0505). <http://wayback.archive-it.org/7993/20171114183724/https://www.fda.gov/Food/IngredientsPackagingLabeling/LabelingNutrition/ucm072926.htm> (accessed September 2017).
36. Mattes RD & Dreher ML (2010) Nuts and healthy body weight maintenance mechanisms. *Asia Pac J Clin Nutr* **19**, 137–141.
37. Mandalari G, Grundy MM, Grassby T *et al.* (2014) The effects of processing and mastication on almond lipid bioaccessibility using novel methods of in vitro digestion modelling and micro-structural analysis. *Br J Nutr* **112**, 1521–1529.
38. Martinez-Lapiscina EH, Pimenta AM, Beunza JJ *et al.* (2010) Nut consumption and incidence of hypertension: the SUN prospective cohort. *Nutr Metab Cardiovasc Dis* **20**, 359–365.
39. Djousse L, Rudich T & Gaziano JM (2009) Nut consumption and risk of hypertension in US male physicians. *Clin Nutr* **28**, 10–14.
40. Mohammadifard N, Salehi-Abargouei A, Salas-Salvado J *et al.* (2015) The effect of tree nut, peanut, and soy nut consumption on blood pressure: a systematic review and meta-analysis of randomized controlled clinical trials. *Am J Clin Nutr* **101**, 966–982.
41. Yu Z, Malik VS, Keum N *et al.* (2016) Associations between nut consumption and inflammatory biomarkers. *Am J Clin Nutr* **104**, 722–728.
42. Mazidi M, Rezaie P, Ferns GA *et al.* (2016) Impact of different types of tree nut, peanut, and soy nut consumption on serum C-reactive protein (CRP): a systematic review and meta-analysis of randomized controlled clinical trials. *Medicine* **95**, e5165.
43. Kahleova H, Levin S & Barnard N (2017) Cardio-metabolic benefits of plant-based diets. *Nutrients* **9**, 8.
44. Lin PH, Allen JD, Li YJ *et al.* (2012) Blood pressure-lowering mechanisms of the DASH dietary pattern. *J Nutr Metabol* **2012**, 472396.
45. Utsugi MT, Ohkubo T, Kikuya M *et al.* (2008) Fruit and vegetable consumption and the risk of hypertension determined by self measurement of blood pressure at home: the Ohasama study. *Hypertens Res* **31**, 1435–1443.
46. Hoshi T, Wissuwa B, Tian Y *et al.* (2013) Omega-3 fatty acids lower blood pressure by directly activating large-conductance Ca(2)(+)-dependent K(+) channels. *Proc Natl Acad Sci USA* **110**, 4816–4821.
47. Del Gobbo LC, Falk MC, Feldman R *et al.* (2015) Effects of tree nuts on blood lipids, apolipoproteins, and blood pressure: systematic review, meta-analysis, and dose-response of 61 controlled intervention trials. *Am J Clin Nutr* **102**, 1347–1356.
48. Sabate J, Oda K & Ros E (2010) Nut consumption and blood lipid levels: a pooled analysis of 25 intervention trials. *Arch Intern Med* **170**, 821–827.