King's Research Portal

DOI:

[10.1017/S0033291711002996](https://doi.org/10.1017/S0033291711002996)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Masillo, A., Day, F., Laing, J., Howes, O., Fusar-Poli, P., Byrne, M., Bhattacharyya, S., Fiori Nastro, P., Girardi, P., McGuire, P. K., & Valmaggia, L. R. (2012). Interpersonal sensitivity in the at-risk mental state for psychosis. *Psychological Medicine*, 42(9), 1835-1845. <https://doi.org/10.1017/S0033291711002996>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Interpersonal sensitivity in the at-risk mental state for psychosis

A. Masillo^{1,2,3}, F. Day^{1,2}, J. Laing¹, O. Howes^{1,2}, P. Fusar-Poli^{1,2}, M. Byrne^{1,2}, S. Bhattacharyya^{1,2}, P. Fiori Nastro⁴, P. Girardi^{3,5}, P. K. McGuire^{1,2} and L. R. Valmaggia^{1,2,6*}

¹ Department of Psychosis Studies, King's College London, King's Health Partners, Institute of Psychiatry, UK

² Outreach and Support in South London (OASIS), South London and Maudsley NHS Trust, UK

³ NESMOS Department (Neurosciences, Mental Health and Sensory Functions), 'Sapienza' University of Rome, 2nd Medical School, Sant'Andrea Hospital, Rome, Italy

⁴ Department of Neurology and Psychiatry, 'Sapienza' University of Rome, 1st Medical School, Rome, Italy

⁵ Department of Neuropsychiatry, Villa Rosa, Suore Ospitaliere of the Sacred Heart of Jesus, Viterbo, Italy

⁶ Department of Psychiatry and Neuropsychology, Maastricht University, The Netherlands

Background. Interpersonal sensitivity is a personality trait described as excessive awareness of both the behaviour and feelings of others. Although interpersonal sensitivity has been found to be one of the vulnerability factors to depression, there has been little interest in its relationship with the prodromal phase of psychosis. The aims of this study were to examine the level of interpersonal sensitivity in a sample of individuals with an at-risk mental state (ARMS) for psychosis and its relationship with other psychopathological features.

Method. Sixty-two individuals with an ARMS for psychosis and 39 control participants completed a series of self-report questionnaires, including the Interpersonal Sensitivity Measure (IPSM), the Prodromal Questionnaire (PQ), the Ways of Coping Questionnaire (WCQ) and the Depression and Anxiety Stress Scale (DASS).

Results. Individuals with an ARMS reported higher interpersonal sensitivity compared to controls. Associations between interpersonal sensitivity, positive psychotic symptoms (i.e. paranoid ideation), avoidant coping and symptoms of depression, anxiety and stress were also found.

Conclusions. This study suggests that being 'hypersensitive' to interpersonal interactions is a psychological feature of the putatively prodromal phase of psychosis. The relationship between interpersonal sensitivity, attenuated positive psychotic symptoms, avoidant coping and negative emotional states may contribute to long-term deficits in social functioning. We illustrate the importance, when assessing a young client with a possible ARMS, of examining more subtle and subjective symptoms in addition to attenuated positive symptoms.

Received 12 May 2011; Revised 18 November 2011; Accepted 5 December 2011; First published online 9 January 2012

Key words: At-risk mental state, coping, depression, early detection, interpersonal sensitivity, prodromal psychotic symptoms.

Introduction

Despite decades of research, schizophrenia and related psychotic disorders remain among the most debilitating disorders in medicine (Tandon *et al.* 2008). Retrospective studies from the 1980s redirected attention to the fact that patients with schizophrenia often showed early, less severe manifestations of the illness for, on average, 5 years before the onset of full psychosis (Häfner *et al.* 1995; Häfner & an der Heiden, 1999). This period has been termed the ultra-high-risk

phase or, retrospectively, the prodromal phase of psychosis (Phillips *et al.* 2002; Yung *et al.* 2003). Recent research has afforded greater importance to this phase. It has been asserted that treatment of the prodrome could prevent onset of the full disorder or ameliorate or delay the onset phase, as claimed by Sullivan in 1927: 'I feel certain that many incipient cases might be arrested before the efficient contact with reality is completely suspended, and a long stay in institutions made necessary' (Sullivan, 1994).

To date, low-intensity or intermittent positive psychotic symptoms are often the most common inclusion criteria for the ultra-high-risk phase (Miller *et al.* 1999, 2002; Broome *et al.* 2005a; Yung *et al.* 2005). Despite the unquestionable importance of these symptoms and their great pragmatic value, many authors have

* Address for correspondence: Dr L. R. Valmaggia, Department of Psychology and Psychosis Studies, King's College London, Institute of Psychiatry, 16 De Crespigny Park, PO box 67, London SE5 8AF, UK.
(Email: lucia.valmaggia@kcl.ac.uk)

stressed the importance of examining psychopathological and phenomenological descriptions for more precise identification of individuals at risk of imminent psychosis (Parnas *et al.* 2005; Davidsen, 2009; Nelson *et al.* 2009a,b; Parnas, 2011; Raballo & Larøi, 2011). This need was explained by Nelson *et al.* (2008) who, in line with Parnas' position (Parnas, 2005) regarding current operational criteria for the assessment of prodromal patients, stated: 'it is not the symptoms as such that put an individual at risk but the underlying or core disturbance of psychotic vulnerability', otherwise it would be like 'predicting extreme heat by an increase in temperature, without identifying the fire that might be causing this change'. To contribute to ongoing research regarding detection of increased risk for psychosis, in this study we aimed to investigate a subtle and subjective psychopathological feature: interpersonal sensitivity.

The importance of interpersonal relationships and their influence on both personality development and psychopathology present challenges to researchers. An outstanding aspect of interpersonal interactions is interpersonal sensitivity, a personality trait described as excessive awareness of both the behaviour and feelings of others (Boyce & Parker, 1989). Highly interpersonally sensitive individuals are extremely sensitive to interpersonal interactions, perceive self-deficiencies in relation to others and behave in such a way as to minimize the risk of negative evaluation (Davidson *et al.* 1989, 1988). High interpersonal sensitivity was also found to be closely linked to low self-confidence, feelings of insecurity, and low self-esteem (Boyce & Parker, 1989). This personality trait was first conceptualized as a set of symptoms occurring both as a consequence of depression and as a vulnerability for the development of depression (Boyce *et al.* 1991; Boyce & Mason, 1996). Early studies indicated high interpersonal sensitivity and problems with self-confidence as being among the subjective symptoms and observable behavioural changes occurring during the prodromal phase of schizophrenia (Subotnik & Nuechterlein, 1988; Häfner *et al.* 1992; Hambrecht *et al.* 1994). More recent studies have confirmed an association between interpersonal sensitivity and persecutory ideations among ultra-high-risk and non-clinical samples (Valmaggia *et al.* 2007; Green *et al.* 2011). Examining interpersonal sensitivity during the prodromal phase of psychosis may also be valuable because of its potential links with dysfunctional coping strategies. Since the early work of Falloon & Talbot (1981), it is generally accepted that coping may serve to diminish threat experiences or other psychotic symptoms and augment controllability. This may only be true for specific coping strategies, such as problem-solving strategies or integration, which can reduce

distress (Dittmann & Schuttler, 1990) and are associated with positive outcome. Less adaptive coping strategies may contribute to negative outcomes such as diminished quality of life and poor social functioning (Tait *et al.* 2004). A functional sense of self or identity may facilitate coping efforts and has been posited as an important resilience factor in recovery from psychosis (Davidson & Strauss, 1992). Feelings of insecurity and negative self-evaluation may encourage development of maladaptive coping strategies (Bernstein *et al.* 1993), such as passivity and avoidance (Tait *et al.* 2004), which could contribute to functional and social deterioration in ultra-high clinical risk individuals. A recent study found that people at ultra-high clinical risk of psychosis showed a greater reliance on maladaptive, passive coping strategies, which are associated with a higher level of negative symptoms, depression and anxiety (Lee *et al.* 2011). In line with these findings, Lin *et al.* (2011) found that emotion-oriented coping (i.e. avoidance, escape) was associated with subclinical psychotic symptoms in a general population adolescent sample.

Greater knowledge concerning interpersonal sensitivity could enhance our understanding of the role of subjective and non-specific symptoms occurring during the prodromal phase of psychosis and their relationship with other psychopathological and behavioural features.

Aims of the study

The aims of the present study were: (i) to assess interpersonal sensitivity in a sample of individuals with an at-risk mental state (ARMS), compared to matched control participants; (ii) to explore, in both samples, the relationship between interpersonal sensitivity and prodromal symptoms of psychosis; (iii) to study, in both samples, the relationship between interpersonal sensitivity and coping; and (iv) to explore, in both samples, the relationship between interpersonal sensitivity and negative emotional states such as depression, anxiety and stress.

On the basis of previous research examining the relationship between interpersonal sensitivity, low self-esteem, feelings of insecurity, dysfunctional sense of identity and mental illness (Davidson *et al.* 1999; Larsen *et al.* 2003; Tait *et al.* 2004), we hypothesized that individuals with an ARMS would report higher interpersonal sensitivity than control participants and that this personality trait would be associated, in both samples, with positive prodromal symptoms, specifically paranoid ideation (Valmaggia *et al.* 2007; Green *et al.* 2011). Moreover, we hypothesized that highly interpersonally sensitive individuals would report increased use of avoidant coping strategies and higher

levels of negative emotional states (depression, anxiety and stress).

Method

Participants

There were 101 participants in this study: 62 with an ARMS and 39 healthy controls. People at high risk for psychosis were recruited through Outreach and Support in South London (OASIS), a clinical service for help-seeking young people, aged 14–35 years, at risk for psychosis (with an ARMS) (Broome *et al.* 2005b). The ARMS was evaluated using the Comprehensive Assessment of At-Risk Mental States (CAARMS; Yung *et al.* 2005). OASIS clients were referred from local general practitioners, schools and colleges, social and faith groups, adolescent and adult mental health services or self-referred. All clients are offered psychological (cognitive behaviour therapy, CBT) and/or pharmacological treatment for a maximum period of 2 years. Healthy control participants, from the same geographic region matched for age, gender and ethnicity to the ARMS group, were recruited using the following methods: searching on the MindSearch research volunteer database (www.mindsearch.iop.kcl.ac.uk); approaching people who had previously taken part in research studies at the Institute of Psychiatry; and asking existing control participants to give details of the study to any friends who might also be interested in taking part. The following inclusion criteria were used: participants aged between 18 and 35 years, lived (or grew up) in South London, and no personal history of mental health problems.

Research ethics approval was obtained from the National Research Ethics Service (Appendix 4.3 Ethics REC no. 08/H0722/45). Participants provided written informed consent prior to commencement of the study.

Measures

Sociodemographic and psychosocial variables were recorded during a clinical assessment using a non-standardized questionnaire modelled on the Census 2001 collection form, named the First Contact with OASIS Questionnaire.

To measure interpersonal sensitivity, we used the Interpersonal Sensitivity Measure (IPSM; Boyce & Parker, 1989), a 36-item self-report questionnaire. Self-statements are rated on a four-point scale (1 = very unlike self, 4 = very like self). The scale generates a total score ranging from 36 to 144, with higher scores indicating greater interpersonal sensitivity, and five

subscales scores: 'Interpersonal awareness' (seven items, range 1–28); 'Need for approval' (eight items, range 8–32); 'Separation anxiety' (eight items, range 8–32); 'Timidity' (eight items, range 8–32) and 'Fragile inner self' (five items, range 5–20). Previous research among a non-clinical sample reported a mean score of 93.2 for the IPSM total score, 18.7 for 'Interpersonal awareness', 26.0 for 'Need for approval', 18.1 for 'Separation anxiety', 20.6 for 'Timidity' and 9.7 for 'Fragile inner self' (Green *et al.* 2011). The IPSM has been found to have good internal consistency (α values from 0.85 to 0.86), test-retest reliability ($r=0.70$) and correlation with clinical judgment ratings of interpersonal sensitivity ($r=0.72$).

To assess prodromal and psychotic symptoms, we used the Prodromal Questionnaire (PQ; Loewy *et al.* 2005), a self-report screening questionnaire that aims to identify individuals who may benefit from a clinical diagnostic interview. The 92 true/false items can be divided into four major subscales: (1) positive symptoms (e.g. unusual thinking and perceptual abnormalities); (2) negative symptoms (e.g. flat affect and social isolation); (3) disorganized symptoms (e.g. odd behaviour); and (4) general symptoms (e.g. depression and diminished role functioning). A score of eight or more positive symptoms on the PQ has been found to differentiate between individuals without an ARMS and those with prodromal or psychotic syndrome diagnoses with 90% sensitivity, 49% specificity, 78% positive predictive value and 69% negative predictive value.

We also used the Ways of Coping Questionnaire (WCQ; Folkman & Lazarus, 1985), a 66-item self-report questionnaire containing a broad range of coping and behavioural strategies that people can use to manage internal or external demands of stressful situations (Folkman *et al.* 1986b). Responses are rated on a four-point Likert scale (0 = not used, 3 = used a great deal). The WCQ comprises eight subscales (Folkman *et al.* 1986a): 'Confrontive coping' (six items, range 0–18); 'Distancing' (seven items, range 0–21); 'Self controlling' (seven items, range 0–21); 'Seeking social support' (six items, range 0–18); 'Accepting responsibility' (four items, range 0–12); 'Escape-Avoidance' (eight items, range 0–24); 'Planful problem-solving' (six items, range 0–18); and 'Positive reappraisal' (seven items, range 0–21). The WCQ has been used extensively in clinical and non-clinical samples, and the stability of its factor structure, its reliability and validity have been the subject of intense scrutiny, indicating good reliability and validity (Parker *et al.* 1993).

Finally, we used the Depression Anxiety Stress Scales (DASS; Lovibond & Lovibond, 1995), a 42-item instrument consisting of three subscales measuring

Table 1. Sociodemographic characteristics of the sample

	ARMS (<i>n</i> = 62)	HC (<i>n</i> = 39)	Test	<i>p</i>
Sex, <i>n</i> (%)				
Male	37 (59.7)	20 (51.3)	$\chi^2(1, 101) = 0.686$	0.407
Female	25 (40.3)	19 (48.7)		
Ethnicity, <i>n</i> (%)				
Black British	10 (16.1)	4 (10.3)	$\chi^2(7, 101) = 7.010$	0.449 (exact)
Black Caribbean	6 (9.7)	5 (12.8)		
Black African	3 (4.8)	0 (0.0)		
White British	21 (33.9)	15 (38.5)		
White Other	11 (17.7)	7 (17.9)		
Asian Oriental	1 (1.6)	2 (5.1)		
Asian Indian	1 (1.6)	3 (7.7)		
Other	9 (14.5)	3 (7.7)		
Employment status, <i>n</i> (%)				
Unemployed	36 (58.1)	3 (7.7)	$\chi^2(2, 101) = 25.646$	0.000
Employed	14 (22.6)	20 (51.3)		
Student	12 (19.4)	16 (41.0)		
Marital status, <i>n</i> (%)				
Single	46 (74.2)	22 (36.4)	$\chi^2(1, 101) = 3.442$	0.064
Relationship	16 (25.8)	17 (43.6)		
Age, mean (s.d.)	22.63 (4.05)	24.03 (4.22)	$t(99) = -1.658$	0.532

ARMS, At-risk mental state; HC, healthy controls; s.d., standard deviation.

Bold values indicate significant results.

current symptoms of depression, anxiety and stress. Each of the subscales consists of 14 items with a 0–3 scale (0 = did not apply at all to me, 3 = applied to me very much). Participants are asked to rate the extent to which they experienced each state over the past week. Higher scores indicate increased levels of emotional distress. Subscale scores range from 0 to 42 and total scores range from 0–126. The scale's reliability and validity has been demonstrated in a large UK non-clinical sample (Crawford & Henry, 2003).

Statistical analysis

Descriptive statistics including mean and standard deviation values for continuous variables and absolute and relative frequencies for categorical variables were calculated. Group differences in categorical variables were examined using the χ^2 test. Mann–Whitney *U* tests were conducted to explore the impact of the ARMS on interpersonal sensitivity (as measured by the IPSM), endorsement of escape/avoidant coping (as measured by the WCQ subscale), negative affectivity (as measured by the DASS) and positive prodromal symptoms (as measured by the PQ positive symptoms subscale).

Spearman's rank-order correlation coefficients (r_s) were calculated to examine associations between

interpersonal sensitivity, positive prodromal symptoms, negative affectivity and escape/avoidance coping according to group membership. Spearman's partial correlations were computed to explore the relationship between interpersonal sensitivity and prodromal positive symptoms of psychosis while controlling for depressive symptoms (as measured by the DASS). The level of statistical difference was set at $p < 0.05$ and all reported significance values were two-tailed. Statistical analyses were performed using SPSS version 18 (SPSS Inc., USA).

Results

A total of 62 individuals with ARMS for psychosis and 39 healthy control participants were included in the present study. Sociodemographic characteristics of the sample are presented in Table 1. There were no significant differences between groups in age, sex, ethnicity and marital status. ARMS employment status differed significantly from healthy controls [$\chi^2(2, 101) = 25.646$, $p < 0.001$]: a greater number of participants with ARMS were unemployed ($n = 36$, 58.1%) relative to control participants ($n = 3$, 7.7%).

As illustrated in Table 2, there were statistically significant differences between groups in IPSM total score ($U = 577.0$, $p < 0.001$), interpersonal awareness

Table 2. Comparisons between participants with ARMS and controls with regard to self-report measures

	ARMS				HC				<i>p</i> ^a
	Mean	S.D.	Median	Min-max	Mean	S.D.	Median	Min-max	
IPSM total score	101.94	17.7	104	60–134	87.67	12.4	89	63–120	0.000
Interpersonal awareness	21.53	4.8	22	10–28	17.31	3.9	17	9–24	0.000
Need for approval	24.05	4.2	25	13–32	25.49	2.5	26	20–31	0.098
Separation anxiety	22.73	5.7	24	8–32	16.31	4.4	16	10–30	0.000
Timidity	21.23	4.8	21	9–31	19.62	3.7	19	13–28	0.078
Fragile inner self	12.4	4.2	13	5–20	8.95	3.2	8	5–18	0.000
PQ-Positive	18.1	11.3	15.5	0–45	4.4	4.2	4	0–15	0.000
DASS-Depression	21.1	12.2	20	0–42	3.3	4.2	1.5	0–16	0.000
DASS-Anxiety	14.4	10.4	12	0–42	2	2.4	1	0–11	0.000
DASS-Stress	20.4	12.1	19	1–42	5.6	5.5	4	0–17	0.000
Escape/avoidance coping	17.6	5.5	17	9–30	13.7	4	12	8–24	0.001

ARMS, At-risk mental state; HC, healthy controls; S.D., standard deviation; IPSM, Interpersonal Sensitivity Measure; PQ, Prodromal Questionnaire; DASS, Depression and Anxiety Stress Scale.

Bold values indicate significant results.

^a Mann-Whitney *U* test.

($U=592.0$, $p<0.001$), separation anxiety ($U=474.5$, $p<0.001$) and fragile inner self ($U=644.5$, $p<0.001$). There were also statistically significant differences between participants at ultra-high clinical risk for psychosis and controls in depression ($U=203.0$, $p<0.001$), anxiety ($U=241.0$, $p<0.001$) and stress ($U=335.5$, $p<0.001$) DASS subscales scores. A significant difference in escape/avoidance WCQ subscale median scores between groups was also found ($U=537.0$, $p=0.001$). Groups significantly differed in median PQ positive symptom subscale scores ($U=314.0$, $p<0.001$), with ARMS reporting significantly higher levels of positive psychotic symptoms (median = 15.5) than control participants (median = 4).

The relationship between symptoms and questionnaire scores are shown by group in Table 3. Higher sensitivity to interpersonal interactions, anxiety about separation from significant others and sense of having an inner or core self that is unlikeable and needs to be hidden from others were all associated with higher numbers of positive prodromal symptoms. Three PQ items specifically address the presence of paranoid ideation and suspiciousness (PQ25: 'I often feel that other people have it in for me'; PQ68: 'I often pick up hidden threats or put-downs from what people say or do'; PQ77: 'I'm often concerned that my closest friends, classmates or co-workers are not really loyal or trustworthy'). The higher the interpersonal awareness ($r_s=0.52$, $p=0.001$), separation anxiety ($r_s=0.71$, $p<0.001$), fragile inner self ($r_s=0.51$, $p<0.001$) and total IPSM ($r_s=0.52$, $p<0.001$) scores among ultra-high clinical risk participants, the higher the level of paranoid ideas and suspiciousness. A significant

Table 3. Correlations between IPSM and PQ positive symptoms subscale (by group)

	ARMS		Controls	
	<i>r_s</i>	Sig.	<i>r_s</i>	Sig.
PQ positive				
IPSM total	0.34	0.006	0.48	0.002
Interpersonal awareness	0.30	0.014	0.43	0.006
Separation anxiety	0.51	0.000	0.51	0.001
Fragile inner-self	0.37	0.003	0.63	0.000
Paranoid ideations/suspiciousness				
IPSM total	0.52	0.000	0.24	0.130
Interpersonal awareness	0.52	0.001	0.21	0.180
Separation anxiety	0.71	0.000	0.32	0.046
Fragile inner-self	0.51	0.000	0.27	0.093

ARMS, At-risk mental state; IPSM, Interpersonal Sensitivity Measure; PQ, Prodromal Questionnaire; Sig., significance.

Bold values indicate significant results.

association between separation anxiety subscale score and paranoid/suspiciousness was also found among control participants ($r_s=0.32$, $p<0.05$).

Spearman's partial correlations were computed to explore the relationship between interpersonal sensitivity and prodromal positive symptoms of psychosis while controlling for depressive symptoms (as measured by the DASS). The degree of association between IPSM total scores and PQ positive symptoms subscale scores were no longer statistically significant after controlling for depression.

Among both at-risk and control participants, statistically significant positive correlations were found between IPSM scores, DASS subscales scores and escape avoidant coping. Among participants with an ARMS, total IPSM score ($r_s=0.40$, $p<0.01$), interpersonal awareness ($r_s=0.34$, $p<0.01$), separation anxiety ($r_s=0.50$, $p<0.01$) and fragile inner self ($r_s=0.35$, $p<0.01$) were significantly positively correlated with escape/avoidance WCQ subscale scores. Only total IPMS score ($r_s=0.38$, $p<0.05$), separation anxiety ($r_s=0.48$, $p<0.01$) and fragile inner self ($r_s=0.37$, $p<0.05$) subscale scores were correlated with escape/avoidance coping among control participants (Table 4).

Among both at-risk and control participants, total IPSM score and interpersonal awareness, fragile inner self and separation anxiety subscales scores were significantly correlated with depression, anxiety and stress measures (Table 5).

Discussion

This study explored interpersonal sensitivity among a sample with an ARMS relative to healthy control participants. Before discussing the results, it is important to note that the study was cross-sectional and therefore it is impossible to infer causality. Without longitudinal follow-up data we cannot draw any conclusion on whether interpersonal sensitivity is a predictive or an independent factor for the transition from an ARMS to first-episode psychosis.

In line with our first research hypothesis, we found that individuals with an ARMS scored higher on all IPSM subscales compared to control participants. IPSM scores of participants with an ARMS were similar to those reported by individuals with major depression (Luty *et al.* 2002). Control participants' IPSM scores were similar to those reported within previous general population studies (Otani *et al.* 2008; Green *et al.* 2011). Interpersonal sensitivity was first proposed as a vulnerability factor for depression (Boyce *et al.* 1991; Boyce & Mason, 1996; Sato *et al.* 2001; Luty *et al.* 2002). Retrospective studies have consistently documented the relatively high frequency of non-specific symptoms, such as depression, anxiety, social isolation and educational difficulties prior to the onset of schizophrenia (Yung & McGorry, 1996a,b; Häfner & an der Heiden, 1999). Lencz *et al.* (2004) conducted a prospective study focusing on negative and non-specific pre-psychotic symptoms showing that individuals at ultra-high clinical risk for psychosis reported depressed mood, anxiety and decline in school functioning about as commonly as positive symptoms. Addington *et al.* (2011) found that a high percentage of individuals with an ARMS had co-morbid diagnoses

Table 4. Correlations between IPSM and escape/avoidance WCQ subscale (by group)

Escape/Avoidance WCQ	ARMS		Controls	
	r_s	Sig.	r_s	Sig.
IPSM total	0.40	0.002	0.38	0.029
Interpersonal awareness	0.34	0.010	0.32	0.064
Separation anxiety	0.50	0.000	0.48	0.005
Fragile inner-self	0.35	0.008	0.37	0.031

ARMS, At-risk mental state; IPSM, Interpersonal Sensitivity Measure; WCQ, Ways of Coping Questionnaire; Sig., significance.

Bold values indicate significant results.

of major depressive disorder and anxiety disorder, suggesting depression as a primary presentation coexistent with and independent from prodromal symptoms of psychosis. Wigman *et al.* (2011) also showed that subclinical psychosis and depression are inter-related phenomena that strongly co-occur in time, but longitudinally; one does not predict change in the other. Other researchers have hypothesized that depressive symptoms in ultra-high-risk individuals may be due to dysphoria and distress secondary to the recent onset of psychotic experiences (Birchwood *et al.* 2000; Demjaha *et al.* 2010). In line with these studies, our results have demonstrated that interpersonal sensitivity, in both participants with an ARMS and healthy controls, is correlated to negative emotional states, such as depression, but also to positive prodromal symptoms. Previous research has demonstrated an association between interpersonal sensitivity and persecutory ideation among non-clinical samples (Free-man *et al.* 2005, 2008; Green *et al.* 2011). Valmaggia *et al.* (2007) also found that paranoid ideation in people with an ARMS was predicted by a high level of interpersonal sensitivity. Similarly, this study found that the higher the sensitivity to interpersonal interactions, anxiety about separation from significant others and the sense of having an inner or core self that is unlikeable and needs to be hidden from others, the higher the level of paranoid ideation. However, as noted above, this research was cross-sectional and therefore it was not possible to evaluate whether interpersonal sensitivity predicted the paranoid thinking in our samples. This finding is in line with cognitive models of positive symptoms of psychosis, in which negative beliefs about the self as fragile and vulnerable to threat may lead to a tendency to attribute experiences as externally caused and in turn facilitate the formation and maintenance of paranoid ideation (Garety *et al.* 2001). It has also been suggested

Table 5. Correlations between IPSM and DASS (by group)

	ARMS		Controls	
	r_s	Sig.	r_s	Sig.
Depression-DASS				
IPSM total	0.56	0.000	0.56	0.000
Interpersonal awareness	0.50	0.000	0.59	0.000
Separation anxiety	0.65	0.000	0.43	0.007
Fragile inner self	0.58	0.000	0.63	0.000
Anxiety-DASS				
IPSM total	0.60	0.000	0.47	0.002
Interpersonal awareness	0.56	0.000	0.50	0.001
Separation anxiety	0.63	0.000	0.32	0.046
Fragile inner self	0.49	0.000	0.53	0.000
Stress-DASS				
IPSM total	0.58	0.000	0.54	0.000
Interpersonal awareness	0.60	0.000	0.75	0.000
Separation anxiety	0.63	0.000	0.40	0.001
Fragile inner self	0.53	0.000	0.63	0.000

ARMS, At-risk mental state; IPSM, Interpersonal Sensitivity Measure; DASS, Depression and Anxiety Stress Scale; Sig., significance.

Bold values indicate significant results.

that paranoid thoughts may build upon interpersonal anxieties and worries such as fear of rejection (Freeman *et al.* 2005). Trower & Chadwick's (1995) model distinguished between two types of paranoia: persecution (or 'poor me') and punishment (or 'bad me') paranoia. Consistent with this theory, individuals who experienced 'bad me' paranoia reported significantly lower self-esteem and increased depression compared to 'poor me' paranoid people (Chadwick *et al.* 2005). An *et al.* (2010) also found a possible association between low self-esteem and depression and severity of paranoia among individuals with an ARMS. The finding of high levels of interpersonal sensitivity and its correlation with negative affectivity and paranoid ideations in the ARMS in the present study is more consistent with 'bad me' paranoia, in which paranoia is based on negative self-evaluation and others' malevolence is seen as a justified and deserved punishment for one's own inadequacies. However, a previous study reported that people with persecutory delusions who are in the initial stages of psychosis do not tend to show 'bad me' paranoia (Fornells-Ambrojo & Garety, 2009). In our study, after controlling for the potential effect of depression, the correlation between interpersonal sensitivity and positive symptoms (including paranoid ideation items) was no longer statistically significant, indicating that there may be a mediating effect of depressive symptoms on the relationship between

interpersonal sensitivity and positive prodromal symptoms. This suggests that the previously reported association between interpersonal sensitivity and paranoid thinking in ARMS (Valmaggia *et al.* 2007) may be explained by the presence of depressive symptoms. We might tentatively conclude that individuals with an ARMS may have a more preserved 'affective core' (i.e. interpersonal sensitivity and depressive symptoms) that may be characterized, at least in part, by negative self-evaluation, personal responsibility for badness/inadequacy and a sense of deserving mistreatment ('bad me' paranoia). Individuals who have crossed the psychosis threshold may show more blunted affectivity and detachment from others, psychopathological features described as 'primary symptoms' of schizophrenia by Bleuler (1911), which may more easily result in 'poor me' persecutory paranoia.

In line with our second research hypothesis we found a significant positive correlation between interpersonal sensitivity and avoidant coping strategies (as measured by the escape/avoidance WCQ subscale), both in participants with an ARMS and healthy controls. These findings are in accordance with previous research reporting an association between interpersonal sensitivity and maladaptive problem-solving styles (McCabe *et al.* 1999). Social withdrawal, habituation or adaptation to illness, and 'self-treatment' with alcohol or drugs may constitute part of avoidant ways of coping used by young people when confronting stressful encounters. Some of the escape/avoidant WCQ subscale items reflected these responses: 'avoided being with people in general'; 'tried to make myself feel better by eating, drinking, smoking, using drugs or medications'; 'refused to believe that it had happened'; 'slept more than usual'. We found that personality traits characterized by negative sense of self and feelings of insecurity were associated with the use of avoidant coping strategies. Examining this association, particularly in relation to changes in social functioning among at-risk individuals, may be a valuable area of future research. It is well known that adolescents often struggle to achieve an integrated and coherent sense of self, consolidating the many different aspects of their private and social persona. The current findings suggest increased levels of inner-self fragility among those with an ARMS. Perceived self-deficiencies in relation to others, feelings of a fragile and bothersome core-self and paranoid ideation may contribute to social anxiety and isolation (Freeman *et al.* 2008). Increased knowledge concerning the possible causes of impaired social functioning is crucial, considering that social withdrawal is the most commonly reported symptom by individuals with an ARMS (Lencz *et al.* 2004), and is an

important factor related to transition from the prodromal phase to frank psychosis (McGlashan *et al.* 2007). Lee *et al.* (2011) argued that individuals who relied on maladaptive coping strategies might be more depressed, anxious and, as a consequence, more likely to avoid social interactions. Avoidant behaviours, such as social withdrawal, may constitute part of the negative symptom profile of psychosis, a consequence of low mood and lack of drive, and/or may result from suspiciousness and paranoia; however, in all of these cases, they may reflect the presence of troubles and difficulties in interpersonal relationships.

With regard to the relationship between interpersonal sensitivity and negative emotional states, we found a statistically significant correlation between fragile inner self, separation anxiety and interpersonal awareness and all three DASS subscales (depression, anxiety, and stress). This result is in line with previous researches (Boyce *et al.* 1991; Boyce & Mason, 1996) that showed that interpersonal sensitivity was both a consequence of depression and a vulnerability for the development of depression. Another study, investigating the relationship between interpersonal sensitivity and anxiety disorders, also found differential associations with specific anxiety disorders (Wilhelm *et al.* 2004). As argued by Clarke & Watson (1991), anxiety and depression share a component called 'negative affectivity', which reflects the experience of subjective distress and unpleasurable engagement, manifested in emotional states such as guilt, anger and nervousness. On the one hand, pervasive feelings of insecurity, low self-esteem and hyper-attentiveness to the reactions and behaviours of others are personality traits that may contribute to emotions such as anger or nervousness and could be related to the development of negative emotions. On the other hand, depression, anxiety or distress could exacerbate the sensitivity to interpersonal interactions and feelings of having a fragile core self. Thus, a vicious cycle may arise between interpersonal sensitivity and negative emotional states that may result, together with avoidant coping strategies, in difficulties in social interactions or social isolation. Assessing levels of interpersonal sensitivity and planning targeted psychotherapeutic interventions during the ARMS for psychosis, focused on producing a more unified and integrated sense of self, as suggested by Nelson *et al.* (2009b), or focused on investigating the deeper nucleus of this self-core fragility may be helpful in combating potential difficulties in interpersonal relationships.

Finally, it is important to underline the possible affinity between the notion of interpersonal sensitivity, the object of our research, and the phenomenological model of self-disturbance, defined by some authors as

the core clinical feature of schizophrenia spectrum disorders (Parnas & Handest, 2003; Sass & Parnas, 2003) and found to be present also in adolescents at risk of psychosis (Davidsen, 2009). In particular, the hyper-reflectivity aspect of this model (defined as 'an increase in the tendency to reflect about one's own thinking, feelings and behaviour, and inability to react and behave spontaneously and carefree; a tendency to excessively monitoring inner life, while at the same time interacting in the world'; Parnas *et al.* 2005) may have some similarities to interpersonal awareness and fragile inner self aspects of interpersonal sensitivity. This is confirmed by a recent study that showed that disturbance of the basic sense of self may underlie the social cognition and interpersonal difficulties present in psychotic disorders (Nelson *et al.* 2009c).

Limitations

Our results should be interpreted in view of the limitations of this study. As mentioned earlier, a major limitation is the lack of follow-up data to evaluate whether interpersonal sensitivity is a predictive, or independent, factor for the transition from an ARMS to first-episode psychosis. In future studies we intend to explore the correlations between baseline interpersonal sensitivity level and long-term outcomes in terms of psychopathology and social functioning. A further limitation was that paranoia and depression were not assessed using specific and diagnostic instruments, but rather using PQ (a screening instrument for prodromal symptoms) and DASS (a measure of current negative emotional states) scores. A final limitation was that all assessments were made by self-reported questionnaires rather than by clinical interviews; this weakens the strength of the results because it is possible that participants misinterpreted some questions.

Conclusions

We found that 'hypersensitivity' to interpersonal interactions was a subjective psychological feature manifest during the ARMS for psychosis and distinguished ultra-high-risk participants from controls. Furthermore, interpersonal sensitivity was related to negative emotional states such as depression, anxiety and stress and avoidant coping strategies, such as social withdrawal and self-treatment with alcohol and drugs. It can be speculated that interpersonal sensitivity may play an active negative role in functional deterioration present in the pre-psychotic phase (Fusar-Poli *et al.* 2010) and contribute to poorer long-term functional outcomes. For this reason, addressing difficulties in interpersonal relationships and offering

targeted psychotherapeutic interventions may usefully be included in early intervention treatment strategies. As the findings of this cross-sectional study were largely correlational, further research will be required to examine the relationship between interpersonal sensitivity and long-term outcomes of individuals at ultra-high clinical risk for psychosis and to assess whether interpersonal sensitivity may predict the clinical features of potential future psychosis.

Acknowledgements

Our special thanks go to the staff and service users of OASIS. Dr L. R. Valmaggia was supported by a National Alliance for Research on Schizophrenia and Depression (NARSAD) Young Investigator Award and by a Peggy Pollack Research Fellowship from the Psychiatry Research Trust. We acknowledge the National Institute for Health Research (NIHR) Biomedical Research Centre for Mental Health at the South London and Maudsley National Health Service (NHS) Foundation Trust and Institute of Psychiatry King's College London for their financial support.

Declaration of Interest

None.

References

- Addington J, Epstein I, Liu L, French P, Boydell KM, Zipursky RBA (2011). Randomized controlled trial of cognitive behavioral therapy for individuals at clinical high risk of psychosis. *Schizophrenia Research* **125**, 54–61.
- An SK, Kang JI, Park JY, Kim KR, Lee SY, Lee E (2010). Attribution bias in ultra-high risk for psychosis and first-episode schizophrenia. *Schizophrenia Research* **118**, 54–61.
- Bernstein DP, Cohen P, Velez N, Schwab-Stone M, Siever L, Shinsato L (1993). Prevalence and stability of the DSM-III-R personality disorders in a community-based sample of adolescents. *American Journal of Psychiatry* **150**, 1237–1243.
- Birchwood M, Meaden A, Trower P, Gilbert P, Plaistow J (2000). The power and omnipotence of voices. *Psychological Medicine* **30**, 337–344.
- Bleuler E (1911). *Dementia Praecox or the Group of Schizophrenias*. Translated by J. Zinkin. International University Press: New York (1950).
- Boyce P, Mason C (1996). An overview of depression-prone personality traits and the role of interpersonal sensitivity. *Australian and New Zealand Journal of Psychiatry* **30**, 90–103.
- Boyce P, Parker G (1989). Development of a scale to measure interpersonal sensitivity. *Australian and New Zealand Journal of Psychiatry* **23**, 341–351.
- Boyce P, Parker G, Barnett B, Cooney M, Smith F (1991). Personality as a vulnerability factor to depression. *British Journal of Psychiatry* **159**, 106–114.
- Broome MR, Woolley JB, Johns LC, Valmaggia LR, Tabraham P, Gafoor R, Bramon E, McGuire PK (2005b). Outreach and support in south London (OASIS): implementation of a clinical service for prodromal psychosis and the at risk mental state. *European Psychiatry* **20**, 372–378.
- Broome MR, Woolley JB, Tabraham P, Johns LC, Bramon E, Murray GK, Pariente C, McGuire PK, Murray RM (2005a). What causes the onset of psychosis? *Schizophrenia Research* **79**, 23–34.
- Crawford JR, Henry JD (2003). The Depression Anxiety Stress Scales (DASS): normative data and latent structure in a large non-clinical sample. *British Journal of Clinical Psychology* **42**, 111–131.
- Davidson KA (2009). Anomalous self-experience in adolescents at risk of psychosis. *Psychopathology* **42**, 361–369.
- Davidson J, Zisook S, Giller E, Helms M (1989). Symptoms of interpersonal sensitivity in depression. *Comprehensive Psychiatry* **30**, 357–368.
- Davidson JR, Giller EL, Zisook S, Overall JE (1988). An efficacy study of isocarboxazid and placebo in depression, and its relationship to depressive nosology. *Archives of General Psychiatry* **45**, 120–127.
- Davidson L, Strauss JS (1992). Sense of self in recovery from severe mental illness. *British Journal of Medical Psychology* **65**, 131–145.
- Davidson M, Reichenberg A, Rabinowitz J, Weiser M, Kaplan Z, Mark M (1999). Behavioral and intellectual markers for schizophrenia in apparently healthy male adolescents. *American Journal of Psychiatry* **156**, 1328–1335.
- Demjaha A, Valmaggia L, Stahl D, Byrne M, McGuire P (2010). Disorganization/cognitive and negative symptom dimensions in the at-risk mental state predict subsequent transition to psychosis. *Schizophrenia Bulletin*. Published online: 12 August 2010. doi:10.1093/schbul/sbq088.
- Dittmann J, Schuttler R (1990). Disease consciousness and coping strategies of patients with schizophrenic psychosis. *Acta Psychiatrica Scandinavica* **82**, 318–322.
- Falloon IR, Talbot RE (1981). Persistent auditory hallucinations: coping mechanism and implications for management. *Psychological Medicine* **11**, 329–339.
- Folkman S, Lazarus RS (1985). If it changes it must be a process: study of emotion and coping during three stages of a college examination. *Journal of Personality and Social Psychology* **48**, 150–170.
- Folkman S, Lazarus RS, Dunkel-Schetter C, DeLongis A, Gruen RJ (1986a). Dynamics of a stressful encounter: cognitive appraisal, coping, and encounter outcomes. *Journal of Personality and Social Psychology* **50**, 992–1003.
- Folkman S, Lazarus RS, Gruen RJ, DeLongis A (1986b). Appraisal, coping, health status, and psychological symptoms. *Journal of Personality and Social Psychology* **50**, 571–579.
- Fornells-Ambrojo M, Garety PA (2009). Understanding attributional biases, emotions and self-esteem in 'poor me' paranoia: findings from an early psychosis sample. *British Journal of Clinical Psychology* **48**, 141–162.
- Freeman D, Dunn G, Garety PA, Bebbington P, Slater M, Kuipers E, Fowler D, Green C, Jordan J, Ray K (2005).

- The psychology of persecutory ideation I: A questionnaire survey. *Journal of Nervous and Mental Disease* **193**, 302–308.
- Freeman D, Gittins M, Pugh K, Antley A, Slater M, Dunn G** (2008). What makes one person paranoid and another person anxious? The differential prediction of social anxiety and persecutory ideation in an experimental situation. *Psychological Medicine* **38**, 1121–1132.
- Fusar-Poli P, Byrne M, Valmaggia L, Day F, Tabraham P, Johns L, McGuire P; OASIS Team** (2010). Social dysfunction predicts two years clinical outcome in people at ultra high risk for psychosis. *Journal of Psychiatric Research* **44**, 294–301.
- Garety PA, Kuipers E, Fowler D, Freeman D, Bebbington PE** (2001). A cognitive model of the positive symptoms of psychosis. *Psychological Medicine* **31**, 189–195.
- Green CE, Freeman D, Kuipers E, Bebbington P, Fowler D, Dunn G, Garety P** (2011). Paranoid explanations of experience: a novel experimental study. *Behavioural and Cognitive Psychotherapy* **39**, 21–34.
- Häfner H, an der Heiden W** (1999). The course of schizophrenia in the light of modern follow-up studies: the ABC and WHO studies. *European Archives of Psychiatry and Clinical Neuroscience* **249**, s14–s26.
- Häfner H, Maurer K, Löffler W** (1995). Onset and early course of schizophrenia. In *Search for the Causes of Schizophrenia* (ed. H. G. W. Häfner), vol. 3, pp. 43–66. Springer-Verlag: Berlin.
- Häfner H, Riecher-Rössler A, Hambrecht M, Maurer K, Meissner S, Schmidtke A, Fätkenheuer B, Löffler W, van der Heiden W** (1992). IRAOS: an instrument for the assessment of onset and early course of schizophrenia. *Schizophrenia Research* **6**, 209–223.
- Hambrecht M, Häfner H, Löffler W** (1994). Beginning schizophrenia observed by significant others. *Social Psychiatry and Psychiatric Epidemiology* **29**, 53–60.
- Larsen TK, Bechdolf A, Birchwood M** (2003). The concept of schizophrenia and phase-specific treatment: cognitive-behavioral treatment in pre-psychosis and in nonresponders. *Journal of the American Academy of Psychoanalysis and Dynamic Psychiatry* **31**, 209–228.
- Lee SY, Kim KR, Park JY, Park JS, Kim B, Kang JI, Lee E, An SK, Kwon JS** (2011). Coping strategies and their relationship to psychopathologies in people at ultra high-risk for psychosis and with schizophrenia. *Journal of Nervous and Mental Disease* **199**, 106–110.
- Lencz T, Smith CW, Auther A, Correll CU, Cornblatt B** (2004). Nonspecific and attenuated negative symptoms in patients at clinical high-risk for schizophrenia. *Schizophrenia Research* **68**, 37–48.
- Lin A, Wigman JT, Nelson B, Vollebergh WA, van Os J, Baksheev G, Ryan J, Raaijmakers QA, Thompson A, Yung AR** (2011). The relationship between coping and subclinical psychotic experiences in adolescents from the general population: a longitudinal study. *Psychological Medicine*. Published online: 28 April 2011. doi:10.1017/S0033291711000560.
- Loewy RL, Bearden CE, Johnson JK, Raine A, Cannon TD** (2005). The prodromal questionnaire (PQ): preliminary validation of a self-report screening measure for prodromal and psychotic syndromes. *Schizophrenia Research* **79**, 117–125.
- Lovibond PF, Lovibond SH** (1995). The structure of negative emotional states: comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behaviour Research and Therapy* **33**, 335–343.
- Luty SE, Joyce PR, Muldera RT, Sullivan PF, McKenzie JM** (2002). The interpersonal sensitivity measure in depression: associations with temperament and character. *Journal of Affective Disorders* **70**, 307–312.
- McCabe RE, Blankstein KR, Mills JS** (1999). Interpersonal sensitivity and social problem-solving: relations with academic and social self-esteem, depressive symptoms, and academic performance. *Cognitive Therapy and Research* **23**, 587–604.
- McGlashan TH, Addington J, Cannon T, Heinimaa M, McGorry P, O'Brien M, Penn D, Perkins D, Salokangas RKR, Walsh B, Woods SW, Yung A** (2007). Recruitment and treatment practices for help-seeking 'prodromal' patients. *Schizophrenia Bulletin* **33**, 715–726.
- Miller TJ, McGlashan TH, Rosen JL, Somjee L, Markovich PJ, Stein K, Woods SW** (2002). Prospective diagnosis of the initial prodrome for schizophrenia based on the Structured Interview for Prodromal Syndromes: preliminary evidence of interrater reliability and predictive validity. *American Journal of Psychiatry* **159**, 863–865.
- Miller TJ, McGlashan TH, Woods SW, Stein K, Driesen N, Corcoran CM, Hoffman R, Davidson L** (1999). Symptom assessment in schizophrenic prodromal states. *Psychiatric Quarterly* **70**, 273–287.
- Nelson B, Fornito A, Harrison BJ, Yücel M, Sass LA, Yung AR, Thompson A, Wood SJ, Pantelis C, McGorry PD** (2009a). A disturbed sense of self in the psychosis prodrome: linking phenomenology and neurobiology. *Neuroscience and Biobehavioural Reviews* **33**, 807–817.
- Nelson B, Sass LA, Skodlar B** (2009b). The phenomenological model of psychotic vulnerability and its possible implications for psychological interventions in the ultra-high risk ('prodromal') population. *Psychopathology* **42**, 283–292.
- Nelson B, Sass LA, Thompson A, Yung AR, Francey SM, Amminger GP, McGorry PD** (2009c). Does disturbance of self underlie social cognition deficits in schizophrenia and other psychotic disorders? *Early Intervention in Psychiatry* **3**, 83–93.
- Nelson B, Yung AR, Bechdolf A, McGorry PD** (2008). The phenomenological critique and self-disturbance: implications for ultra-high risk ('Prodrome') research. *Schizophrenia Bulletin* **34**, 381–392.
- Otani K, Suzuki A, Ishii G, Matsumoto Y, Kamata M** (2008). Relationship of interpersonal sensitivity with dimensions of the Temperament and Character Inventory in healthy subjects. *Comprehensive Psychiatry* **49**, 184–187.
- Parker JDA, Endler NS, Bagby RM** (1993). If it changes, it might be unstable: examining the factor structure of the Ways of Coping Questionnaire. *Psychological Assessment* **5**, 361–368.
- Parnas J** (2005). Clinical detection of schizophrenia-prone individuals: critical appraisal. *British Journal of Psychiatry* **48**, s111–s112.

- Parnas J** (2011). A disappearing heritage: the clinical core of schizophrenia. *Schizophrenia Bulletin* **37**, 1121–1130.
- Parnas J, Handest J** (2003). Phenomenology of anomalous self-experience in early schizophrenia. *Comprehensive Psychiatry* **44**, 121–134.
- Parnas J, Møller P, Kircher T, Thalbitzer J, Jansson L, Handest P, Zahavi D** (2005). EASE: Examination of Anomalous Self-Experience. *Psychopathology* **38**, 236–258.
- Phillips LJ, Leicester SB, O'Dwyer LE, Francey SM, Koutsogiannis J, Abdel-Baki A, Kelly D, Jones S, Vay C, Yung AR, McGorry PD** (2002). The PACE clinic: identification and management of young people at 'ultra' high risk of psychosis. *Journal of Psychiatric Practice* **8**, 255–269.
- Raballo A, Larøi F** (2011). Psychosis risk syndrome and DSM-5: time for a dimensional approach to at-risk mental states? *Clinical Schizophrenia and Related Psychoses* **5**, 155–158.
- Sass LA, Parnas J** (2003). Schizophrenia, consciousness, and the self. *Schizophrenia Bulletin* **29**, 427–444.
- Sato T, Narita T, Hirano S, Kusunoki K, Sakado K, Uehara T** (2001). Is interpersonal sensitivity specific to non-melancholic depressions? *Journal of Affective Disorders* **64**, 133–144.
- Subotnik KL, Nuechterlein KH** (1988). Prodromal signs and symptoms of schizophrenic relapse. *Journal of Abnormal Psychology* **97**, 405–412.
- Sullivan HS** (1994). The onset of schizophrenia. 1927. *American Journal of Psychiatry* **151**, s135–s139.
- Tait L, Birchwood M, Trower P** (2004). Adapting to the challenge of psychosis: personal resilience and the use of sealing-over (avoidant) coping strategies. *British Journal of Psychiatry* **185**, 410–415.
- Tandon R, Keshavan MS, Nasrallah HA** (2008). Schizophrenia, 'Just the Facts': what we know in 2008 Part 1: Overview. *Schizophrenia Research* **100**, 4–19.
- Trower P, Chadwick PDJ** (1995). Pathways to the defence of the self: a theory of two types of paranoia. *Clinical Psychology: Science and Practice* **2**, 263–278.
- Valmaggia LR, Freeman D, Green C, Garety P, Swapp D, Antley A, Prescott C, Fowler D, Kuipers E, Bebbington P, Slater M, Broome M, McGuire P** (2007). Virtual reality and paranoid ideations in people with an 'at-risk mental state' for psychosis. *British Journal of Psychiatry* **51**, s63–s68.
- Wigman JTW, Lin A, Vollebergh WAM, van Os J, Raaijmakers QAW, Nelson B, Baksheev G, Yung AR** (2011). Subclinical psychosis and depression: co-occurring phenomena that do not predict each other over time. *Schizophrenia Research* **130**, 277–281.
- Wilhelm K, Boyce P, Brownhill S** (2004). The relationship between interpersonal sensitivity, anxiety disorders and major depression. *Journal of Affective Disorders* **79**, 33–41.
- Yung AR, McGorry PD** (1996a). The prodromal phase of first-episode psychosis: past and current conceptualizations. *Schizophrenia Bulletin* **22**, 353–370.
- Yung AR, McGorry PD** (1996b). The initial prodrome in psychosis: descriptive and qualitative aspects. *Australian and New Zealand Journal of Psychiatry* **30**, 587–599.
- Yung AR, Phillips LJ, Yuen HP, Francey SM, McFarlane CA, Hallgren M, McGorry PD** (2003). Psychosis prediction: 12-month follow up of a high-risk ('prodromal') group. *Schizophrenia Research* **60**, 21–32.
- Yung AR, Yuen HP, McGorry PD, Phillips LJ, Kelly D, Dell'Olio M, Francey SM, Cosgrave EM, Killackey E, Stanford C, Godfrey K, Buckby J** (2005). Mapping the onset of psychosis: the Comprehensive Assessment of At-Risk Mental States. *Australian and New Zealand Journal of Psychiatry* **39**, 964–971.