King's Research Portal

DOI:

[10.1111/camh.12034](https://doi.org/10.1111/camh.12034)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Ames, C. S., Richardson, J., Payne, S., Smith, P., & Leigh, E. (2014). Mindfulness-based cognitive therapy for depression in adolescents. *Child and Adolescent Mental Health*, 19(1), 74-78.
<https://doi.org/10.1111/camh.12034>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Innovations in Practice: Mindfulness-based cognitive therapy for depression in adolescents

Catherine S. Ames^{1,3}, Jessica Richardson¹, Susanna Payne², Patrick Smith^{1,3} & Eleanor Leigh^{1,3}

¹National and Specialist Child and Adolescent Mental Health Service, South London and Maudsley NHS Foundation Trust Michael Rutter Centre, Maudsley Hospital, London, SE5 8AZ, UK. E-mail: catherine.ames@kcl.ac.uk

²Bethlem Adolescent Unit, South London and Maudsley NHS Foundation Trust Bethlem Royal Hospital, Beckenham, UK

³Psychology Department, Institute of Psychiatry, King's College London, London, UK. E-mail: catherine.ames@kcl.ac.uk

Background: Mindfulness-Based Cognitive Therapy (MBCT) has been shown to be effective in preventing relapse of depression in adults, but has not previously been applied to adolescents who have residual symptoms of depression following treatment. **Method:** An 8-week MBCT group was adapted for adolescents, and evaluated using qualitative and quantitative measures. **Results:** Participants report high levels of satisfaction with the group intervention. Qualitative analysis of semi-structured interviews provides areas for future development of this intervention. Pilot data indicate reductions in depressive symptoms, alongside positive change in mindfulness skills, quality of life and rumination. **Conclusions:** Preliminary evidence for the use of MBCT with this group of adolescents is provided.

Key Practitioner Message

- It is known that the course of depression from childhood is characterized by relapse and continuity into adulthood.
- There is a need to develop interventions that ameliorate the course of this disorder.
- Mindfulness-based approaches have good evidence in adults in reducing symptoms of depression and in relapse reduction. They have also been seen to be acceptable to children and adolescents, though not yet with a clinically referred group of adolescents with symptoms of depression.
- A group programme of MBCT for adolescents with symptoms of low mood is presented here.
- This research demonstrates the feasibility of this approach.

Keywords: Depression; Mindfulness-based cognitive therapy; adolescents

Introduction

NICE guidelines recommend psychological therapy as the first-line treatment of depression in young people (National Institute for Health and Clinical Excellence, 2005, 2011). However, the evidence for the efficacy of psychological therapy, particularly Cognitive Behavioural Therapy, among children and young people is discordant, and in the main, disappointing (Klein, Jacobs, & Reinecke, 2007; Weisz, McCarty, & Valeri, 2006). Many remain symptomatic at the end of treatment (Wood, Harrington, & Moore, 1996) and even among those who respond to psychological therapies most gains are not maintained at 1-year follow-up (Wood et al., 1996).

Identifying effective interventions to reduce psychological vulnerability to future depressive episodes is a priority in the overall management of the disorder (National Institute for Health and Clinical Excellence, 2005) as the long-term prognosis for young people with depression includes high rates of recurrence (Birmaher, Arbelaez, & Brent, 2002), and continuity of depression into adulthood (Costello et al., 2002). However, although NICE

currently recommends psychological therapy for relapse prevention with young people, the form of this therapy is as yet unspecified. Individual CBT or group-based Mindfulness Behavioural Cognitive Therapy (MBCT) is recommended for adults at significant risk of relapse (National Institute for Health and Clinical Excellence, 2009) and meta-analyses have indicated that MBCT is effective in relapse reduction (Piet & Hougaard, 2011) and in the reduction in mood and anxiety symptoms (Hofmann, Sawyer, Witt, & Oh, 2010). MBCT is proposed to reduce risk in part through increasing awareness of the ruminative cycles and thinking styles associated with relapse vulnerability (Teasdale et al., 2000). Rumination has also been identified in adolescence and research evidence suggests that it may play a comparably toxic role in depression amongst young people (Abela, Brozina, & Haigh, 2002). Rumination as a potential target of treatment points to the potential value of MBCT among this population.

MBCT has been extended and adapted for use with children and adolescents and there is growing evidence of its acceptability and efficacy (for reviews see

Burke, 2009; Greco & Hayes, 2008; Black, Milam, & Sussman, 2009). The evidence base has developed from early case series reports targeting anxiety and rumination in school-based samples (Mendelson et al., 2010; Semple, Reid, & Miller, 2005) to more recent RCTs in a community child sample (Semple, Lee, Rosa, & Miller, 2009) and a sample of adolescents referred for a range of psychiatric difficulties (Biegel, Brown, Shapiro, & Schubert, 2009). These have demonstrated preliminary indications of benefits for a range of emotional and behavioural outcomes. However, there has been no published evaluation of an MBCT group intervention for referred adolescents with residual symptoms of depression.

This study sought to evaluate the feasibility of developing MBCT provision for young people, who had received individual psychological therapy and who continued to have residual symptoms of depression. In addition, the study provided the opportunity to conduct a pilot evaluation of the efficacy of MBCT in reducing symptoms of depression in adolescents, and to measure the impact of MBCT on cognitive factors implicated in depressive relapse. This study has been designed in line with the Medical Research Council Framework for the Development of Complex Interventions (2008) recommendation that feasibility and pilot studies are undertaken prior to controlled studies. The study aims to evaluate the feasibility and acceptability of the qualitative and quantitative measures and of the MBCT intervention to guide future controlled evaluations.

Method

Participants

Young people (aged 12–18), who had received psychological treatment for a mood disorder or low mood alongside an anxiety disorder, but continued to present with residual symptoms of depression were recruited by care coordinators. Those that presented with a high level of risk; who were experiencing an acute episode of depression; with active substance misuse difficulties; or who had not completed one course of psychological therapy were excluded. Young people prescribed antidepressant medication were not excluded from the group, but at the time of intervention none were prescribed.

The data presented here is derived from two MBCT groups. The first was conducted at the Mood Disorder Clinic (National & Specialist CAMHS service, Maudsley Hospital) and the second in Lewisham CAMHS. This research received ethical approval from the South West London REC 3 NHS National Research Ethics Committee (ref: 10/H0803/47). Informed consent to participate in this research was given by both young people and their parents.

Treatment protocol

The full 8-week protocol (see Table 1) was based primarily on the 8 week adult MBCT programme (Williams, Teasdale, Segal, & Kabat-Zinn, 2007). It was also informed by the 12-week MBCT-Children protocol (MBCT-C, Semple & Lee, 2008) and the 8-week adolescent programme developed by Bögels, Hoogstad, van Dun, de Schutter, and Restifo (2008) to ensure it was developmentally appropriate. The protocol used here was designed to make mindfulness as accessible as possible to young people through our choice of simple sensory exercises, for example, the use of visual illusions to illustrate processes of 'describing' and 'judging' as well as directing attention. In separate precourse sessions for participants and parents the rationale of the treatment and attendance and homework expectations were outlined and suitability for MBCT confirmed.

Measures

Measures of acceptability. Acceptability was assessed using semistructured interviews. Transcripts were analysed using Interpretive Phenomenological Analysis (Smith & Osborn, 2003) by one of the group facilitators experienced in this approach (JR). This researcher has a personal and professional interest in MBCT. In line with principles for evaluating qualitative research by Elliott, Fischer, and Rennie (1999) a clinical psychology colleague, unconnected to this project, audited parts of the data by checking that themes corresponded with the raw data. Questions focused on perceived benefits of the intervention, useful and unhelpful aspects of the course, barriers to attendance and ideas for adaptations for future groups.

Measures of efficacy. The effects of the MBCT intervention were examined using a battery of well-validated self-report measures at pretreatment, end-treatment and at 1 month follow-up. These assessed depression (Moods and Feelings Questionnaire; Angold et al., 1995), rumination (Child Response Style Questionnaire; Meiser-Stedman, Dalgleish, Smith, Yule, & Glucksman, 2007), worry (Penn State Worry Questionnaire; Meyer, Miller, Metzger, & Borkovec, 1990), mindfulness skills (Child Acceptance and Mindfulness Measure; Greco, Dew, & Ball, 2005) and perceived quality of life (Paediatric Quality of Life Enjoyment and Satisfaction Questionnaire; Endicott, Nee, Yang, & Wohlberg, 2006) as well as the overall impact of symptoms and emotional distress (Strengths and difficulties Questionnaire; Goodman, 1997).

Results

Eleven (10 female adolescents, 1 male adolescent) young people met inclusion criteria at the time of recruitment. Seven completed the course (7 female adolescents, aged 12–18) and four did not (aged 14–17). Of the seven completers four attended 100% of sessions, two attended 75% and one attended 50%. Three participants left the study in the first 3 weeks, two following a relapse and one because of a complex family situation. The only male adolescent did not specify his reason for non-completion.

Acceptability

Analysis of the semistructured interviews with IPA generated 6 themes, with two superordinate themes. These were conducted with all young people who completed the course.

I. Superordinate Theme: Mindfulness Helps. Theme 1: Increased awareness

All participants talked about increased awareness of thoughts and actions.

It's like your life is in slow motion and you see everything you're doing and understand it more...life is more exciting knowing mindfulness exists (P4).

Theme 2: Relating differently to thoughts and feelings

All participants described decentred awareness enabling them to relate to their thoughts and/or feelings differently.

I try to sort of accept my thought and its there and go back to what I am doing (P5).

Three participants described thinking more positively. Two reported stepping back from strong emotions and consequently enjoying and engaging in life more.

Table 1. MBCT adolescent 8-week session outline

	Aims	Agenda	Exercises	Home practice
1. Tuning in	Introduction and orientation Introducing regular formal practice	Group rules Orientation to mindfulness Information about homework and barriers to practice Mindful daily activities	Eating a strawberry Awareness of the breath: 3-min breathing space	Mindful daily activity 3-min breathing space
2. Two sides to the same coin	Help YP identify thoughts, feelings, behaviour Encouraging present orientation (in the body)	Differentiating Thoughts, Feelings, and Behaviours—Where would MBCT fit it? Introduction to body scan Introduction to pleasant event recording	Body scan	Pleasant events diary Body scan
3. Listen up	Mindfulness in all the senses Full review of present orientation in the body	Mindful listening (holding on/letting go) Review of body scan Introduction of unpleasant event recording	Receptive listening Body scan	Unpleasant and pleasant events diary 3-min breathing space
4. Seeing is believing	Mindfulness in all the senses Continuing regular formal practice	Mindful seeing and smelling Practising directing attention Differentiating judging from describing	Optical illusions Mindful smelling	Pleasant and unpleasant events diary Body scan
5. The five senses	Mindfulness in all the senses Encouraging present orientation and acceptance	Mindful movement Staying in the present, accepting difficult feelings	Mindful walking Exploration of photographs of other people's reactions to an aversive stimulus (bees)	Pleasant and unpleasant events diary Mindful walking
6. Thoughts are not facts	Discriminating thoughts from facts Accepting difficult thoughts/feelings in the present	Noticing and staying with difficult thoughts/feelings in the present	Breath and body sitting meditation Discussion: automatic thoughts	Mindful awareness of negative event 3-min Breathing space Breathing space for coping.
7. Putting mindfulness into practice	Building mindfulness into everyday life	Discuss how to integrate mindfulness into everyday life Review of previous sessions Acceptance of experiences and mindfulness 3-min breathing space to choose what to do. Identifying signs of relapse (action plan)	30 min sitting practice, introduce a difficulty Pleasure/mastery activities Small groups—action plans	Experiment with alternative daily exercises Action plan
8. Moving on mindfully	Concluding	Generalizing mindfulness to everyday life Exploring and sharing reflections Graduation ceremony Stone meditation	Individual 1-month plan	

Note: From week 2 every session started with a breathing space and enquiry and finished with another breathing space without enquiry.

I've been thinking don't panic if you haven't done every bit of homework, you go into school and at least you don't miss more stuff (P2).

Theme 3: Using mindfulness when distressed
Most participants talked about mindfulness as a tool to manage distress.

Before if I was feeling in a low mood then I wouldn't want to do anything... but now after the mindfulness... I think about all the things we talked about and I put on the breathing space and it makes me feel a bit better (P7).

However, some of the same young people said that when feeling very distressed they did not believe mindfulness would work and therefore did not practice.

II. Superordinate Theme: The Group Experience. Theme 4: Working it out together

Three participants gave feedback on the size of the group, with two finding a smaller group more comfortable. Two participants thought the age range was too broad. Several participants said they enjoyed being in the group and sharing the experience with others.

It was nice listening to others' problems and how I could relate to them was useful (P6).

Theme 5: Formal meditation versus simple sensory exercises

Three participants reported finding formal meditation too long or difficult.

I find I sometimes get a bit lost in the body scan, maybe I don't concentrate so much (P2).

Table 2. Descriptive statistics for outcome measures

Measure Mean (SD)	Pre n = 7	End n = 6	Follow-up n = 6	Pre-end d	Pre-FU d
MFQ	36.57 (11.18)	27.17 (8.52)	23.67 (9.52)	.79	.77
CRSQ	35.29 (11.95)	31.83 (8.61)	24.33 (13.17)	.22	.48
PSWQ	22.86 (8.28)	23.17 (7.41)	19.50 (5.13)	.02	.41
CAMM	46.14 (9.53)	47.33 (3.56)	49.00 (4.90)	.11	.33
PQ-LES-Q	35.29 (10.69)	42.33 (3.98)	44.83 (8.52)	.59	.56
SDQ emotional	7.57 (1.90)	7.33 (1.21)	6.50 (1.38)	.15	.35
SDQ impact	5.86 (1.22)	5.00 (2.00)	3.33 (1.21)	.51	1.13

MFQ, Moods and Feelings Questionnaire; CAMM, Child Acceptance and Mindfulness Measure; PSWQ, Penn State Worry Questionnaire; CRSQ, Child Response Style Questionnaire; PQ-LES-Q, Paediatric Quality of Life Enjoyment and Satisfaction Questionnaire; SDQ, Strengths and difficulties Questionnaire.

There were several comments about not prioritizing or remembering home practice. However, all participants said they most enjoyed the sensory exercises and reported applying these to their everyday life.

Theme 6: Keeping mindfulness going

There was a wish to keep going with 'mindfulness' alongside recognition that this was challenging. Three participants wanted more group sessions and one wanted a longer term plan.

Even when I become aware of the fact that I want to be mindful at a particular point it's really hard to keep that going (P3).

Efficacy

Analysis of outcome measures is limited by the small number of participants. Effect sizes of change in measures are reported in Table 2. Large effect sizes are seen in the reduction in depression and the impact of symptoms. Modest decreases in worry and rumination, increase in quality of life, and a smaller increase in mindfulness are evident.

Discussion

This study has indicated that MBCT with adolescents, targeted at symptoms of low mood, is feasible. The retention rate is reasonable and attendance rates were good. The qualitative assessment indicates that MBCT was acceptable to this group. Participants reported enjoying the group experience. Participants were positive about the premise of MBCT and the material and format of the sessions. They valued the range of material, indicating that the combination of protocols was appropriate. Following the CONSORT Group (Consolidated Standards of Reporting Trials, Moher, Schulz, and Altman 2001) guidance for clinical trials it is clear that at this stage, the primary outcome for this study is feasibility. The secondary findings reported below are offered as indications of areas for further development in future research.

Participants described how mindfulness had benefited them through increasing their awareness and facilitating distance from thoughts and strong emotions. Participants found the 3-min breathing space helpful, requested having a longer course and support in continuing their practice. However, they also indicated difficulties with home practice and longer formal meditations. The development of mindfulness relies on the encountering of one's changing mind states, recognizing them and

disengaging from them (Segal, Williams, & Teasdale, 2002). These can be encountered in regular meditation and also 'practiced in all situations' (Segal et al., 2002, p. 76). When working with adolescent groups greater emphasis might be put on practice in everyday life.

Our analysis of the efficacy of MBCT for this group is limited by the small number of participants and the absence of a control group. It is also limited by the use of self-report measures and the absence of triangulation of information from other informants. However, we can tentatively conclude that the intervention was associated with a reduction in symptom and impact scores and an increase in perceived quality of life.

These conclusions are made with a significant caveat, which is that with this low number of participants, the absence of a control group or full assessment of non-completers, and follow-up only at 1 month, the data may reflect natural recovery rather than a meaningful and lasting response to treatment. Furthermore, future research might also consider the impact of mindfulness intervention with a greater number of male participants (given the predominance of female adolescents in this study) and deploy measures of treatment integrity. Therefore, these encouraging findings are seen to provide justification for future research and are informative about how best to adapt the adult MBCT course to suit the needs of young people.

Acknowledgements

This study received no external funding support. The authors thank Mark Williams for his generosity in encouraging us to use the adult MBCT protocol and materials, Jane Hutton for her support and encouragement with the development of the protocol, Wendy Geraghty for clinical supervision and Adenique Galloway for conducting and transcribing interviews. The authors have declared that they have no competing or potential conflicts of interest.

References

- Abela, J.R.Z., Brozina, K., & Haigh, E.P. (2002). An examination of the response styles theory of depression in third- and seventh-grade children: A short-term longitudinal study. *Journal of Abnormal Child Psychology*, 30, 515–527.
- Angold, A., Costello, E.J., Messer, S.C., Pickles, A., Winder, F., & Silver, D. (1995). The development of a short questionnaire for use in epidemiological studies of depression in children and adolescents. *International Journal of Methods in Psychiatric Research*, 5, 237–249.
- Biegel, G.M., Brown, K.W., Shapiro, S.L., & Schubert, C.A. (2009). Mindfulness-Based Stress Reduction for the Treat-

- ment of Adolescent Psychiatric Outpatients: A Randomized Clinical Trial. *Journal of Consulting and Clinical Psychology*, 77, 855–866.
- Birmaher, B., Arbelaez, C., & Brent, D. (2002). Course and outcome of child and adolescent major depressive disorder. *Child and Adolescent Psychiatric Clinics of North America*, 11, 619–637.
- Black, D.S., Milam, J., & Sussman, S. (2009). Sitting-meditation interventions among youth: A review of treatment efficacy. *Pediatrics*, 124(3), e532–e541.
- Bögels, S., Hoogstad, B., van Dun, L., de Schutter, S., & Restifo, K. (2008). Mindfulness training for adolescents with externalizing disorders and their parents. *Behavioural and Cognitive Psychotherapy*, 36, 193–209.
- Burke, C.A. (2009). Mindfulness-based approaches with children and adolescents: A preliminary review of current research in an emergent field. *Journal of Family and Child Studies*, doi:10.1007/s10826-009-9282-x.
- CONSORT Group (Consolidated Standards of Reporting Trials), Moher, D., Schulz, K.F., & Altman, D.G. (2001). The CONSORT statement: Revised recommendations for improving the quality of reports of parallel-group randomized trials. *Journal of the American Podiatric Medical Association*, 91, 437–442.
- Costello, E.J., Pine, D.S., Hammen, C., March, J.S., Plotsky, P.M., Weissman, M.M., ... & Leckman, J.F. (2002). Development and natural history of mood disorders. *Biological Psychiatry*, 52, 529–542.
- Elliott, R., Fischer, C.T., & Rennie, D.L. (1999). Evolving guidelines for publication of qualitative research studies in psychology and related fields. *British Journal of Clinical Psychology*, 38, 215–229.
- Endicott, J., Nee, J., Yang, R., & Wohlberg, C. (2006). Paediatric quality of life enjoyment and satisfaction questionnaire (PQLES-Q): Reliability and validity. *Journal of the American Academy of Child and Adolescent Psychiatry*, 45, 401–407.
- Goodman, R. (1997). The strengths and difficulties questionnaire: A research note. *Journal of Child Psychology and Psychiatry*, 38, 581–586.
- Greco, L.A., Dew, S.E., & Ball, S. (2005). Acceptance, mindfulness, and related processes in childhood: Measurement issues, clinical relevance, and future directions. In S.E. Dew & R. Baer (Chairs), *Measuring acceptance, mindfulness, and related processes: Empirical findings and clinical applications across child, adolescent, and adult samples*. Washington, DC: Symposium presented at the Association for Behavior and Cognitive Therapies.
- Greco, L.A., & Hayes, S.C. (2008). *Acceptance and mindfulness interventions for children, adolescents, and families*. Oakland, CA: Context Press/New Harbinger Publications.
- Hofmann, S.G., Sawyer, A.T., Witt, A.A., & Oh, D. (2010). The effect of mindfulness-based therapy on anxiety and depression: A meta-analytic review. *Journal of Consulting and Clinical Psychology*, 78, 169–183.
- Klein, J.B., Jacobs, R.H., & Reinecke, M.A. (2007). Cognitive-behavioural therapy for adolescent depression: A meta-analytic investigation of changes in effect-size estimates. *Journal of the American Academy of Child and Adolescent Psychiatry*, 46, 1403–1413.
- Medical Research Council (2008). *Developing and evaluating complex interventions: New guidance*. London: Medical Research Council.
- Meiser-Stedman, R., Dalgleish, T., Smith, P., Yule, W., & Glucksman, E. (2007). Diagnostic, demographic, memory quality, and cognitive variables associated With acute stress disorder in children and adolescents. *Journal of Abnormal Psychology*, 116, 65–79.
- Mendelson, T., Greenberg, M.T., Dariotis, J.K., Gould, L.F., Rhoades, B.L., & Leaf, P.J. (2010). Feasibility and preliminary outcomes of a school-based mindfulness intervention for urban youth. *Journal of Abnormal Child Psychology*, 38, 985–994.
- Meyer, T.J., Miller, M.L., Metzger, R.L., & Borkovec, T.D. (1990). Development and validation of the Penn State Worry Questionnaire. *Behavioural Research and Therapy*, 28, 487–495.
- National Institute for Health and Clinical Excellence (2005). *Depression in Children and Young People*. NICE. Available at: <http://www.nice.org.uk/nicemedia/pdf/CG028NICEguideline.pdf> (accessed 15 January 2010).
- National Institute for Health and Clinical Excellence (2009). *Depression*. NICE. Available at: <http://www.nice.org.uk/nicemedia/pdf/CG90NICEguideline.pdf> (accessed 15 January 2010).
- National Institute for Health and Clinical Excellence (2011). *CG28 Depression in children and young people: review decision*. NICE. Available at: <http://guidance.nice.org.uk/CG28/ReviewDecision/pdf/English> (accessed 8 August 2011).
- Piet, J., & Hougaard, E. (2011). The effect of mindfulness-based cognitive therapy for prevention of relapse in recurrent major depressive disorder: A systematic review and meta-analysis. *Clinical Psychology*, 31, 1032–1040.
- Segal, Z.V., Williams, J.M.G., & Teasdale, J.D. (2002). *Mindfulness-based cognitive therapy for depression: A new approach to preventing relapse*. New York: Guilford Press.
- Semple, R.J., & Lee, J. (2008). Treating anxiety with mindfulness: Mindfulness-based cognitive therapy for children. In L.A. Greco & S.C. Hayes (Eds.), *Acceptance and mindfulness interventions for children, adolescents, and families* (pp. 94–134). Oakland, CA: Context Press/New Harbinger Publications.
- Semple, R.J., Lee, J., Rosa, D., & Miller, L.F. (2009). A randomized trial of mindfulness-based cognitive therapy for children: Promoting mindful attention to enhance social-emotional resiliency in children. *Journal of Child and Family Studies*, doi:10.1007/s10826-10009-19301-y.
- Semple, R.J., Reid, E.F., & Miller, L. (2005). Treating anxiety with mindfulness: An open trial of mindfulness training for anxious children. *Journal of Cognitive Psychotherapy*, 19, 379–392.
- Smith, J.A., & Osborn, M. (2003). Interpretive phenomenological analysis. In J.A. Smith (Ed.), *Qualitative Psychology: A practical guide for research methods* (pp. 51–80). London: Sage.
- Teasdale, J.D., Segal, Z.V., Williams, J.M.G., Ridgeway, V.A., Soulsby, J.M., & Lau, M.A. (2000). Prevention of relapse/recurrence in major depression by mindfulness-based cognitive therapy. *Journal of Consulting and Clinical Psychology*, 68, 615–623.
- Weisz, J.R., McCarty, C.A., & Valeri, S.M. (2006). Effects of psychotherapy for depression in children and adolescents: A meta-analysis. *Psychological Bulletin*, 132, 132–149.
- Williams, J., Teasdale, J., Segal, Z., & Kabat-Zinn, J. (2007). *The mindful way through depression, freeing yourself from chronic unhappiness*. New York: Guilford Press.
- Wood, A., Harrington, R., & Moore, A. (1996). Controlled trial of a brief cognitive-behavioural intervention in adolescent patients with depressive disorders. *Journal of Child Psychology and Psychiatry*, 37, 737–746.

Accepted for publication: 31 May 2013
Published online: 28 August 2013