King's Research Portal

DOI:

[10.1016/j.foodchem.2020.127671](https://doi.org/10.1016/j.foodchem.2020.127671)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Khoja, K. K., Aslam, M. F., Sharp, P. A., & Latunde-dada, G. O. (2021). In vitro bioaccessibility and bioavailability of iron from fenugreek, baobab and moringa. *FOOD CHEMISTRY*, 335, 127671. [127671]. <https://doi.org/10.1016/j.foodchem.2020.127671>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

1
2
3 **In Vitro Bioaccessibility and Bioavailability of Iron from**
4 **Fenugreek, Baobab and Moringa**

5 Kholoud K. Khoja, Mohamad F. Aslam, Paul A. Sharp, and Gladys O. Latunde-Dada*

6 Department of Nutritional Sciences, School of Life Course Sciences, King's College London. Franklin-Wilkins-Building,
7 150 Stamford Street, London, SE1 9NH, UK.

8 * Corresponding Author: yemisi.latunde-dada@kcl.ac.uk.

9
10
11
12 **Abstract**

13 Iron deficiency anaemia (IDA) is a common nutritional disorder worldwide. Sustainable food-based
14 approaches are being advocated to use high and bioavailable dietary iron sources to prevent iron
15 deficiency. The study investigated the bioaccessibility and bioavailability of iron from some plant
16 products.

17 Total iron levels in the samples were measured by inductively coupled plasma optical emission
18 spectrometry (ICP-OES). Fractionation of the iron from the digested extracts was carried out by
19 centrifugation and ultrafiltration. Iron bioavailability was determined using an in vitro simulated
20 peptic-pancreatic digestion, followed by measurement of ferritin in Caco-2 cells.

21 The highest amount of bioaccessible iron was obtained from moringa leaves ($9.88\% \pm 0.45$ and 8.44
22 ± 0.01 mg/100 g), but the highest percentage bioavailability was from baobab fruit pulp ($99.7\% \pm 0.13$
23 and 1.74 ± 0.01 mg/100 g) respectively. All the plant products, except for baobab, significantly
24 inhibited iron uptake from FeSO_4 and FAC, with fenugreek sprout being the most inhibitory.

25
26 **Key words: plants, in vitro, iron, availability, Caco-2**
27
28

1. Introduction

Iron deficiency anaemia (IDA) is a nutritional disorder affecting several population groups in the world (WHO). This disorder is common amongst vulnerable infants, adolescent girls, pregnant women and the elderly in most countries. The increased physiological requirements of iron for growth and reproduction, within these population groups, is exacerbated by low intake and poor iron bioavailability from foods (Aspuru, Villa, Bermejo, Herrero, & López, 2011). This is particularly evident in populations subsisting predominantly on vegetables or plants for their sources of iron. Substantial evidence has revealed that IDA has deleterious consequences on cognition, mental function, work performance, and pregnancy outcomes (Muñoz & Humeres, 2012). Consequently, iron supplementation and fortification of staple foods in different countries (Gera, Sachdev, & Boy, 2012) have been practical approaches to alleviate this critical nutritional disorder.

However, oral iron supplements are associated with gastrointestinal irritations and inflammation, while highly bioavailable iron salts used in food fortification cause adverse sensory changes in foods (Saini, Manoj, Shetty, Srinivasan, & Giridhar, 2014). Food fortification also poses a significant challenge in developing economies as it relies on mass processing and distribution of staple foods (Gómez-Galera et al., 2010). Consequently, sustainable food-based approaches are being advocated to increase the intake of food with high iron content and bioavailability (Bouis & Saltzman, 2017). In general, dietary guidelines for the sustainability of the planet advocate the consumption of predominantly plant foods to meet the nutritional requirements of the world's population (Lang & Barling, 2012). Recently, the EAT-Lancet Commission on Food, Planet, Health submitted that 'A diet rich in plant-based foods and with fewer animal source foods confers both improved health and environmental benefits' and assures sustainability of natural resources' (Willett et al., 2019). Consequently, there is a surge in the use of plant parts and plant products in health care and medicinal purposes (Mbah, Eme, & Ogbusu, 2012).

In recent years, several tropical and sub-tropical plants have gained popularity in Western countries in health food shops as novel foods with nutritional potential as sources of minerals and vitamin C (Magaia, Uamusse, Sjöholm, & Skog, 2013). The legume plant, *Trigonella foenum-graecum* (Fenugreek) has long been used for culinary and medicinal purposes in countries such as India, China, Egypt and across the Middle East where it is widely cultivated (Petropoulos, 2003). Fenugreek seeds and leaves have been reported as good sources of iron, calcium and zinc (Goswami, 2012). Baobab, *Adansonia digitata*, a tropical African fruit, is used for food and beverages and medical purposes. However, it is also inherently high in polyphenols (Coe, Clegg, Armengol, & Ryan, 2013) and dietary fibre (Magaia et al., 2013), which are natural inhibitors of mineral absorption. *Moringa oleifera* (moringa) is reputed as a herb that cures a myriad of ailments and diseases which include anaemia,

asthma and skin infections in Africa, South East Asia and South America. Moringa leaves contain high levels of minerals, vitamins and phytochemicals (Anwar et al., 2007). The hematinic potential of moringa leaves has been reported to be comparable with ferric citrate in haemoglobin (Hb) repletion study in rats (Saini et al., 2014). Consequently, this current study has evaluated levels of iron in fenugreek, baobab and moringa samples and also determined the release of soluble iron following *in vitro* digestion and iron uptake in Caco-2 cells.

2. Materials and Methods

2.1. Reagents and Chemicals

Unless otherwise stated, all the reagents and chemicals used in this study were purchased from Sigma-Aldrich Ltd. (Dorset, U.K.). Solutions of enzymes were all prepared freshly just before use.

2.2. Plant Samples

Fenugreek sprout and fenugreek seeds were acquired from Plant Organic UK. Baobab fruit pulp powder was purchased from Aduna Ltd., London. Moringa dried leaf powder was acquired from Mother Nature's Garden Co., London.

2.3. Moisture Analysis

The moisture content of the samples was determined according to the AOAC (2002) method. Briefly, 2 g of samples were weighed and placed in an oven (Gallenkamp model IH-100) at 100 °C to dry for 24–48 h until constant weights were achieved. Afterwards, the percentage of moisture content was calculated for each sample. Samples (50 g) of each vegetable were ground in a classic Moulinex AR1043 grinder to fine powders and stored in sealed bags at –70 °C before analysis.

2.4. Determination of Mineral Content in Plant Products

Samples were processed using the MARS 6 Microwave digestion system. Samples (0.5 g of starting material or 5 ml of digest) and 5 ml of 15.8 M nitric acid were added into reaction vessels and placed into the microwave digester. Digestion of the samples was carried out for an hour. The contents were then transferred into Falcon tubes containing 140 µl of 100 ppm yttrium internal standard, and the volume was made to 14 ml with deionised water. Iron in the samples was read using the inductively coupled plasma optical emission spectrometry - ICP-OES (Thermo ICAP 6000).

Fractionation of the bioaccessible iron into percentages, and low-molecular-weight Fe, in the digested extracts, were carried out by centrifugation (Eppendorf microcentrifuge 5417) and ultrafiltration as described by Powell et al. (2014). Aqueous suspensions (0.5 ml) were centrifuged (110 g, 5 min), and the supernatant represents the total bioaccessible iron (TBF) released during *in vitro* digestion. To separate the low-molecular-weight Fe fraction (LMW), a fraction of the supernatant was ultrafiltered through AMICON ULTRA 3 kDa molecular weight cut-off columns (Merck-UFC500396) (110 g rpm, 5 min). Iron concentrations of samples were determined in ICP-OES (Thermo ICAP 6000). The TBF and the LMW were calculated as follows:

$$[(\%) \text{ Total bioaccessible Fe}] = [(\text{Fe}_{\text{supernatant after digestion}}) / \text{Total Fe}] \times 100$$

$$[(\%) \text{ Fe low-molecular-weight fraction}] = [(\text{Fe}_{\text{ultrafiltrate}}) / \text{Total Fe}] \times 100$$

2.5. Peptic-pancreatic *in vitro* digestion

Simulated gastrointestinal digestion was performed on the samples using a procedure described previously (Glahn, Lee, Yeung, Goldman, & Miller, 1998). Briefly, in dark tubes, 0.5 g samples were mixed with 10 ml of saline solution (140 mmol/L NaCl and 5 mmol/L KCl) and left for 5 min. Then, the pH was adjusted to 2.0, using 1 M HCl. Afterwards, 0.5 ml of pepsin (Sigma-P7000) (16 mg/mL) was added. Samples were incubated at 37°C on a rocking platform (150 rpm) for 75 min. Following this, the pH of the samples was adjusted to pH 5.5 using solid NaHCO₃. Bile extract and pancreatin (8.5 mg/ml bile extract and 1.4 mg/ml pancreatin) were added, and the pH adjusted again to pH 7.0. The solution was made up to 30 ml with saline solution, and the samples were incubated at 37°C for 2 hours. At the end of the incubation period, samples were centrifuged at 5000 rpm for 10 min, and the supernatants were decanted and used for the determination of TBF and LMW fractions. Furthermore, the digested extracts were applied to Caco-2 cells to estimate iron uptake.

2.6. Phytic Acid Analysis

Phytic acid content (total phosphorus) was measured by using a kit (Megazyme- K-PHYT, Bray, Ireland) using the protocol described by the manufacturer (McKie & McCleary, 2016). Briefly, acid extracts of inositol phosphates from the samples were digested with phytase and alkaline phosphatase suspension was used to release phosphate from all the myo-inositol phosphate forms. The total phosphate released was measured using a modified colourimetric method, and was calculated as grams of phosphorus per 100 g of sample material.

123 **2.7. Cell culture**

124 Human Caco-2 cell line was obtained from American Type Culture Collection [ATCC] at passage
125 40 and used in experiments at passage 45. Cells were sub-cultured in a 75 cm² flask to 70-80%
126 confluence. The growth medium contained Dulbecco's Modified Eagle Medium [DMEM], high
127 glucose with glutamine , 10 % fetal calf serum (FBS), 1 % Penicillin-streptomycin (100X) , 1 %
128 L-glutamine (100X) and 1 % MEM non-essential amino acids in an incubator at 37°C, 5% CO₂
129 and 95% oxygen.

130 **2.8. Cell Viability Studies**

131 Caco-2 cells were seeded at a density of 1×10^4 cells/cm² in 96-well plates. After 14 days of
132 differentiation, the medium was discarded, and the cells were washed twice with sterile phosphate
133 buffer saline [PBS] and then incubated with 100 µl of the digested extracts of fenugreek sprouts,
134 seeds, baobab or moringa for 2 h. Following this, 100 µl of fresh Modified Eagle's medium
135 [DME] along with 10 µl of Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide [MTT]
136 sterile solution (5 mg/ml MTT in PBS) were added to each well. After incubating for 3 h in the
137 dark at 37 °C, 100 µl of a solubilisation buffer in dimethyl sulfoxide [DMSO] was added and
138 incubated for 15 minutes at room temperature. To determine the MTT reaction in the cells, optical
139 density was read in a microplate reader (Bio-Tek ELx800) at 490 nm. Cell viability was expressed
140 as a percentage of the controls.

141 **2.9. Iron availability**

142 Human Caco-2 cell line was used to evaluate the bioavailability of iron. These cells are derived
143 from colon adenocarcinoma and are used as a surrogate for enterocytes in the small intestine. This
144 model compares well with human studies and is commonly used to analyse iron bioavailability
145 from various food types. (Glahn et al., 1998). Caco-2 cells were trypsinised and cultured in 6-
146 well plates for 14 days to allow them to differentiate, and the medium was changed every 2 days.
147 Before experiments, cells in 6-well plates were treated with 2 ml serum-free medium [SFM] MEM
148 for 24 hours. Sample digests were centrifuged and heated at 100°C for 5 min to inactivate the
149 digestive enzymes. Serum-free medium (1 ml) was added to cells, followed by the addition of 1
150 ml of sample digest, and these were incubated in a rotating shaker for 2 h. Following this, 1 ml of
151 MEM was added, and samples were incubated at 37°C for a further 22 hours for ferritin synthesis.
152 After the incubation, the culture medium was discarded, and cells were washed with versene (PBS
153 + EDTA). Afterwards, 100 µl of mammalian protein extraction reagent (MPER, Thermo
154 Scientific, UK) was added to wells and left on a shaker for 15 min for cell lysis.

Ferritin ELISA kit, Spectro Ferritin MT (Ramco Laboratories Inc., USA) was used to determine ferritin content in the cells according to the manufacturer's protocol.

2.10. Statistical Analysis

Experiments were performed in 3-6 replicates and data are shown as mean $\pm$ standard error of the means. Comparisons of iron content, solubility and ferritin concentrations in Caco-2 cells were analysed using one-way or two-way ANOVA followed by Tukey's post hoc test where appropriate, using GraphPad Prism software. The significance level was at $P \leq 0.05$.

3. Results

3.1. Mineral content and moisture in plant products

Levels of iron (and other minerals) in the plant samples were measured by ICP-OES (Supplementary Table 1). Iron content was highest in moringa (85.44 mg/100 g) and lowest in baobab (1.67 mg/100 g). Fenugreek sprouts (19.85 mg/100 g) contained more than twice the amount of iron present in fenugreek seeds (7.75 mg/100 g). Moisture content did not vary between plant food samples (range 92.5 – 95.7 %).

3.2. The bioaccessible and fractional low-molecular-weight iron content of the digested extracts of the vegetables

Percentage of TBF and LMW fraction are shown in Table 1. Although baobab fruit pulp had the highest TBF (99.7%), the absolute amount of Fe from baobab was the lowest (1.74 mg/ 100 g). In contrasts, the percentage of TBF in moringa was the lowest, but the absolute quantity LMW was the highest (Table 1). Percentage of fenugreek seeds LMW Fe was significantly ($P \leq 0.05$) higher than the sprouts; nevertheless, the quantity of Fe in both products was comparable. Interestingly, the trend of the LMW fractions from the plant products was similar to the TBF.

Table 1: Total bioaccessible and low-molecular-weight iron in digest samples.

Sample	Bioaccessible	Low-Molecular-Weight
	%	%
Fenugreek sprout	17.18 $\pm$ 0.04 (3.41 $\pm$ 0.01 mg/100 g) a	9.16 $\pm$ 0.09 (1.81 $\pm$ 0.01 mg/100 g) a
Fenugreek Seeds	42.71 $\pm$ 0.22 (3.31 $\pm$ 0.02 mg/100 g) b	25.18 $\pm$ 0.14 (1.95 $\pm$ 0.02 mg/100 g) b
Baobab fruit pulp	99.7 $\pm$ 0.13 (1.74 $\pm$ 0.01 mg/100 g) c	97.8 $\pm$ 0.01 (1.70 $\pm$ 0.01 mg/100 g) c

Moringa leaves	9.88±0.45 (8.44±0.01 mg/100 g)	d	1.96±0.04 (1.68±0.03 mg/100 g)	d
-----------------------	-----------------------------------	---	-----------------------------------	---

Data are presented as means ± SEM of n = 3 measurements. Different letters indicate significant difference between groups (P≤0.05). Mean values represent total bioaccessible iron for each sample category while the lower type phase in bracket are the absolute quantities of iron in the samples.

3.3. The total bioaccessible and fractional low-molecular-weight iron content of the digested extracts of the vegetables with added ascorbic acid

Adding ascorbic acid (AA) to the plant products during digestion decreased TBF and the LMW (Table.2) significantly (P≤0.05) compared with the digested sample without AA. Percentage TBF was the highest from fenugreek seeds. Moringa was the lowest of the plant products in TBF percentage although moringa yielded the highest absolute Fe. The LMW fractions from the plant digests with added AA varied significantly, and the relative proportions differ from the TBF. However, enhancing the effect of ascorbic acid became evident only when it was added along with the digests to Caco-2 cells during the iron uptake study.

Table 2: Total bioaccessible and low-molecular-weight iron when ascorbic acid was added during the digestion of the samples.

Sample	Bioaccessible		Low-Molecular-Weight	
	%		%	
Fenugreek sprout	15.4±0.13 (3.06±0.026 mg/100 g)	a	1.6±0.03 (0.32±0.007 mg/100 g)	a
Fenugreek Seeds	24.3±0.11 (1.88±0.009 mg/100 g)	b	2.7±0.09 (0.21±0.008 mg/100 g)	b
Baobab fruit pulp	19.2±0.9 (0.32±0.015 mg/100 g)	c	15.1±1.15 (0.25±0.019 mg/100 g)	c
Moringa leaves	8.1±0.02 (6.88±0.016 mg/100 g)	d	0.2±0.01 (0.16±0.009 mg/100 g)	d

Data are presented as means ± SEM of n = 3 measurements. Different letters indicate significant difference between groups (P≤ 0.05). Mean values represent total bioaccessible iron for each sample category while the lower type phase in bracket are the absolute quantities of iron in the samples.

3.4. Cell viability of Caco-2 cells after exposure to digested vegetable samples

To ascertain whether the digested extracts from the plants were cytotoxic to Caco-2 cells; the MTT viability assay was performed. The application of the boiled samples of the digested extracts of fenugreek sprouts and seeds, baobab or moringa, did not adversely affect the viability of Caco-2 cells. Indeed, in some cases, there was a significant effect on cell growth as judged by an increase in viability (Supplementary Figure 3).

211
212
213
214
215
216
217
218

219

220
221
222
223
224

225
226
227
228
229

230

231

3.5. *In vitro* accessibility of iron from fenugreek sprouts, seeds, baobab or moringa in Caco-2 cells

To estimate the accessibility of iron from the samples, an *in vitro* simulated peptic–pancreatic digestion was carried out followed by ferritin analysis (a surrogate marker for iron absorption) in Caco-2 cells. In contrast to the bioaccessible iron profile of the samples, fenugreek sprouts exhibited comparatively higher iron available ($P \leq 0.05$) than the seeds (Figure 1). Iron availability from the baobab fruit pulp was the lowest. The absorption of FeSO_4 and FAC served as positive controls.

Figure 1: Iron uptake by Caco-2 cells from plant food digests. “Control” indicates digest system components only (i.e., pepsin, pancreatin, bile extract). Values are presented as means $n = 6 \pm \text{SEM}$. Data were analysed using a two-way ANOVA. Different letters indicate significant difference between groups ($P \leq 0.05$).

To enhance iron extraction during the peptic-pancreatic digestion of the plant products, ascorbic acid was added to the samples during the digestion process. Iron availability from fenugreek sprouts, fenugreek seeds, baobab or moringa was not significantly influenced when ascorbic acid was included in the digestion medium (Figure 2). Ascorbic acid, however, enhanced iron availability from the plant products when added to the digest during exposure to Caco-2 cells during the iron uptake stage.

Figure 2. Iron uptake by Caco-2 cells from vegetable samples with added ascorbic acid (AA) during digestion (a) and (b.) vegetable samples with AA added during exposure to cells. The control sample contained 50 μ M FeSO₄ and 250 μ M AA. Values are presented as means $\pm$ SEM, n = 6. Data were analysed using a two-way ANOVA. Different letters indicate significant difference between vegetables ($P \leq 0.05$). The differences between groups (a) and (b) are denoted (* $P \leq 0.05$, *** $P \leq 0.001$ and **** $P \leq 0.0001$).

3.6. Modulating effects of fenugreek sprouts, fenugreek seeds, baobab or moringa on iron accessibility from iron salts in Caco-2 cells

We next explored the interactions of the plant products on the accessibility of iron salts. FeSO₄ and FAC had comparable iron availability for uptake in Caco-2 cells (Figure 3). Except for baobab fruit pulp, the other plant products significantly ($P \leq 0.05$) reduced iron availability from both FeSO₄ and FAC in Caco-2 cells. Moreover, inhibition of iron from FAC (Fe(III)) was significantly higher than from FeSO₄ (Fe(II)) for fenugreek seeds ($P \leq 0.01$) and moringa leaves ($P \leq 0.0001$) respectively.

253

254 **Figure 3:** Iron uptake by Caco-2 cells from vegetable digests. Cells were exposed to the digest
255 samples with added FeSO₄ or FAC (50 μ M). Values are presented as means $\pm$ SEM, n = 6. Data were
256 analysed using a two-way ANOVA between FeSO₄ and FAC groups and one-way ANOVA amongst
257 the treatment groups. Different letters indicate significant difference between vegetables ($P \leq 0.05$).
258 The differences between added FeSO₄ and added FAC groups are denoted (* $P \leq 0.05$, *** $P \leq 0.001$
259 and **** $P \leq 0.0001$).

260

261 **3.7. Phytic acids levels in fenugreek sprouts, seeds, baobab and moringa**

262 Phytic acid is a major inhibitory component in plant foods. Figure 4 shows a significant level of
263 variations in the phytate content of the plant products. Both fenugreek samples exhibited the highest
264 levels of phytic acid and may in part explain why both plant products inhibited FeSO₄ and FAC
265 uptake by Caco-2 cells the most (Figure 4).

Figure 4: Phytic acid content (g/100 g) of the vegetable samples. Values are presented as means $\pm$ SEM, n=6. Data were analysed using a two-way ANOVA test. Different letters indicate significant difference between groups ($P \leq 0.05$).

4. Discussion

A sustainable food-based approach to tackle iron deficiency implies the consumption of plant foods that have high iron levels and that have been processed to reduce inhibitors and facilitate the natural release of enhancers of iron bioavailability in meals (Gibson, Perlas, & Hotz, 2006). The current study, therefore, investigated the potentials of fenugreek sprouts, fenugreek seeds, baobab fruit pulp, and moringa leaves as sources of minerals and of accessible iron using *in vitro* methods of analysis.

The levels of iron varied between these plant products and it was clear that they could contribute to daily intake significantly depending on dietary formulations. The differences in the mineral levels from plant products, when compared to published data could be accounted for by factors such as the seed variety, geographical location, processing types, storage conditions and analytical methods (Aslam, Anwar, Nadeem, Rashid, Kazi, & Nadeem, 2005; Chadare, Linnemann, Hounhouigan, Nout, & Van Boekel, 2008).

The iron content of moringa in the current study is higher than the published data (Price, 2007). The mineral profile of moringa leaves has been reported to be superior nutritionally to those of amaranth, mushrooms, taro leaves, cassava leaves and pumpkin seeds (Owusu, Ellis, & Oduro, 2008). All daily

287 calcium and 75% of iron requirements could be met in children aged 1-3 years from 100 g of fresh
288 moringa leaves (Price, 2007). In comparison, an equivalent weight of baobab pulp will provide 30%
289 of calcium and 23% of iron daily needs in children 4-13 years, and about 29% of the calcium
290 requirement of pregnant women (Magaia et al., 2013). Furthermore, van der Merwe et al. (2019)
291 reported that fortification of pearl millet in a cereal-based meal with moringa leaf powder, roselle
292 calyces or baobab fruit pulp contributed about 3 times more iron and zinc than from pearl millet
293 porridge fortified with provitamin A source alone. The added plant food fortificants contributed about
294 28% and 41% of the absolute iron and zinc requirements, respectively, of women of reproductive
295 age from a single meal. A recent study (Shija, Rumisha, Oriyo, Kilima, & Massaga, 2019) also
296 reported that daily supplementation of 25 g moringa flour to food intake for 6 months significantly
297 reduced anaemia prevalence in children whose average age was about one year old. Improvement in
298 haemoglobin levels for these children were significant irrespective of the degree of anaemia with
299 moringa intervention plus nutrition education compared with the control that had nutrition education
300 alone. Iron in plant foods exists predominantly as ferric non-haem iron and might be complexed with
301 other organic such as phytate and polyphenols or inorganic molecules in the matrices of fenugreek
302 sprouts, fenugreek seeds, baobab fruit pulp and moringa leaves. Iron, though relatively abundant in
303 the earth crust and most foods, is characterised by poor availability from plant sources.

304 The digested extract solution in the current study was subsequently fractionated with a 3 kDa column
305 to exclude the agglomerated particulate component, and the soluble fractions in the samples ranged
306 between 9.88 and 99.7 % (Tables 1&2). Consequently, the peptic-pancreatin digestion extracts from
307 the plant products after centrifugation consisted of micronised insoluble particulates and non-
308 particulate soluble matter comprising aggregates and agglomerates of iron chelates of diverse sizes
309 and bioaccessibility (Anderson & Frazer, 2017). Digested extracts from baobab fruit pulp had the
310 highest percentage of bioaccessible iron, which however, did not translate into enhanced iron uptake
311 by Caco-2 cells, thus indicating that some components of iron chelates or ligands though soluble
312 might not be bioavailable (Miller & Berner, 1989).

313 Fenugreek sprouts had the highest available iron in Caco-2 cells but paradoxically, exhibited a high
314 level of phytic acid. This was an unexpected finding as the process of sprouting is known to activate
315 phytase (Ou, Cheng, Xing, Lin, Nout, & Liang, 2011) that enzymatically degrades phytate, thereby
316 enhancing iron bioavailability (Hurrell, 2004). In contrast to the current study, previous work reported
317 increased iron availability in germinated fenugreek seeds (Hooda & Jood, 2003). The ratio of phytate
318 to iron has been shown to correlate with iron dialyzability (Glahn, Wortley, South, & Miller, 2002).
319 The phytate to iron ratios were 39.66, 115.07 137.35 and 4.37, while the calcium to iron ratios were
320 9.23, 22.30, 187.25 and 27.13 respectively for fenugreek sprouts, fenugreek seeds, baobab and

321 moringa. Vitamin C in plant products in sequential order were 2 (Ahmed, 2014), 19.55 (Sharara,
322 2017), 266 and 0.11 (Sankhyan, Sharma, Seth, Chauhan, & Kulshrestha, 2013) mg/100 g. Hence,
323 baobab is abundant in both potent inhibitors and an enhancer of iron availability. In general, the
324 comparatively low solubility of iron from these plant products may be due to chemical complexation
325 with a range of inhibitory factors including phytate, polyphenols or fibre (Chadare et al., 2008).

326 The exploitation of the synergy among dietary components to enhance iron absorption promotes the
327 use of foods that are rich in ascorbic acid, a potent enhancer of iron absorption in composite meals.
328 Consequently, ascorbic acid was added to the plant products during the simulated digestion or added
329 to the digested extracts before the application to Caco-2 cells in the current study. Iron availability
330 was significantly enhanced from fenugreek sprouts, fenugreek seeds, baobab pulp and moringa leaves
331 when ascorbic acid was added to the digest before application onto Caco-2 cells but not when it was
332 added during the peptic-pancreatic digestion process. This might be due to the oxidation of the Fe (II)
333 species by ascorbic acid during heat denaturation of the digestive enzymes in the digests before the
334 application onto Caco-2 cells (Vikram, Ramesh & Prapulla, 2005). Moreover, enhanced iron uptake
335 in Caco-2 cells could be due to an augmentation of Dcytb-dependent, ascorbate-mediated
336 ferrereduction (Luo, Hill, Johnson, & Latunde-Dada, 2014). Alternatively, it could be due to the
337 complexation of iron, which is known to form insoluble complexes when the pH level is higher than
338 5.3, thus decreasing the availability and absorption of iron (Scheers, Andlid, Alming, & Sandberg,
339 2010). In support of the latter possibility, previous studies showed that there was no effect of ascorbic
340 acid on iron availability from pork meat at pH 7 during peptic and pancreatic digestion (Sørensen &
341 Bukhave, 2010). Furthermore, the study highlighted the pH dependence of iron uptake from foods
342 and revealed the significance of separating pepsin-digested and pepsin + pancreatin-digested proteins
343 during in vitro studies on iron availability. This exposes a technical limitation of the *in vitro* method
344 as the spatial interactions of dietary constituents in the food matrix, enzymatic digestive processes,
345 bioaccessibility and absorption of iron occur concurrently in the gastrointestinal tract of the
346 organisms.

347 Three indices, namely total bioaccessible iron fraction (TBF), low molecular-weight (LMW) iron
348 fraction and iron uptake in Caco-2 cells were employed in the current study to estimate iron
349 availability from fenugreek sprouts, fenugreek seeds, baobab or moringa. Caco-2 cells have been
350 proven for assessing iron uptake because the cells express intestinal microvilli, enzymes and
351 differentiation markers typical of human small intestine enterocytes (Glahn et al., 1998; Sharp, 2005).
352 There was a positive correlation between TBF and LMW for all samples. However, no correlation
353 was evident between TBF, or LMW and iron uptake in Caco-2 cells. Although iron must be soluble
354 to be absorbable, variables such as the chemical nature of the soluble ligand, molecular weights and

the composition of the digest matrix could confound, in some cases, extrapolation to *in vivo* iron absorption studies (Miller & Berner, 1989). Furthermore, clathrin-mediated endocytosis and micropinocytosis have been shown to play a role in the uptake of nanoparticulate iron complexes in intestinal cells (Latunde-Dada et al., 2014).

In vitro assay of ferritin formation in Caco-2 cells after exposure to an iron source was used as a surrogate marker of iron uptake (Yun, Habicht, Miller, & Glahn, 2004). Variations of the original protocol (Glahn et al., 1998) abound in the literature, and it might now be necessary for a review to enable a form of standardisation of the methodology across different laboratories. Nonetheless, *in vitro* systems of estimating iron solubility are useful for predicting the trends of absorption or relative bioavailability which can, therefore, be used to screen or compare large quantities of food types and different varieties of plant products (Sharp, 2005).

Conclusion

Iron content of the dried vegetable samples (mg/100g) are 1.67 ± 0.05 , (baobab fruit pulp), 7.75 ± 0.04 , (fenugreek seeds), 19.85 ± 0.33 (fenugreek sprout) and 85.44 ± 1.22 (moringa leaves). These underutilised plant foods with variable iron content which upon processing or incorporation in composite cuisines containing rich sources of ascorbic acid, could potentially be employed to treat improve and maintain iron homeostasis in groups at risk of iron deficiency or anaemia. Maximising the potentials of these plant products requires further research in food processing, dietary formulation and nutrition education of the populace.

Funding: This research received no external funding. Kholoud Khoja receives financial sponsorship from King Abdullah Scholarship Programme, Saudi Arabia.

5. References

- Ahmed, A. Y. (2014). Effect of Spraying Fenugreek Seed Sprout and Some Nutrients on Fruiting of Keitte Mango Trees Grown Under Aswan Region Conditions.
- Anderson, G. J., & Frazer, D. M. (2017). Current understanding of iron homeostasis. *The American journal of clinical nutrition*, 106(suppl_6), 1559S-1566S.
- Anwar, F., Latif, S., Ashraf, M., & Gilani, A. H. (2007). Moringa oleifera: a food plant with multiple medicinal uses. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*, 21(1), 17-25.
- Aslam, M., Anwar, F., Nadeem, R., Rashid, U., Kazi, T., & Nadeem, M. (2005). Mineral composition of Moringa oleifera leaves and pods from different regions of Punjab, Pakistan. *Asian J. Plant Sci*, 4(4), 417-421.

Aspuru, K., Villa, C., Bermejo, F., Herrero, P., & López, S. G. (2011). Optimal management of iron deficiency anemia due to poor dietary intake. *International journal of general medicine*, 4, 741.

Bouis, H. E., & Saltzman, A. (2017). Improving nutrition through biofortification: a review of evidence from HarvestPlus, 2003 through 2016. *Global food security*, 12, 49-58.

Chadare, F., Linnemann, A., Hounhouigan, J., Nout, M., & Van Boekel, M. (2008). Baobab food products: a review on their composition and nutritional value. *Critical Reviews in Food Science and Nutrition*, 49(3), 254-274.

Coe, S. A., Clegg, M., Armengol, M., & Ryan, L. (2013). The polyphenol-rich baobab fruit (*Adansonia digitata* L.) reduces starch digestion and glycemic response in humans. *Nutrition research*, 33(11), 888-896.

Feldsine, P., Abeyta, C., & Andrews, W. H. (2002). AOAC International methods committee guidelines for validation of qualitative and quantitative food microbiological official methods of analysis. *Journal of AOAC International*, 85(5), 1187-1200.

Gera, T., Sachdev, H. S., & Boy, E. (2012). Effect of iron-fortified foods on hematologic and biological outcomes: systematic review of randomized controlled trials. *The American journal of clinical nutrition*, 96(2), 309-324.

Gibson, R. S., Perlas, L., & Hotz, C. (2006). Improving the bioavailability of nutrients in plant foods at the household level. *Proceedings of the Nutrition Society*, 65(2), 160-168.

Glahn, R. P., Lee, O. A., Yeung, A., Goldman, M. I., & Miller, D. D. (1998). Caco-2 cell ferritin formation predicts nonradiolabeled food iron availability in an in vitro digestion/Caco-2 cell culture model. *The Journal of nutrition*, 128(9), 1555-1561.

Glahn, R. P., Wortley, G. M., South, P. K., & Miller, D. D. (2002). Inhibition of iron uptake by phytic acid, tannic acid, and ZnCl₂: studies using an in vitro digestion/Caco-2 cell model. *Journal of agricultural and food chemistry*, 50(2), 390-395.

Gómez-Galera, S., Rojas, E., Sudhakar, D., Zhu, C., Pelacho, A. M., Capell, T., & Christou, P. (2010). Critical evaluation of strategies for mineral fortification of staple food crops. *Transgenic research*, 19(2), 165-180.

Goswami, T. (2012). A review on the functional properties, nutritional content, medicinal utilization and potential application of fenugreek. *Journal of Food Processing & Technology*.

Hooda, S., & Jood, S. (2003). Effect of soaking and germination on nutrient and antinutrient contents of fenugreek (*Trigonella foenum graecum* L.). *Journal of Food Biochemistry*, 27(2), 165-176.

Hurrell. (2004). Phytic acid degradation as a means of improving iron absorption. *International journal for vitamin and nutrition research*, 74(6), 445-452.

Lang, T., & Barling, D. (2012). Food security and food sustainability: reformulating the debate. *The Geographical Journal*, 178(4), 313-326.

Latunde-Dada, G. O., Pereira, D. I., Tempest, B., Ilyas, H., Flynn, A. C., Aslam, M. F., Simpson, R.J., & Powell, J. J. (2014). A nanoparticulate ferritin-core mimetic is well taken up by HuTu 80 duodenal cells and its absorption in mice is regulated by body iron. *The Journal of Nutrition*, 144(12), 1896-1902.

Luo, X., Hill, M., Johnson, A., & Latunde-Dada, G. O. (2014). Modulation of Dcytb (Cybrd 1) expression and function by iron, dehydroascorbate and Hif-2 α in cultured cells. *Biochimica et Biophysica Acta (BBA)-General Subjects*, 1840(1), 106-112.

Magaia, T., Uamusse, A., Sjöholm, I., & Skog, K. (2013). Dietary fiber, organic acids and minerals in selected wild edible fruits of Mozambique. *SpringerPlus*, 2(1), 88.

Mbah, B., Eme, P., & Ogbusu, O. (2012). Effect of cooking methods (boiling and roasting) on nutrients and anti-nutrients content of Moringa oleifera seeds. *Pakistan Journal of Nutrition*, 11(3), 211.

McKie, V. A., & McCleary, B. V. (2016). A novel and rapid colorimetric method for measuring total phosphorus and phytic acid in foods and animal feeds. *Journal of AOAC International*, 99(3), 738-743.

- Miller, D., & Berner, L. (1989). Is solubility in vitro a reliable predictor of iron bioavailability? *Biological trace element research*, 19(1-2), 11.
- Muñoz, P., & Humeres, A. (2012). Iron deficiency on neuronal function. *Biometals*, 25(4), 825-835.
- Ou, K., Cheng, Y., Xing, Y., Lin, L., Nout, R., & Liang, J. (2011). Phytase activity in brown rice during steeping and sprouting. *Journal of food science and technology*, 48(5), 598-603.
- Owusu, D., Ellis, W. O., & Oduro, I. (2008). Nutritional potential of two leafy vegetables: *Moringa oleifera* and *Ipomoea batatas* leaves.
- Petropoulos, G. A. (2003). *Fenugreek: the genus Trigonella*: London/New York: Taylor and Francis.CRC Press.
- Powell, J. J., Bruggraber, S. F., Faria, N., Poots, L. K., Hondow, N., Pennycook, T. J., . . . Pereira, D. I. (2014). A nano-disperse ferritin-core mimetic that efficiently corrects anemia without luminal iron redox activity. *Nanomedicine: Nanotechnology, Biology and Medicine*, 10(7), 1529-1538.
- Price, M. (2007). The moringa tree. ECHO Technical Note. North Fort Myers, USA.
- Saini, R., Manoj, P., Shetty, N., Srinivasan, K., & Giridhar, P. (2014). Dietary iron supplements and *Moringa oleifera* leaves influence the liver hepcidin messenger RNA expression and biochemical indices of iron status in rats. *Nutrition research*, 34(7), 630-638.
- Sankhyan, N., Sharma, A., Seth, C. A., Chauhan, A., & Kulshrestha, S. (2013). Determination and comparison of Vitamin C content from *Moringa oleifera* by different methods. *International Journal of Agriculture Science and Research*, 3(2), 67-70.
- Scheers, N., Andlid, T., Alminger, M., & Sandberg, A. S. (2010). Determination of Fe²⁺ and Fe³⁺ in aqueous solutions containing food chelators by differential pulse anodic stripping voltammetry. *Electroanalysis: An International Journal Devoted to Fundamental and Practical Aspects of Electroanalysis*, 22(10), 1090-1096.
- Sharara, M. S. (2017). Effect of Germination and Heat Treatment on Chemical Composition and Bioactive Components of Fenugreek Seeds. *World Journal of Dairy & Food Sciences*, 12(1), 33-41.
- Sharp, P. (2005). Methods and options for estimating iron and zinc bioavailability using Caco-2 cell models: benefits and limitations. *International journal for vitamin and nutrition research*, 75(6), 413-421.
- Shija, A. E., Rumisha, S. F., Oriyo, N. M., Kilima, S. P., & Massaga, J. J. (2019). Effect of *Moringa Oleifera* leaf powder supplementation on reducing anemia in children below two years in Kisarawe District, Tanzania. *Food science & nutrition*, 7(8), 2584-2594.
- Sørensen, A. D., & Bukhave, K. (2010). Iron uptake by Caco-2 cells following in vitro digestion: Effects of heat treatments of pork meat and pH of the digests. *Journal of Trace Elements in Medicine and Biology*, 24(4), 230-235.
- van der Merwe, R., Kruger, J., Ferruzzi, M. G., Duodu, K. G., & Taylor, J. R. (2019). Improving iron and zinc bioaccessibility through food-to-food fortification of pearl millet with tropical plant foodstuffs (moringa leaf powder, roselle calyces and baobab fruit pulp). *Journal of food science and technology*, 56(4), 2244-2256.
- Vikram, V. B., Ramesh, N.M., & Prapulla, S. G. (2005). Thermal degradation kinetics of nutrients in orange juice heated by electromagnetic and conventional methods. *Journal of Food Engineering*, 69, 31-40.
- Willett, W., Rockström, J., Loken, B., Springmann, M., Lang, T., Vermeulen, S., Garnett, T., Tilman, D., DeClerck F., Wood, A., Jonell, M., Clark, M., Gordon L.J., Fanzo, J., Hawkes, C., Zurayk, R., Rivera, J.A., Vries, W.D., Sibanda, L.M., Afshin, A., Chaudhary, A., Herrero, M., Agustina, R., Branca, F., Lartey, A., Fan, S., Crona, B., Fox, E., Bignet, V., Troell, M., Lindahl, T., Singh, S., Cornell, S.E., Reddy, K.S., Narain, S., Nishtar, S., Murray, C.J.L & Wood, A. (2019). Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *The Lancet*, 393(10170), 447-492.

495
496 World Health Organization. (2015). The global prevalence of anaemia in 2011. World Health
497 Organization. <https://apps.who.int/iris/handle/10665/177094> .
498
499 Yun, S., Habicht, J.-P., Miller, D. D., & Glahn, R. P. (2004). An in vitro digestion/Caco-2 cell culture
500 system accurately predicts the effects of ascorbic acid and polyphenolic compounds on iron
501 bioavailability in humans. *The Journal of nutrition*, 134(10), 2717-2721.
502
503

6. Supplementary

Figure 1 Schematic diagram of the in-vitro digestion procedure and iron availability determinations

Table 1 Moisture and Mineral content of the dried vegetable samples (mg/100g)

Sample	Moisture content (%)	Fe	Ca	Cu	Mg	Mn	Zn
Fenugreek sprouts	95.77 ± 0.034	19.85 ± 0.33 a	183.26 ± 0.61 a	2.67 ± 0.01 a	258.0 ± 0.30 a	2.42 ± 0.01 a	2.42 ± 0.01 a
Fenugreek seeds	92.55 ± 0.052	7.75 ± 0.04 b	172.91 ± 3.22 b	2.33 ± 0.03 a	152.89 ± 1.04 b	1.61 ± 0.06 a	1.59 ± 0.05 a
Baobab fruit pulp	94.1 ± 0.098	1.67 ± 0.05 c	313.05 ± 4.13 c	1.77 ± 0.03 a	172.33 ± 2.21 c	1.07 ± 0.14 a	1.079 ± 0.14 a
Moringa leaves	93.2 ± 0.014	85.44 ± 1.22 d	2318.27 ± 23.20 d	1.54 ± 0.03 a	639.50 ± 6.20 d	6.13 ± 0.07 b	6.13 ± 0.06 b

Mineral contents, calcium (Ca), copper (Cu), iron (Fe), magnesium (Mg) manganese (Mn) and zinc (Zn) in 100 g of sample. Values are presented as means ± SEM, n=4. Data were analysed using a two-way ANOVA test. Different letters indicate significant difference between groups ($P \leq 0.05$).

Figure 2: Mineral bioaccessible from digested vegetable. Values are presented as means $\pm$ SEM, n = 6. Data were analysed using a two-way ANOVA. The differences between vegetables are denoted (* $P \leq 0.05$, *** $P \leq 0.001$ and **** $P \leq 0.0001$).

Figure 3 Cell viability of Caco-2 after exposure to digest for 2 h. Values are presented as means $n = 4 \pm$ SEM. Data were analysed using a one-way ANOVA test. Different letters indicate significant difference between groups ($P < 0.05$).