



King's Research Portal

DOI:

[10.1136/bmjopen-2019-035779](https://doi.org/10.1136/bmjopen-2019-035779)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Gungabissoon, U., Perera, G., Galwey, N. W., & Stewart, R. (2020). The association between dementia severity and hospitalisation profile in a newly assessed clinical cohort: The South London and Maudsley case register. *BMJ Open*, *10*(4), [e035779]. <https://doi.org/10.1136/bmjopen-2019-035779>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

BMJ Open The association between dementia severity and hospitalisation profile in a newly assessed clinical cohort: the South London and Maudsley case register

Usha Gungabissoon ^{1,2}, Gayan Perera ², Nicholas W Galwey,³ Robert Stewart^{2,4}

To cite: Gungabissoon U, Perera G, Galwey NW, *et al.* The association between dementia severity and hospitalisation profile in a newly assessed clinical cohort: the South London and Maudsley case register. *BMJ Open* 2020;**10**:e035779. doi:10.1136/bmjopen-2019-035779

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2019-035779>).

Received 15 November 2019
Revised 07 February 2020
Accepted 20 March 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY. Published by BMJ.

¹Epidemiology (Value Evidence and Outcomes), GSK, Brentford, UK

²Department of Psychological Medicine, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK

³Target Sciences, GSK, Stevenage, UK

⁴NIHR Maudsley Biomedical Research Centre, South London and Maudsley NHS Foundation Trust, London, UK

Correspondence to

Usha Gungabissoon;
usha.2.gungabissoon@gsk.com

ABSTRACT

Objectives To evaluate the risk and common causes of hospitalisation in patients with newly diagnosed dementia and variation by severity of cognitive impairment.

Setting We used data from a large London mental healthcare case register linked to a national hospitalisation database.

Participants Individuals aged ≥65 years with newly diagnosed dementia with recorded cognitive function and the catchment population within the same geography.

Outcome measures We evaluated the risk and duration of hospitalisation in the year following a dementia diagnosis. In addition we identified the most common causes of hospitalisation and calculated age-standardised and gender-standardised admission ratios by dementia severity (mild/moderate/severe) relative to the catchment population.

Results Of the 5218 patients with dementia, 2596 (49.8%) were hospitalised in the year following diagnosis. The proportion of individuals with mild, moderate and severe dementia who had a hospital admission was 47.9%, 50.8% and 51.7%, respectively ($p=0.097$). Duration of hospital stay increased with dementia severity (median 2 days in mild to 4 days in severe dementia, $p<0.0001$). After excluding readmissions for the same cause, the most common primary hospitalisation discharge diagnoses among patients with dementia were urinary system disorders, pneumonia and fracture of femur, accounting for 15%, 10% and 6% of admissions, respectively. Overall, patients with dementia were hospitalised 30% more than the catchment population, and this trend was observed for most of the discharge diagnoses evaluated. Standardised admission ratios for urinary and respiratory disorders were higher in those with more severe dementia at diagnosis.

Conclusions Individuals with a dementia diagnosis were more likely to be hospitalised than individuals in the catchment population. The length of hospital stay increased with dementia severity. Most of the common causes of hospitalisation were more common than expected relative to the catchment population, but standardised admission ratios only varied by dementia stage for certain groups of conditions.

Strengths and limitations of this study

- We evaluated hospitalisation in a large and representative sample of newly diagnosed cases of dementia in a London mental healthcare case register linked to a national hospitalisation database providing near-complete outcome ascertainment.
- Severity of dementia at the time of diagnosis was determined from the Mini-Mental State Examination (MMSE) score or Health of the Nation Outcome Scales impairment score, which has acceptable/good psychometric properties and correlates with MMSE score measurement.
- We obtained the analysed samples from a single service provider in a single London catchment, although highly socially diverse, which may limit the generalisability of our findings to other settings.
- The analyses did not account for comorbidities, use of medications, institutional residence or socioeconomic status, aetiology of dementia or hospitalisation rates prior to the diagnosis of dementia, all of which may influence the risk of hospitalisation.

INTRODUCTION

In the United Kingdom, there are around 850 000 people living with dementia and the associated annual healthcare costs are around £4.3 billion; major contributing factors to costs include disease severity and hospitalisation.¹ It follows that the associated healthcare costs are expected to increase in line with the increasing prevalence of dementia as a result of population ageing.² Compared with age-matched controls, people with dementia are more likely to require an acute hospital admission,^{3,4} more hours of care and longer hospital stays.⁵ Hospitalisation also often represents a pivotal event for people with dementia due to increased risk of admission to long-term institutional care, functional decline, mortality, loss of independence and impact

on caregivers.⁶ To date most published studies examining hospital admissions of people with dementia have evaluated prevalence samples of people with dementia.⁷⁻⁹ Other studies have been small,^{3 10 11} or have not taken into account the severity of dementia.^{3 8 9 12} Some have ascertained admission information from family carers^{10 13} which is prone to recall bias.

There is increasing focus on improving interventions and causes of hospitalisation which may be prevented by optimised primary care in the UK for those with dementia.¹⁴ Identifying common causes of hospitalisation in this population could help guide development of preventive strategies to reduce the risk of unnecessary events and/or to avoid lengthy and disruptive admission episodes. However, it is highly likely that the profile of hospitalisations and underlying causal pathways will vary according to the severity of dementia, given the substantial changes that occur during the course of the disease in level of frailty and ability to self-care, not to mention direct effects of neuropsychiatric symptoms on physical health risk. Despite the impact of hospitalisation and the potential for preventative intervention, we are not aware of any previous study that has evaluated the risk and identified the most common causes of hospital admission in newly diagnosed dementia and the extent to which this varies with dementia severity.

In this study, we sought to evaluate the frequency and common causes of hospitalisation in a newly diagnosed cohort of people with a dementia diagnosis relative to the catchment population, and also the potential association between this relative likelihood of cause-specific hospitalisation and severity of cognitive impairment, using a large cohort drawn from a mental healthcare case register linked to national hospitalisation episodes.

METHODS

Study design and setting

This retrospective observational cohort study involved analysis of newly presenting patients who received a dementia diagnosis. We evaluated hospital episodes in the 12-month period following this index diagnosis date.

The Clinical Record Interactive Search (CRIS) data resource was used to identify dementia cases. CRIS provides research access to anonymised electronic health records (EHR) from the South London and Maudsley NHS Foundation Trust (SLaM). SLaM provides mental healthcare, including dementia assessment and management, for a south London catchment containing approximately 1.2 million residents; EHRs were implemented across all SLaM services from 2006.¹⁵ CRIS data are linked to mortality records and national Hospital Episode Statistics (HES) described below. The Oxfordshire Research Ethics Committee C (reference 18/SC/0372) has approved CRIS at the Maudsley as a data resource for secondary analysis.

Routine diagnoses recorded in SLaM are structured according to the WHO International Classification of

Diseases, 10th edition (ICD-10) and are supplemented in CRIS, by a natural language processing (NLP) algorithm ascertaining diagnoses recorded in correspondence and other text fields.^{15 16} Dementia was defined on the presence of F00*, F01*, F02* or F03* ICD diagnosis codes recorded up to the sixth position in the structured data, or F00, F01, F02, F03 dementia, Alzheimer's disease, vascular dementia and mixed dementia from NLP. Hospital outcomes (hospital admissions and its cause) for patients with dementia were obtained from the HES-CRIS linkage. HES contains details of all inpatient admissions at NHS hospitals in England.¹⁷ Discharge diagnoses are recorded as ICD-10 codes and are available for each hospitalisation episode. Additionally, a subset of the HES database detailing hospitalisations for all residents within SLaM's catchment was used in this analysis to generate expected rates for standardisation.

Study participants

We aimed to identify people with newly diagnosed dementia and retrieved records from CRIS for patients with a first recorded diagnosis of dementia between 1 January 2008 and 31 December 2012. We restricted the sample to individuals aged ≥ 65 years at the time of dementia diagnosis, to those with a measure of cognition within 6 months of their dementia diagnosis, and to those with a Mini-Mental State Examination (MMSE) score of < 28 or Health of the Nation Outcome Scales (HoNOS) cognitive impairment score > 1 . Patients who were receiving care from an acute hospital liaison service at the time of the initial diagnosis were excluded since they reflected cases whose dementia diagnosis might have been precipitated by a hospitalisation and who would have an accompanying higher risk of further events.

Covariates

Age at diagnosis (in 5 year bands), sex, ethnicity and dementia severity (within 6 months of dementia diagnosis) were extracted from CRIS. Dementia severity was estimated primarily from the MMSE score recorded closest to the dementia diagnosis date. If no MMSE score was present within 6 months of the diagnosis date, the closest-recorded cognitive impairment subscale of HoNOS was used if data on this were available within the 6-month period around dementia diagnosis. HoNOS is a clinician-rated instrument usually completed at first assessment, which contains subscales rated 0 (no problem) to 4 (severe or very severe problem), has acceptable/good psychometric properties and correlates with MMSE measurement.^{18 19} Dementia severity was categorised as mild, moderate or severe based on MMSE scores of 21–27, 10–20 and 0–9 which is similar to cut-offs used by the National Institute for Health and Care Excellence (NICE), or a HoNOS cognitive impairment subscale score of 2, 3 or 4, respectively.²⁰

All acute general hospital inpatient admissions (both elective and unplanned) for cases were obtained for the 12-month period following the date of dementia diagnosis.




Figure 1 Identification of the case sample. HoNoS, Health of the Nation Outcome Scales; MMSE, Mini Mental State Examination.

Follow-up was censored at the earliest of 31 March 2013 or date of death. Hospitalisations were defined from HES episodes, combining contiguous episodes (ie, where start and end dates were on the same day). Length of hospital stay (LOS) was defined as the number of days from admission to discharge. The three-digit primary ICD-10 discharge diagnosis was obtained for cases and the catchment population for each hospitalisation. All diagnoses were also grouped at the highest level (letter) ICD-10 code, in line with ICD-10 chapter classifications, broadly related to the body system affected.

Hospitalisations for the catchment population from January 2008 to March 2012 were extracted, and the age and sex profile of the catchment was obtained from the UK Census data for the same time frame. **Figure 1** shows the derivation of the case sample.

Statistical analyses

Microsoft Excel for Office 365 ProPlus and Stata V.13.0 were used for analyses. Following a descriptive analysis, we calculated the cumulative incidence of general hospital admissions in the dementia cohort accounting for person-years (py) of follow-up. The association of dementia severity with risk of hospitalisation and length of stay was evaluated by χ^2 test and Kruskal-Wallis tests, respectively.

Age and sex standardised admission ratios (SARs) with 95% CI were calculated overall and for each of the 20 most frequent causes of hospital admission for the 12-month period from the dementia diagnosis based on the three-digit ICD-10 codes. The SAR was defined as the observed number of cases in the dementia cohort admitted with a given cause, divided by the number expected, the second estimate derived from age-specific and sex-specific rates

for admissions with that cause in the SLam catchment. SARs were also calculated for diagnoses at ICD-10 chapter level. SARs were then calculated separately for each category of dementia severity at the time of dementia diagnosis.

To describe variation in the SAR for each specific cause of admission of interest (three-digit ICD-10 code or ICD-10 chapter) by severity of dementia at diagnosis, linear regression models, with SAR as the dependent variable and dementia stage (mild, moderate or severe) as the exposure variable, were fitted. The slope of the regression (direction and magnitude) was used to describe the trend across dementia stages for each specific cause as a way in which to quantify whether SARs tended to increase or decrease with dementia severity. In a sensitivity analysis, we excluded repeat admissions (defined as repeat admissions for the same three-digit ICD-10 code in the 12-month period of interest).

RESULTS

The mean age (SD) of the included patients in the case sample ($n=5218$) was 82.2 (7.0) years, and almost two-thirds were women ($n=3338$; 64.0%); descriptive characteristics of the analysed cohort by the severity of dementia are displayed in **table 1**. At the time of dementia diagnosis, 39.4% were classified as mild, 50.1% as moderate and 10.4% as severe. The 12-month mortality rate from dementia diagnosis was 15.6% overall: 10.9% for mild severity, 17.5% for moderate and 24.6% for severe. Approximately half of the patients with dementia were hospitalised during this period (**table 1**). While the mean number of admissions did not differ substantially between the dementia severity groups, the median duration of inpatient stay (for all admissions in the 12-month period) increased with higher severity of dementia at diagnosis ($p < 0.0001$). Although the proportion of patients with at least one admission increased with dementia severity, this did not reach statistical significance ($p < 0.097$). Patients with dementia were hospitalised 30% more than the catchment population (SAR: 1.3, 95% CI 1.2 to 1.3) during the 12-month follow-up and this SAR was similar across dementia severities: mild 1.2 (95% CI 1.2 to 1.2), moderate 1.3 (95% CI 1.3 to 1.3) and severe 1.3 (95% CI 1.2 to 1.3).

The 20 most common three-digit primary discharge diagnoses for hospitalisation episodes in the year following dementia diagnosis are shown in **table 2**. The most common postdementia discharge diagnosis was chronic renal failure, accounting for 27% of the admissions among those hospitalised, followed by disorders of the urinary system (21%). When readmissions for the same cause were excluded, the most common causes of hospitalisation were disorders of the urinary system, pneumonia and fracture of the femur, accounting for 15%, 10% and 6% of admissions, respectively (online supplementary table 1). Considering primary discharge diagnoses at ICD-10 chapter level, the highest proportions of admissions

**Table 1** Descriptive characteristics of analysis population

	Dementia severity			
	Mild n=2057	Moderate n=2616	Severe n=545	All n=5218
Mean age (SD)	81.7 (6.7)	82.6 (7.1)	82.4 (7.5)	82.2 (7.0)
Sex, N (%)				
Male	765 (37.2)	926 (35.4)	189 (34.7)	1880 (36.0)
Female	1292 (62.8)	1690 (64.6)	356 (65.3)	3338 (64.0)
Ethnicity, N (%)				
Afro-Caribbean	219 (10.6)	400 (15.3)	87 (16)	706 (13.5)
Asian	87 (4.2)	105 (4.0)	30 (5.5)	222 (4.3)
European	1683 (81.8)	2015 (77.0)	394 (72.3)	4092 (78.4)
Other	68 (3.3)	96 (3.7)	34 (6.2)	198 (3.8)
Hospitalisation (12 months after diagnosis)				
Number of patients with ≥1 inpatient admission (%)	986 (47.9)	1328 (50.8)	282 (51.7)	2596 (49.8)
Number of patients per 100 py (95% CI)*	54.6 (51.3 to 58.1)	60.0 (56.9 to 63.3)	63.3 (56.4 to 71.2)	58.1 (55.9 to 60.4)
Including readmissions for the same cause				
Number of hospitalisations†	2413	3201	664	6278
Mean number of admissions (SD)	2.4 (5.9)	2.4 (5.9)	2.3 (4.6)	2.4 (5.8)
Median (IQR) inpatient stay	2 (1 to 10)	3 (1 to 13)	4 (1 to 14)	3 (1 to 12)
After excluding readmissions for the same cause				
Number of hospitalisations†	1868	2461	508	4837
Mean number of admissions (SD)	1.9 (1.4)	1.9 (1.3)	1.8 (1.2)	1.9 (1.3)
Median (IQR) inpatient stay	3 (1 to 12)	5 (1 to 15)	6 (2 to 17)	4 (1 to 14)

*P value= 0.097 (χ^2 test).†P value= 0.0001 (Kruskal-Wallis tests).
py, person years.

were observed in specific diseases of the genitourinary, circulatory and respiratory systems, accounting for 54%, 25% and 22% of the hospitalisation episodes, respectively (table 3); however, the broader chapters for 'other symptoms/signs' and 'injury/external cause events' were also common, accounting for 35% and 22% of admissions, respectively. After exclusion of repeat admissions (online supplementary table 2), the highest proportions of admissions at a chapter level were in specific diseases of the genitourinary, circulatory and respiratory systems, each accounting for around 20% of the hospitalisation episodes. The broader chapters for 'other symptoms/signs' and 'injury/external cause events', accounted for 31% and 21% of admissions, respectively, after exclusion of repeat admissions at a three-digit ICD level.

SARs for each discharge diagnosis by the severity of dementia at diagnosis are also shown in tables 2 and 3, and are illustrated in figures 2 and 3. For the cohort as a whole, patients with dementia were hospitalised more than the catchment population (SAR significantly greater than 1) for most of the 20 most common discharge diagnoses, the exceptions being chronic renal failure (N18), other chronic obstructive pulmonary disease (J44), heart failure (I50, SARs not significantly >1) and other cataracts (H26, SAR <1). The trends across

dementia severity groups were positive for 17 of the 20 conditions (increased SARs with increasing severity, indicated by a positive regression coefficient); however, only the trend for the R41 code ('other symptoms and signs involving cognitive function and awareness') was statistically significant. Considering ICD chapters, in most groups, patients with dementia were hospitalised more than the catchment population (table 3). The exceptions were cancers, benign neoplasms and diseases of the eye (SAR <1). Of the 18 ICD-10 chapters, trends across dementia severity groups were positive in 10; however, only those for mental and behavioural disorders (F), diseases of the respiratory system (J) and diseases of the genitourinary system (N) were statistically significant. No chapters showed significantly decreasing SARs with increasing dementia severity, although the negative trend for diseases of the digestive system (K) was close to significance. After excluding repeat admissions for the same cause, similar trends were observed at the three-digit ICD and chapter level, although an increasing trend of SAR with dementia severity reached or approached statistical significance for disorders of the urinary system and pneumonia with an increasing trend with dementia severity, and a decreasing trend, which reached statistical significance, for pain in throat

Table 2 SARs by primary discharge diagnosis (20 most common ICD-10 three-digit code) for hospitalisations in the 12-month period following a dementia diagnosis

Three-digit ICD primary discharge diagnosis	Number of hospitalisations with this discharge diagnosis	Proportion (%) of episodes with this discharge diagnosis	SAR (95% CI) by dementia severity at diagnosis				Regression coefficient*	P value*
			All cases	Mild n=986	Moderate n=1328	Severe n=282		
Chronic renal failure (N18)	710	27.3	1.0 (0.9–1.0)	0.8 (0.7–0.9)	1.0 (0.9–1.1)	2.2 (1.9–2.6)	0.71	0.26
Other disorders of urinary system (N39)	532	20.5	2.7 (2.5–2.7)	2.4 (2.1–2.8)	2.9 (2.6–3.2)	4.8 (3.9–5.8)	1.18	0.21
Pneumonia, organism unspecified (J18)	263	10.1	1.9 (1.7–2.2)	1.4 (1.1–1.8)	2.1 (1.8–2.5)	4.5 (3.5–5.7)	1.56	0.2
Fracture of femur (S72)	156	6.0	2.4 (2.0–2.8)	2.4 (1.8–3.1)	2.3 (1.9–2.9)	3.1 (1.9–4.6)	0.32	0.39
Senility (R54)	155	6.0	2.9 (2.5–3.4)	2.9 (2.2–3.7)	2.8 (2.2–3.5)	4.6 (3.0–6.7)	0.86	0.36
Syncope and collapse (R55)	133	5.1	2.9 (2.4–3.4)	3.0 (2.3–3.9)	2.8 (2.2–3.6)	4.6 (2.9–6.9)	0.77	0.4
Other cataract (H26)	132	5.1	0.6 (0.5–0.7)	0.5 (0.4–0.7)	0.6 (0.5–0.8)	0.6 (0.3–1.0)	0.05	0.58
Unspecified acute lower respiratory infection (J22)	103	4.0	1.8 (1.5–2.2)	1.4 (1.0–2.0)	1.9 (1.5–2.5)	3.9 (2.5–5.8)	1.24	0.21
Cerebral infarction (I63)	100	3.9	1.7 (1.4–2.1)	1.4 (1.0–2.0)	1.8 (1.3–2.3)	3.4 (2.1–5.2)	1.00	0.23
Other symptoms and signs involving cognitive function and awareness (R41)	96	3.7	5.2 (4.2–6.3)	4.2 (2.9–6.1)	5.9 (4.5–7.7)	8.1 (4.6–13.2)	1.93	0.05
Pain in throat and chest (R07)	91	3.5	1.4 (1.1–1.7)	1.7 (1.2–2.3)	1.3 (1.0–1.8)	1.2 (0.5–2.4)	–0.24	0.16
Other chronic obstructive pulmonary disease (J44)	87	3.4	1.0 (0.8–1.2)	1.0 (0.7–1.4)	1.0 (0.8–1.4)	0.8 (0.3–1.6)	–0.13	0.32
Open wound of head (S01)	78	3.0	3.0 (2.4–3.8)	3.2 (2.1–4.5)	2.8 (2.0–3.9)	4.5 (2.4–7.7)	0.65	0.47
Unspecified dementia (F03)	77	3.0	8.3 (6.6–10.4)	5.0 (2.9–7.9)	10.6 (7.9–14.0)	12.7 (6.8–21.7)	3.87	0.17
Superficial injury of head (S00)	73	2.8	3.6 (2.8–4.5)	3.6 (2.4–5.3)	3.3 (2.3–4.5)	6.9 (3.9–11.2)	1.64	0.39
Heart failure (I50)	61	2.3	0.8 (0.6–1.1)	0.6 (0.4–1.0)	1.0 (0.7–1.4)	0.8 (0.3–1.7)	0.08	0.76
Acute renal failure (N17)	61	2.3	2.1 (1.6–2.8)	1.6 (1.0–2.6)	2.2 (1.5–3.1)	5.7 (3.3–9.2)	2.05	0.25

Continued



Table 2 Continued

Three-digit ICD primary discharge diagnosis	Number of hospitalisations with this discharge diagnosis	Proportion (%) of episodes with this discharge diagnosis	SAR (95% CI) by dementia severity at diagnosis				Regression coefficient*	P value*
			All cases	Mild n=986	Moderate n=1328	Severe n=282		
Alzheimer's disease (G30)	60	2.3	10.4 (8.0–13.4)	7.0 (3.9–11.5)	10.8 (7.4–15.2)	31.9 (19.5–49.2)	12.46	0.24
Other functional intestinal disorders (K59)	59	2.3	1.8 (1.4–2.3)	2.6 (1.8–3.6)	1.5 (1.0–2.2)	0.9 (0.2–2.5)	–0.85	0.09
Pneumonitis due to solids and liquids (J69)	52	2.0	3.0 (2.2–3.9)	1.7 (0.8–3.0)	3.5 (2.4–5.0)	7.9 (4.4–13.1)	3.14	0.15

*The regression coefficient indicates the increase or decrease in SAR associated with a one-step increase in severity of dementia. See Statistical analyses section for the linear model relating to these variables. The p value is for a two-sided test of the null hypothesis regression coefficient=0.

ICD-10, International Classification of Diseases, 10th edition; SARs, standardised admission ratios.

and chest (online supplementary tables 3–4, figures 2 and 3).

DISCUSSION

In this retrospective cohort study of people with a dementia diagnosis, over half had at least one hospitalisation in the 12 months after diagnosis. The risk of hospitalisation and the LOS increased with higher dementia severity, although risk of hospitalisation did not reach statistical significance. Most of the common reasons for hospitalisation, based on the primary discharge diagnosis, were also more common than expected in cases relative to the catchment population. Differences in SARs by baseline dementia severity, where present, were more evident for discharge diagnoses grouped at ICD-10 chapter level than at the level of specific three-digit ICD-10 codes.

The hospitalisation rate for at least one inpatient admission in the 12-month period after dementia diagnosis was 58 per 100 py, slightly higher than reported elsewhere.^{21 22} This may reflect more recent data (given that hospitalisation rates are increasing), the near-complete data on hospitalisation in our study and/or the fact that hospitalisation is free at the point of delivery in the UK context.⁷ Increasing likelihood and LOS with increasing dementia severity is consistent with other reports, although differences between admissions from community and institutional facilities have been described, with reduced healthcare use following relocation to institutional care, potentially reflecting higher support.^{23–25} This, as well as survival effects, may account for our finding that the most marked difference in hospitalisation was between mild and moderate dementia rather than between moderate and severe dementia.

The most common causes of hospital admission, after excluding repeat admissions for the same cause, were disorders of the urinary system and pneumonia, followed by fracture of femur, senility and syncope and collapse; this is similar to the findings reported elsewhere.^{4 9 26–28} Most of the common causes of admission identified in people with dementia were more frequent than expected relative to the catchment population (SAR >1).

Higher risk of syncope and collapse, and of other diagnoses related to fractures and falls, has been previously reported in patients with dementia,^{4 9 26} and may reflect age-related autonomic dysfunction, comorbid disorders and polypharmacy,²⁹ impaired visuospatial functions and gait instability.³⁰ A review of published clinical trials of atypical antipsychotic medication use did not find evidence of increased injury, falls or syncope associated with their use,³¹ nor did a recent study of hospitalised falls/fractures in a CRIS-derived cohort of people with dementia.³² Significant predictors included social/demographic factors, physical health problems and previous episode; no associations were found with neuropsychiatric symptoms, cognitive (MMSE) scores or functional problems.³² Inpatient admissions due to infection were also more common than expected, both at the level of specific disorders and at the ICD-10 chapter level. This may reflect more rapid age-related decline in immune function,³³ problems with mobility associated with urinary incontinence and subsequent infection³⁴ or inactivity and dysphagia causing respiratory tract infection.³⁵ Acute renal failure is reported to be associated with a high risk of hospital readmission.^{36 37} In our study it remained a common cause of admission among individuals with dementia even after excluding repeat admissions. Risk factors for acute renal failure include hospitalisation,

Table 3 SARs (95% CI) by discharge diagnosis (ICD-10 chapter) for hospitalisations in the 12-month period after a dementia diagnosis

Discharge diagnosis by ICD-10 chapter		Number of hospitalisations with this discharge diagnosis	Proportion (%) of episodes with this discharge diagnosis	SAR (95% CI) by dementia severity at diagnosis				Regression coefficient*	P> T *
				All cases	Mild n=986	Moderate n=1328	Severe n=282		
A	Infectious and parasitic diseases	62	2.4	1.5 (1.2–2.0)	1.0 (0.6–1.7)	1.8 (1.3–2.5)	2.1 (0.9–3.9)	0.51	0.16
B	Infectious and parasitic diseases	11	0.4	1.6 (0.8–2.9)	0.8 (0.1–2.8)	2.1 (0.8–4.2)	2.8 (0.3–10.0)	0.99	0.11
C	Cancers	167	6.4	0.3 (0.3–0.4)	0.3 (0.3–0.4)	0.4 (0.3–0.5)	0.1 (0.1–0.3)	–0.1	0.47
D	Benign neoplasms or diseases of the blood	158	6.1	0.8 (0.7–1.0)	0.9 (0.7–1.1)	0.9 (0.7–1.1)	0.3 (0.1–0.7)	–0.3	0.31
E	Endocrine, nutritional and metabolic diseases	138	5.3	1.9 (1.6–2.3)	1.9 (1.5–2.5)	1.9 (1.4–2.4)	2.1 (1.2–3.5)	0.09	0.54
F	Mental and behavioural disorders	194	7.5	5.7 (4.9–6.6)	4.7 (3.6–6.1)	6.1 (5–7.4)	7.4 (4.9–10.8)	1.35	0.01
G	Diseases of the nervous system	183	7.0	2.4 (2.1–2.8)	2.1 (1.6–2.7)	2.4 (2.0–3.0)	3.4 (2.3–5.0)	0.65	0.18
H	Diseases of the eye	209	8.1	0.6 (0.5–0.7)	0.6 (0.5–0.8)	0.7 (0.5–0.8)	0.3 (0.2–0.6)	–0.13	0.43
I	Diseases of the circulatory system	638	24.6	1.3 (1.2–1.3)	1.4 (1.2–1.5)	1.2 (1.1–1.4)	1.0 (0.8–1.4)	–0.16	0.06
J	Diseases of the respiratory system	567	21.8	1.6 (1.5–1.6)	1.3 (1.1–1.5)	1.7 (1.5–1.9)	2.2 (1.8–2.7)	0.46	0.03
K	Diseases of the digestive system	448	17.3	1.0 (0.9–1.0)	1.2 (1.1–1.4)	1.0 (0.8–1.1)	0.6 (0.4–0.9)	–0.3	0.06
L	Diseases of the skin	105	4.0	1.2 (1.0–1.5)	1.2 (0.9–1.6)	1.1 (0.8–1.4)	1.9 (1.1–3.0)	0.33	0.41
M	Diseases of the musculoskeletal system	239	9.2	0.9 (0.8–1.1)	1.1 (0.9–1.3)	0.8 (0.7–1.0)	0.9 (0.6–1.4)	–0.07	0.64
N	Diseases of the genitourinary system	1389	53.5	1.3 (1.2–1.3)	1.1 (1.0–1.1)	1.4 (1.3–1.4)	1.7 (1.4–1.9)	0.28	0.01
R	Symptoms and sign not elsewhere classified	913	35.2	1.9 (1.8–1.9)	2 (1.8–2.2)	1.9 (1.7–1.9)	1.7 (1.3–2.1)	–0.15	0.18
S	Injury, poisoning and certain other external causes	574	22.1	2.3 (2.1–2.3)	2.3 (2.0–2.7)	2.2 (2.0–2.5)	2.4 (1.9–3.1)	0.05	0.69
T	Injury, poisoning and certain other external causes	120	4.6	1.6 (1.3–1.9)	1.6 (1.2–2.1)	1.8 (1.4–2.3)	0.7 (0.2–1.5)	–0.46	0.47
Z	External causes	161	6.2	1.1 (1.0–1.3)	1.6 (1.3–1.9)	0.9 (0.7–1.1)	0.8 (0.4–1.4)	–0.4	0.25
Total		2596	–	1.3 (1.2–1.3)	1.2 (1.2–1.2)	1.3 (1.3–1.3)	1.3 (1.2–1.3)	0.03	0.3

*The regression coefficient indicates the increase or decrease in SAR associated with a one-step increase in severity of dementia. See Statistical analyses section for the linear model relating to these variables. The p value is for a two-sided test of the null hypothesis regression coefficient=0.

ICD-10, International Classification of Diseases, 10th edition; SARs, standardised admission ratios.




Figure 2 SARs by three-digit ICD code for mild, moderate and severe dementia populations (excludes repeat admissions for the same cause at a three-digit ICD level). ICD, International Classification of Diseases; SARS, standardised admission ratios.

infection, diabetes, pre-existing renal impairment and advanced age,³⁸ which themselves may be associated with dementia severity.

A lower than expected frequency of hospitalisation (SAR <1) was observed for heart failure, cancers, benign neoplasms and diseases of the eye, consistent with other

reports^{21 28} and possibly reflecting diagnostic delay. For example, an association of dementia with later-stage cancer at diagnosis has been reported,³⁹ as well as unexpected cases being identified at autopsy⁴⁰; furthermore, patients with dementia may be less likely to undergo invasive diagnostic tests and receive fewer treatment




Figure 3 SARs by ICD chapter for mild, moderate and severe dementia populations (excludes repeat admissions for the same cause at a three-digit ICD level). ICD, International Classification of Diseases; SARs, standardised admission ratios.

interventions, potentially explaining a shorter duration of survival.⁴¹

While rates of healthcare utilisation and costs, or limited specified causes of admission,^{23 24} have been reported in relation to severity of dementia, we are not aware of any evaluation of common causes of hospital admission by dementia stage. Most of the SARs for specific causes of hospitalisation were positively associated with dementia severity, although most not to a significant extent. Associations, where significant, were more often with discharge diagnoses categorised by ICD-10 chapter rather than three-character code. This may have arisen because of higher statistical power for the chapter-level comparisons but might also reflect differences in the accuracy and/or variability of code assignments in routine administrative data; for example, there may be differences between hospitals in which specific three-character code to assign to a particular hospitalisation profile but stronger agreement on the broader chapter heading.

As well as hospitalisations occurring as a direct result of dementia, with worsening cognitive impairment there may be an increase in admissions to acute care for exacerbation, because of an individual's decreasing ability to manage existing comorbidity. Similarly, if identification of new comorbidity is delayed at earlier stages of dementia, this may increase the risk of hospitalisations later in its course due to underrecognised and/or undertreated disorders, although there is also evidence for an increased risk of some causes of hospitalisation even prior to a dementia diagnosis.⁴² An increased LOS is consistent with the higher rate of complications seen in hospitalised patients with dementia, as well as with increasing severity of dementia.⁴³ Increasing respiratory and genitourinary disease admissions most likely represent infections, and diseases of the nervous system may reflect cases where the dementia itself has been provided as a primary diagnosis, or else comorbidities such as depression, psychosis or other behavioural/psychological manifestations. Risk for diseases of the digestive system diminished across the severity groups, but this may reflect a diminished ability to communicate non-specific symptoms rather than a reduced risk of defined disorders, as indicated by the negative coefficient for 'other functional intestinal disorders' (K59) in [table 2](#), which was the most common digestive system diagnosis.

Strengths of the study included the large and representative sample of diagnosed cases of dementia and the linkage to a national hospitalisation database providing near-complete outcome ascertainment. Patients who were receiving care from acute hospital liaison services at the time of their dementia diagnosis were excluded from the study since these may reflect individuals whose diagnosis might have been precipitated by the hospitalisation, and yet who would also have an increased risk of rehospitalisation by virtue of their status, thus biasing the association of interest. Considering limitations, the analysed sample was from a single service and also only included cases with dementia diagnosed in specialist services; however,

estimated proportions of people with dementia in the SLAM catchment who receive a specialist diagnosis is relatively high at 75.2%, compared with 67.6% nationally.⁴⁴ This study was descriptive in nature and used all available data; as such it was not generated with a specific power calculation for the associations of interest, and the relatively small number of cases with severe dementia at diagnosis will have limited the detection of differences across the three groups. Furthermore, unrecognised dementia cases omitted from observed admission rates will have biased findings towards the null, as would healthy survivor effects (which may also be particularly an issue for the severe dementia group). We used the discharge diagnosis recorded during the last episode of the hospital spell, and it is possible that the primary discharge diagnosis code may reflect complications that arose during the hospital stay rather than the reason for initial hospital admission. Although we compared SARs across dementia severities for a given discharge diagnosis, due to potential differences in the underlying age structure between severity categories SARs are not strictly comparable with each other. However, as the mean age was broadly similar between dementia severities, this mitigates this concern in part. Finally, this analysis did not attempt to account for comorbidity, medication use, institutional residence or socioeconomic status, did not subclassify dementia and was not able to investigate or take into account clustering by admission units.

Understanding the factors influencing health service utilisation by patients with dementia is necessary to inform care needs and to guide future healthcare resource planning and allocation. Our findings are of importance given that hospitalisation is a significant element in the cost of dementia care and that the prevalence of dementia is increasing. Our study highlights the need to develop specific strategies for those causes of hospitalisation that are most amenable to prevention in dementia. Further research on factors influencing patterns of healthcare use over time and severity would be useful in this context.

Acknowledgements The authors would like to acknowledge Hitesh Shetty for his contributions to the study.

Contributors UG was responsible for the design, analysis, interpretation and drafting of the manuscript. GP contributed to the design and analysis of the study. NWG provided statistical input and contributed to the drafting of the manuscript. RS contributed to the design, interpretation and drafting of the manuscript.

Funding This work was supported by NIHR Biomedical Research Centre at the South London and Maudsley NHS Foundation Trust and King's College London. RS is additionally part funded by a Medical Research Council Mental Health Data Pathfinder Award to King's College London, and an NIHR Senior Investigator Award.

Disclaimer The funder of the study had no role in the study design, data collection, data analysis, data interpretation or writing of the report. UG, GP and RS had access to the anonymised data. The corresponding author had the final responsibility for the decision to submit for publication.

Competing interests UG and NWG are employees of GSK, hold stock and receive a salary from GSK. RS has received research funding in the last 5 years from Roche, Janssen, Takeda and GSK.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data may be obtained from a third party and are not publicly available. Because of their nature and to comply with their ethical approval, CRIS data are required to remain within the firewall of the South London and Maudsley NHS Foundation Trust (SLaM). Access to the data used for this study can be facilitated by the CRIS Oversight Committee on application and with appropriate SLaM affiliation, details of which can be obtained from cris.administrator@slam.nhs.uk.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Usha Gungabissoon <http://orcid.org/0000-0002-2040-1763>

Gayana Perera <http://orcid.org/0000-0002-3414-303X>

REFERENCES

- Alzheimer's Society. *Dementia UK update*, 2014.
- Ahmadi-Abhari S, Guzman-Castillo M, Bandosz P, et al. Temporal trend in dementia incidence since 2002 and projections for prevalence in England and Wales to 2040: modelling study. *BMJ* 2017;358:j2856.
- Phelan EA, Borson S, Grothaus L, et al. Association of incident dementia with hospitalizations. *JAMA* 2012;307:165–72.
- Natalwala A, Potluri R, Uppal H, et al. Reasons for hospital admissions in dementia patients in Birmingham, UK, during 2002–2007. *Dement Geriatr Cogn Disord* 2008;26:499–505.
- Mukadam N, Sampson EL. A systematic review of the prevalence, associations and outcomes of dementia in older General Hospital inpatients. *Int Psychogeriatr* 2011;23:344–55.
- Lehmann J, Michalowsky B, Kaczynski A, et al. The impact of hospitalization on readmission, institutionalization, and mortality of people with dementia: a systematic review and meta-analysis. *J Alzheimers Dis* 2018;64:735–49.
- Bernardes C, Massano J, Freitas A. Hospital admissions 2000–2014: a retrospective analysis of 288 096 events in patients with dementia. *Arch Gerontol Geriatr* 2018;77:150–7.
- Tolppanen A-M, Taipale H, Purmonen T, et al. Hospital admissions, outpatient visits and healthcare costs of community-dwellers with Alzheimer's disease. *Alzheimers Dement* 2015;11:955–63.
- Draper B, Karmel R, Gibson D, et al. The hospital dementia services project: age differences in hospital stays for older people with and without dementia. *Int Psychogeriatr* 2011;23:1649–58.
- Voisin T, Andrieu S, Cantet C, et al. Predictive factors of hospitalizations in Alzheimer's disease: a two-year prospective study in 686 patients of the REAL.FR study. *J Nutr Health Aging* 2010;14:288–91.
- Russ TC, Parra MA, Lim AE, et al. Prediction of General Hospital admission in people with dementia: cohort study. *Br J Psychiatry* 2015;206:153–9.
- Joyce AT, Zhao Y, Bowman L, et al. Burden of illness among commercially insured patients with Alzheimer's disease. *Alzheimers Dement* 2007;3:204–10.
- Andrieu S, Reynish E, Nourhashemi F, et al. Predictive factors of acute hospitalization in 134 patients with Alzheimer's disease: a one year prospective study. *Int J Geriatr Psychiatry* 2002;17:422–6.
- Department of Health. *Living well with dementia: a national dementia strategy*. Department of Health, 2009.
- Perera G, Broadbent M, Callard F, et al. Cohort profile of the South London and Maudsley NHS Foundation trust biomedical research centre (SLaM BRC) case register: current status and recent enhancement of an electronic mental health Record-derived data resource. *BMJ Open* 2016;6:e008721.
- Jackson RG, Patel R, Jayatilake N, et al. Natural language processing to extract symptoms of severe mental illness from clinical text: the clinical record interactive search comprehensive data extraction (CRIS-CODE) project. *BMJ Open* 2017;7:e012012.
- NHS. Hospital episode statistics (hES) 2019, 2019. Available: <https://digital.nhs.uk/data-and-information/data-tools-and-services/data-services/hospital-episode-statistics> [Accessed 13 Jan 2019].
- Pirkis JE, Burgess PM, Kirk PK, et al. A review of the psychometric properties of the health of the nation outcome scales (HoNOS) family of measures. *Health Qual Life Outcomes* 2005;3:76.
- Canuto A, Weber K, Gold G, et al. Structured assessment of mental health status in psychogeriatrics: validity of the French HoNOS65+. *Can J Psychiatry* 2007;52:37–45.
- NICE. Donepezil, galantamine, rivastigmine and memantine for the treatment of dementia. In: *Health technology appraisal 2018 update*. NICE, 2018.
- Tuppin P, Kusnik-Joinville O, Weill A, et al. Primary health care use and reasons for hospital admissions in dementia patients in France: database study for 2007. *Dement Geriatr Cogn Disord* 2009;28:225–32.
- Chen L, Reed C, Happich M, et al. Health care resource utilisation in primary care prior to and after a diagnosis of Alzheimer's disease: a retrospective, matched case-control study in the United Kingdom. *BMC Geriatr* 2014;14:76.
- Lin P-J, Rane PB, Fillit HM, et al. O2-11-01: national estimates of potentially avoidable hospitalizations among Medicare beneficiaries with Alzheimer's disease and related dementias. *Alzheimer's & Dementia* 2016;12:P253–4.
- Zhu CW, Cosentino S, Ornstein K, et al. Use and cost of hospitalization in dementia: longitudinal results from a community-based study. *Int J Geriatr Psychiatry* 2015;30:833–41.
- Fillenbaum G, Heyman A, Peterson BL, et al. Use and cost of hospitalization of patients with AD by stage and living arrangement: CERAD XXI. *Neurology* 2001;56:201–6.
- Zhao Y, Kuo T-C, Weir S, et al. Healthcare costs and utilization for Medicare beneficiaries with Alzheimer's. *BMC Health Serv Res* 2008;8:108.
- Rao A, Suliman A, Vuik S, et al. Outcomes of dementia: systematic review and meta-analysis of hospital administrative database studies. *Arch Gerontol Geriatr* 2016;66:198–204.
- Guijarro R, San Román CM, Gómez-Huelgas R, et al. Impact of dementia on hospitalization. *Neuroepidemiology* 2010;35:101–8.
- O' Brien H, Anne Kenny R, Kenny A. Syncope in the elderly. *Eur Cardiol* 2014;9:28–36.
- Sheridan PL, Hausdorff JM. The role of higher-level cognitive function in gait: executive dysfunction contributes to fall risk in Alzheimer's disease. *Dement Geriatr Cogn Disord* 2007;24:125–37.
- Schneider LS, Dagerman K, Insel PS. Efficacy and adverse effects of atypical antipsychotics for dementia: meta-analysis of randomized, placebo-controlled trials. *Am J Geriatr Psychiatry* 2006;14:191–210.
- Sharma S, Mueller C, Stewart R, et al. Predictors of falls and fractures leading to hospitalization in people with dementia: a representative cohort study. *J Am Med Dir Assoc* 2018;19:607–12.
- Janssens J-P. Pneumonia in the elderly (geriatric) population. *Curr Opin Pulm Med* 2005;11:405–9.
- Schnelle JF, Leung FW. Urinary and fecal incontinence in nursing homes. *Gastroenterology* 2004;126:S41–7.
- Palmer JL, Metheny NA. Preventing aspiration in older adults with dysphagia. *Am J Nurs* 2008;108:40–8. quiz 49.
- Jencks SF, Williams MV, Coleman EA. Rehospitalizations among patients in the Medicare fee-for-service program. *N Engl J Med* 2009;360:1418–28.
- Silver SA, Harel Z, McArthur E, et al. 30-Day readmissions after an acute kidney injury hospitalization. *Am J Med* 2017;130:163–72.
- Hilton R. Acute renal failure. *BMJ* 2006;333:786–90.
- Gorin SS, Heck JE, Albert S, et al. Treatment for breast cancer in patients with Alzheimer's disease. *J Am Geriatr Soc* 2005;53:1897–904.
- Gupta SK, Lamont EB. Patterns of presentation, diagnosis, and treatment in older patients with colon cancer and comorbid dementia. *J Am Geriatr Soc* 2004;52:1681–7.
- Libert Y, Dubruielle S, Borghgraef C, et al. Vulnerabilities in older patients when cancer treatment is initiated: does a cognitive impairment impact the two-year survival? *PLoS One* 2016;11:e0159734.
- Desai U, Kirson NY, Ye W, et al. Trends in health service use and potentially avoidable hospitalizations before Alzheimer's disease diagnosis: a matched, retrospective study of US Medicare beneficiaries. *Alzheimers Dement* 2019;11:125–135.
- Möllers T, Stocker H, Wei W, et al. Length of hospital stay and dementia: a systematic review of observational studies. *Int J Geriatr Psychiatry* 2019;34:8–21.
- NHS England. *Dementia diagnosis monthly workbook: secondary dementia diagnosis monthly workbook*, 2017.