King's Research Portal

DOI:

[10.1097/ACI.0000000000000719](https://doi.org/10.1097/ACI.0000000000000719)

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Santos, A. F. (2020). Prevention of food allergy: can we stop the rise of IgE mediated food allergies? *Current Opinion in Allergy & Clinical Immunology*, *Publish Ahead of Print*.
<https://doi.org/10.1097/ACI.0000000000000719>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Title page

Title: Prevention of food allergy: can we stop the rise of IgE mediated food allergies?

Author: Alexandra F. Santos^{1,2,3,4}

Affiliations:

¹Department of Women and Children's Health (Paediatric Allergy), School of Life Course Sciences, Faculty of Life Sciences and Medicine, King's College London, London, United Kingdom

²Peter Gorer Department of Immunobiology, School of Immunology and Microbial Sciences, King's College London, London, United Kingdom

³Children's Allergy Service, Guy's and St Thomas' Hospital, London, United Kingdom

⁴Asthma UK Centre for Allergic Mechanisms in Asthma, London, United Kingdom

Corresponding author:

Alexandra F. Santos, MD PhD

alexandra.santos@kcl.ac.uk

+44 (0) 20 7188 6424

Funding: A.F.Santos is funded by the Medical Research Council (MRC Transition support MR/T032081/1, MRC Clinician Scientist Fellowship MR/M008517/1, MRC Centenary Early Career Award and MRC Clinical Research Training Fellowship G0902018), Asthma UK (AUK-BC-2015-01) and by the UK National Institute for Health Research comprehensive Biomedical Research Centre award to Guy's and St Thomas' National Health Service (NHS)

Foundation Trust, in partnership with the King's College London and King's College Hospital
NHS Foundation Trust.

Structured Abstract

Purpose of review: Food allergy has become more prevalent in recent decades. Without a
curative treatment for food allergy, prevention is key. Can we intervene and halt the food
allergy epidemic?

Recent findings: There are three main hypotheses to explain the rise in food allergy: the dual-
allergen exposure hypothesis, the hygiene hypothesis and the vitamin D hypothesis. In a recent
systematic review of randomized controlled trials, only introduction of allergenic foods,
namely egg and peanut, in the diet at the time of weaning and avoidance of temporary
supplementation with cow's milk formula in the first few days of life showed low to moderate
evidence of a preventative effect.

Summary: For primary prevention, introduction of allergenic foods at the time of weaning and
avoidance of temporary supplementation with cow's milk formula in the first few days of life
has been recommended. Introduction of foods once allergy has been excluded may be
beneficial for sensitized subjects (secondary prevention). Once food allergy has been
established, it is important to minimise complications (tertiary prevention) through allergen
avoidance, timely treatment of allergic reactions, control of atopic co-morbidities and dietetic
and psychological support, as appropriate. Immunomodulatory treatments can potentially be
disease-modifying and require further research.

Abstract word count: 199

Keywords: Food allergy, prevention, treatment, IgE, eczema, anaphylaxis

Word count: 2547

Introduction

Food allergy affects about 8% of children in Western countries (1, 2) and its prevalence is increasing in the adult age group (3), probably reflecting both an increase in incidence and in life-time prevalence as children grow older. Food allergy has also been rising in other parts of the world, like South East Asia and Africa, particularly in urban centres (4, 5). Hospital admissions for food allergy and severity of food allergy have also increased, with a recent US study showing that about 40% of food allergic children have multiple food allergies, severe food allergies and have been prescribed an adrenaline auto-injector (1). Although many food allergies resolve spontaneously, others can be life-long. The 4-year follow-up of the Healthnuts cohort from Melbourne in Australia showed that the prevalence of food allergy decreased from 11% at 1 year to 3.8% at 4 years of age (6). The Schoolnuts study, in the same geographical area, showed a prevalence of food allergy of 4.5% at 10-14 years of age (7). The natural history of food allergy seems to be changing over time with more recent studies reporting a later acquisition of tolerance than previously, although such reports can be influenced by the population studied, the type of food allergy (e.g. whether IgE or non-IgE-mediated) and the approach adopted to assess resolution. Over the past few decades, we have witnessed an increase in the prevalence, severity and persistence of food allergy overall.

Why is the world becoming more allergic to food?

There are 3 main working hypotheses to explain the food allergy epidemic (**Figure 1**):

- Dual allergen exposure hypothesis
- Hygiene hypothesis
- Vitamin D hypothesis

Dual allergen exposure hypothesis

Atopic eczema is probably the main risk factor for food allergy, particularly early onset moderate-severe eczema (8, 9). The dual allergen exposure hypothesis, proposed by Gideon Lack in 1996, postulates that the development of food allergy results from the combination of time, route and dose of exposure to food allergens in early life (10). The balance between exposure through the gastro-intestinal tract, which is tolerogenic, particularly in high doses, and exposure through the skin, particularly to low doses of allergen through abraded inflamed skin affected with eczema, determines food allergy risk. When exposure via the gastrointestinal tract happens before exposure through the skin, tolerance is established; conversely, when exposure via the skin happens before the food is introduced in the diet, food allergy can develop. The dual allergen exposure hypothesis was tested in the Learning Early About Peanut Allergy (LEAP) randomized controlled trial (RCT), which demonstrated that early peanut consumption, from age 4-11 months, prevented the development of peanut allergy by the age of 5 years with 81% relative reduction in the prevalence of peanut allergy (11). This effect was sustained after 1 year of peanut avoidance, suggesting that the preventative effects were long-lasting (12). The higher prevalence of peanut allergy in the avoidance arm was evident immunologically with peanut avoiders having larger wheals on skin prick testing to peanut, higher prevalence of high peanut and Ara h 2-specific IgE levels and higher proportion of activated basophils, with basophil activation being inversely correlated with ratios of IgG4/IgE to peanut (11-13).

Hygiene hypothesis

The hygiene hypothesis was proposed by David Strachan in the 1989 following the observation that children from larger families had less asthma and allergic rhinitis. For food allergies, number of siblings and dog ownership have been identified as protective factors, probably

related to greater microbial exposure (14, 15). The gut microbiome has been implicated in the development of food allergy and available evidence has recently been reviewed (16). With oral tolerance developing in the gut, commensal microbiota can modulate immune development and are required for a healthy immune response to foods (17). Observational studies in humans have showed different microbiota between food allergic and non-allergic individuals and between food allergic children who resolve their food allergies and children with persistent food allergies (18-21). Studies in mouse models have demonstrated that intestinal microbiota can transfer susceptibility or resistance to the development of food allergy, supporting its implication in this condition, and have shed light on the mechanisms of their effects (17, 22). Such mechanistic studies have allowed to understand the modifications to the immune system induced by the microbiota and to identify key regulatory cells induced in the GALT as a consequence of the introduction of solid foods in the diet, confirming that both the microbiota and allergen exposure are required for the establishment of oral tolerance to foods.

Vitamin D hypothesis

Observational studies have documented a greater prevalence of reported food allergies, of emergency visits for food allergic reactions and of prescriptions of adrenaline auto-injectors in geographical areas with lower sun exposure (23). In the Healthnuts study, vitamin D insufficiency was more common in infants with food allergy, with or without eczema, compared to infants with eczema only and to infants with neither eczema nor food allergy (24). Vitamin D insufficiency was also more common in children with multiple food allergies compared to children with a single food allergy (24). However, vitamin D levels can be influenced by various factors, including vitamin D binding protein levels and genetic polymorphisms influencing vitamin D metabolism, which can explain the conflict with results of other studies (25). A recent systematic review of the literature failed to find evidence to

support vitamin D supplementation in pregnancy (1 RCT and 1 observational study), lactation (0 studies), infancy (1 observational study) and childhood (0 studies) for primary food allergy prevention (26). Further studies are needed, possibly with patient stratification according to genotype and taking into account other factors affecting vitamin D levels (23).

Can we stop the food allergy epidemic?

With the alarming increase in prevalence, severity and persistence of food allergy and in the absence of a curative treatment, the hope of changing the current burden of food allergy is on prevention. There are three levels of prevention to consider:

- Primary prevention, i.e. reducing food allergy before allergic sensitisation develops;
- Secondary prevention, i.e. reducing food allergy after allergic sensitisation but before clinical food allergy develops;
- Tertiary prevention, i.e. minimising the consequences of clinical food allergy.

Primary prevention

Recently, a task force of the European Academy of Allergy and Clinical Immunology (EAACI) published a systematic review of RCT covering various interventions that could possibly modify the food allergy risk in the first year of life (27), namely food allergen ingestion during pregnancy or lactation, breastfeeding, milk formulas (standard cow's milk formula, hydrolysed formulas, soy-based formulas), introduction of complementary foods, vitamin supplementation, ingestion of fish oil, prebiotics, probiotics and synbiotics, application of emmollients, prophylactic allergen immunotherapy and BCG vaccination. From the available evidence, the only interventions that showed an effect with low to moderate certainty were the ones related to the introduction of allergenic foods in the diet. Introducing cooked egg and peanut in the diet at weaning (cooked egg from 6 months and peanut from median 7.8 months)

led to an absolute reduction in the prevalence of egg (29%) and peanut (12-23%) allergies with a moderate certainty of effect (28). Avoiding temporary supplementation with standard cow's milk formula in the first week of life was estimated to reduce in 6% the prevalence of cow's milk allergy with low certainty of effect (28).

Based on this systematic review, the EAACI will be soon publishing the guidelines for the prevention of food allergy. Recommendations (**Table 1**) include:

- No avoidance of food allergens during pregnancy or breastfeeding;
- Avoidance of temporary supplementation with cow's milk formula in the first week of life;
- Introduction of well-cooked egg in the infant's diet during weaning;
- Introduction of peanut in the infant's diet during weaning in populations with a high prevalence of peanut allergy.

No recommendations could be made about vitamin or fish oil supplementation, the use of prebiotics, probiotics or synbiotics, emmollients, prophylactic use of allergen-specific immunotherapy or BCG vaccination with the view of preventing the development of food allergy. Other systematic reviews support these recommendations (29, 30). Various studies are ongoing to try to gather high-quality evidence on the effect of interventions on food allergy risk, including RCT of high-allergen diet in the second half of pregnancy and first 4 months of lactation (e.g. peanut and egg in PrEggNut RCT), perinatal administration of probiotics (NCT00200954), ceramide-based emollient (e.g. PEBBLES RCT, NCT03667651), proactive treatment of atopic eczema (e.g. SEAL RCT), introduction of cashew and other tree nuts in the infants' diet (e.g. Cashew RCT, TreEat RCT), vitamin D supplementation (e.g. Vitality RCT, NCT02112734), whole cell pertussis vaccine instead of acellular pertussis vaccine (e.g. Optimum RCT) and BCG vaccination compared to placebo (e.g. MIS BAIR RCT).

Secondary prevention

What can be done to prevent food allergy in infants and children who are already sensitized? Considering the current recommendations in, for instance, Europe (27), USA (31) and Australia (32) (**Table 1**), the main intervention to prevent food allergy in sensitized infants and children is the introduction of allergenic foods in the diet, in the case that they are not allergic to them. In the LEAP study (11), the preventative effect of early peanut consumption was indeed observed in the SPT-positive stratum, although with a lower relative risk reduction compared to the SPT-negative stratum (70% versus 86%, respectively). However, data is absent for infants with SPT of 5 mm or greater as they were excluded from the LEAP study at screening, as they were likely to already have developed peanut allergy (33).

Most guidelines do not recommend allergy testing prior to introduction of the food, due to the eventual need for supervised challenges and the fact that limited resources could cause a delay in the food introduction, which could be counterproductive due to rapid immune changes in the first few months of life and consequent risk of developing food allergy (34, 35). However, the US guidelines recommend testing of high-risk infants prior to peanut introduction and also underscore the importance of being particularly proactive in this subgroup of infants, recommending peanut consumption from 4-6 months of age, whereas for the low risk group the introduction of peanut could be left to after 6 months, depending on the importance of peanut in the family's diet (31).

From the Allergist's perspective, for the family's that seek our advice prior to food introduction, it is justified to test for the foods that are commonly involved in food allergy in children and that the infant or child as yet to be exposed to via the oral route. This is particularly important for peanut in children with early onset severe eczema and/or egg allergy, as we have high-quality evidence with the LEAP study that early consumption of peanut is protective. The use of SPT and specific IgE testing should be based on a consciously defined pre-test

probability and interpreted with caution, based on existing studies on the predictive values of SPT and specific IgE in infants and young children as much as possible (**Table 2**). In children younger than 2 years of age, lower levels of IgE and smaller weals on SPT are more likely to be clinically relevant as reflected by the lower levels for 95% PPV cut-offs and the shift in the curves for the probability of clinical reactivity for a given IgE level compared to older children (36-38). As new modalities of testing are being developed like the basophil activation test, optimal cut-offs seem to be different in this age group and further research is needed (Santos AF, unpublished). Allergy testing prior to weaning should be done with the possibility of offering families supervised introduction of foods, in the office or the hospital setting, depending on the risk of reactivity. This is particularly important in this age group as immunological changes can happen quite rapidly in the first few months of life and the risk of food allergy can change from one assessment to the other. For this reason, ideally, there would be priority access for food challenges for infants as part of specialized Allergy services.

Tertiary prevention

In the cases where food allergy could not be prevented, it is important to minimise the impact of food allergy in the lives of children and their families, the risk of accidental allergic reactions and the progression and complications that could arise from food allergy and accompanying atopic comorbidities (39-41). Following an accurate and definitive diagnosis of food allergy, it is important to recommend appropriate allergen avoidance and emergency medication to treat allergic reactions in the context of accidental allergen exposure. Keeping good control of asthma (and also allergic rhinitis and eczema) is very important as more severe reactions can develop in food allergic children with uncontrolled asthma. Education of patients and family about the nature of food allergies, reading labels and good communication with school, peers and family is paramount. The support of a specialised dietitian is invaluable to advise on the

practical management of food allergy, appropriate replacement foods and adequate nutrient intake. Support from a clinical psychologist may also be beneficial in case of significant anxiety, depression or other mental health issues of child and/or family related with food allergies.

A definitive treatment of food allergy is highly expected. Immunomodulatory treatments have been investigated to try to modify the course of the disease and shut down the aberrant IgE-mediated immune response to foods. Allergen immunotherapy (AIT) has shown to increase the threshold of clinical reactivity while on treatment (desensitisation) with a subset of patients sustaining this effect following discontinuation of treatment (sustained unresponsiveness) (42-44). The foods that have been studied the most are peanut, egg and cow's milk. As AIT involves the administration of allergen to allergic patients, it incurs on a significant risk of allergic reactions during up-dosing and also during the maintenance phase, usually in the presence of co-factors, such as exercise, infection, menstruation and sleep deprivation (45). Some patients do not tolerate the side-effects and end up abandoning the treatment. Epicutaneous have a better safety profile than sublingual oral immunotherapy but the dose of allergen administered and the dose tolerated increase in reverse order. The administration of anti-IgE omalizumab concomitantly with AIT reduces the rate of adverse events and allows more rapid up dosing. Active research on omalizumab (e.g. OUtMATCH, NCT03881696) and other anti-IgE molecules, dupilumab (anti-IL4R – e.g. NCT03793608, NCT03682770, NCT04148352) and etokimab (anti-IL33) (46) in food allergy is being pursued, at different stages of development. Long-lasting disease-modifying effect of immunomodulatory treatments are yet to be demonstrated.

Conclusion:

Food allergy is a significant public health problem and its increase seems to be related to environmental factors related to the Western modern life-style. The main hypothesis to explain this increase in food allergy prevalence, severity and persistence are the timing, dosing and route of exposure to food allergens in infancy, microbial exposure in early life and levels of vitamin D and possibly other factors. High quality evidence is lacking for many possible interventions to modify the food allergy risk. The main recommendation currently to prevent food allergy is to introduce allergenic foods in the infant's diet alongside with breastfeeding as exposure to food allergens via the oral route is necessary for the establishment of oral tolerance and can be protective. Future research needs to focus on the search not only for evidence of interventions for food allergy prevention but also for definitive treatments for food allergy.

Key points:

- Food allergy affects 8% of children and 11% of adults and its prevalence, severity and persistence has increased over time.
- For primary prevention of food allergy, introduction of allergenic foods, namely cooked egg and peanut, during weaning alongside breastfeeding is the main recommendation.
- Immunomodulatory treatments, such as allergen-specific immunotherapy and biologics, have been explored and further research into curative treatments for food allergy is needed.

Acknowledgements: none.

Conflicts of interest: Dr. Santos reports grants and personal fees from Medical Research Council (MR/M008517/1); grants from Asthma UK and the NIHR through the Biomedical Research Centre (BRC) award to Guy's and St Thomas' NHS Foundation Trust, during the conduct of the study; grants from Immune Tolerance Network/National Institute of Allergy and Infectious Diseases (NIAID, NIH), grants from Asthma UK; personal fees from Thermo

Scientific, Nutricia, Infomed, Novartis, Allergy Therapeutics, Buhlmann, as well as research support from Buhlmann and Thermo Fisher Scientific through a collaboration agreement with King's College London.

Tables

Table 1. Recommendations of major Allergy societies for primary prevention of food allergy.

Geographical location	Recommendations	Reference
Europe	<ul style="list-style-type: none"> No avoidance of allergenic foods during pregnancy or breastfeeding. Avoidance of temporary administration of standard cow's milk formula in the first days of life. Introduction of cooked egg and peanut (the latter in populations with high prevalence of peanut allergy) during weaning. 	(27)
USA	Introduction of peanut during weaning: <ul style="list-style-type: none"> for infants with severe eczema and/or egg allergy at 4-6 months with consideration given to prior SPT/specific IgE testing for infants with mild/moderate eczema around 6 months of age for infants without eczema or food allergy, according to family's diet and preferences 	(31)
Australia	<ul style="list-style-type: none"> Weaning between 4-6 months, starting with iron-rich foods, while continuing breast-feeding. Introduction of allergenic foods during weaning for all infants, including peanut butter, cooked egg, dairy, and wheat. Hydrolyzed (partially or extensively) infant formula is not recommended for the prevention of allergic disease. 	(32)

Table 2. Positive predictive value (PPV) cut-offs for skin prick test (100% PPV) and specific IgE (95% PPV) to diagnose common food allergies in infants and young children (38). ND, not determined.

Food	SPT	Specific IgE
Cow's milk	6 mm	5 KU/L
Egg	4 mm	2 KU/L
Peanut	4 mm	ND

295 **Figure Legends:**

296

297 **Figure 1.** Main hypotheses to explain the rise in food allergy in recent decades

298

299

Reference

1. Gupta RS, Warren CM, Smith BM, Blumenstock JA, Jiang J, Davis MM, et al. The Public Health Impact of Parent-Reported Childhood Food Allergies in the United States. *Pediatrics*. 2018;142(6).
2. Perkin MR, Logan K, Tseng A, Raji B, Ayis S, Peacock J, et al. Randomized Trial of Introduction of Allergenic Foods in Breast-Fed Infants. *N Engl J Med*. 2016;374(18):1733-43.
3. Gupta RS, Warren CM, Smith BM, Jiang J, Blumenstock JA, Davis MM, et al. Prevalence and Severity of Food Allergies Among US Adults. *JAMA Netw Open*. 2019;2(1):e185630.
4. Tham EH, Leung ASY, Pacharn P, Lee S, Ebisawa M, Lee BW, et al. Anaphylaxis - Lessons learnt when East meets West. *Pediatr Allergy Immunol*. 2019;30(7):681-8.
5. Botha M, Basera W, Facey-Thomas HE, Gaunt B, Gray CL, Ramjith J, et al. Rural and urban food allergy prevalence from the South African Food Allergy (SAFFA) study. *J Allergy Clin Immunol*. 2019;143(2):662-8 e2.
6. Peters RL, Koplin JJ, Gurrin LC, Dharmage SC, Wake M, Ponsonby AL, et al. The prevalence of food allergy and other allergic diseases in early childhood in a population-based study: HealthNuts age 4-year follow-up. *J Allergy Clin Immunol*. 2017;140(1):145-53 e8.
7. Sasaki M, Koplin JJ, Dharmage SC, Field MJ, Sawyer SM, McWilliam V, et al. Prevalence of clinic-defined food allergy in early adolescence: The SchoolNuts study. *J Allergy Clin Immunol*. 2018;141(1):391-8 e4.
8. Grimshaw KEC, Roberts G, Selby A, Reich A, Butiene I, Clausen M, et al. Risk Factors for Hen's Egg Allergy in Europe: EuroPrevall Birth Cohort. *J Allergy Clin Immunol Pract*. 2020;8(4):1341-8 e5.
9. Martin PE, Eckert JK, Koplin JJ, Lowe AJ, Gurrin LC, Dharmage SC, et al. Which infants with eczema are at risk of food allergy? Results from a population-based cohort. *Clin Exp Allergy*. 2015;45(1):255-64.
10. Brough HA, Nadeau KC, Sindher SB, Alkotob SS, Chan S, Bahnson HT, et al. Epicutaneous sensitization in the development of food allergy: What is the evidence and how can this be prevented? *Allergy*. 2020;75(9):2185-205.
11. Du Toit G, Roberts G, Sayre PH, Bahnson HT, Radulovic S, Santos AF, et al. Randomized trial of peanut consumption in infants at risk for peanut allergy. *N Engl J Med*. 2015;372(9):803-13.
12. Du Toit G, Sayre PH, Roberts G, Sever ML, Lawson K, Bahnson HT, et al. Effect of Avoidance on Peanut Allergy after Early Peanut Consumption. *N Engl J Med*. 2016;374(15):1435-43.
13. Santos AF, Du Toit G, O'Rourke C, Becares N, Couto-Francisco N, Radulovic S, et al. Biomarkers of severity and threshold of allergic reactions during oral peanut challenges. *J Allergy Clin Immunol*. 2020;146(2):344-55.
14. Koplin JJ, Dharmage SC, Ponsonby AL, Tang ML, Lowe AJ, Gurrin LC, et al. Environmental and demographic risk factors for egg allergy in a population-based study of infants. *Allergy*. 2012;67(11):1415-22.
15. Marrs T, Logan K, Craven J, Radulovic S, McLean W, Lack G, et al. Dog ownership at three months of age is associated with protection against food allergy. *Allergy*. 2019;74(11):2212-9.
16. Bunyavanich S, Berin MC. Food allergy and the microbiome: Current understandings and future directions. *J Allergy Clin Immunol*. 2019;144(6):1468-77.

17. Kim KS, Hong SW, Han D, Yi J, Jung J, Yang BG, et al. Dietary antigens limit mucosal immunity by inducing regulatory T cells in the small intestine. *Science*. 2016;351(6275):858-63.
18. Fazlollahi M, Chun Y, Grishin A, Wood RA, Burks AW, Dawson P, et al. Early-life gut microbiome and egg allergy. *Allergy*. 2018;73(7):1515-24.
19. Bunyavanich S, Shen N, Grishin A, Wood R, Burks W, Dawson P, et al. Early-life gut microbiome composition and milk allergy resolution. *J Allergy Clin Immunol*. 2016;138(4):1122-30.
20. Savage JH, Lee-Sarwar KA, Sordillo J, Bunyavanich S, Zhou Y, O'Connor G, et al. A prospective microbiome-wide association study of food sensitization and food allergy in early childhood. *Allergy*. 2018;73(1):145-52.
21. Feehley T, Plunkett CH, Bao R, Choi Hong SM, Culleen E, Belda-Ferre P, et al. Healthy infants harbor intestinal bacteria that protect against food allergy. *Nat Med*. 2019;25(3):448-53.
22. Abdel-Gadir A, Stephen-Victor E, Gerber GK, Noval Rivas M, Wang S, Harb H, et al. Microbiota therapy acts via a regulatory T cell MyD88/RORgammat pathway to suppress food allergy. *Nat Med*. 2019;25(7):1164-74.
23. Hawrylowicz CM, Santos AF. Vitamin D: can the sun stop the atopic epidemic? *Curr Opin Allergy Clin Immunol*. 2020;20(2):181-7.
24. Allen KJ, Koplin JJ, Ponsonby AL, Gurrin LC, Wake M, Vuillermin P, et al. Vitamin D insufficiency is associated with challenge-proven food allergy in infants. *J Allergy Clin Immunol*. 2013;131(4):1109-16, 16 e1-6.
25. Koplin JJ, Suaini NH, Vuillermin P, Ellis JA, Panjari M, Ponsonby AL, et al. Polymorphisms affecting vitamin D-binding protein modify the relationship between serum vitamin D (25[OH]D3) and food allergy. *J Allergy Clin Immunol*. 2016;137(2):500-6 e4.
26. Yepes-Nunez JJ, Brozek JL, Fiocchi A, Pawankar R, Cuello-Garcia C, Zhang Y, et al. Vitamin D supplementation in primary allergy prevention: Systematic review of randomized and non-randomized studies. *Allergy*. 2018;73(1):37-49.
27. de Silva D, Halken S, Singh C, Antonella M, Angier E, Arasi S, et al. Preventing food allergy in infancy and childhood: systematic review of randomised controlled trials. *Pediatr Allergy Immunol*. 2020.
28. Urashima M, Mezawa H, Okuyama M, Urashima T, Hirano D, Gocho N, et al. Primary Prevention of Cow's Milk Sensitization and Food Allergy by Avoiding Supplementation With Cow's Milk Formula at Birth: A Randomized Clinical Trial. *JAMA Pediatr*. 2019.
29. Burgess JA, Dharmage SC, Allen K, Koplin J, Garcia-Larsen V, Boyle R, et al. Age at introduction to complementary solid food and food allergy and sensitization: A systematic review and meta-analysis. *Clin Exp Allergy*. 2019;49(6):754-69.
30. Roberts G, Grimshaw K, Beyer K, Boyle R, Lack G, Austin M, et al. Can dietary strategies in early life prevent childhood food allergy? A report from two iFAAM workshops. *Clin Exp Allergy*. 2019;49(12):1567-77.
31. Togias A. Addendum guidelines for the prevention of peanut allergy in the United States: Report of the National Institute of Allergy and Infectious Diseases-sponsored expert panel. *J Allergy Clin Immunol*. 2017.
32. Netting MJ, Campbell DE, Koplin JJ, Beck KM, McWilliam V, Dharmage SC, et al. An Australian Consensus on Infant Feeding Guidelines to Prevent Food Allergy: Outcomes From the Australian Infant Feeding Summit. *J Allergy Clin Immunol Pract*. 2017;5(6):1617-24.

33. Du Toit G, Roberts G, Sayre PH, Plaut M, Bahnson HT, Mitchell H, et al. Identifying infants at high risk of peanut allergy: the Learning Early About Peanut Allergy (LEAP) screening study. *J Allergy Clin Immunol*. 2013;131(1):135-43 e1-12.
34. Muraro A, Agache I, Clark A, Sheikh A, Roberts G, Akdis CA, et al. EAACI food allergy and anaphylaxis guidelines: managing patients with food allergy in the community. *Allergy*. 2014;69(8):1046-57.
35. Boyce JA, Assa'ad A, Burks AW, Jones SM, Sampson HA, Wood RA, et al. Guidelines for the Diagnosis and Management of Food Allergy in the United States: Summary of the NIAID-Sponsored Expert Panel Report. *J Allergy Clin Immunol*. 2010;126(6):1105-18.
36. Santos AF, Brough HA. Making the Most of In Vitro Tests to Diagnose Food Allergy. *J Allergy Clin Immunol Pract*. 2017;5(2):237-48.
37. Foong RX, Santos AF. Biomarkers of diagnosis and resolution of food allergy. *Pediatr Allergy Immunol*. 2020.
38. R.X. F, J. D, R. W, A.F. S. Improving diagnostic accuracy in food allergy. *J Allergy Clin Immunol Pract*. 2020.
39. Muraro A, Hoffmann-Sommergruber K, Holzhauser T, Poulsen LK, Gowland MH, Akdis CA, et al. EAACI Food Allergy and Anaphylaxis Guidelines. Protecting consumers with food allergies: understanding food consumption, meeting regulations and identifying unmet needs. *Allergy*. 2014;69(11):1464-72.
40. Muraro A, Dubois AE, DunnGalvin A, Hourihane JO, de Jong NW, Meyer R, et al. EAACI Food Allergy and Anaphylaxis Guidelines. Food allergy health-related quality of life measures. *Allergy*. 2014;69(7):845-53.
41. Muraro A, Werfel T, Hoffmann-Sommergruber K, Roberts G, Beyer K, Bindslev-Jensen C, et al. EAACI food allergy and anaphylaxis guidelines: diagnosis and management of food allergy. *Allergy*. 2014;69(8):1008-25.
42. Kulis MD, Patil SU, Wambre E, Vickery BP. Immune mechanisms of oral immunotherapy. *J Allergy Clin Immunol*. 2018;141(2):491-8.
43. Pajno GB, Fernandez-Rivas M, Arasi S, Roberts G, Akdis CA, Alvaro-Lozano M, et al. EAACI Guidelines on allergen immunotherapy: IgE-mediated food allergy. *Allergy*. 2018;73(4):799-815.
44. Nurmatov U, Dhami S, Arasi S, Pajno GB, Fernandez-Rivas M, Muraro A, et al. Allergen immunotherapy for IgE-mediated food allergy: a systematic review and meta-analysis. *Allergy*. 2017;72(8):1133-47.
45. Chu DK, Wood RA, French S, Fiocchi A, Jordana M, Waserman S, et al. Oral immunotherapy for peanut allergy (PACE): a systematic review and meta-analysis of efficacy and safety. *Lancet*. 2019;393(10187):2222-32.
46. Chinthrajah S, Cao S, Liu C, Lyu SC, Sindher SB, Long A, et al. Phase 2a randomized, placebo-controlled study of anti-IL-33 in peanut allergy. *JCI Insight*. 2019;4(22).

Notes on references:

Gupta 2018

**Recent questionnaire-based study performed in the USA providing recent data for prevalence and severity of food allergy in children.*

Gupta 2019

**Recent questionnaire-based study performed in the USA providing recent data for prevalence of food allergy in adults.*

Grimshaw 2020

**Data from Europrevall birth cohort studies establishing moderate-severe eczema as a major risk factor for food allergy.*

Marrs 2019

**Data from the EAT study in the UK demonstrating that dog ownership is a protective factor for the development of food allergy.*

B 2019

**Excellent recent review on the evidence for the role of the microbiome in food allergy.*

Feehley 2019

***Study in mouse models colonised with microbiome from healthy infants and infants with cow's milk allergy related specific bacteria in the ileum with protective and regulatory effects.*

Abdel-Gadir 2019

**Study using mouse models identified a population of regulatory T cells induced as a result of bacteriotherapy to reverse dysbiosis and suppress food allergy.*

Silva 2020

***Systematic review of randomized controlled trials of interventions aiming to prevent the development of food allergy, which formed the basis to the soon-to-be-published EAACI Guidelines for food allergy prevention.*

U 2019

***Randomized controlled trial demonstrating that supplementation with cow's milk formula in the first few days of life increases the risk of cow's milk allergy.*

Roberts 2019

***Meta-analyses of studies of early introduction of allergenic foods demonstrates the benefits of early introduction of egg and peanut (but not milk) in the prevention of food allergy.*

Pajno 2018

***EAACI Guidelines on allergen-specific immunotherapy for food allergy recommending oral immunotherapy for peanut, egg and cow's milk for children older than 4 years of age with persistent allergies to induce desensitization, which should be undertaken only in specialized centres with the appropriate resources to treat anaphylaxis and ensure patient safety during treatment.*

Chu 2019

***Systematic review and meta-analyses showing that patients submitted to oral peanut immunotherapy suffer more allergic reactions and anaphylaxis than those treated with placebo.*