



## King's Research Portal

DOI:

[10.1161/CIRCRESAHA.120.316748](https://doi.org/10.1161/CIRCRESAHA.120.316748)

*Document Version*

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Floyd, C. N., Shahed, F., Ukah, F., McNeill, K., O'Gallagher, K., Mills, C. E., Evangelopoulos, D., Lim, S., Mudway, I., Barratt, B., Walton, H., & Webb, A. J. (2020). Acute Blood Pressure-Lowering Effects of Nitrogen Dioxide Exposure from Domestic Gas Cooking via Elevation of Plasma Nitrite Concentration in Healthy Individuals. *Circulation Research*, 127(6), 847-848. <https://doi.org/10.1161/CIRCRESAHA.120.316748>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# Acute Blood Pressure-Lowering Effects of Nitrogen Dioxide Exposure From Domestic Gas Cooking Via Elevation of Plasma Nitrite Concentration in Healthy Individuals

Christopher N. Floyd<sup>1</sup>, Fawzia Shahed, Frances Ukah, Karen McNeill, Kevin O'Gallagher, Charlotte E. Mills<sup>1</sup>, Dimitris Evangelopoulos, Shanon Lim, Ian Mudway, Benjamin Barratt, Heather Walton, Andrew J. Webb<sup>1</sup>

**A**ir pollution is a major cause of cardiovascular and all-cause mortality. Disentangling the relative contributions of pollutants is challenging, as epidemiological data measuring exposure to one (eg, nitrogen dioxide [NO<sub>2</sub>]) is inevitably confounded by exposure to others (eg, particulate matter). Animal studies suggest that inhaled NO<sub>2</sub> has the potential to increase plasma [nitrite]<sup>1</sup>; a chemical originally considered to be physiologically inert before we found that its reduction to nitric oxide protects the myocardium against ischemia-reperfusion injury and lowers blood pressure in humans.<sup>2</sup>

We conducted an acute, randomized, controlled, crossover study to assess the impact of 90 minutes exposure to NO<sub>2</sub> (from sitting next to a domestic gas cooker with gas hobs lit and uncovered) versus control (room air) on plasma [nitrite] (primary end point) and blood pressure (secondary end points) in 12 healthy participants. All underwent both interventions/visits (interval 7–108 days) in a computer-generated randomized order. Baseline characteristics (mean±SD): 26±4 years, 10/12 female, body mass index 21.9±3.0 kg/m<sup>2</sup>, systolic blood pressure 113.8±7.9 mmHg, diastolic blood pressure 72.8±5.7 mmHg. The exposure phase was followed by a 90 minutes washout phase at background [NO<sub>2</sub>]. Participants fasted for 12 hours before each visit and received 250 mL low-nitrate water at time 0 h/1.5 h. The study was powered for a difference in plasma [nitrite] of 27±40 nmol/L on repeated-measures, 2-way ANOVA

( $\alpha$ , 0.05 and  $\beta$ , 0.2) following D'Agostino-Pearson normality-confirmation, with Sidak post-test (GraphPad Prism v8.2.1).<sup>3</sup>

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Relative to control, exposure increased ambient [NO<sub>2</sub>]: 276.3±38.5 versus 27.6±2.8 ppb ( $P<0.001$ ). Plasma [nitrite] was increased through both the 90 minutes NO<sub>2</sub> exposure and 90 minutes washout ( $P<0.001$ ; Figure [A]). NO<sub>2</sub> exposure decreased both systolic blood pressure and diastolic blood pressure (both  $P<0.001$ ; Figure [B] and [C]). The largest decrease in systolic blood pressure relative to control occurred at 45 minutes (4.6 mmHg [95% CI, 0.2–8.9];  $P=0.032$ ) and 90 minutes (5.5 mmHg [95% CI, 1.2–9.9];  $P=0.005$ ). The effect of NO<sub>2</sub> on diastolic blood pressure was maximal at 45 minutes (5.7 mmHg [95% CI, 0.9–10.5];  $P=0.009$ ).

The temporal relationship between the increase in plasma [nitrite] and systolic blood pressure/diastolic blood pressure reduction ( $\approx 5$  mmHg) is consistent with studies investigating dietary nitrate.<sup>2</sup> Furthermore, whilst the level of NO<sub>2</sub> exposure (276.3±38.5 ppb) was  $\approx 2.5$ -fold greater than recommended limits for exposure (eg, World Health Organization guideline 105 ppb 1-hour mean), it is less than that recorded adjacent to busy roads or in some domestic kitchens ( $\approx 2000$  ppb).


**Key Words:** air pollution ■ blood pressure ■ nitrite ■ nitrogen dioxide ■ particulate matter

Meet the First Author, see p 706

Correspondence to: Andrew Webb, MBBS, PhD, King's College London, St Thomas' Hospital, London, United Kingdom, SE1 7EH. Email [andrew.1.webb@kcl.ac.uk](mailto:andrew.1.webb@kcl.ac.uk)  
For Sources of Funding and Disclosures, see page 848.

© 2020 The Authors. *Circulation Research* is published on behalf of the American Heart Association, Inc., by Wolters Kluwer Health, Inc. This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution, and reproduction in any medium, provided that the original work is properly cited.

*Circulation Research* is available at [www.ahajournals.org/journal/res](http://www.ahajournals.org/journal/res)



**Figure.** Effect of nitrogen dioxide (NO<sub>2</sub>) exposure on plasma [nitrite] and blood pressure.

Plasma [nitrite] (A), systolic blood pressure ( $\Delta$ SBP; B), diastolic blood pressure ( $\Delta$ DBP; C). Comparison between experimental and control shown as ††† $P < 0.001$  and individual timepoints as \* $P < 0.05$  and \*\* $P < 0.01$ . Data expressed as mean  $\pm$  SEM ( $n = 12$ ).

Our model is, therefore, conservative relative to real-world exposure.

Previously, an increase in plasma [nitrite] at 2 hours following diesel exhaust inhalation was thought to be due to particulate matter-mediated induction of inflammatory pathways.<sup>3</sup>

However, our data suggest a more rapid increase in plasma [nitrite] which favors chemical conversion from NO<sub>2</sub> (eg, via a nitrous acid intermediary) and presents a plausible mechanism through which inhaled NO<sub>2</sub> increases plasma [nitrite].<sup>1</sup> This novel ecophysiological

NOx cycle may directly feed into the established nitrate-nitrite-nitric oxide pathway and contribute nitric oxide-mediated cardiovascular effects.<sup>2</sup> Adverse respiratory effects of inhaled NO<sub>2</sub> were not investigated here.<sup>1</sup>

These data must be considered in the context of the strong epidemiological association between NO<sub>2</sub> exposure and cardiovascular mortality.<sup>1</sup> However, particulate matter-free NO<sub>2</sub> does not appear to impair either vascular function, fibrinolysis, or affect heart rate variability in patients with coronary heart disease: parameters adversely affected by increased ambient NO<sub>2</sub> exposure in epidemiological studies.<sup>4,5</sup> This study expands our understanding of how inhaled NO<sub>2</sub> might impact the cardiovascular system, and the role of diet in disease: it is not just what you eat, but how you cook it that matters.

## ARTICLE INFORMATION

### Affiliation

From King's College London (KCL) British Heart Foundation (BHF) Centre, School of Cardiovascular Medicine and Sciences, Department of Clinical Pharmacology, National Institute for Health Research (NIHR) Biomedical Research Centre (BRC), Clinical Research Facility (CRF), Guy's and St Thomas' NHS Foundation Trust (GSTFT), London, UK (C.N.F., F.S., F.U., K.M., K.O.G., A.J.W.); Nutritional Sciences, School of Life Course Sciences, King's College London, UK, Food and Nutritional Sciences, School of Chemistry, Food and Pharmacy, University of Reading, UK (C.E.M.); and NIHR Health Protection Research Unit (HPRU) on Health Impacts of Environmental Hazards at KCL in Partnership with Public Health England (PHE) and Imperial College London (ICL), UK (D.E., S.L., I.M., B.B., H.W.).

### Sources of Funding

King's Together Multi and Interdisciplinary Research Scheme (Wellcome Trust Institutional Strategic Support Fund [204823/Z/16/Z]). H. Walton and D. Evangelopoulos's posts part-funded by NIHR-HPRU on Health Impacts of Environmental Hazards at KCL in Partnership with PHE and ICL, London, UK. Internal infrastructure financial support: KCL-BHF Centre, NIHR-CRF/NIHR-BRC at GSTFT/KCL. Views expressed are authors' (not necessarily NHS/NIHR/DHSC/PHE).

### Disclosures

None.

## REFERENCES

- World Health Organisation. *Review of evidence on health aspects of air pollution – REVIHAAP project: final technical report*. 2013. <https://www.euro.who.int/en/health-topics/environment-and-health/air-quality/publications/2013/review-of-evidence-on-health-aspects-of-air-pollution-revihaap-project-final-technical-report>.
- Khatri J, Mills CE, Maskell P, Odongel C, Webb AJ. It is rocket science - why dietary nitrate is hard to 'beet'! Part I: twists and turns in the realization of the nitrate-nitrite-NO pathway. *Br J Clin Pharmacol*. 2017;83:129–139. doi: 10.1111/bcp.12913
- Langrish JP, Unosson J, Bosson J, Barath S, Muala A, Blackwell S, Söderberg S, Pourazar J, Megson IL, Treweeke A, et al. Altered nitric oxide bioavailability contributes to diesel exhaust inhalation-induced cardiovascular dysfunction in man. *J Am Heart Assoc*. 2013;2:e004309. doi: 10.1161/JAHA.112.004309
- Langrish JP, Lundbäck M, Barath S, Söderberg S, Mills NL, Newby DE, Sandström T, Blomberg A. Exposure to nitrogen dioxide is not associated with vascular dysfunction in man. *Inhal Toxicol*. 2010;22:192–198. doi: 10.3109/08958370903144105
- Scaife A, Barclay J, Hillis GS, Srinivasan J, Macdonald DW, Ross JA, Ayres JG. Lack of effect of nitrogen dioxide exposure on heart rate variability in patients with stable coronary heart disease and impaired left ventricular systolic function. *Occup Environ Med*. 2012;69:587–591. doi: 10.1136/oemed-2011-100126