



King's Research Portal

DOI:

[10.1016/j.biopsych.2014.05.021](https://doi.org/10.1016/j.biopsych.2014.05.021)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Ekonomou, A., Savva, G. M., Brayne, C., Forster, G., Francis, P. T., Johnson, M., Perry, E. K., Attems, J., Somani, A., Minger, S. L., Ballard, C. G., & The Medical Research Council Cognitive Function and Ageing Neuropathology Study (2015). Stage-Specific Changes in Neurogenic and Glial Markers in Alzheimer's Disease. *Biological psychiatry*, 77(8), 711-719. <https://doi.org/10.1016/j.biopsych.2014.05.021>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

28 **Abstract**

29 **Background:** Reports of altered endogenous neurogenesis in people with Alzheimer's disease (AD)
30 and transgenic AD models have suggested that endogenous neurogenesis may be an important
31 treatment target, but there is considerable discrepancy between studies. We examined endogenous
32 neurogenesis and glia changes across the range of pathological severity of AD in people with/without
33 dementia to address this key question.

34 **Methods:** Endogenous neurogenesis and glia in the subventricular zone and dentate gyrus
35 neurogenic niches were evaluated using single and double immunohistochemistry and a validated
36 antibody selection for stage-and-type-specific markers in autopsy tissue from a representative cohort
37 of 28 participants in the MRC Cognitive Function and Ageing study (MRC CFAS). Immunopositive
38 cells were measured blinded to diagnosis, using brightfield and fluorescent microscopy.

39 **Results:** The number of newly-generated neurons significantly declined only in the dentate gyrus of
40 patients with severe tau pathology. No other changes in other neurogenic markers were observed in
41 either of the neurogenic niches. Interestingly, alterations in astrocytes and microglia were also
42 observed in the dentate gyrus across the different stages of tau pathology. No change in any of the
43 markers was observed in individuals who died with dementia compared to those who did not.

44 **Conclusions:** Alterations in endogenous neurogenesis appear to be confined to a reduction in the
45 generation of new neurons in the dentate gyrus of AD patients with severe neurofibrillary tangle
46 pathology and were accompanied by changes in the glia load. These data suggest that intervention
47 enhancing endogenous neurogenesis may be a potential therapeutic target in AD.

48

49 **Introduction**

50 Dementia currently affects more than 34 million people worldwide, with estimations that more than
51 110 million people will be affected by dementia in 2050 (1). Alzheimer's disease (AD), the most
52 common form of dementia, causes enormous personal, social and financial burdens on the patients,
53 their caregivers and society. Current pharmacological treatments offer symptomatic benefits, thus
54 effective disease-modifying therapies are urgently needed. As AD is a neurodegenerative disease,
55 cell replacement strategies are a potential target for therapeutic intervention; such as promoting
56 endogenous neurogenesis.

57 Endogenous neurogenesis is evident in two areas of the brain: the hippocampal dentate gyrus (DG)
58 and the wall of the lateral ventricles (subventricular zone–SVZ) (2-5). In mammals, neural
59 progenitors at the base of the DG granular layer (subgranular layer, SGL) give rise to neurons that
60 can be functionally integrated in the granular cell layer, whereas the SVZ neural progenitors follow a
61 distinct pathway, the Rostral Migratory Stream (RMS), to the olfactory bulb (OB) where they create
62 interneurons. In the healthy adult brain, SVZ neurogenesis maintains cellular turnover in the OB,
63 contributing to olfactory adaptation and learning (6-8), whereas in DG endogenous neurogenesis is
64 crucial for the hippocampal-dependent spatial learning and memory throughout life (8-11).

65 Groundbreaking work over the last two decades has demonstrated the presence of the same
66 neurogenic niches in the adult human brain, including the temporal horn of the lateral ventricles,
67 located adjacent to the hippocampal formation (12-15). Consequently, there has been evolving
68 interest in the therapeutic potential of strategies that aim to enhance endogenous neurogenesis. Many
69 available compounds, of which some are already in clinical use, such as retinoid agonists,
70 cannabinoids, Selective Serotonin Reuptake Inhibitors (SSRIs), cholinesterase inhibitors and certain
71 hormones have a substantial positive impact on neurogenesis in animals, by either stimulating
72 proliferation of endogenous neural stem cells and/or increasing their differentiation into neurons (for
73 reviews see 16 and 17).

74 However, the potential clinical relevance for AD patients is less clear, with contradictory results
75 from the small number of human autopsy studies that have been undertaken. Ziabreva et al. (2006)
76 and Perry et al. (2012) identified an increase at the proliferation stages of neurogenesis in the anterior
77 SVZ (18) and the temporal horn SVZ and DG (19) respectively, but a reduction in the early stage
78 neural progenitors in the SVZ of AD patients compared to age-matched controls (18). In a previous
79 study from our group focussing on a different cohort of AD patients, including people with
80 concurrent cerebrovascular disease, no statistically significant difference was observed in early
81 neuronal marker immunoreactivity between AD and controls (19). In contrast, in another study
82 increased numbers of neural progenitors were detected in the DG of AD patients, which resulted in
83 an unsuccessful maturation to newly-generated neurons (20). In a report focusing on younger
84 patients with AD, increased glial proliferation was reported in the SGL, but no alteration in
85 neurogenesis was identified (21). Other studies suggested that both concurrent cerebrovascular
86 pathology (22-26) and the severity of cortical cholinergic system deficits (18, 19, 27) are likely to
87 represent key mediating factors in either increasing or decreasing endogenous neurogenesis,
88 respectively. Hence, the influence of age-associated neuropathological changes on neurogenesis is
89 not fully elucidated, in particular with respect to the early stages of the AD process.

90 Similar to the often contradictory data from studies on human tissue, studies investigating
91 neurogenesis in transgenic animal models carrying the human mutations for APP and/or PS1 or PS2
92 and/or tau proteins reported increased, decreased or unchanged progenitor activity (28-32).

93 To further elucidate the role of neurogenesis in AD, we examined *post mortem* brain tissue from a
94 subset of participants of the Medical Research Council Cognitive Function and Ageing Study (MRC
95 CFAS), including individuals who died with and without dementia, and who showed all
96 neuropathological stages of AD-associated tau pathology (i.e., Braak stages 0 to VI), without any
97 other neuropathology such as cerebrovascular disease. For the first time, the levels of astrocytic and
98 microglia cell numbers were also identified in the different Braak stages. Our study primarily aimed

to identify alterations in the various phases of endogenous neurogenesis in relation to dementia and AD-associated pathology.

Methods and Materials

Tissue was obtained from brains donated to the UK MRC CFAS. Details of the study have been described elsewhere (33, 34) and can also be found at the website (www.cfes.ac.uk). In brief, MRC CFAS included an initial cohort of 13,004 individuals, representative of the population aged 65 and over recruited from general practice lists in five areas of England and Wales. The cohort for the current study consists of 28 brain samples selected from those participants of the MRC CFAS who agreed to donate their brain on death and among whom a successful autopsy was performed. At the time of sampling, 456 brain donations had been made to the study, 114 of which were available and had sufficient tissue for the current analysis (those from the Cambridgeshire and Newcastle centres). Those who received a neuropathological diagnosis of ‘normal brain’, ‘possible AD’, ‘probable AD’ or ‘definite AD’ were considered for selection into the study. Those with any diagnosis of Lewy body disease, cerebrovascular disease or other neuropathology were excluded. Neurofibrillary tangle (NFT) pathology was assessed using Braak staging (35-37) after histochemistry, by experienced neuropathologists working in the MRC CFAS study, blinded to clinical findings (supplementary figure S1). Neurofibrillary Braak stages are stated based on the topographical distribution of neurofibrillary tangles and neuropil threads which are neuropathological hallmark lesions of AD; at Braak stages I and II, NFTs are confined mainly to the transentorhinal region of the brain; at stages III and IV they are also found in limbic regions such as the hippocampus, and at the severe stages V and VI they are extensively located in other brain areas, including the neocortex (35, 36). The neuropathological diagnosis of AD was done according to internationally accepted criteria that include the assessment of amyloid- β pathology, which progresses spatially and temporally differently than the tau pathology. Of note, similar to many other studies, we used Braak stages to

indicate the overall severity of AD pathology, but not to directly compare the severity of tau pathology with neurogenesis; in this study neither tau nor amyloid- β pathology was directly compared with neurogenesis in the same topographic locations.

Diagnosis of dementia

Dementia status at death was determined based on interviews during the last years of life, including the full GMS-AGECAT diagnostic algorithm that was equivalent to that in the *Diagnostic and Statistical Manual of Mental Disorders*, third edition, revised (DSM-III-R), interviews with informants after the respondent's death when this was possible and death certification (37). Of the 28 individuals included in the present study, 13 received a study diagnosis of dementia at death. Demographic data are shown in table 1.

Immunohistochemistry

8 μ m thick, paraffin- embedded sections were obtained at the level of basal ganglia, including the anterior SVZ, and also at the level of hippocampus, including the temporal horn of the SVZ. Slides were processed for immunohistochemistry and double immunofluorescence (described in supplementary *Methods and Materials*), according to previously published procedures (22, 23, 26, 27).

Cell counts

Cell counting was performed twice, blind to the clinical and neuropathological diagnosis, using a Nikon Eclipse E800 microscope and the NIS elements software for bright field microscopy (version. 2.3, both from Nikon Europe, Netherlands) and a Carl Zeiss Apotome Axioplan 2 microscope and the AxioVision software for the immunofluorescence (version 4.7.2, all from Carl Zeiss, UK). The length of the hippocampus and that of three areas of the SVZ and the neighbouring ependymal cell layer in either the anterior ventricle horn or the temporal ventricle horn were measured on each slide section under a very low magnification and using the relevant microscope software tools. Results

were expressed as the number of immunopositive cells for each antibody/mm of length for all the markers, in order to adjust for variances due to different lengths of the areas measured.

Statistical analysis

For all markers except doublecortin, negative binomial regression analysis was used to model the difference in the number of immunopositive cells per mm length across groups defined by Braak stage. Regression models were adjusted for age at death and for gender. Differences in cell counts per mm between those with and without dementia at autopsy were assessed by negative binomial regression adjusting for age at autopsy, gender and Braak stage. Negative binomial regression is a count based regression model, and allows a number of events (immunopositive cells) to be modelled in terms of covariates of interest (Braak stage and dementia diagnosis), as well as potentially confounding covariates (age and sex), while also taking into account differences in an ‘exposure’ variable (length of tissue sample being examined). Negative binomial regression is a generalisation of Poisson regression in that it allows for heterogeneity in the number of cells per mm across individuals within a group. Poisson regression assumes an even distribution of immunopositive cells across samples within groups and would lead to Type 1 errors since this assumption is highly unlikely to be met. Negative binomial regression results in an estimate of ‘rate’ (cells per mm) in each group standardised to remove the effect of any differences in age and sex across groups and an estimate of difference in terms of an ‘incident rate ratio’ corresponding to the ratio of the rates across groups, assuming all other covariates held constant. It should be noted that the rate estimate is very similar to the raw count of cells/mm in each case, suggesting that the effect of any difference in age and sex across groups in this analysis is minimal.

As a secondary albeit more conventional analysis, we conducted an analysis of variance (ANOVA) across groups treating total cells per mm as a continuous outcome. This led to substantively similar results, however it is likely that the outcome measure of cells per mm violates the assumptions of an

172 ANOVA (normally distribution with equal variances across groups) and so we prefer the negative
173 binomial regression as the primary analysis.

174 The numbers of cells positive for doublecortin were too small for meaningful multivariate analysis or
175 analysis using a count-based regression model, and so for each brain area the proportion of cases
176 with any doublecortin positive cells was compared across groups using Fisher's exact tests.

177 An initial p value of $p < 0.05$ was set for statistical significance, however owing to the large number
178 of hypotheses considered, we subsequently corrected for multiple testing using the method of
179 Benjamini and Hochberg (38) by setting a false discovery rate of $q = 0.05$, leading to a revised critical
180 value of $p < 0.0016$ for each individual hypothesis. Finally, Spearman's correlation coefficients were
181 estimated to identify any association between the detected changes. For the data analysis, IBM SPSS
182 (version 19) and STATA (version 12.1) statistical software were used.

183

184 **Results**

185 Demographic data for the cohort are presented in Table 1. Across the entire study cohort, the mean
186 age at death was 84.8 years (± 8.6 , range 71-103) and 50% of the participants were female (for
187 details see table 1). Statistical analysis showed that the post mortem delay was not related either to
188 Braak Stage and dementia diagnosis ($p > 0.05$ for both) or to the cell counts of any of the markers
189 examined (data not shown).

190 The pattern of immunoreactivity of the neural stem/progenitor cells and their progeny were
191 consistent with our previous descriptions (22, 23). Nestin and doublecortin immunoreactivity was
192 observed in cellular somata and processes, although in the case of doublecortin, immunoreactivity in
193 the processes was rarely observed, possibly due to the post mortem delay, as described previously
194 (21). As doublecortin is also expressed in astrocytes (39), any doublecortin-positive cells with
195 astrocytic appearance (multiple processes) were excluded from counting. PCNA and HuC/D
196 immunoreactivity were detected in the cell nucleus. Figure 1 (A and B) shows the different cell

types, as detected by DAB immunohistochemistry and double immunofluorescence in the hippocampal DG and the subventricular zone.

Dentate gyrus

The estimated number of immunopositive cells per mm length in the DG and the estimated frequency of cells across Braak stages, standardised to the average age at the time of autopsy and for gender is shown in figure 2 (A and B) for both neurogenic markers and glia. There was a lower number of HuC/D-positive post mitotic early neurons in the DG in individuals with Braak staging V-VI (severe neuropathology) compared to those with Braak staging 0-II ($p=0.032$, Figures 1B and 2A).

A significantly higher number of microglia cells, as identified by Iba1 immunohistochemistry, were detected in individuals with Braak stage III-IV compared to the other two groups ($p=0.033$, Figures 2B and 3). However, the differences in early neurons and microglia cells detected were not statistically significant after adjusting for multiple testing.

There was a significantly lower number of GFAP-positive cells (identifying both neural stem cells and astrocytes) among those with Braak stage III-IV (IRR=0.6; 95% CI=0.4-0.9) compared to those with Braak stage 0-II, but a higher number among those in those in Braak stage V-VI (IRR=1.9; 95% CI=1.3-2.8), and this difference across Braak groups remained statistically significant after correction for multiple testing ($p<0.0001$, Figures 2B and 3).

Table 2 shows the difference in cell numbers per mm between those with and without dementia, and the rate ratio standardised for age, gender and Braak stage. There is some evidence of an increase in HuC/D-positive cell numbers in those with dementia (adjusted IRR=2.1; 95% CI=1.0-4.3; $p=0.05$). As the number of doublecortin-positive cells was very low in the DG, a different statistical analysis was performed, as described in *Methods and Materials* section. Table 3 shows the proportion of cases with doublecortin-immunopositive cells in each brain area examined, with some evidence that

these are more commonly found in the DG of those with higher Braak stages (Fisher's exact test $p=0.05$). All of the samples with doublecortin immunopositive cells in the DG were from participants who died with dementia (table 3, Fisher's exact test $p=0.04$), however owing to small numbers of samples and the large number of hypotheses considered these findings should be treated with caution.

In order to examine potential correlations between the neurogenic and glial markers we performed Spearman's correlation (ρ) analysis. There was a significant positive association between newly-generated neurons and the activated microglia ($R=0.52$, $p=0.005$), but negatively associated with the changes of astrocytic cell numbers ($R=-.396$, $p=0.045$) in the DG (Table 4).

SVZ and ependymal cell layer (anterior and temporal horn)

The neurogenic markers remained unchanged at the anterior (at the level of basal ganglia) and temporal (at the level of hippocampus) horns of the SVZ and the ependymal cell layer adjacent to those when analysed by Braak stage and by the presence of dementia (supplemental table S1 and table 3). PCNA-HuC/D double and HuC/D single immunolabelled cells were not detected in these areas.

The astrocytic and microglia markers also remained unchanged in the SVZ neurogenic niche and neighbouring ependymal cell layer in Braak stages 0 to VI (supplemental table S1).

Discussion

A detailed analysis of endogenous neurogenesis in both neurogenic niches at various stages of tau pathology in 28 people with and without AD was undertaken from a representative community cohort of patients followed to autopsy (MRC CFA Study). This is a population-based representative pilot study of endogenous neurogenesis in AD using human autopsy tissue, and focuses on the full range of pathological disease severity, the cognitive status and both neurogenic niches from each individual, compared to our previous studies (18, 19). Furthermore, the study has the methodological

248 advantage of excluding patients with concurrent cerebrovascular pathology, Lewy body disease and
249 any other pathology that have been found to affect adult endogenous neurogenesis (22-27, 40), as
250 opposed to previous studies of endogenous neurogenesis in AD (20, 21), including our previous
251 studies focussing on different patient cohorts (18, 19) where the percentage area of immunoreactivity
252 was measured rather than number of immunopositive cells.

253 There were limited but statistically significant changes in one of the markers used in this study in
254 individuals with dementia or at the different Braak stages. Specifically, we observed a significant
255 reduction in newly generated neurons, as determined by single HuC/D immunohistochemistry, but
256 not neural progenitors as determined by nestin, doublecortin and double PCNA-HuC/D
257 immunolabelling in people with severe tau pathology (Braak stage V and VI), compared to those
258 with no significant tau changes (Braak stage 0-II). Interestingly and consistent with other recent
259 studies about the role of microglia in modulating endogenous neurogenesis, there was a significant
260 relationship with activated microglia.

261 There were no changes in neural stem cells or progenitors in the anterior and temporal horn of the
262 SVZ, which is in contrast with the increased numbers of neural stem cells and progenitors observed
263 in the SVZ of patients suffering from stroke, vascular dementia, dementia with Lewy bodies and
264 small vessel disease, as our previous studies have shown (22, 23, 26, 27, 40). Markers of
265 neurogenesis did not vary between those with and without dementia after adjusting for Braak stage,
266 thus the reduced numbers of new neurons were specifically associated with the severe Alzheimer
267 pathology.

268 Earlier studies examining endogenous neurogenesis in the course of AD have produced conflicting
269 results, possibly because of the predominant focus on more severe AD, variable concurrent vascular
270 changes and the limited focus on the later stages of neurogenesis, i.e. the newly-generated neurons.

271 The only other study to examine early neurons, as well as progenitors, reported a significant

reduction on their maturation as marked by the the decreased levels of MAP2a and MAP2b isoforms in the DG of AD patients (20), consistent with our current findings.

The present study exhibits some limitations, mainly due to methodology. As there are no reliable and/or applicable markers to “visualise” and follow the fate of neural stem cells in the adult brain, it is impossible to draw any conclusions about adult endogenous neurogenesis during the lifespan of the participants in the study, so all our results represent adult neural stem cells and their progeny at a single time point, that of autopsy. Optimistically, the development of new technologies will facilitate this and clarify more the role of adult neurogenesis and its involvement in cognitive decline in ageing and AD.

As exclusion criteria were applied to the cases in reference to the presence of other neurodegenerative diseases, our sample was modest (n=28), but it was a population-representative it is pilot study of endogenous neurogenesis focussing upon “pure” AD in human autopsy tissue, including individuals with different stages of AD pathology.

The use of antibodies as markers of endogenous neurogenesis on human autopsy tissue represents a challenge, but we employed a validated battery of antibodies for the identification of progenitors and newly-developed neurons at the various stages of neurogenesis in human autopsy tissue. Long post-mortem delay, quite common factor for obtaining human tissue, has been shown to alter but not eliminate the immunostaining pattern for doublecortin, with similar overall levels of staining but reduced staining within the soma. For example, Boekhoorn et al. (2006) have shown that post mortem delay reduced immunoreactivity within the dendrites of doublecortin-positive neuroblasts (21). A similar pattern was seen for doublecortin in our study, but importantly the changes in the overall pattern of staining by dementia stage were similar for other markers of neuroblasts/immature neurons (HuC/D with PCNA). As it has been suggested that DCX can also be expressed in astrocytes (39) and in dormant cells in non neurogenic areas (41, 42), the use of HuC/D as an additional marker for neuronal progenitors/early neurons is important and limits the possibility of over-

297 interpreting results obtained from DCX immunohistochemistry. In addition, post-mortem delay was
298 not significantly correlated with the overall level of staining. Many hypotheses were explored and
299 when a correction was applied to ensure a false discovery rate of less than 0.05 only the changes
300 detected in the numbers of GFAP-immunopositive cells in the DG remained still statistically
301 significant across groups.

302 As there is an age-related decline in neurogenesis observed also in humans (43), our estimates have
303 also been adjusted for age, which further confirms the validity of our results. Physical activity and
304 certain pharmacological treatments such as SSRIs (for a review see 16, 17 and references therein)
305 can have an impact on neurogenesis in rodent models (44-51 and for a review 16, 17). These were
306 not specifically examined in the current study, but it is unlikely that the magnitude of these effects
307 would be sufficient to confound the analysis.

308 The PCNA immunohistochemistry (Figure 1B) also revealed a number of cells that are not co-
309 expressing the neuronal fate marker HuC/D (green only cells in figure 1B). Although these cells
310 were not counted for the study, we can speculate that they could represent astrocytic, microglial, or
311 endothelial cell proliferating progenitors or cells re-entering the cell cycle according to Yang et al.
312 (2003) (52), although that was not observed in the DG.

313 Our results still showed some evidence that although the early stages of endogenous neurogenesis
314 remained unchanged throughout the different Braak stages, severe AD pathology had a detrimental
315 effect on the numbers of newly-generated neurons in the DG of the affected individuals. In contrast,
316 endogenous neurogenesis at all stages and areas remained unchanged in individuals with dementia
317 compared to people without, suggesting that severe AD pathology impaired only the production of
318 new neurons.

319 We used Braak staging, a neurofibrillary tangle-based staging system, to describe the overall severity
320 of AD. Amyloid and tau pathology both increase with increasing disease severity, and many other
321 concurrent pathways related to a broad range of changes including inflammation and mitochondrial

function are activated. The specific mechanisms associated with the reduced production of new neurones in people with severe AD are therefore difficult to unravel from the results of the current study. Previous work has however suggested that tau transgenic mice (29, 53) do have reduced neurogenesis, supporting the potential role of tau pathology as a contributor to this effect.

There was a significant positive correlation between the cell numbers of activated microglia and those of the newly-generated neurons in the DG, suggesting that the reduction in activated microglia in people with Braak stage V- VI tangle pathology may be a key driver for the decline in the newly-produced neurons in these individuals, along with the presence of tangles. Microglia have an important role in adult neurogenesis in the healthy brain, as it has been demonstrated to control the numbers of newly-produced neuron in the hippocampus through apoptosis (54) and can have both pro- and anti- neurogenic effects, finely “tuning” adult neurogenesis (for a review see 55).

For the first time, changes in GFAP-positive astrocytes have been examined at the various Braak stages. Though we have no knowledge of the causal factors of these changes, one can hypothesize that as astrocytes have a significant role in the support and protection of neurons in the healthy brain, the decrease identified at Braak stage III-IV, below the levels seen in individuals with Braak stage 0-II, could be another contributing factor to the disease progression and pathology. The two-fold increase above the levels observed in the healthy brain in astrocytes numbers in the Braak stage V-VI could have a detrimental effect on the diseased brain, as it has been described for certain neurodegenerative diseases, including dementia (for a review see 56). A separate study further investigating this hypothesis could clarify that observation and the underlying mechanism(s).

Our study examined the fate of neuronal progenitors and their progeny at the DG and the anterior and temporal horn SVZ and adjacent ependymal layers in various stages of AD, without any effect from concurrent cerebrovascular or other neuropathology, and showed that specific and significant reductions in newly-generated neurons were detected only in the DG of those with severe AD pathology, and were associated to the microglial load of the area. Previous studies (57-59)

identifying that abnormal endogenous neurogenesis relates to age-related learning impairment have indicated that the manipulation of endogenous neurogenesis may be a potential treatment target in people with AD. As a cross-sectional autopsy study, our results have to be interpreted cautiously, but our findings do support the concept for an enhancement of aspects of endogenous neurogenesis, as a possible treatment target in AD.

Acknowledgements

The authors would like to thank Mr Carl Hobbs and Dr David Howlett at the Wolfson Centre for Age-Related Diseases, King's College London for technical assistance and advice with aspects of the immunohistochemical procedures. This study was supported by Research Into Ageing-Age UK through a Research Fellowship to A.E (Grant ref. no. 309). The CFAS study is supported by the Department of Health and the Medical Research Council (grants MRC/G9901400 and MRC U.1052.00.0013); the UKNIHR Biomedical Research Centre for Ageing and Age - related Disease Award to the Newcastle up Tyne Hospitals Foundation Trust; the Cambridge Brain Bank is supported by the NIHR Cambridge Biomedical Research Centre; The Cambridgeshire and Peterborough NIHR CLAHRC; Nottingham University Hospitals NHS Trust; University of Sheffield and the Sheffield Teaching Hospitals NHS Foundation Trust; The Thomas Willis Oxford Brain Collection, supported by the Oxford Biomedical Research Centre; The Walton Centre NHS Foundation Trust, Liverpool; and Brains for Dementia Research. . C.G.B. and P.T.F. would like to thank the National Institute for Health Research (NIHR) Mental Health Biomedical Research Centre and Dementia Unit at South London and Maudsley NHS Foundation Trust and [Institute of Psychiatry] King's College London. We would like to acknowledge the essential contribution of the liaison officers, the general practitioners, their staff, and nursing and residential home staff. We are grateful to our respondents and their families for their generous gift to medical research, which has made this study possible.

372

373 Corporate CFAS authorship: Professor Carol Brayne (University of Cambridge); Dr. Thais Minett
374 (University of Cambridge); Dr. Fiona Matthews (University of Cambridge); Sally Hunter (University
375 of Cambridge), Dr Tuomo Polvikoski (Newcastle University); Professor Paul Ince (University of
376 Sheffield); Professor Steve Wharton (University of Sheffield); Gill Forster (University of Sheffield);
377 Dr. Cherie McCracken (University of Liverpool); Simon Harrison (Exeter); Dr. Hannah Keage
378 (University of South Australia).

379

380

381 **Financial Disclosures**

382 Within the last three years, Prof. Clive Ballard has received research grants from Lundbeck
383 pharmaceutical company and fees for consultancy or speaking from Lundbeck, Acadia, Bristol-Myer
384 Squibb, Bial and Novartis pharmaceutical companies. None of this work is however directly relevant
385 to the submitted manuscript. Within the last three years, Prof. Paul Francis has received research
386 grants from Lundbeck pharmaceutical company and fees for consultancy or speaking from Lundbeck
387 and Novartis pharmaceutical companies. He has also received payment related to expert witness
388 testimony related to cases involving Novartis and Janssen Alzheimer Immunotherapy. None of this
389 work is however directly relevant to the submitted manuscript.

390 All other authors report no biomedical financial interests or potential conflicts of interest.

391

392 **Table 1: Demographic data for the cases used in the present study**

	Braak stage		
	0-II	III-IV	V-VI
Age (years) ± SD	n=12 80.3±8.4	n=11 88.9±8.2	n=5 86.8±5.3
Gender	Female: 5; Male: 7	Female: 8; Male: 3	Female: 1; Male: 4
Dementia	n=3	n=5	n=5
Gender	Female: 2; Male: 1	Female: 4; Male: 1	Female: 1; Male: 4
PM delay in hours median (IQR)	17.5 (12-28)	25 (7-27)	17.5 (9.5-33)

393

394 Data represent the mean or median in each group. SD: standard deviation, PM delay: Post-mortem
 395 delay, IQR: InterQuartile Range.

Table 2: Cell numbers and area lengths (mm) for the neurogenic markers in the human dentate gyrus by study diagnosis of dementia at death.

Antibody	No dementia (n=15)			Dementia (n=13)		
	Cells/mm (raw data)	Adjusted Cells/mm	IRR	Cells/mm (raw data)	Adjusted Cells/mm	IRR
Nestin	0.094	0.13 (0.08)	Ref.	0.10	0.09 (0.05)	0.7 (0.1-4.0)
HuC/D-PCNA	0.88	1.0 (0.3)	Ref.	0.61	0.6 (0.2)	0.6 (0.2-1.6)
HuC/D	0.74	0.6 (0.1)	Ref.	0.73	1.3 (0.4) *	2.1 (1.0-4.3)
GFAP	4.79	5.2 (0.7)	Ref.	2.46	4.9 (0.5)	0.9 (0.7-1.4)
Iba1	7.82	7.3 (1.9)	Ref.	5.57	7.0 (2.2)	1.0 (0.4-2.3)

Rates and incident rate ratios for the difference in cell density in the dementia group compared to the no dementia group are estimated by negative binomial regression, adjusted for Braak stage. Cells per mm are standardised to the sample age and gender and numbers in parentheses represent standard error; IRR = Incidence rate ratio adjusted for age and gender, numbers in parentheses represent 95% confidence interval. *: p=0.045.

Table 3: Occurrence of Doublecortin immunoreactivity in the adult human brain neurogenic niches

Antibody	Braak 0-II (n=12)	Braak III-IV (n=11)	Braak V-VI (n=5)	Fisher's exact test p-value	No dementia (n=15)	Dementia (n=13)	Fisher's exact test p-value
DG	0 (0)	2 (20)	2 (40)	0.05	0 (0)	4 (31)	0.04
SVZ	1 (8.3)	2 (20)	0 (0)	0.57	1 (7)	2 (15)	0.58
EP	1 (8.3)	1 (10)	0 (0)	1.00	1 (7)	1 (8)	1.00
SVZ BG	4 (33)	1 (9)	0 (0)	0.22	2 (13)	3 (23)	0.64
EP BG	7 (58)	6 (54)	1 (20)	0.47	9 (60)	5 (38)	0.45

The numbers represent number of samples with doublecortin immunopositive cells in each brain area by Braak stage and dementia status where number in parentheses show the % of the same samples in each subgroup by Braak stage and dementia status. SVZ BG: Subventricular Zone at the level of basal ganglia (anterior horn). EP BG: Ependymal cell layer at the level of basal ganglia, adjacent to SVZ BG.

Table 4: Correlation analysis among the observed changes in the dentate gyrus

			GFAP DG	HuC/D DG	Iba1 DG
Spearman's rho	GFAP DG	<i>R</i>	1.000	-.396*	-.476*
	HuC/D DG	<i>R</i>	-.396*	1.000	.524**
	Iba1 DG	<i>R</i>	-.476*	.524**	1.000

Statistical analysis was performed using a two-tailed Spearman's rho correlation analysis. *R*: correlation coefficient. Statistical significance: *: $p < 0.05$ and **: $p < 0.01$.

References

1. Alzheimer's Disease International: World Alzheimer report 2009.
<http://www.alz.co.uk/research/world-report>.
2. Lois C, Garcia-Verdugo JM , Alvarez-Buylla A (1996): Chain migration of neuronal precursors. *Science* 271: 978-981.
3. Doetsch F, Garcia-Verdugo JM , Alvarez-Buylla A (1997): Cellular composition and three-dimensional organization of the subventricular germinal zone in the adult mammalian brain. *J Neurosci* 17: 5046-5061.
4. Palmer TD, Takahashi J , Gage FH (1997): The adult rat hippocampus contains primordial neural stem cells. *Mol Cell Neurosci* 8: 389-404.
5. Ma DK, Bonaguidi MA, Ming GL , Song H (2009): Adult neural stem cells in the mammalian central nervous system. *Cell Res* 19: 672-682.
6. Rochefort C, Gheusi G, Vincent JD , Lledo PM (2002): Enriched odor exposure increases the number of newborn neurons in the adult olfactory bulb and improves odor memory. *J Neurosci* 22: 2679-2689.
7. Magavi SS, Mitchell BD, Szentirmai O, Carter BS , Macklis JD (2005): Adult-born and preexisting olfactory granule neurons undergo distinct experience-dependent modifications of their olfactory responses in vivo. *J Neurosci* 25: 10729-10739.
8. Imayoshi I, Sakamoto M, Ohtsuka T, Takao K, Miyakawa T, Yamaguchi M, *et al.* (2008): Roles of continuous neurogenesis in the structural and functional integrity of the adult forebrain. *Nat Neurosci* 11: 1153-1161.
9. Ge S, Yang CH, Hsu KS, Ming GL , Song H (2007): A critical period for enhanced synaptic plasticity in newly generated neurons of the adult brain. *Neuron* 54: 559-566.

10. Kee N, Teixeira CM, Wang AH , Frankland PW (2007): Preferential incorporation of adult-generated granule cells into spatial memory networks in the dentate gyrus. *Nat Neurosci.* 10: 355-362.
11. Tashiro A, Makino H , Gage FH (2007): Experience-specific functional modification of the dentate gyrus through adult neurogenesis: a critical period during an immature stage. *J Neurosci.* 27: 3252-3259.
12. Eriksson PS, Perfilieva E, Bjork-Eriksson T, Alborn AM, Nordborg C, Peterson DA *et al.* (1998): Neurogenesis in the adult human hippocampus. *Nat Med* 4: 1313-1317.
13. Sanai N, Tramontin AD, Quinones-Hinojosa A, Barbaro NM, Gupta N, Kunwar S, *et al.* (2004): Unique astrocyte ribbon in adult human brain contains neural stem cells but lacks chain migration. *Nature* 427: 740-744.
14. Quinones-Hinojosa A, Sanai N, Soriano-Navarro M, Gonzalez-Perez O, Mirzadeh Z, Gil-Perotin S, *et al.* (2006): Cellular composition and cytoarchitecture of the adult human subventricular zone: a niche of neural stem cells. *J Comp Neurol* 494: 415-434.
15. Curtis MA, Kam M, Nannmark U, Anderson MF, Axell MZ, Wikkelsso C, *et al.* (2007): Human neuroblasts migrate to the olfactory bulb via a lateral ventricular extension. *Science* 315: 1243-1249.
16. Abrous DN, Koehl M , Le Moal M (2005): Adult neurogenesis: from precursors to network and physiology. *Physiol Rev* 85: 523-569.
17. Vaidya VA, Vadodaria KC, Jha S (2007): Neurotransmitter regulation of adult neurogenesis: putative therapeutic targets. *CNS Neurol Disord Drug Targets* 6(5): 358-74.
18. Ziabreva I, Perry E, Perry R, Minger SL, Ekonomou A, Przyborski S , Ballard C (2006): Altered neurogenesis in Alzheimer's disease. *J Psychosom Res* 61: 311-316.

19. Perry EK, Johnson M, Ekonomou A, Perry RH, Ballard C, Attems J (2012): Neurogenic abnormalities in Alzheimer's disease differ between stages of neurogenesis and are partly related to cholinergic pathology *Neurobiol Dis* 47(2): 155-162.
20. Li B, Yamamori H, Tatebayashi Y, Shafit-Zagardo B, Tanimukai H, Chen S, *et al.* (2008): Failure of neuronal maturation in Alzheimer disease dentate gyrus. *J Neuropathol Exp Neurol* 67: 78-84.
21. Boekhoorn K, Joels M, Lucassen PJ (2006): Increased proliferation reflects glial and vascular-associated changes, but not neurogenesis in the presenile Alzheimer hippocampus. *Neurobiol Dis* 24: 1-14.
22. Ekonomou A, Ballard CG, Pathmanaban ON, Perry RH, Perry EK, Kalaria RN, Minger SL (2011): Increased neural progenitors in vascular dementia. *Neurobiol Aging* 32: 2152-2161.
23. Ekonomou A, Johnson M, Perry RH, Perry EK, Kalaria RN, Minger SL, Ballard CG (2012): Increased neural progenitors in individuals with cerebral small vessel disease. *Neuropathol Appl Neurobiol* 38(4): 344-353.
24. Jin K, Wang X, Xie L, Mao XO, Zhu W, Wang Y, *et al.* (2006): Evidence for stroke-induced neurogenesis in the human brain. *Proc Natl Acad Sci USA* 103: 13198-13202.
25. Macas J, Nern C, Plate KH, Momma S (2006): Increased generation of neuronal progenitors after ischemic injury in the aged adult human forebrain. *J Neurosci* 26: 13114-13119.
26. Minger SL, Ekonomou A, Carta EM, Chinoy A, Perry RH, Ballard CG (2007): Endogenous neurogenesis in the human brain following cerebral infarction. *Regen Med* 2: 69-74.
27. Johnson M, Ekonomou A, Hobbs C, Ballard CG, Perry RH, Perry EK (2011): Neurogenic marker abnormalities in the hippocampus in dementia with Lewy bodies. *Hippocampus* 21: 1126-1136.

28. Demars M, Hu YS, Gadadhar A , Lazarov O (2010): Impaired neurogenesis is an early event in the etiology of familial Alzheimer's disease in transgenic mice. *J Neurosci Res* 88: 2103-2117.
29. Hamilton LK, Aumont A, Julien C, Vadnais A, Calon F , Fernandes KJ (2010): Widespread deficits in adult neurogenesis precede plaque and tangle formation in the 3xTg mouse model of Alzheimer's disease. *Eur J Neurosci* 32: 905-920.
30. Jin K, Galvan V, Xie L, Mao XO, Gorostiza OF, Bredesen DE , Greenberg DA (2004): Enhanced neurogenesis in Alzheimer's disease transgenic (PDGF-APP^{Sw,Ind}) mice. *Proc Natl Acad Sci U S A* 101: 13363-13367.
31. Wen PH, Hof PR, Chen X, Gluck K, Austin G, YOUNKIN SG, *et al.* (2004): The presenilin-1 familial Alzheimer disease mutant P117L impairs neurogenesis in the hippocampus of adult mice. *Exp Neurol* 188: 224-237.
32. Ermini FV, Grathwohl S, Radde R, Yamaguchi M, Staufenbiel M, Palmer TD, Jucker M (2008): Neurogenesis and alterations of neural stem cells in mouse models of cerebral amyloidosis. *Am J Pathol* 172: 1520-1528.
33. Brayne C, McCracken C, Matthews FE (2006): Cohort profile: the Medical Research Council Cognitive Function and Ageing Study (CFAS). *Int J Epidemiol* 35: 1140-1145.
34. Matthews FE, Brayne C, Lowe J, McKeith I, Wharton SB , Ince P (2009): Epidemiological pathology of dementia: attributable-risks at death in the Medical Research Council Cognitive Function and Ageing Study. *PLoS Med* 6: e1000180.
35. Braak, H. and Braak, E. (1991): "Neuropathological staging of Alzheimer-related changes". *Acta Neuropathol* 82 (4): 239–59.
36. Braak H, Alafuzoff I, Arzberger T, Kretschmann H, Del Tredici K. (2006): Staging of Alzheimer disease-associated neurofibrillary pathology using paraffin sections and immunohistochemistry. *Acta Neuropathol* 112: 389–404.

37. Savva GM, Wharton SB, Ince PG, Forster G, Matthews FE, Brayne C; Medical Research Council Cognitive Function and Ageing Study (2009): Age, neuropathology and dementia. *N Engl J Med* 360(22): 2302-2309.
38. Benjamini Y and Hochberg Y (1995): Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society, Series B*, 57: 289-300.
39. Verwer RW, Sluiter AA, Balesar RA, Baayen JC, Noske DP, Dirven CM, *et al.* (2007): Mature astrocytes in the adult human neocortex express the early neuronal marker doublecortin. *Brain*. 130(Pt 12):3321-35.
40. Ziabreva I, Ballard C, Johnson M, Larsen JP, McKeith I, Perry R, *et al.* (2007): Loss of Musashi1 in Lewy body dementia associated with cholinergic deficit. *Neuropathol Appl Neurobiol* 33(5): 586-590.
41. Kremer T, Jagasia R, Herrmann A, Matile H, Borroni E, Francis F, *et al.*, (2013): Analysis of adult neurogenesis: evidence for a prominent "non-neurogenic" DCX-protein pool in rodent brain. *PLoS One*, 8(5):e59269
42. Martí-Mengual U, Varea E, Crespo C, Blasco-Ibáñez JM, Nacher J. (2013): Cells expressing markers of immature neurons in the amygdala of adult humans. *Eur J Neurosci*. 37(1):10-22.
43. Kempermann G, Gast D, Gage FH (2002): Neuroplasticity in old age: sustained fivefold induction of hippocampal neurogenesis by long-term environmental enrichment. *Ann Neurol* 52(2): 135-43.
44. van Praag H, Shubert T, Zhao C, Gage FH. (2005): Exercise enhances learning and hippocampal neurogenesis in aged mice. *J Neurosci*. 25(38): 8680-5.
45. Leem YH, Lim HJ, Shim SB, Cho JY, Kim BS, Han PL (2009): Repression of tau hyperphosphorylation by chronic endurance exercise in aged transgenic mouse model of tauopathies. *J Neurosci Res*. 87(11): 2561-70.

46. Marlatt MW and Lucassen PJ (2010): Neurogenesis and Alzheimer's disease: Biology and pathophysiology in mice and men. *Curr Alzheimer Res* 7(2): 113-25.
47. Belarbi K, Burnouf S, Fernandez-Gomez FJ, Laurent C, Lestavel S, Figeac M, *et al.* (2011): Beneficial effects of exercise in a transgenic mouse model of Alzheimer's disease-like Tau pathology. *Neurobiol Dis* 43(2): 486-94.
48. García-Mesa Y, López-Ramos JC, Giménez-Llort L, Revilla S, Guerra R, Gruart A, Laferla FM, *et al.* (2011): Physical exercise protects against Alzheimer's disease in 3xTg-AD mice. *J Alzheimers Dis* 24(3): 421-54.
49. Marlatt MW, Potter MC, Lucassen PJ, van Praag H (2012): Running throughout middle-age improves memory function, hippocampal neurogenesis, and BDNF levels in female C57BL/6J mice. *Dev Neurobiol.* 72(6): 943-52.
50. Marlatt MW, Potter MC, Bayer TA, van Praag H, Lucassen PJ. (2013): Prolonged running, not fluoxetine treatment, increases neurogenesis, but does not alter neuropathology, in the 3xTg mouse model of Alzheimer's disease. *Curr Top Behav Neurosci.* 15:313-40.
51. Voss MW, Vivar C, Kramer AF, van Praag H. (2013): Bridging animal and human models of exercise-induced brain plasticity. *Trends Cogn Sci.*, 17(10): 525-44.
52. Yang Y, Mufson EJ, Herrup K (2003): Neuronal cell death is preceded by cell cycle events at all stages of Alzheimer's disease. *J Neurosci.* 23(7):2557-63.
53. Hong XP, Peng CX, Wei W, Tian Q, Liu YH, Yao XQ, Zhang Y, Cao FY, Wang Q, Wang JZ (2010): Essential role of tau phosphorylation in adult hippocampal neurogenesis. *Hippocampus* 20(12):1339-49.
54. Sierra A, Encinas JM, Deudero JJ, Chancey JH, Enikolopov G, Overstreet-Wadiche LS, *et al.* (2010): Microglia shape adult hippocampal neurogenesis through apoptosis-coupled phagocytosis. *Cell Stem Cell.* 7: 483-495.

55. Ekdahl CT (2012): Microglial activation - tuning and pruning adult neurogenesis. *Front Pharmacol* 3: 411-419.
56. Verkhratsky A, Olabarria M, Noristani HN, Yeh CY and Rodriguez JJ (2012): Astrocytes in Alzheimer's disease. *Neurotherapeutics*. 7(4): 399-412.
57. Nyffeler M, Yee BK, Feldon J, Knuesel I. (2010): Abnormal differentiation of newborn granule cells in age-related working memory impairments. *Neurobiol Aging*. 31(11): 1956-74.
58. Kim SE, Ko IG, Kim BK, Shin MS, Cho S, Kim CJ, et al. (2010): Treadmill exercise prevents aging-induced failure of memory through an increase in neurogenesis and suppression of apoptosis in rat hippocampus. *Exp. Gerontol*. 45(5): 357-65.
59. Déry N, Pilgrim M, Gibala M, Gillen J, Wojtowicz JM, Macqueen G, Becker S. (2013): Adult hippocampal neurogenesis reduces memory interference in humans: opposing effects of aerobic exercise and depression. *Front Neurosci*. 7: 66.

Figure legends

Figure 1: Immunohistochemistry showing neurogenesis in the dentate gyrus (DG) and the subventricular zone (SVZ) of the adult human brain. **A:** DAB immunohistochemistry for nestin (top left, arrows) and doublecortin (top right, arrow) depicting neural stem cell/progenitors and late neural progenitors, respectively in both neurogenic niches. Scale bar: 5um. **B:** Double immunofluorescence for HuC/D (red) and PCNA (green) in severe AD patients (Braak stage V-VI) and their age-matched controls (Braak stage 0-II). There are significantly less postmitotic immature neurons (arrows, HuC/D-immunopositive cells) in the DG of severe AD patients compared to their age-matched controls. Yellow arrowheads indicate proliferating neuronal progenitors/neuroblasts immunopositive for both PCNA (a proliferating marker) and the HuC/D antigen. Images **i** and **ii** show higher magnification of double labelled cells at Braak stage 0-II and V-VI, respectively. Scale bar: 20um. **C:** Schematic representation of the markers used for the characterization of the different stages of neurogenesis in the present study. **Gr:** dentate gyrus granular layer, **h:** hilus, **LV:** lateral ventricle, **Ep:** Ependymal cell layer.

Figure 2: Graph bars represent the adjusted mean number (\pm standard error) of immunopositive cells per mm of length of dentate gyrus for neurogenic (A) and glial (B) markers in AD patients at different Braak stages, adjusted for age and dementia status. *: $p < 0.05$, **: $p < 0.001$.

Figure 3: Immunofluorescence for astrocytes (GFAP) and microglia (Iba1) in age-matched individuals at Braak stage 0-II and patients with moderate (Braak stage III-IV) and severe (Braak stage V-VI) tangle pathology. **DG:** dentate gyrus, **h:** hilus. Scale bar: 50 μ m.

Figure 1
[Click here to download high resolution image](#)




Figure 2
[Click here to download high resolution image](#)

Dentate Gyrus





Figure 3
[Click here to download high resolution image](#)



Stage-Specific Changes in Neurogenic and Glial Markers in Alzheimer's Disease

Supplemental Information

Supplementary Materials and Methods

Immunohistochemistry

After deparaffinization, rehydration and microwave pressure cooking antigen retrieval using 10 mM citrate buffer, pH 6, slides were processed for immunohistochemistry using DAB staining and double immunofluorescence. A validated battery of antibodies was used in order to identify the type of neural stem cells and progenitors as marked by nestin (1/200, Chemicon) (1) and doublecortin (1/200, Santa Cruz) (2) and immature neurons (HuC/D, 1/1000, Invitrogen) (3). In order to identify proliferating neuronal progenitors, double immunofluorescence was performed with HuC/D and an antibody against the proliferating cell nuclear antigen (1/1000, DAKO). Astrocytes were quantified using an antibody against glial fibrillary acidic protein (GFAP, 1/6000, Dako), which also marks adult neural stem cells. Although all GFAP-positive cells are astrocytes, only some of them are the adult neural stem cells. Microglial cells were labelled using the Iba1 antibody (1/1500, Wako, Japan). Counterstaining was performed using either haematoxylin (DAB staining) or Hoescht33258 (immunofluorescence). The endogenous lipofuscin autofluorescence was removed by a 10-minute incubation of the sections in a 0.5% Sudan Black B (SIGMA, UK) in 70% methanol solution, as previously described (4). Adjacent sections were incubated in the absence of the primary or secondary antibodies in order to determine non-specific antibody binding, and they were devoid of immunoreactivity (Figure S2B and S2C).

AT8 immunohistochemistry




Figure S1. Examples for semi-quantitative scoring using immunohistochemistry for the AT8 antibody against hyperphosphorylated tau in hippocampal sections. **(A)** Low (arrow: neurofibrillary tangle; arrowhead: neuropil thread), **(B)** moderate and **(C)** high immunopositivity in the CA1 region of the human hippocampus. A denotes sparse pathology. Scale bar: 500 μm, valid for all photomicrographs. DG, dentate gyrus.




Figure S2. (A) DAB immunohistochemistry for nestin and doublecortin (DCX), arrows, in the hippocampal dentate gyrus (DG) under X63 magnification. (B) Fluorescence immunohistochemistry in the hippocampal DG after omission of the primary antibodies HuC/D and proliferating cell nuclear antibody (PCNA) and in the presence of fluorescence-conjugated donkey anti-mouse secondary antibodies Alexa[®]Fluor488 and AlexaFluor[®]568 antibodies under 20X magnification. (C) DAB immunohistochemistry after omission of the primary antibodies nestin and doublecortin in the presence of HRP-conjugated anti-rabbit and anti-goat secondary antibodies. h, hilus; LV, lateral ventricle; Ep, endepymal cell layer; SVZ, subventricular zone.

Table S1. Neurogenic and glia markers at the anterior and temporal horn of the subventricular zone

			Stage 0-2 (<i>n</i> = 12)			Stage 3-4 (<i>n</i> = 11)			Stage 5-6 (<i>n</i> = 5)		
			Cells/mm (raw data)	Adjusted Cells/mm	IRR	Cells/mm (raw data)	Adjusted Cells/mm	IRR	Cells/mm (raw data)	Adjusted Cells/mm	IRR
Iba1	SVZ	BG	8.28	7.5 (2.3)	Ref.	8.82	9.6 (3.3)	1.3 (0.5-3.4)	3.64	5.0 (2.9)	0.7 (0.2-2.4)
Iba1	EP	BG	1.83	2.6 (1.3)	Ref.	2.24	1.6 (0.7)	0.6 (0.1-2.4)	00.30	0.5 (0.6)	0.2 (0.02-2.3)
GFAP	SVZ	BG	4.58	5.6 (2.9)	Ref.	4.27	4.4 (2.1)	0.8 (0.2-2.7)	5.28	5.1 (3.6)	0.9 (0.2-5.0)
GFAP	EP	BG	23.3	38.8 (20.1)	Ref.	5.73	4.5 (2.6)	0.1 (0.02-0.8)	24.17	20.6 (16.5)	0.5 (0.05-5.3)
Nestin	SVZ	BG	22.12	20.9 (7.0)	Ref.	12.59	14.3 (5.9)	0.7 (0.2-2.1)	13.94	11.9 (5.7)	0.6 (0.2-1.7)
Nestin	EP	BG	93.77	86.3 (26.3)	Ref.	51.29	49.1 (15.8)	0.6 (0.2-1.5)	95.46	120.6 (54.4)	1.4 (0.5-3.8)
Iba1	SVZ		11.63	11.4 (2.2)	Ref.	21.49	21.4 (3.9)	1.9 (1.1-3.3)	16.88	19.3 (5.3)	1.7 (0.9-0.6)
Iba1	EP		1.83	2.1 (0.6)	Ref.	4.37	4.0 (0.9)	1.9 (0.9-4.1)	2.19	2.1 (0.9)	1.0 (0.4-2.7)
GFAP	SVZ		15.12	14.8 (2.7)	Ref.	19.10	19.7 (3.8)	1.3 (0.8-2.3)	17.5	17.3 (4.6)	1.2 (0.6-2.2)
GFAP	EP		10	11.3 (3.4)	Ref.	11.67	10.4 (3.1)	0.9 (0.3-2.2)	12.19	14.8 (7.0)	1.3 (0.5-3.8)
Nestin	SVZ		15.93	16.9 (2.8)	Ref.	20.67	22.1 (3.6)	1.3 (0.8-2.1)	17.81	14.2 (3.3)	0.8 (0.5-1.5)
Nestin	EP		13.58	14.6 (4.5)	Ref.	9.2	9.3 (3.1)	0.6 (0.3-1.6)	13.13	11.5 (5.0)	0.8 (0.3-2.2)

Cells/mm and incident rate ratios (IRR) (compared to the group with Braak stage 0-2 as a reference) are estimated by negative binomial regression. Adjusted cells per mm are standardized to the sample age and gender and numbers in parentheses represent standard error. None of the differences across groups are statistically significant. GFAP, glial fibrillary acidic protein; SVZ BG, subventricular zone at the level of basal ganglia (anterior horn); EP BG, ependymal cell layer at the level of basal ganglia, adjacent to SVZ BG.

Supplemental References

1. Lendahl U, Zimmerman LB, McKay RD (1990): CNS stem cells express a new class of intermediate filament protein. *Cell* 60: 585-595.
2. Brown JP, Couillard-Despres S, Cooper-Kuhn CM, Winkler J, Aigner L, Kuhn HG (2003): Transient expression of doublecortin during adult neurogenesis. *J Comp Neurol* 467: 1-10.
3. Grandel H, Kaslin J, Ganz J, Wenzel I, Brand M (2006): Neural stem cells and neurogenesis in the adult zebrafish brain: origin, proliferation dynamics, migration and cell fate. *Dev Biol* 295(1): 263-277.
4. Schnell SA, Staines WA, Wessendorf MW (1999): Reduction of lipofuscin-like autofluorescence in fluorescently labeled tissue. *J Histochem Cytochem* 47: 719–730.