King's Research Portal

DOI:

[10.1136/bmj.h5903](https://doi.org/10.1136/bmj.h5903)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Møller, H. (2015). Author's reply to Keatinge and Wong and Cruickshank. *BMJ (Clinical research ed.)*, 351, [h5903]. <https://doi.org/10.1136/bmj.h5903>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

LETTERS

URGENT CANCER REFERRAL PATHWAY

Author's reply to Keatinge and Wong and Cruickshank

Henrik Møller *professor of cancer epidemiology*

Cancer Epidemiology and Population Health, King's College London, London SE1 9RT, UK

We appreciate the interest and debate that our article has generated in rapid responses and conferences and workshops in the past year.¹⁻³ We too thought that the press headlines (mainly arising from a single press agency) were sensationalist and unhelpful.

The concern about lead time contributing to the observed association between urgent referral and patient survival is valid and is covered in the paper's discussion. Lead time is unlikely to be the single cause of the survival difference, and there is evidence that the use of urgent referral is associated with a favourable stage distribution.⁴ This implies that GPs' awareness of cancer and their propensity to use urgent referral is important; this is also illustrated by our finding (web appendix) that cancer patients from practices with many old patients have better age adjusted survival than those from practices with younger patients.

We agree that the results of the overall analysis of all types of cancer do not necessarily apply to each of the less common types of cancer.

Prostate cancer kills more than 10 000 UK men each year and the importance of referral and early diagnosis should not be dismissed.

Our conclusion that "General practices that consistently have a low propensity to use urgent referrals could consider increasing their use of this pathway, thereby plausibly increasing the survival of their patients with cancer" was not meant to advocate excessive use of urgent referral. The statement is consistent with the recently updated NICE guidance.⁵ A single research article cannot be authoritative regarding the use of urgent referral—GPs should look to the NICE guidance for such advice.

Competing interests: see www.bmj.com/content/351/bmj.h5102.

- 1 Møller H, Gildea C, Meechan D, et al. Use of the English urgent referral pathway for suspected cancer and mortality in patients with cancer: cohort study. *BMJ* 2015;351:h5102. (13 October.)
- 2 Keatinge RM. Article on urgent cancer referral pathway did not take lead time bias into account. *BMJ* 2015;351:h5880.
- 3 Wong W, Cruickshank H. Urgent cancer referral pathway: the difficulty in diagnosing head and neck cancers. *BMJ* 2015;351:h5881.
- 4 Maclean R, Jeffreys M, Ives A, et al. Primary care characteristics and stage of cancer at diagnosis using data from the national cancer registration service, quality outcomes framework and general practice information. *BMC Cancer* 2015;15:500.
- 5 National Institute for Health and Care Excellence. Suspected cancer: recognition and referral. (NICE guidance 12.) 2015. www.nice.org.uk/guidance/NG12.

Cite this as: *BMJ* 2015;351:h5903

© BMJ Publishing Group Ltd 2015