King's Research Portal

DOI:

[10.1016/j.chemphyslip.2016.02.004](https://doi.org/10.1016/j.chemphyslip.2016.02.004)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Pal, S., Chakraborty, H., Bandari, S., Yahioğlu, G., Suhling, K., & Chattopadhyay, A. (2016). Molecular Rheology of Neuronal Membranes Explored using A Molecular Rotor: Implications for Receptor Function. *Chemistry and Physics of Lipids*. <https://doi.org/10.1016/j.chemphyslip.2016.02.004>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Title: Molecular Rheology of Neuronal Membranes Explored using A Molecular Rotor: Implications for Receptor Function

Author: Sreetama Pal Hirak Chakraborty Suman Bandari
Gokhan Yahioğlu Klaus Suhling Amitabha Chattopadhyay

PII: S0009-3084(16)30016-0
DOI: <http://dx.doi.org/doi:10.1016/j.chemphyslip.2016.02.004>
Reference: CPL 4451

To appear in: *Chemistry and Physics of Lipids*

Received date: 14-12-2015
Revised date: 23-1-2016
Accepted date: 10-2-2016

Please cite this article as: Pal, Sreetama, Chakraborty, Hirak, Bandari, Suman, Yahioğlu, Gokhan, Suhling, Klaus, Chattopadhyay, Amitabha, Molecular Rheology of Neuronal Membranes Explored using A Molecular Rotor: Implications for Receptor Function. *Chemistry and Physics of Lipids* <http://dx.doi.org/10.1016/j.chemphyslip.2016.02.004>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Short Communication

Molecular Rheology of Neuronal Membranes Explored using A Molecular Rotor: Implications for Receptor Function

Sreetama Pal^{‡,¶}, Hirak Chakraborty^{†,#}, Suman Bandari[†], Gokhan Yahiloglu[§], Klaus Suhling[§]
and Amitabha Chattopadhyay^{†,¶,*}

[†]CSIR-Centre for Cellular and Molecular Biology, Uppal Road, Hyderabad 500 007, India;

[‡]CSIR-Indian Institute of Chemical Technology, Uppal Road, Hyderabad 500 007, India;

[#]School of Chemistry, Sambalpur University, Burla, Odisha 768 019, India;

[¶]Academy of Scientific and Innovative Research, New Delhi, India; [§]Antikor Biopharma Ltd., Stevenage Bioscience Catalyst, Gunnels Wood Road, Stevenage SG1 2FX, Herts, UK;

[§]Department of Physics, King's College London, Strand, London WC2R 2LS, UK

*To whom correspondence should be addressed. Tel.: +91-40-2719-2578;

Fax: +91-40-2716-0311; E-mail: amit@ccmb.res.in

Graphical abstract

Highlights

- Fluorescent molecular rotors are characterized by an excited state process dependent on intramolecular rotational motion
- BODIPY-C₁₂ allows measurement of membrane viscosity by its characteristic viscosity-sensitive fluorescence depolarization
- Specific agonist binding by the serotonin_{1A} receptor exhibits close correlation with hippocampal membrane viscosity
- The role of membrane fluidity in GPCR function has important physiological implications

Abstract

The role of membrane cholesterol as a crucial regulator in the structure and function of membrane proteins and receptors is well documented. However, there is a lack of consensus on the mechanism for such regulation. We have previously shown that the function of an important neuronal receptor, the serotonin_{1A} receptor, is modulated by cholesterol in hippocampal membranes. With an overall objective of addressing the role of

membrane physical properties in receptor function, we measured the viscosity of hippocampal membranes of varying cholesterol content using a *meso*-substituted fluorophore (BODIPY-C₁₂) based on the BODIPY probe. BODIPY-C₁₂ acts as a fluorescent molecular rotor and allows measurement of hippocampal membrane viscosity through its characteristic viscosity-sensitive fluorescence depolarization. A striking feature of our results is that specific agonist binding by the serotonin_{1A} receptor exhibits close correlation with hippocampal membrane viscosity, implying the importance of global membrane properties in receptor function. We envision that our results are important in understanding GPCR regulation by the membrane environment, and is relevant in the context of diseases in which GPCR signaling plays a major role and are characterized by altered membrane fluidity.

Abbreviations: 2-AS, 2-(9-anthroyloxy)stearic acid; 12-AS, 12-(9-anthroyloxy)stearic acid; BODIPY, 4,4-difluoro-4-bora-3a,4a-diaza-*s*-indacene; GPCR, G protein-coupled receptor; 8-OH-DPAT, 8-hydroxy-2-(di-*N*-propylamino)tetralin; 5-PC, 1-palmitoyl-2-(5-doxy)stearoyl-*sn*-glycero-3-phosphocholine; 12-PC, 1-palmitoyl-2-(12-doxy)stearoyl-*sn*-glycero-3-phosphocholine; DMPC, 1,2-dimyristoyl-*sn*-glycero-3-phosphocholine; DOPC, 1,2-dioleoyl-*sn*-glycero-3-phosphocholine; M β CD, methyl- β -cyclodextrin; Tempo-PC, 1,2-dioleoyl-*sn*-glycero-3-phosphotempocholeline

Keywords: Molecular rotor; hippocampal membrane; viscosity; cholesterol; neuronal receptor

1. Introduction

Biological membranes are complex, highly organized, two-dimensional, supramolecular assemblies of a diverse variety of lipids and proteins. The function of membranes is to allow cellular compartmentalization, and impart an identity to individual cells and organelles, besides providing an appropriate environment for proper functioning of membrane proteins. Interestingly, cellular membranes in the nervous system are characterized by very high concentration and remarkable diversity of lipids, and these are correlated with increased complexity in the function of the nervous system (Sastry, 1985; Wenk, 2005). In this context, cholesterol represents an important lipid since brain cholesterol has been implicated in a number of neurological disorders (Chattopadhyay and Paila, 2007; Martín et al., 2014), some of which share a common etiology of defective cholesterol

metabolism in the brain (Porter and Herman, 2011). More importantly, the function of neuronal receptors depends on cholesterol (Pucadyil and Chattopadhyay, 2006; Allen et al., 2007; Paila and Chattopadhyay, 2010; Jafurulla and Chattopadhyay, 2013), which affects neurotransmission, resulting in mood and anxiety disorders (Papakostas et al., 2004). In spite of these important functional correlates, the organization and dynamics of neuronal membranes as a consequence of alterations in membrane cholesterol is only beginning to be understood.

We have earlier established native hippocampal membranes as a convenient natural source for exploring the interaction of neuronal receptors such as the serotonin_{1A} receptor, with membrane lipids of neuronal origin (Pucadyil and Chattopadhyay, 2004). In this overall context, we have demonstrated the requirement of membrane cholesterol in modulating the function of the serotonin_{1A} receptor (Pucadyil and Chattopadhyay, 2004, 2006; Paila et al., 2008; Paila and Chattopadhyay, 2010; Shrivastava et al., 2010; Jafurulla and Chattopadhyay, 2013). A continuing effort in our laboratory has been to explore how to correlate these cholesterol-dependent functional changes of the serotonin_{1A} receptor with alterations in membrane organization and dynamics (Mukherjee et al., 2007; Saxena et al., 2008; Singh et al., 2012).

In this work, we have monitored the change in viscosity associated with cholesterol depletion in hippocampal membranes using viscosity-dependent fluorescence depolarization of BODIPY-C₁₂ (Levitt et al., 2009), a *meso*-substituted molecular rotor based on the BODIPY (4,4-difluoro-4-bora-3a,4a-diaza-*s*-indacene) structure (see Fig. 1a). BODIPY is a popular fluorescent probe and is characterized by high extinction coefficient, quantum yield and photostability (Johnson et al., 1991). Fluorescent molecular rotors are interesting reporter molecules that are generally characterized by an excited state process that is dependent on an intramolecular rotational motion (Haidekker et al., 2010; Levitt et al., 2011; Kuimova, 2012). A fluorescent molecular rotor typically consists of an electron donor moiety covalently linked to an electron acceptor unit *via* a spacer that conjugates the donor and the acceptor units, and facilitates electron movement between the two. The donor and acceptor groups assume a planar or near-planar conformation (with respect to each other) in the ground state. Electronic excitation triggers an intramolecular charge transfer from the donor to the acceptor *via* a twisting of the bond connecting them. This leads to a non-planar excited state conformation termed as twisted internal charge transfer (TICT) state characterized by a lower energy. The radiative relaxation from the TICT state is associated with a red-shifted fluorescence emission. The relaxation of these rotors on photoexcitation is

therefore coupled to rotations of the donor and acceptor units relative to each other. The rotation of the donor and acceptor groups in the excited state is extremely sensitive to the viscosity of the environment. This property makes molecular rotors superior viscosity probes.

A major application of molecular rotors is to measure microviscosity in biological systems since the intramolecular rotation depends on viscosity of the immediate environment in which the molecular rotor is localized (Haidekker and Theodorakis, 2007, 2010; Kuimova et al., 2008; Levitt et al., 2009; Wu et al., 2013). Our results using BODIPY-C₁₂ show that the viscosity of hippocampal membranes exhibits reduction under conditions of cholesterol depletion. More importantly, receptor activity (measured as specific binding of radioligand) exhibits a close correlation with hippocampal membrane viscosity, thereby implying the importance of global membrane properties in receptor activity.

2. Materials and methods

2.1. Materials

1,2-Dioleoyl-*sn*-glycero-3-phosphocholine (DOPC), 1,2-dioleoyl-*sn*-glycero-3-phosphotempocholine (Tempo-PC), 1-palmitoyl-2-(5-doxyl)stearoyl-*sn*-glycero-3-phosphocholine (5-PC), and 1-palmitoyl-2-(12-doxyl)stearoyl-*sn*-glycero-3-phosphocholine (12-PC) were obtained from Avanti Polar Lipids (Alabaster, AL). 1,2-Dimyristoyl-*sn*-glycero-3-phosphocholine (DMPC), methyl- β -cyclodextrin (M β CD), EDTA, EGTA, MgCl₂, MnCl₂, iodoacetamide, phenylmethylsulfonyl fluoride (PMSF), sucrose, Na₂HPO₄, sodium azide, Tris, and sodium acetate were obtained from Sigma Chemical Co. (St. Louis, MO). [³H]8-OH-DPAT (sp. activity 134.1Ci/mmol) was purchased from MP Biomedicals (Santa Ana, CA). Bicinchoninic acid (BCA) reagent for protein estimation was from Pierce (Rockford, IL). *n*-AS probes (2- and 12-(9-anthroyloxy)stearic acid) and Amplex Red cholesterol assay kit were from Molecular Probes/Invitrogen (Eugene, OR). BODIPY-C₁₂ was synthesized as described previously (Levitt et al., 2009). DOPC was checked for purity by thin layer chromatography on silica gel precoated plates obtained from Merck (Darmstadt, Germany) in chloroform/methanol/water (65:35:5, v/v/v) and was found to give only one spot with a phosphate-sensitive spray and upon subsequent charring (Baron and Coburn, 1984). Concentrations of stock solutions of *n*-AS probes and BODIPY-C₁₂ in methanol were estimated using the molar extinction coefficients (ϵ_m) of 8000 and 82100 M⁻¹cm⁻¹ at 361 nm (Haugland, 1996) and 492 nm (López Arbeloa et al., 1999), respectively. Solvents used were of spectroscopic grade. Water was purified through a Millipore (Bedford, MA) Milli Q

system and used throughout. Fresh bovine brains were obtained from a local slaughterhouse within 10 min of death, and the hippocampal region was carefully dissected out. The hippocampi were immediately flash frozen in liquid nitrogen and stored at -80 °C till further use.

2.2. *Methods*

2.2.1. *Estimation of phospholipids*

The concentration of DOPC was determined by phosphate assay subsequent to total digestion by perchloric acid (McClare, 1971). DMPC was used as an internal standard to assess lipid digestion. Samples without perchloric acid digestion produced negligible readings. The phosphate content of native and cholesterol-depleted hippocampal membranes was estimated by phosphate assay.

2.2.2. *Depth measurements using the parallax method*

The actual spin (nitroxide) contents of the spin-labeled phospholipids (Tempo-, 5- and 12-PC) were assayed using fluorescence quenching of anthroyloxy-labeled fatty acids (2- and 12-AS) as described previously (Abrams and London, 1993). For depth measurements, unilamellar vesicles (ULV) of DOPC containing 10% (mol/mol) spin-labeled phospholipids (Tempo-, 5- or 12-PC) labeled with 1% (mol/mol) of BODIPY-C₁₂ were prepared by the ethanol injection method (Kremer et al., 1977). This method produces predominantly unilamellar vesicles of ~100 nm diameter. For this, 160 nmol of lipid (90% DOPC/10% Tempo- or 5- or 12-PC, mol/mol) and 1.6 nmol of BODIPY-C₁₂ was co-dried under a steady stream of nitrogen with gentle warming (~35 °C), followed by further drying under a high vacuum for at least 3 h. The dried lipids were then dissolved in ethanol to give a final concentration of 40 mM. The ethanolic lipid solution was then injected into 1 ml of 10 mM sodium acetate, 150 mM sodium chloride, pH 5 buffer while vortexing to give a final concentration of 0.16 mM lipid in the buffer. Duplicate samples were prepared in each case except samples lacking the quencher for which triplicates were prepared. Background samples lacking BODIPY-C₁₂ were prepared in all experiments, and their fluorescence intensity was subtracted from the respective sample fluorescence intensity in each case. Samples were kept in the dark for 12 h before fluorescence measurements.

2.2.3. *Steady state fluorescence measurements*

Steady state fluorescence measurements was carried out at room temperature (~25 °C) with a Hitachi F-7000 spectrofluorometer (Tokyo, Japan) using 1 cm path length quartz cuvettes. Fluorescence anisotropy measurements were performed at room temperature (~25 °C) with a Hitachi F-7000 spectrofluorometer (Tokyo, Japan). Fluorescence anisotropy values were calculated using the following equation (Lakowicz, 2006):

$$r = \frac{I_{VV} - GI_{VH}}{I_{VV} + 2GI_{VH}} \quad (1)$$

where I_{VV} and I_{VH} are the measured fluorescence intensities (after appropriate background subtraction) with the excitation polarizer vertically oriented and emission polarizer vertically and horizontally oriented, respectively. G is the grating correction factor and is the ratio of the efficiencies of the detection system for vertically and horizontally polarized light, and is equal to I_{HV}/I_{HH} . The excitation wavelength was set at 490 nm, and emission was monitored at 510 nm. Excitation and emission slits with slit widths of 3 nm were used for all measurements. The optical density of the samples measured at 490 nm was less than 0.15. Data shown in Fig. 2b represent means $\pm$ SE of at least three independent measurements.

2.2.4. *Generation of calibration curve for fluorescence anisotropy with viscosity*

The calibration curve for fluorescence anisotropy with viscosity was generated by measuring fluorescence anisotropy of BODIPY- C_{12} (~2 μ M) in glycerol-methanol mixtures of varying ratio and plotting anisotropy as a function of viscosity. The viscosity values of glycerol-methanol mixtures were taken from previous literature (Levitt et al., 2009, 2011). The following hyperbolic equation was fitted to the data points:

$$r = a - b/(1 + c\eta)^{1/d} \quad (2)$$

where r is fluorescence anisotropy and η is viscosity, while a , b , c and d are fitting parameters.

2.2.5. *Hippocampal membrane preparation and cholesterol depletion*

Native hippocampal membranes were prepared from frozen hippocampal tissue, as described previously (Jafurulla et al., 2014). The final pellet (native membranes) was suspended in a minimum volume of buffer C (50 mM Tris, pH 7.4), homogenized using a hand-held Dounce homogenizer, flash frozen in liquid nitrogen and stored at -80 °C. Protein concentration was assayed using the BCA assay (Smith et al., 1985). Native hippocampal

membranes were depleted of cholesterol using M β CD as described previously (Pucadyil and Chattopadhyay, 2004). Briefly, membranes resuspended at a protein concentration of ~2 mg/ml were treated with different concentrations of M β CD in buffer C at room temperature (~25 °C) with constant shaking for 1 h. Membranes were then spun down at 50,000xg for 5 min, washed once with buffer C and resuspended in the same buffer. Cholesterol was estimated using the Amplex Red cholesterol assay kit (Amundson and Zhou, 1999). Radioligand binding assays were carried out as described previously (Jafurulla et al., 2014).

Native and cholesterol-depleted hippocampal membranes containing 50 nmol of total phospholipids were suspended in 1.5 ml of buffer C and used for fluorescence anisotropy experiments. The amount of probe added was such that the final probe concentration was 0.25 mol% with respect to the total phospholipid content. This ensures that the fluorescence anisotropy is not affected due to probe-probe interaction (Wu et al., 2013). Membranes were vortexed for 1 min after addition of the probe and kept in the dark overnight before measurements. Background samples were prepared the same way except that the probe was omitted. The final probe concentration was 0.08 μ M in all cases and the methanol content was always low (0.01%, v/v). Control experiments showed that at this concentration of methanol, the ligand binding properties of the receptor are not altered (Pucadyil and Chattopadhyay, 2004). Viscosity values for native and cholesterol-depleted hippocampal membranes were obtained by extrapolation from the calibration plot (Fig. 2b).

2.2.6. Statistical analysis

Significance levels were estimated using Student's two-tailed unpaired *t*-test using Graphpad Prism version 4.0 (San Diego, CA). The correlation between agonist binding activity of the serotonin_{1A} receptor with membrane viscosity was analyzed using the same software with 95% confidence interval. Plots were generated using Microcal Origin version 6.0 (OriginLab, Northampton, MA).

3. Results and discussion

Knowledge of the penetration depth of the fluorescent moiety in a membrane probe is important because it allows to correlate the fluorescence parameters of the probe with a defined location in the membrane (Halder et al., 2012; Saxena et al., 2014). The penetration depth of the BODIPY fluorophore of BODIPY-C₁₂ in DOPC membranes was determined by the parallax method (Chattopadhyay and London, 1987) using the equation:

$$z_{cF} = L_{c1} + \{[(-1/\pi C) \ln (F_1/F_2) - L_{21}^2]/2 L_{21}\} \quad (3)$$

where z_{cF} is the depth of the fluorophore from the center of the bilayer, L_{c1} is the distance of the center of the bilayer from the shallow quencher (Tempo-PC), L_{21} is the difference in depth between the two quenchers (*i.e.*, the vertical distance between the shallow and deep quenchers), and C is the two-dimensional quencher concentration in the plane of the membrane (molecules/Å²). Here, F_1/F_2 is the ratio of F_1/F_0 and F_2/F_0 , in which F_1 and F_2 are the fluorescence intensities in the presence of the shallow quencher (Tempo-PC) and the deep quencher (5-PC), respectively, both at the same quencher concentration (C); F_0 is the fluorescence intensity in the absence of any quencher. All the bilayer parameters used were the same as described previously (Chattopadhyay and London, 1987). Analysis of quenching data using Eq. 3 shows that the depth of penetration of the BODIPY group in BODIPY-C₁₂ is ~20 Å from the center of the bilayer (see Table 1 and Fig. 1b). These results suggest that the BODIPY moiety of the probe is localized at the interfacial region of the membrane, a region characterized by unique dynamics and dielectric properties (Haldar et al., 2011).

We have previously shown that cholesterol content in hippocampal membranes can be efficiently modulated using MβCD as it forms a selective inclusion complex with cholesterol by including it in the central nonpolar cavity (Pucadyil and Chattopadhyay, 2004). Fig. 2a shows that cholesterol content in hippocampal membranes gets considerably reduced upon treatment with increasing concentrations of MβCD. The extent of cholesterol depletion was highest when 40 mM MβCD was used, which resulted in cholesterol content being reduced to ~10% of the control. The membrane phospholipid level did not display significant change under these conditions (see inset in Fig. 2a).

We utilized the viscosity-sensitive fluorescence depolarization of BODIPY-C₁₂, a fluorescent molecular rotor, for estimating viscosity of native and cholesterol-depleted hippocampal membranes. For this purpose, we generated a calibration curve (see Fig. 2b) by plotting the variation in steady state fluorescence anisotropy of BODIPY-C₁₂ as a function of viscosity using methanol-glycerol mixtures of varying composition (viscosity values of glycerol-methanol mixtures were taken from the literature) (Levitt et al., 2009, 2011). The figure shows that at the higher end of the viscosity range, the fluorescence anisotropy approaches r_0 , *i.e.*, the limiting (fundamental) anisotropy of the BODIPY in the absence of any depolarizing processes such as rotational diffusion (Karolin et al., 1994). An advantage of fluorescence anisotropy measurements is that anisotropy is generally independent of fluorophore concentration.

We measured fluorescence anisotropy of BODIPY-C₁₂ in hippocampal membranes and utilized the calibration plot (shown in Fig. 2b) to obtain an estimate of apparent membrane viscosity. The apparent viscosity of native hippocampal membranes was found to be ~197 cP, which is in the same range as previously reported for natural membranes from the nervous system (Cone, 1972; Poo and Cone, 1974). The apparent viscosity of native and hippocampal membranes with progressive cholesterol depletion (corresponding to increasing concentrations of M β CD) was estimated from the calibration plot (see Fig. 2c). The figure shows that there was an overall decrease in membrane viscosity with cholesterol depletion. For example, viscosity was reduced by ~45% (to ~108 cP), relative to control, when 40 mM M β CD was used.

Membrane cholesterol has previously been shown to be a crucial modulator of the function of membrane receptors (Burger et al., 2000; Pucadyil and Chattopadhyay, 2006; Paila and Chattopadhyay, 2010; Oates and Watts, 2011; Jafurulla and Chattopadhyay, 2013). In this context, we have previously shown that the specific ligand binding activity of the serotonin_{1A} receptor is dependent on hippocampal membrane cholesterol concentration (Pucadyil and Chattopadhyay, 2004). The specific agonist ([³H]8-OH-DPAT) binding was reduced to ~60% of the original binding observed (in control membranes in the absence of M β CD) when 40 mM M β CD was used for cholesterol depletion (see Fig. 3a). In order to gain insight into cholesterol-dependent changes in the function of the serotonin_{1A} receptor in terms of changes in hippocampal membrane viscosity, we plotted the specific agonist binding of the serotonin_{1A} receptor (taken from Fig. 3a) as a function of hippocampal membrane viscosity (from Fig. 2c). This plot is shown in Fig. 3b. Interestingly, a linear regression analysis between specific agonist binding and hippocampal membrane viscosity resulted in a correlation coefficient (*r*) of ~0.84. The 95% confidence intervals contained all the data points, thereby implying that the close correlation among these parameters is significant. Such a close correlation indicates that membrane viscosity plays a crucial role in the function of the serotonin_{1A} receptor in hippocampal membranes. The role of membrane fluidity in GPCR function has important physiological implication (Escribá et al., 2007). Certain diseases in which GPCR signaling is known to play a crucial role, such as hypertension and Alzheimer's disease, have been reported to be characterized by altered membrane fluidity (Roth et al., 1995; Zicha et al., 1999).

Although the role of membrane cholesterol in the function of membrane receptors is well documented (Burger et al., 2000; Pucadyil and Chattopadhyay, 2006; Paila and Chattopadhyay, 2010; Oates and Watts, 2011; Jafurulla and Chattopadhyay, 2013), zeroing in

on the mechanism underlying the effect of membrane cholesterol on the structure and function of membrane proteins and receptors has proved to be challenging (Paila and Chattopadhyay, 2009, 2010; Paila et al., 2009; Lee, 2011). A possible mechanism is centered on change in membrane physical properties (such as viscosity, thickness and curvature) that would give rise to altered protein function (Brown, 2012; daCosta et al., 2013; Soubias et al., 2014). An alternate mechanism could involve a direct (specific) interaction between cholesterol and the given membrane protein that triggers subtle conformational changes in the receptor, giving rise to altered function (). An emerging theme in this area is that these mechanisms need not be mutually exclusive, and the possibility of such cholesterol-dependent membrane protein function could be due to a combination of both.

In case of the serotonin_{1A} receptor, previous work from our laboratory has comprehensively demonstrated the requirement of membrane cholesterol in the function of the receptor (reviewed in Pucadyil and Chattopadhyay, 2006; Paila and Chattopadhyay, 2010; Jafurulla and Chattopadhyay, 2013). An important aspect of our results is that the interaction between cholesterol and the serotonin_{1A} receptor appears to be stringent, since immediate biosynthetic precursors of cholesterol (differing with cholesterol merely in a double bond) were not able to maintain receptor function (Paila et al., 2008; Singh et al., 2009). In addition, we recently showed that the requirement of cholesterol for receptor function is diastereospecific, but not enantiospecific (Jafurulla et al., 2014). This difference in sterol stereospecificity for receptor function appears to be related to membrane dipole potential (Bandari et al., 2014). We have also shown by coarse-grain molecular dynamics simulations that membrane cholesterol has higher occupancy in certain sites on the serotonin_{1A} receptor (Sengupta and Chattopadhyay, 2012). Taken together, these results tend to support a specific mechanism for the requirement of membrane cholesterol for the function of the serotonin_{1A} receptor.

On the other hand, we have shown that the serotonin_{1A} receptor is more compact (Paila et al., 2011) and stable (Saxena and Chattopadhyay, 2012; Patra et al., 2015) in the presence of membrane cholesterol. More importantly, we previously reported that the agonist binding of the serotonin_{1A} receptor in hippocampal membranes was reduced in the presence of capsaicin, which changes the physical properties of membranes by increasing the membrane elasticity (Prasad et al., 2009). In the overall context of these results, our present observation of the close correlation between membrane viscosity and receptor activity assumes relevance. We conclude that lipid effects on membrane protein structure and function are complex and

multifaceted, and adequate caution should be exercised while interpreting experimental results.

Conflict of Interest

The authors declare no conflict of interest

Acknowledgments

This work was supported by the Council of Scientific and Industrial Research, Govt. of India (A.C.). S.P. thanks the University Grants Commission for the award of a Junior Research Fellowship. H.C. acknowledges the Council of Scientific and Industrial Research (Govt. of India) for the award of Senior Research Associateship and the University Grants Commission for UGC-Assistant Professor position. A.C. is an Adjunct Professor of Tata Institute of Fundamental Research (Mumbai), RMIT University (Melbourne, Australia), Indian Institute of Technology (Kanpur), and Indian Institute of Science Education and Research (Mohali). A.C. gratefully acknowledges J.C. Bose Fellowship (Department of Science and Technology, Govt. of India). We thank Md. Jafurulla, Sandeep Shrivastava and Pavan Chodisetti for help with experiments and useful discussion, and members of the Chattopadhyay laboratory for critically reading the manuscript.

References

- Abrams, F.S., London, E., 1993. Extension of the parallax analysis of membrane penetration depth to the polar region of model membranes: use of fluorescence quenching by a spin-label attached to the phospholipid polar headgroup. *Biochemistry* 32, 10826-10831.
- Allen, J.A., Halverson-Tamboli, R.A., Rasenick, M.M., 2007. Lipid raft microdomains and neurotransmitter signalling. *Nat. Rev. Neurosci.* 8, 128-140.
- Amundson, D.M., Zhou, M., 1999. Fluorometric method for the enzymatic determination of cholesterol. *J. Biochem. Biophys. Methods* 38, 43-52.
- Bandari, S., Chakraborty, H., Covey, D.F., Chattopadhyay, A., 2014. Membrane dipole potential is sensitive to cholesterol stereospecificity: implications for receptor function. *Chem. Phys. Lipids* 184, 25-29.
- Baron, C.B., Coburn, R.F., 1984. Comparison of two copper reagents for detection of saturated and unsaturated neutral lipids by charring densitometry. *J. Liq. Chromatogr.* 7, 2793-2801.
- Brown, M.F., 2012. Curvature forces in membrane lipid-protein interactions. *Biochemistry*, 51, 9782-9795.
- Burger, K., Gimpl, G., Fahrenholz, F., 2000. Regulation of receptor function by cholesterol. *Cell. Mol. Life Sci.* 57, 1577-1592.
- Chattopadhyay, A., London, E., 1987. Parallax method for direct measurement of membrane penetration depth utilizing fluorescence quenching by spin-labeled phospholipids. *Biochemistry* 26, 39-45.
- Chattopadhyay, A., Paila, Y.D., 2007. Lipid-protein interactions, regulation and dysfunction of brain cholesterol. *Biochem. Biophys. Res. Commun.* 354, 627-633.
- Cone, R.A., 1972. Rotational diffusion of rhodopsin in the visual receptor membrane. *Nat. New Biol.* 236, 39-43.
- daCosta, C.J.B., Dey, L., Therien, J.P.D., Baenziger, J.E., 2013. A distinct mechanism for activating uncoupled nicotinic acetylcholine receptors. *Nat. Chem. Biol.*, 9, 701-707.
- Escribá, P.V., Wedegaertner, P.B., Goñi, F.M., Vögler, O., 2007. Lipid-protein interactions in GPCR-associated signaling. *Biochim. Biophys. Acta* 1768, 836-852.
- Haidekker, M.A., Nipper, M., Mustafic, A., Lichlyter, D., Dakanali, M., Theodorakis, E. A., 2010. Dyes with segmental mobility: molecular rotors. In: Demchenko, A.P. (Ed.), *Advanced Fluorescence Reporters in Chemistry and Biology I: Fundamentals and Molecular Design*; Vol. 8, Springer-Verlag: Heidelberg, pp 267-308.

- Haidekker, M.A., Theodorakis, E.A., 2007. Molecular rotors-fluorescent biosensors for viscosity and flow. *Org. Biomol. Chem.* 5, 1669-1678.
- Haidekker, M.A., Theodorakis, E.A., 2010. Environment-sensitive behavior of fluorescent molecular rotors. *J. Biol. Eng.* 4, 11.
- Haldar, S., Chaudhuri, A., Chattopadhyay, A., 2011. Organization and dynamics of membrane probes and proteins utilizing the red edge excitation shift. *J. Phys. Chem. B* 115, 5693-5706.
- Haldar, S., Kanaparthi, R.K., Samanta, A., Chattopadhyay, A., 2012. Differential effect of cholesterol and its biosynthetic precursors on membrane dipole potential. *Biophys. J.* 102, 1561-1569.
- Haugland, R.P., 1996. In: *Handbook of Fluorescent Probes and Research Chemicals*, 6th ed.; Molecular Probes Inc: Eugene, OR.
- Jafurulla, M., Chattopadhyay, A., 2013. Membrane lipids in the function of serotonin and adrenergic receptors. *Curr. Med. Chem.* 20, 47-55.
- Jafurulla, M., Rao, B.D., Sreedevi, S., Ruyschaert, J.-M., Covey, D.F., Chattopadhyay, A., 2014. Stereospecific requirement of cholesterol in the function of the serotonin_{1A} receptor. *Biochim. Biophys. Acta* 1838, 158-163.
- Johnson, I.D., Kang, H.C., Haugland, R.P., 1991. Fluorescent membrane probes incorporating dipyrrometheneboron difluoride fluorophores. *Anal. Biochem.* 198, 228-237.
- Karolin, J., Johansson, L.B.-A., Strandberg, L., Ny, T., 1994. Fluorescence and absorption spectroscopic properties of dipyrrometheneboron difluoride (BODIPY) derivatives in liquids, lipid membranes, and proteins. *J. Am. Chem. Soc.* 116, 7801-7806.
- Kremer, J.M.H., Esker, M.W.J., Pathmamanoharan, C., Wiersema, P.H., 1977. Vesicles of variable diameter prepared by a modified injection method. *Biochemistry* 16, 3932-3935.
- Kuimova, M.K., 2012. Mapping viscosity in cells using molecular rotors. *Phys. Chem. Chem. Phys.* 14, 12671-12686.
- Kuimova, M.K., Yahioglu, G., Levitt, J.A., Suhling, K., 2008. Molecular rotor measures viscosity of live cells via fluorescence lifetime imaging. *J. Am. Chem. Soc.* 130, 6672-6673.
- Lakowicz, J.R., 2006. *Principles of fluorescence spectroscopy*, 3rd ed.; Springer: New York.
- Lee, A.G., 2011. Biological membranes: the importance of molecular detail. *Trends Biochem. Sci.* 36, 493-500.

- Levitt, J.A., Chung, P.-H., Kuimova, M.K., Yahiloglu, G., Wang, Y., Qu, J., Suhling K., 2011. Fluorescence anisotropy of molecular rotors. *ChemPhysChem* 12, 662-672.
- Levitt, J.A., Kuimova, M.K., Yahiloglu, G., Chung, P.-H., Suhling, K., Phillips, D., 2009. Membrane-bound molecular rotors measure viscosity in live cells via fluorescence lifetime imaging. *J. Phys. Chem. C* 113, 11634-11642.
- López Arbeloa, F., López Arbeloa, T., López Arbeloa, I., 1999. Electronic spectroscopy of pyrromethene 546. *J. Photochem. Photobiol. A Chem.* 121, 177-182.
- Martín, M.G., Pfrieger, F., Dotti, C.G., 2014. Cholesterol in brain disease: sometimes determinant and frequently implicated. *EMBO Rep.* 15, 1036-1052.
- McClare, C.W.F., 1971. An accurate and convenient organic phosphorus assay. *Anal. Biochem.* 39, 527-530.
- Mukherjee, S., Kombrabail, M., Krishnamoorthy, G., Chattopadhyay, A., 2007. Dynamics and heterogeneity of bovine hippocampal membranes: role of cholesterol and proteins. *Biochim. Biophys. Acta* 1768, 2130-2144.
- Oates, J., Watts, A., 2011. Uncovering the intimate relationship between lipids, cholesterol and GPCR activation. *Curr. Opin. Struct. Biol.* 21, 802-807.
- Paila, Y.D., Chattopadhyay, A., 2009. The function of G-protein coupled receptors and membrane cholesterol: specific or general interaction? *Glycoconj. J.* 26, 711-720.
- Paila, Y.D., Chattopadhyay, A., 2010. Membrane cholesterol in the function and organization of G-protein coupled receptors. *Subcell. Biochem.* 51, 439-466.
- Paila, Y.D., Murty, M.R.V.S., Vairamani, M., Chattopadhyay, A., 2008. Signaling by the human serotonin_{1A} receptor is impaired in cellular model of Smith-Lemli-Opitz Syndrome. *Biochim. Biophys. Acta* 1778, 1508-1516.
- Paila, Y.D., Tiwari, S., Chattopadhyay, A., 2009. Are specific nonannular cholesterol binding sites present in G-protein coupled receptors? *Biochim. Biophys. Acta* 1788, 295-302.
- Paila, Y.D., Tiwari, S., Sengupta, D., Chattopadhyay, A., 2011. Molecular modeling of the human serotonin_{1A} receptor: role of membrane cholesterol in ligand binding of the receptor. *Mol. Biosyst.* 7, 224-234.
- Papakostas, G.I., Öngür, D., Iosifescu, D.V., Mischoulon, D., Fava, M., 2004. Cholesterol in mood and anxiety disorders: review of the literature and new hypotheses. *Eur. Neuropsychopharmacol.* 14, 135-142.

- Patra, S.M., Chakraborty, S., Shahane, G., Prasanna, X., Sengupta, D., Maiti, P.K., Chattopadhyay, A., 2015. Differential dynamics of the serotonin_{1A} receptor in membrane bilayers of varying cholesterol content revealed by all atom molecular dynamics simulation. *Mol. Membr. Biol.* 32, 127-137.
- Poo, M., Cone, R.A., 1974. Lateral diffusion of rhodopsin in the photoreceptor membrane. *Nature* 247, 438-441.
- Porter, F.D., Herman, G.E., 2011. Malformation syndromes caused by disorders of cholesterol synthesis. *J. Lipid Res.* 52, 6-34.
- Prasad, R., Singh, P., Chattopadhyay, A., 2009. Effect of capsaicin on ligand binding activity of the hippocampal serotonin_{1A} receptor. *Glycoconj. J.* 26, 733-738.
- Pucadyil, T.J., Chattopadhyay, A., 2004. Cholesterol modulates ligand binding and G-protein coupling to serotonin_{1A} receptors from bovine hippocampus. *Biochim. Biophys. Acta* 1663, 188-200.
- Pucadyil, T.J., Chattopadhyay, A., 2006. Role of cholesterol in the function and organization of G-protein coupled receptors. *Prog. Lipid Res.* 45, 295-333.
- Roth, G.S., Joseph, J.A., Mason, R.P., 1995. Membrane alterations as causes of impaired signal transduction in Alzheimer's disease and aging. *Trends Neurosci.* 18, 203-206.
- Sastry, P.S., 1985. Lipids of nervous tissue: composition and metabolism. *Prog. Lipid Res.* 24, 69-176.
- Saxena, R., Chattopadhyay, A., 2012. Membrane cholesterol stabilizes the human serotonin_{1A} receptor. *Biochim. Biophys. Acta* 1818, 2936-2942.
- Saxena, R., Shrivastava S., Chattopadhyay, A., 2008. Exploring the organization and dynamics of hippocampal membranes utilizing pyrene fluorescence. *J. Phys. Chem. B* 112, 12134-12138.
- Saxena, R., Shrivastava, S., Haldar, S., Klymchenko, A.S., Chattopadhyay, A., 2014. Location, dynamics and solvent relaxation of a Nile red-based phase-sensitive fluorescent membrane probe. *Chem. Phys. Lipids* 183, 1-8.
- Sengupta, D., Chattopadhyay, A., 2012. Identification of cholesterol binding sites in the serotonin_{1A} receptor. *J. Phys. Chem. B* 116, 12991-12996.
- Shrivastava, S., Pucadyil, T.J., Paila, Y.D., Ganguly, S., Chattopadhyay, A., 2010. Chronic cholesterol depletion using statin impairs the function and dynamics of human serotonin_{1A} receptors. *Biochemistry* 49, 5426-5435.
- Singh, P., Saxena, R., Paila, Y.D., Jafurulla, M., Chattopadhyay, A., 2009. Differential effects of cholesterol and desmosterol on the ligand binding function of the

- hippocampal serotonin_{1A} receptor: implications in desmosterolosis. *Biochim. Biophys. Acta* 1788, 2169-2173.
- Singh, P., Tarafdar, P.K., Swamy, M.J., Chattopadhyay, A., 2012. Organization and dynamics of hippocampal membranes in a depth-dependent manner: an electron spin resonance study. *J. Phys. Chem. B* 116, 2999-3006.
- Smith, P.K., Krohn, R.I., Hermanson, G.T., Mallia, A.K., Gartner, F.H., Provenzano, M.D., Fujimoto, E.K., Goeke, N.M., Olson, B.J., Klenk, D.C., 1985. Measurement of protein using bicinchoninic acid. *Anal. Biochem.* 150, 76-85.
- Soubias, O., Teague, W.E., Hines, K.G., Gawrisch, K., 2014. The role of membrane curvature elastic stress for function of rhodopsin-like G protein-coupled receptors. *Biochimie* 107, 28-32.
- Wenk, M.R., 2005. The emerging field of lipidomics. *Nat. Rev. Drug Discov.* 4, 594-610.
- Wu, Y., Štefl, M., Olżyńska, A., Hof, M., Yahioğlu, G., Yip, P., Casey, D.R., Ces, O., Humpolíčková, J., Kuimova, M.K., 2013. Molecular rheometry: direct determination of viscosity in L_o and L_d lipid phases via fluorescence lifetime imaging. *Phys. Chem. Chem. Phys.* 15, 14986-14993.
- Zicha, J., Kunes, J., Devynck, M.-A., 1999. Abnormalities of membrane function and lipid metabolism in hypertension: a review. *Am. J. Hypertens.* 12, 315-331.

Figure Legends

Fig. 1. (a) The chemical structure of the fluorescent molecular rotor BODIPY- C_{12} used in this study. The free rotation of the C-C single bond connecting the BODIPY moiety with the benzene ring (marked by a curved arrow) depends on viscosity of the medium and this phenomenon modulates the photophysical properties of the probe. The hydrophobic C_{12} chain ensures that the molecule partitions into the membrane. See text for other details. (b) A schematic representation of one-half of the membrane bilayer showing the localization of the BODIPY fluorophore in phosphatidylcholine membranes. The horizontal line at the bottom indicates the center of the bilayer. See Table 1 for other details.

Fig. 2. (a) Effect of increasing concentration of M β CD on cholesterol content of bovine hippocampal membranes. Values are expressed as percentage of cholesterol content for native hippocampal membranes in the absence of M β CD treatment. Data represent means $\pm$ SE of at least three independent measurements (***) corresponds to significant ($p < 0.001$) difference in cholesterol content of M β CD treated hippocampal membranes relative to native membranes). The inset shows phospholipid content of hippocampal membranes upon cholesterol depletion. (b) Steady state anisotropy of BODIPY- C_{12} in methanol-glycerol mixtures with increasing viscosity. The viscosity values corresponding to various compositions of methanol and glycerol were taken from Levitt et al., 2009, 2011. Data represent means $\pm$ SE of at least three independent measurements. See Experimental Section for other details. (c) Effect of cholesterol depletion on viscosity of hippocampal membranes. Data represent means $\pm$ SE of at least three independent measurements (*, ** and *** correspond to significant ($p < 0.05$, $p < 0.01$, and $p < 0.001$, respectively) difference in viscosity of cholesterol-depleted hippocampal membranes relative to native membranes). See Section 2 for more details.

Fig. 3. (a) Effect of increasing concentration of M β CD on specific [3 H]8-OH-DPAT binding to serotonin $_{1A}$ receptors in hippocampal membranes. Values are expressed as percentage of specific binding for native hippocampal membranes in the absence of M β CD treatment. Data represent means $\pm$ SE of at least three independent measurements (* and *** correspond to significant ($p < 0.05$ and $p < 0.001$, respectively) difference in specific [3 H]8-OH-DPAT binding in M β CD treated hippocampal membranes relative to native membranes). (b) Correlation of agonist binding activity of the serotonin $_{1A}$ receptor with membrane viscosity. Specific [3 H]8-OH-DPAT binding to serotonin $_{1A}$ receptors (values taken from Fig. 3a) and corresponding values of membrane viscosity (from Fig. 2c) are plotted. Linear regression analysis yielded a correlation coefficient (r) ~ 0.84 . The significance of the correlation is apparent from the 95% confidence band (plotted as dashed lines). The close correlation between membrane viscosity and receptor activity is noteworthy. See text for more details.

Figure 1
Pal *et al.*

Figure 2
Pal *et al.*

Figure 3
Pal *et al.*

Table 1. Average membrane penetration depth of BODIPY-C₁₂^a

Spin-labeled PC pair used for quenching analysis	Calculated distance from the bilayer center z _{CF} (Å)
Tempo-PC/5-PC	~ 20

^aDepths were calculated from fluorescence quenching obtained with samples containing 10 mol% of Tempo- and 5-PC and using Eq. 3. Samples were excited at 490 nm and emission was collected at 510 nm. The ratio of BODIPY-C₁₂ to total lipid was 1:100 (mol/mol) and the concentration of BODIPY-C₁₂ was 1.6 μM in all cases. See Experimental Section for more details.