King's Research Portal

DOI:

[10.1093/qjmed/hcv224](https://doi.org/10.1093/qjmed/hcv224)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Webb, J., Gemmell, M., Al-Fakih, K., & Chiribiri, A. (2016). Medical treatment of left ventricular pseudoaneurysms. *QJM*, 109(3), 213-214. <https://doi.org/10.1093/qjmed/hcv224>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Medical treatment of Left Ventricular Pseudoaneurysms

Jessica Webb BM BCh, MA (Cantab), MRCP FHEA¹, Rebecca M Gemmell MBBS, MA (Cantab)², Khaled Al-Fakih ³, Amedeo Chiribiri¹

¹ Division of Imaging Sciences and Biomedical Engineering, King's College London, SE1 7EH, United Kingdom

² Department of Cardiology, Princess Royal University Hospital, BR6 8ND, United Kingdom

³ Department of Cardiology, Lewisham and Greenwich NHS Trust Hospital, SE13 6LH, United Kingdom

Address for correspondence

Dr Jessica Webb Division of Imaging Sciences and Biomedical Engineering, 4th floor Lambeth Wing, King's College London, SE1 7EH, United Kingdom

Tel: +44 7970440562

Email: Jessica.webb@kcl.ac.uk

Word count: 376

Clinical picture

A 59-year-old male experienced severe central chest pain for one hour with no associated breathlessness. His medical history included rheumatoid arthritis for which he took methotrexate. His 12 lead electrocardiogram confirmed sinus rhythm with a heart rate of 68 bpm, Q waves in the inferior leads and persisting ST elevation in leads V3-V6. Blood tests confirmed increased troponin and an echocardiogram confirmed severely impaired LV function. Angiography noted an occluded right coronary artery with plaque in the left anterior descending artery but no targets for revascularisation. As he had no further chest pain he was discharged home with follow up at one month to consider whether he was suitable for an implantable cardioverter defibrillator.

Cardiac Magnetic Resonance Imaging (MRI) after one week confirmed a dilated left ventricle with an ejection fraction of 35%. There was a transmural myocardial infarction with a pseudoaneurysm of the basal inferior and inferolateral wall, contained by the pericardium (Figure 1a). Late gadolinium enhancement imaging demonstrated full thickness infarction in this region (Figure 1b) and infarction in the apex. This patient was treated medically.

A pseudoaneurysm is a contained rupture of a blood vessel or of the myocardial wall contained by pericardium, thrombus, or adhesions. An aneurysm, in contrast, results from a weakness in the wall with an outer layer than contains all layers of the myocardium. Differentiating these is clinically relevant as pseudoaneurysms have a greater risk of rupture and historically have had surgical treatment. The incidence of left ventricular aneurysms in patients with Q wave myocardial infarction is thought to

be 8-15%¹. The natural history of untreated ventricular pseudoaneurysm in asymptomatic patients is not clearly defined and the evidence is based largely on retrospective single centre case series. The risk of rupture had been thought to be as high as 30-46%² although advances in imaging have increased the detection of 'incidental' pseudoaneurysms in asymptomatic patients, possibly diluting this rupture risk. In a series published in 1998, the ten patients with ventricular pseudoaneurysms who did not undergo surgery, did not rupture over the four year follow up³. A recent publication described a good outcome in a similar patient not treated surgically⁴. Treatment of blood pressure, optimising heart failure medications and consideration of anticoagulation are undoubtedly important in these patients.

Figure Legends

Figure 1a

Cardiac magnetic resonance imaging (MRI) Left ventricular outflow tract confirming pseudoaneurysm in the basal inferolateral segment. A aorta, LA left atrium, LV left ventricle, PA pseudoaneurysm

Figure 1b

Late gadolinium enhancement of the left ventricular outflow tract confirming full thickness infarction in the basal inferolateral segment (white arrow) and in apex. Normal myocardium appears black.

References

1. Glower DG, Lowe EL. Left ventricular aneurysm. In: Cardiac Surgery in the Adult, Edmunds LH (Ed), McGraw-Hill, New York 1997. p.677.
2. Frances C, Romero A, Grady D Left ventricular pseudoaneurysm. J Am Coll Cardiol. 1998; 32:557-561
3. Yeo TC, Malouf JF, Oh JK, Seward JB Clinical profile and outcome in 52 patients with cardiac pseudoaneurysm. Ann intern Med. 1998;128:299-305
4. Hulten EA, Blankstein R Psuroaneurysms of the Heart. Circulation, 2012;125:1920-1925

Abbreviations

MRI Magnetic Resonance Imaging

This is a pre-copyedited, author-produced PDF of an article accepted for publication in QJM: An International Journal of Medicine, following peer review. The version of record Webb J, et al. Medical treatment of left ventricular pseudoaneurysms. QJM Dec 2015, DOI: 10.1093/qjmed/hcv224 is available online at:

<https://qjmed.oxfordjournals.org/content/qjmed/early/2016/01/05/qjmed.hcv224.full.pdf>