King's Research Portal

DOI:

[10.1111/bjd.14615](https://doi.org/10.1111/bjd.14615)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Rashidghamat, E., Hsu, C-K., Nanda, A., Liu, L., Al-Ajmi, H., & McGrath, J. A. (2016). Incontinentia pigmenti in a father and daughter. *British Journal of Dermatology*. <https://doi.org/10.1111/bjd.14615>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Received Date : 10-Nov-2015

Revised Date : 08-Feb-2016

Accepted Date : 08-Mar-2016

Article type : Research Letter

Incontinentia pigmenti in a father and daughter

E. Rashidghamat,¹ C.K. Hsu,^{1,2} A. Nanda,³ L. Liu,⁴ H. Al-Ajmi,³ J.A. McGrath¹

¹St. John's Institute of Dermatology, King's College London, London, UK; ²Department of Dermatology, National Cheng Kung University Hospital, College of Medicine, National Cheng Kung University, Tainan, Taiwan; ³As'ad Al-Hamad Dermatology Center, Al-Sabah Hospital, Kuwait; ⁴Viapath, St Thomas's Hospital, London, UK.

Correspondence: John McGrath, Dermatology Research Laboratories, Floor 9 Tower Wing, Guy's Hospital, Great Maze Pond, London SE1 9RT, United Kingdom. Tel 44-20-71886409; Fax 44-20-71888050; E-mail john.mcgrath@kcl.ac.uk

Incontinentia pigmenti (IP) is a rare multi-system X-linked dominant genetic disorder caused by mutations in *IKBKG*, encoding inhibitor of nuclear factor kappa-B kinase subunit gamma (IKK- γ).

Functionally, the encoded IKK- γ protein participates in nuclear factor kappa-light-chain-enhancer of activated B cell (NF- κ B) signalling to regulate inflammatory and immune responses and prevent

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/bjd.14615

This article is protected by copyright. All rights reserved.

apoptosis. Loss-of-function mutations in *IKBK*G, as occurs in IP, leave mutant cells vulnerable to apoptosis when exposed to tumour necrosis factor alpha.¹ Females with IP who inherit germline mutations in *IKBK*G usually survive because of skewed X-inactivation favouring expression of the wild-type allele, whereas most males with IP die *in utero*. However, possible explanations for rare examples of male IP survival include: Klinefelter syndrome (47XXY), hypomorphic *IKBK*G mutations, and post-zygotic mutations in *IKBK*G leading to somatic mosaicism.² Here, we describe an adult male with IP, with survival being attributed to tissue mosaicism, who then fathered a female child with IP affecting her germline.

A 29-year-old Kuwaiti man was born with severe linear blistering followed by hyperpigmentation and hyperkeratotic plaques in a Blaschkoid distribution on the head, trunk, upper and lower limbs (Figure 1a and b). He had linear facial asymmetry and blindness of the left eye. His dentition was abnormal, necessitating dentures. The rest of his neurological examination was unremarkable. His one-year-old daughter was born at term via normal vaginal delivery. There was no history of consanguinity in the family. Soon after birth she had linear, streaky erythema and vesicles, with subsequent hyperpigmentation and hyperkeratotic plaques; lesions were noted on the upper and lower limbs along lines of Blaschko (Figure 1c). Other systemic abnormalities were excluded on clinical examination and developmental parameters were normal. No neurological abnormalities have been noted during her first 12 months.

Following informed consent, a three-primer PCR protocol was used to screen for the most common recurrent pathogenic mutation in *IKBK*G that underlies ~80% of IP cases worldwide, an intragenic deletion from within intron 3 to within intron 10 that removes all of exons 4-10.¹ Such a PCR protocol is necessary because of a non-functional copy of *IKBK*G running in the opposite

Accepted Article

direction. We used a recently reported method,³ that represents an improvement over other protocols because of its capacity to non-preferentially amplify both wild-type and deletion mutant *IKBKG* alleles (c.f. some other protocols that favour amplification of the wild-type allele because of GC nucleotide mismatch).

This PCR protocol (Figure 2) failed to demonstrate presence of the common deletion in *IKBKG* in the father's genomic DNA extracted from his peripheral blood. However, the deletion was clearly identified in his lesional skin DNA and, to a much lesser extent, in his non-lesional skin DNA. In contrast to the father, the deletion was present in peripheral blood genomic DNA from his daughter. Presence of all deletions was confirmed by Sanger sequencing (data not shown). The clinical phenotype in this family infers that the father must be a tissue mosaic for the IP deletion mutation that has unexpectedly also affected the gonads and hence been transmitted to his daughter as a germline abnormality. Her clinical signs reflect skewed X-inactivation, resulting in survival, as is the case for most affected females.

To our knowledge, there are only three previous reports of father-to-daughter transmission of IP. Two cases were similar to ours, with a history of probable tissue mosaicism in the fathers,^{4,5} although the clinical descriptions predated the discovery of *IKBKG* mutations in IP and thus neither of these cases had supportive molecular pathology. One other report described a clinically unaffected father having two daughters with IP (born to different mothers).⁶ The lack of clinical abnormalities in the father in that case is consistent with gonadal mosaicism, although again the report was published before the discovery of *IKBKG* as the IP gene. Thus our fourth case represents the first example of father to daughter transmission of IP in which a pathogenic mutation in *IKBKG* has been demonstrated.

Acknowledgements

This study received funding/support from the National Institute for Health Research (NIHR) Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London

References

- [1] Smahi A, Courtois G, Vabres P *et al.* Genomic rearrangement in NEMO impairs NF-kappa-B activation and is a cause of incontinentia pigmenti. The International Incontinentia Pigmenti Consortium. *Nature* 2000; **405**: 466-72.
- [2] Fusco F, Paciolla M, Conte M *et al.* Incontinentia pigmenti: report on data from 2000 to 2013. *Orphanet J Rare Dis* 2014; **9**: 93.
- [3] Guevara BEK, Hsu CK, Liu L *et al.* Improved molecular diagnosis of the common recurrent intragenic deletion mutation in *IKBKG* in a Filipino family with incontinentia pigmenti. *Australas J Dermatol* 2015; in press.
- [4] Sommer A, Liu PH. Incontinentia pigmenti in a father and his daughter. *Am J Med Genet* 1984; **17**: 655-9.
- [5] Emery MM, Siegfried EC, Stone MS *et al.* Incontinentia pigmenti: transmission from father to daughter. *J Am Acad Dermatol* 1993; **29**: 368-72.
- [6] Kirchman TT, Levy ML, Lewis RA *et al.* Gonadal mosaicism for incontinentia pigmenti in a healthy male. *J Med Genet* 1995; **32**: 887-90.

Figure legends

Figure 1. Clinical features (a) Blaschkoid distribution of hyperpigmentation with hyperkeratotic plaques, with some pallor and atrophy, affecting the abdomen in the father; (b) Similar Blaschko-linear changes are evident on his right leg; (c) Lower limbs of his daughter showing Blaschko-linear vesicles and early verrucous inflammatory plaques.

Figure 2. A modified PCR amplification method identifies the recurrent intragenic deletion of *IKBKG* in the DNA samples from the father's affected skin and the proband's blood, but not from the father's blood (see ref. 1 for detailed methodology). Only a very faint band was detected in the DNA from the father's unaffected skin. For these specific PCR conditions, the upper 1045-bp band indicates the presence of the deletion, whereas the 733-bp band is wild-type. C = control DNA; MW = molecular weight ladder.