King's Research Portal

DOI:

[10.1016/j.clnu.2016.04.022](https://doi.org/10.1016/j.clnu.2016.04.022)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Weekes, C. E., Baldwin, C., Munk, T., & Beck, A. M. (2016). Are oral nutritional supplements more effective than dietary advice in malnourished care home residents? *CLINICAL NUTRITION*, 35(4), 984.
<https://doi.org/10.1016/j.clnu.2016.04.022>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Are oral nutritional supplements more effective than dietary advice in malnourished care home residents?

C. Elizabeth Weekes, PhD RD, Senior Consultant Dietitian, Christine Baldwin, PhD RD, Lecturer in Nutrition & Dietetics, Tina Munk, PhD RD, Head of Nutritional Unit, Anne Marie Beck, PhD RD, Senior Researcher, Dietician

PII: S0261-5614(16)30068-1

DOI: [10.1016/j.clnu.2016.04.022](https://doi.org/10.1016/j.clnu.2016.04.022)

Reference: YCLNU 2812

To appear in: *Clinical Nutrition*

Received Date: 2 March 2016

Accepted Date: 20 April 2016

Please cite this article as: Weekes CE, Baldwin C, Munk T, Beck AM, Are oral nutritional supplements more effective than dietary advice in malnourished care home residents?, *Clinical Nutrition* (2016), doi: 10.1016/j.clnu.2016.04.022.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Are oral nutritional supplements more effective than dietary advice in malnourished care home residents?

Dear Editor,

With the high prevalence of malnutrition in care homes¹, it is important to determine the most effective strategies for improving nutritional status and outcomes in this population. A recent paper² attempts to address this by comparing the impact of oral nutritional supplements (ONS) with dietary advice (DA) on quality of life. The authors are to be commended for conducting this trial however we feel that some aspects of their paper require clarification.

The study was conducted in 104 care home residents identified as at risk of malnutrition and randomised to receive either ONS or DA. Nutritional status and quality of life were measured at baseline and at six and twelve weeks.

Our first concern relates to the amount of detail provided on the interventions. While acknowledging the often strict word limits for journal articles, authors should provide sufficient information for their study to be replicated and the comparability of the interventions to be assessed. Parsons *et al.* state that the ONS group were 'given access to a range of ONS... so they could take them *ad libitum*' while those in the DA group 'were given a specially designed diet sheet encouraging intake of high energy foods, and drinks and snacks'. Provision of information alone does not constitute DA yet there is no description of how the residents were able to act on the information provided in the diet sheet. While it seems that the dietitian discussed the intervention with both groups (DA and ONS), and care and catering staff were informed of the intervention, it is not clear if the DA group were given access to the high energy foods, snacks and drinks they required in the same way as the ONS were available to the other group.

Our second concern relates to the appropriateness of DA, as described in this study, for this population. DA has been described as a "cognitive intervention"³ and requires the participants to have adequate cognitive function to implement suggestions. The Alzheimer's Society estimates that 80% of people in residential care have a form of dementia or severe memory problems, and only 44% of people with dementia in the UK receive a diagnosis⁴. Parsons *et al.* state that participants "*without obvious dementia*" were selected but this does not rule out the possibility that some of the participants were cognitively impaired and thus may not have had the capacity to act on the advice provided. This might explain the high drop-out rate due to confusion. Furthermore, by selecting only those without cognitive impairment the authors have limited the generalisability of their findings to a minority of the care home population.

35 The title of the study suggests a failure of DA in managing malnourished residents in care
36 homes, whereas we suspect that the findings are more likely to be related to the
37 inappropriateness of the intervention in this population and/or a failure of implementation.

38

39 References

- 40 1. Valentini L, Schindler K, Schlaffer R, Bucher H, Mouhieddine M, Steininger K,
41 Tripamer J, Handschuh M, Christian S, Volkert D, Lochs H, Sieber C.C, Hiesmayr M,
42 The first nutritionDay in nursing homes: participation may improve malnutrition
43 awareness. Clin Nutr 28 (2009) 109-116
44 2. Parsons EL, Stratton RJ, Cawood AL, Smith TR, Elia M. Clin Nutr. Oral nutritional
45 supplements in a randomised trial are more effective than dietary advice at improving
46 quality of life in malnourished care home residents. 2016 pii: S0261-5614(16)00003-
47 0. doi: 10.1016/j.clnu.2016.01.002. [Epub ahead of print]
48 3. Silver HJ (2009) Nestlé Nutr Inst Workshop Ser Clin Perform Program, vol. 12, pp
49 79–93
50 4. The Alzheimer's Society
51 (https://www.alzheimers.org.uk/site/scripts/documents_info.php?documentID=341)