King's Research Portal

DOI:

[10.1016/S2215-0366\(15\)00491-5](https://doi.org/10.1016/S2215-0366(15)00491-5)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Fusar-Poli, P., Schultze-Lutter, F., & Addington, J. (2016). Intensive community outreach for those at ultra high risk of psychosis: Dilution, not solution. *The Lancet Psychiatry*, 3(1). [https://doi.org/10.1016/S2215-0366\(15\)00491-5](https://doi.org/10.1016/S2215-0366(15)00491-5)

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

INTENSIVE COMMUNITY OUTREACH MAY DILUTE TRANSITION TO PSYCHOSIS IN SUBJECTS AT HIGH CLINICAL RISK

Letter

Fusar-Poli P^{1,2, 31}, Schultze-Lutter F⁴, Addington J⁵.

1. King's College London, Institute of Psychiatry, London, United Kingdom;
2. OASIS service, South London and the Maudsley NHS Foundation Trust, London, United Kingdom;
3. Department of Brain and Behavioural Sciences, University of Pavia, Italy;
4. University Hospital of Child and Adolescent Psychiatry and Psychotherapy, University of Bern, Bern, Switzerland;
5. Department of Psychiatry, University of Calgary, Calgary, Alberta, Canada.

Endnote : CAARMS_SIPS

Word count: 497

Refs: 5

Figures/Tables: 0

Financial support: Department of Psychosis Studies, Institute of Psychiatry Psychology and Neuroscience, King's College London and South London and SLaM National Health Service Foundation Trust.

Conflict of Interests: None

¹¹ Corresponding author Dr. Paolo Fusar-Poli, Department of Psychosis Studies, Institute of Psychiatry Psychology and Neuroscience PO63, De Crespigny Park, SE58AF London UK. Phone ++44 (0) 20 7848 0900; e-mail: paolo.fusar-poli@kcl.ac.uk

Dear Editor,

The Liaison and Education in General Practices (LEGs) cluster-randomised controlled trial¹ investigated whether increased liaison between primary and secondary care improves the detection of primary care patients at ultra-high risk (UHR) of developing psychosis or first episode-psychosis (FEP) and concluded that it was both clinically and cost effective. However, the ratio of true positives (UHR or FEP) and false positives (neither UHR nor FEP) referrals from general practices was similarly (approximately 1:1) across all the three treatment conditions (high intensity: 2.2:2.3, low intensity: 1.1:0.9, practice as usual: 0.6:0.8)¹. Thus, there is no evidence of the high-intensity intervention specifically improving diagnostic accuracy of general practitioners in UHR or FEP identification, which is the key outcome of interest. Furthermore, UHR true/false positives were defined as new referrals “who met criteria for high risk [...] according to CAARMS (psychosis true positives) and those who did not fulfill the criteria (false positives)”¹. This is problematic as, commonly, “psychosis true positive” is prognostically defined by the proportion of baseline subjects who actually transition to psychosis (i.e. positive predictive value). Indeed, it is not only CAARMS criteria that determine the actual risk of psychosis-transition but also the underlying prevalence of psychosis-risk in the population referred. In a recent meta-analysis, psychosis risk in populations seeking help at high risk services was on average 0.15 (at 38 months)². This is significantly higher than the 0.001 risk of psychosis within 38 months (incidence of psychosis 0.0317 per 100 person-years 95%CI: 0.025-0.041)³ in the general population. Specifically, there is meta-analytical evidence that intensive community outreach campaigns promoting referrals to high risk services may actually dilute the risk enrichment of populations undergoing risk (e.g. CAARMS) assessment, resulting in low transitions. Non-specific referrals in Perez’ study¹ could lead to such a dilution of risk enrichment and transitions. We have hypothetically tested this hypothesis. We used the pre-test probability of psychosis estimated in primary care services, reported by the UK Health Improvement Network (THIN) epidemiological primary care database⁴. Based on more than 10 million longitudinal patient records from 437 general practices a rate of first diagnosis of

schizophrenia or non-organic psychoses of 31.3 per 100,000 person years is indicated (corresponding to a 0.00099 risk of psychosis over 38 months)⁴. Using our meta-analytical probability modifying plot (positive likelihood ratio=1.82, negative likelihood ratio=0.09²), we estimated the hypothetical positive and negative post-test probabilities (at 38-month follow-up) associated with a CAARMS assessment in populations referred from UK GP's. The post-test probability for developing psychosis over 38 months was less than 0.001 in case of a negative risk assessment (e.g. CAARMS negative) and only 0.002 in case of a positive risk assessment (e.g. CAARMS positive). This suggests a very limited clinical utility of in-depth risk assessments for psychosis in GP's practices. However, these numbers are only hypothetical and might not hold against real-life numbers that we hope the authors will provide by following-up their sample, thus dismissing our concerns by showing a clinically significant real risk of high transitions in their population.

References

1. Perez J, Jin H, Russo D, et al. Clinical effectiveness and cost-effectiveness of tailored intensive liaison between primary and secondary care to identify individuals at risk of a first psychotic illness (the LEGs study): a cluster-randomised controlled trial. *Lancet Psychiatry* 2015.
2. Fusar-Poli P, Cappucciati M, Rutigliano G, et al. At risk or not at risk? Meta-analysis of the prognostic accuracy of psychometric interviews for psychosis prediction. *World Psychiatry* 2015: in press.
3. Kirkbride JB, Errazuriz A, Croudace TJ, et al. Incidence of schizophrenia and other psychoses in England, 1950-2009: a systematic review and meta-analyses. *PloS one* 2012; **7**(3): e31660.
4. Hardoon S, Hayes JF, Blackburn R, et al. Recording of severe mental illness in United Kingdom primary care, 2000-2010. *PloS one* 2013; **8**(12): e82365.