King's Research Portal

DOI:

[10.1016/j.jcpo.2016.05.004](https://doi.org/10.1016/j.jcpo.2016.05.004)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Harding, R., Luyirika, E., & Sleeman, K. E. (2016). Palliative care: when and how, and what are the implications for global cancer policy? *Journal of Cancer Policy*. <https://doi.org/10.1016/j.jcpo.2016.05.004>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Title: Palliative care: when and how, and what are the implications for global cancer policy?

Author: Richard Harding Emmanuel Luyirika Katherine E Sleeman

PII: S2213-5383(16)30021-2
DOI: <http://dx.doi.org/doi:10.1016/j.jcpo.2016.05.004>
Reference: JCPO 68

To appear in:

Received date: 19-4-2016
Accepted date: 30-5-2016

Please cite this article as: Richard Harding, Emmanuel Luyirika, Katherine E Sleeman, Palliative care: when and how, and what are the implications for global cancer policy?, Journal of Cancer Policy <http://dx.doi.org/10.1016/j.jcpo.2016.05.004>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Palliative care: when and how, and what are the implications for global cancer policy?

Richard Harding (corresponding author)

King's College London

Cicely Saunders Institute

Department of Palliative Care, Policy & Rehabilitation

Bessemer Road

London SE5 9PJ

Richard.harding@kcl.ac.uk

+44 (0) 207 848 5518

Emmanuel Luyirika

African Palliative care Association

Kampala, Uganda

Katherine E Sleeman

King's College London

Cicely Saunders Institute

Highlights

- Palliative care is an essential component of health services. It improves outcomes and saves cost. Despite increasing need for palliative care in all regions of the world, provision is woefully inadequate.
- Despite the needs and benefits for palliative care, investment is poor even in high income countries.
- Integration of palliative care, and quality of death, is greatest in high income countries.
- Some low and middle income countries have shown regional leadership by including palliative care in the National health Plans, and by supporting policy to enhance provision in primary, secondary and tertiary sectors.

The concept and practice of palliative care

Palliative care is defined by the World Health Organization (WHO) as "an approach that improves quality of life of patients and their families facing the problems associated with life-threatening illness, through the prevention and relief of suffering by means of early identification and impeccable assessment and treatment of pain and other problems, including physical, psychosocial, and spiritual".¹ The word palliative comes from the Latin *palliare*, meaning *to cloak*. The goal of palliative care is to ease the symptoms and other problems associated with living with an incurable illness, rather than to try to extend the amount of time that the patient has left to live.

In this article the term palliative care will refer to care delivered by health care professionals working in specialist palliative care teams. Specialist palliative care teams are multi-professional, including doctors, nurses, social workers, and other allied health professionals for whom palliative care is their core daily work. Specialist palliative care teams deliver care directly to patients, and indirectly by supporting other professionals to deliver care.²

The term hospice is context dependent. In the UK, hospice most often refers to an institution within which the health care professionals are specialists in palliative care. Hospices in the UK usually have inpatient beds, as well as community teams who see patients in their place of residence. In the United States, hospice care is the term used for a Medicare benefit available to people with a prognosis of six months or less.³ US hospice care is most often provided in the community, with inpatient units being less common.

Since the development of the modern hospice movement by Dame Cicely Saunders in the UK in the 1960's, it has become a global movement (for example the first hospice in Africa opened in Zimbabwe in 1979), and is often recognised as a medical and nursing specialty. Given the advances in the policy, practice and research of palliative care in the UK, in this article we will largely draw on the UK context. However, as an academic collaborative that has focused on palliative care as a global health need, we will also draw on further contexts and lessons learned from around the world. In 2004 the World Health Assembly passed its first resolution on palliative care and so now Governments have a responsibility to train health care professionals in palliative care, deliver appropriate and effective palliative care, and ensure an evidence base.

The benefits of palliative care

There is now a body of evidence from controlled trials that specialist palliative care teams provide improved symptom control, better satisfaction with care, and improved quality of life for patients, and better outcomes for care-givers.⁴⁻⁷ Systematic reviews have shown that receipt of community-based palliative care is associated with double the odds of dying at home,⁸ and half the odds of cancer patients attending Emergency Departments in their last month of life.⁹ In the United States, among people with cancer, hospice care was associated with lower rates of hospitalization, intensive care unit admission, and invasive procedures close to the end-of-life.¹⁰ In Canada, community-based specialist palliative care is associated with reduced risk of acute hospital care use, and a higher rate of out-of-hospital death.¹¹

Paradoxically, although the aim of palliative care is not to lengthen life, there is early but growing evidence that early palliative care may also improve survival among people with cancer.^{12,13} The mechanism of this is unclear but it may be a result of patients receiving less chemotherapy near to the end-of-life, having fewer hospital admissions, or due to improvement in patients' depression.¹⁴ In the UK, trial evidence has shown that a brief palliative care consultant advisory service can reduce symptoms and caregiver burden¹⁵ and costs of care¹⁶. A further UK trial to integrate palliative care into a novel multiprofessional brief service to manage breathlessness found In addition to self-mastery of breathlessness, it also has a positive survival effect¹⁷. In an RCT evaluation from Africa, training and use of a basic assessment tool with a care plan in palliative care delivered by existing clinic nurses alongside treatment improved mental health and psychosocial problems¹⁸.

Palliative care reduces the high costs of end-of-life care

Care costs for people in the last year of life account for a large proportion of total health care expenditure. In the UK, around 20% of hospital bed days are related to the last year of life.¹⁹ In the USA, 25% of total health care expenditure is used in the last year of life.²⁰

Annual direct costs for cancer care are projected to rise in the USA from \$104 billion in 2006 to \$173 billion in 2020, driven by dramatic rises in the costs of therapy and the extent of care.²¹ These are unsustainable increases. Better integration of

palliative care has been suggested as a potential solution to reducing costs without depriving patients of benefit.²¹

Evidence on the cost effectiveness of palliative care has lagged behind evidence on its effectiveness with respect to patient and family outcomes. This is due to the complexity of capturing social care and informal care costs in addition to health care related costs. Emerging evidence suggests that palliative care is likely to be cost-saving overall. In a systematic review, Smith et al examined 46 studies that reported on costs of palliative care.²² They found a wide variety of study types, characteristics, and quality. Most of these studies were from the United States, and they included both cancer and non-cancer conditions. Despite these variations, they found a consistent pattern: palliative care was frequently found to be less costly than comparator groups, and in most cases this difference was statistically significant. This indicates that the expense incurred by having an extra team involved in the patient's care is offset by fewer tests and treatments, and less hospital-based care.

End-of-life communication

It has been said that if palliative care was a drug, patients and policy makers alike would demand it. Unfortunately, societal attitudes towards death and dying, and a reluctance to speak about end-of-life care issues, mean that palliative care does not always seem palatable. The 2013 Neuberger review called for a National conversation about death and dying, to dispel some of the taboos that still surround it.²³ Initiatives such as the Dying Matters coalition in the UK, the Death Café movement, and the US Conversation Project encourage the normalization of conversations about dying. However, many people still have a 'not yet' approach to talking about their own mortality.

This is compounded by health care professionals' reluctance to initiate conversations about end-of-life care preferences with their patients. A US study showed that 69% of patients with lung cancer (and 81% with colorectal cancer) were not aware that the intention of their palliative chemotherapy was not to cure their disease.²⁴ In one study, around half of medical oncologists preferred not to discuss resuscitation, advance directives, or palliative care until there were no more treatment options available.²⁵ Perhaps not surprisingly, cancer patients who have late end-of-life conversations are more likely to receive overly aggressive care.²⁶ In the US, recent provision of reimbursement to physicians who have end-of-life conversations with their patients is likely to lead to such conversations becoming more mainstream.

Good end-of-life communication must be initiated by health care professionals, who in turn require adequate education and training. Remarkably, given its ubiquity, it is only relatively recently that caring for people who are dying has become a mainstream part of medical education. While palliative care is slowly gaining prominence on the medical curriculum, still the average time devoted to palliative care is 20 hours over five years.²⁷

Providing enough palliative care: a lack of investment

While there is now convincing evidence that palliative care can improve patients' quality of life and death, and frequently at lower costs, investment in palliative care services has been slow and often inadequate. In the UK, palliative care developed from the hospice sector, and historically much of its funding has been private or charitable. Currently, the charitable sector spends £950 million on hospice care annually, with state funding of around £460 million.²⁸

There is widespread geographical variation in investment in specialist palliative care services: in 2010/11 the average spend by Primary Care Trusts on specialist palliative care varied from £186 to £6213 per death.²⁸ Inequalities in palliative care provision are widespread, with evidence of disparity according to diagnosis, age and relative affluence in addition to geographical location.²⁹

Improving care for people approaching the end of their lives, and providing care that reflects their individual choices and preferences, has been cited as a priority for the UK Government. The 2015 Choice review calculated that a relatively modest investment - £130 million – could provide a meaningful level of service improvement and greater choice for patients approaching the end-of-life.³⁰ However, despite the government's commitment to respond to the Choice review by the end of 2015, it is yet to publish its position.

Global palliative care provision

Access to palliative care is recognized as a human right claimable within existing global covenants^{31,32}. Palliative care may be even more necessary in low and middle income countries where there is poorer access to health services, later presentation, and fewer curative treatment options.

In 2010, and again in 2015, the Economist Intelligence Unit attempted to rank quality of end-of-life care in 80 countries worldwide according to five indicators: the palliative and healthcare environment, human resources, the affordability of care, the quality of care and the level of community engagement. Not surprisingly, rich nations tend to rank highest in terms of quality of end-of-life care. The report showed that countries with a high quality of end-of-life care shared the following qualities:

- A strong and effectively implemented national palliative care policy framework;
- High levels of public spending on healthcare services;
- Extensive palliative care training resources for general and specialised medical workers;
- Generous subsidies to reduce the financial burden of palliative care on patients;
- Wide availability of opioid analgesics;
- Strong public awareness of palliative care.

The WHO global atlas of palliative care has categorized every country according to the level of palliative care integration and provision³³. While 74 low and middle income countries fall within the worst category (i.e. “no known activity”, e.g. Afghanistan, Bhutan, Senegal, Uzbekistan) it is notable that several African countries fall in the top two categories (“Preliminary integration” and “Advanced integration”, e.g. Kenya, Tanzania, Zambia, Zimbabwe, Uganda and South Africa). Sub-Saharan Africa has achieved much in advancing the policy, training, practice and research of palliative care. African advocacy bodies and services have worked hard to roll out opioid availability in districts^{34,35}, to develop and implement simple outcome measures to guide and evaluate care³⁶, and to undertake research on physical³⁷, social³⁸ spiritual^{39,40} and psychological aspects⁴¹ of palliative care. The WHO foundation measures were designed to assist countries develop a public health approach to cancer palliative care⁴²: policy, education, drug availability and implementation. However, these foundation measures must be underpinned by local research that determines *what* the policy should advocate, *how and what* should be taught, *which* drugs need to be available, and *how best* to implement⁴³. African has been an exemplar in advancing the science of palliative care, generating the original research urgently needed in the emerging field of global health palliative care⁴⁴. The

African example shows how clinical cultures and public understanding can be changed to endure effective, compassionate and appropriate end-of-life care. Such change is urgently needed in many parts of the world, for example research in Cuba found that advanced cancer patients were often unaware of their diagnosis or prognosis, and that this was associated with greater distress⁴⁵.

Drug availability for the management of advanced cancer pain is a key challenge in many regions⁴⁶. Very economically poor terminal cancer patients and their families in Bangladesh describe spending their few resources on futile treatments, yet not being able to access cheap and effective opioids^{47,48}. The patchy coverage of palliative care highlighted in the WHO Global Atlas underlines the challenge of unequal access even within countries. An example of this is India, where the Kerala region has become well known for good availability of community-based palliative care, while other states have very little coverage at all⁴⁹. Similar inequity exists in Africa⁵⁰, and the challenge remains that we must expand global coverage while maintaining quality.

Cancer palliative care policy in Africa

According to the **World Bank Cancer Mortality report of 2012** low- income countries largely in sub-Saharan African contributed a total of 350,000 deaths to cancer mortality of which 70,000 tobacco related cancers and 90,000 deaths were infection related cancers.

Sub-Saharan Africa has a unique cancer picture compared to the rest of the world. It is the only continent where among women cervical cancer is equivalent to breast cancer in prevalence. According to Global Cancer Report 2014 the prevalence of cervical cancer among women in sub-Saharan Africa is 25.2% while breast cancer is 25.5%. Both these are the top cancers for women while prostate and liver cancers are top cancers in men (World Cancer Report 2014).

The dilemma facing most cancer patients are many in Africa is lack of appropriate and accessible services. Most cancer patients in sub Saharan Africa either present late for treatment, or go to facility where appropriate cancer treatment does not exist. There are many cancer patients who never access any form of western medicine services (World Cancer Report 2014).

The situation is changing slowly in terms of cancer policy and cancer control plans in Africa. Zimbabwe and Kenya are model countries that have now adopted model cancer control plans that cover the whole continuum from primary prevention, early detection, diagnosis and treatment, support and rehabilitation as well as palliative care (<http://www.iccp-portal.org/sites/default/files/plans>) and <http://www.iccp-portal.org/cancer-plans>)

Six African countries in namely Swaziland, Mozambique, Malawi, Zimbabwe, Tanzania and Rwanda have national overarching standalone palliative care policies with clear national strategies. Uganda and Botswana have completed national palliative care policy documents which are awaiting approval.

Evidence based policy

As described above, an accumulation of research evidence in recent decades has shown that palliative care is an essential component of cancer services, improves outcomes for patients and their family care givers (who also have high care needs⁵¹), and reduces health care costs. However, further research is necessary to understand when the 'right' time to introduce palliative care is, and which models of care work best in different settings. Late referral is common, and this means that patients and their families are less able to access low cost, effective care.

In order to answer these questions, investment in research is necessary. Currently in the UK, less than 0.5% of the cancer research budget is allocated to research that aims to improve end-of-life cancer care.⁵² In 2010 this money was allocated in 48 separate awards, providing an average of just £25,000 per award. The danger of this piecemeal approach to palliative care funding is to produce a mass of small-scale studies that have little impact, while national policies are developed without an adequate evidence base.

The importance of evidence based policy making in care of the dying is exemplified by the Liverpool Care Pathway.⁵³ Developed in the late 1990s, the LCP aimed to improve care for people dying outside specialist palliative care settings. Greeted with enthusiasm by policy makers, it was rapidly rolled out nationally, and was endorsed in the National End-of-life Care Strategy. However, following sustained media focus

on its potential harms an independent review in 2013 found that there was a lack of convincing evidence of its benefits, and recommended that the LCP should be phased out.²³

It is essential that future policies aiming to improve care for people who are dying are based on appropriate evidence, and sustained investment in research is necessary to deliver this.

Conclusions

The nature of illness is changing. As a population, we are living longer and survival from disease is prolonged. This, together with a projected increase in the annual number of deaths,⁵⁴ presents an unprecedented challenge for health care systems.

While the search for a cure for cancer occupies many newspaper headlines, the fact that roughly 50% of people diagnosed with cancer will die from it is barely acknowledged. Our societal focus on cure must not come at the expense of openness about what to do when cure is no longer an option. When Aneurin Bevan founded the UK's National Health Service in 1948, his vision was of a cradle to grave health service, freely available to all. However, the huge progress in maternity care in the intervening decades has not been matched by progress in palliative and end-of-life care. Investment in palliative care has the potential to transform the quality of life and death for patients, and lead to improved outcomes in bereavement for their carers, while potentially avoiding burdensome (and expensive) hospital care. As the world's population ages, and costs of care become an important component of health care planning, policy makers must turn their attention to care of the dying. The question is no longer why, but how, effective palliative and end-of-life care is provided.

Reference

1. World Health Organisation, pp
<http://www.who.int/cancer/palliative/definition/en/>
2. Murtagh F: Can palliative care teams relieve some of the pressure on acute services? *Bmj* 348:g3693, 2014
3. Harris PS, Stalam T, Ache KA, et al: Can hospices predict which patients will die within six months? *J Palliat Med* 17:894-8, 2014
4. Higginson IJ, Finlay I, Goodwin DM, et al: Do hospital-based palliative teams improve care for patients or families at the end of life? *J Pain Symptom Manage* 23:96-106, 2002
5. Abernethy AP, Currow DC, Fazekas BS, et al: Specialized palliative care services are associated with improved short- and long-term caregiver outcomes. *Support Care Cancer* 16:585-97, 2008
6. Garcia-Perez L, Linertova R, Martin-Olivera R, et al: A systematic review of specialised palliative care for terminal patients: which model is better? *Palliat Med* 23:17-22, 2009
7. Dionne-Odom JN, Azuero A, Lyons KD, et al: Benefits of Early Versus Delayed Palliative Care to Informal Family Caregivers of Patients With Advanced Cancer: Outcomes From the ENABLE III Randomized Controlled Trial. *J Clin Oncol* 33:1446-52, 2015
8. Gomes B, Calanzani N, Curiale V, et al: Effectiveness and cost-effectiveness of home palliative care services for adults with advanced illness and their caregivers. *Cochrane Database Syst Rev* 6:CD007760, 2013
9. Henson LA, Gao W, Higginson IJ, et al: Emergency department attendance by patients with cancer in their last month of life: a systematic review and meta-analysis. *J Clin Oncol* 33:370-6, 2015
10. Obermeyer Z, Makar M, Abujaber S, et al: Association between the Medicare hospice benefit and health care utilization and costs for patients with poor-prognosis cancer. *Jama* 312:1888-96, 2014
11. Seow H, Brazil K, Sussman J, et al: Impact of community based, specialist palliative care teams on hospitalisations and emergency department visits late in life and hospital deaths: a pooled analysis. *Bmj* 348:g3496, 2014
12. Temel JS, Greer JA, Muzikansky A, et al: Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 363:733-42, 2010
13. Bakitas MA, Tosteson TD, Li Z, et al: Early Versus Delayed Initiation of Concurrent Palliative Oncology Care: Patient Outcomes in the ENABLE III Randomized Controlled Trial. *J Clin Oncol* 33:1438-45, 2015
14. Irwin KE, Greer JA, Khatib J, et al: Early palliative care and metastatic non-small cell lung cancer: potential mechanisms of prolonged survival. *Chron Respir Dis* 10:35-47, 2013
15. Edmonds P, Hart S, Wei G, et al: Palliative care for people severely affected by multiple sclerosis: evaluation of a novel palliative care service. *Mult Scler* 16:627-36, 2010
16. Higginson IJ, McCrone P, Hart SR, et al: Is short-term palliative care cost-effective in multiple sclerosis? A randomized phase II trial. *J Pain Symptom Manage* 38:816-26, 2009

17. Higginson IJ, Bausewein C, Reilly CC, et al: An integrated palliative and respiratory care service for patients with advanced disease and refractory breathlessness: a randomised controlled trial. *Lancet Respir Med* 2:979-87, 2014
18. Lowther K, Selman L, Simms V, et al: **A randomised controlled trial of a nurse-led palliative care intervention for HIV positive patients on ART in Kenya.** *Lancet HIV*, 2015
19. Evi Hatziaandreu FA, Andrew Daly: The potential cost savings of greater use of home and hospice-based end of life care in England. Cambridge: RAND Corporation, 2008
20. Riley GF, Lubitz JD: Long-term trends in Medicare payments in the last year of life. *Health Serv Res* 45:565-76, 2010
21. Smith TJ, Hillner BE: Bending the cost curve in cancer care. *N Engl J Med* 364:2060-5, 2011
22. Smith S, Brick A, O'Hara S, et al: Evidence on the cost and cost-effectiveness of palliative care: a literature review. *Palliat Med* 28:130-50, 2014
23. More care, less pathway: a review of the Liverpool Care Pathway, Department of Health, 2013
24. Weeks JC, Catalano PJ, Cronin A, et al: Patients' expectations about effects of chemotherapy for advanced cancer. *N Engl J Med* 367:1616-25, 2012
25. Keating NL, Landrum MB, Rogers SO, Jr., et al: Physician factors associated with discussions about end-of-life care. *Cancer* 116:998-1006, 2010
26. Mack JW, Cronin A, Keating NL, et al: Associations between end-of-life discussion characteristics and care received near death: a prospective cohort study. *J Clin Oncol* 30:4387-95, 2012
27. Field D, Wee B: Preparation for palliative care: teaching about death, dying and bereavement in UK medical schools 2000-2001. *Med Educ* 36:561-7, 2002
28. Palliative Care Funding Review, 2011
29. Sleeman KE, Davies JM, Verne J, et al: The changing demographics of inpatient hospice death: Population-based cross-sectional study in England, 1993-2012. *Palliat Med*, 2015
30. What's important to me. A Review of Choice in End of Life Care. The Choice in End of Life Care Programme Board, 2015
31. Brennan F: Palliative care as an international human right. *Journal of Pain and Symptom Management* 33:494-499, 2007
32. Gwyther L, Brennan F, Harding R: Advancing palliative care as a human right. *J Pain Symptom Manage.* 38:767-774, 2009
33. Organisation WH: Global Atlas of Palliative Care at the End of Life, in Connor S, Sepulveda C (eds). Geneva, 2014
34. Merriman A, Harding R: Pain control in the African context: the Ugandan introduction of affordable morphine to relieve suffering at the end of life. *Philos Ethics Humanit Med* 5:10, 2010
35. Logie DE, Harding R: An evaluation of a morphine public health programme for cancer and AIDS pain relief in Sub-Saharan Africa. *BMC Public Health* 5:82, 2005
36. Harding R, Selman L, Agupio G, et al: Validation of a core outcome measure for palliative care in Africa: the APCA African Palliative Outcome Scale. *Health Qual Life Outcomes* 8:10, 2010

37. Harding R, Selman L, Agupio G, et al: The prevalence and burden of symptoms amongst cancer patients attending palliative care in two African countries. *Eur J Cancer* 47:51-6, 2011
38. Stanford J, Sandberg DM, Gwyther L, et al: Conversations worth having: the perceived relevance of advance care planning among teachers, hospice staff, and pastors in Knysna, South Africa. *J Palliat Med* 16:762-7, 2013
39. Selman L, Siegert RJ, Higginson IJ, et al: The "Spirit 8" successfully captured spiritual well-being in African palliative care: factor and Rasch analysis. *J Clin Epidemiol* 65:434-43, 2012
40. Selman L, Harding R, Higginson I, et al: Spiritual wellbeing in sub-Saharan Africa: the meaning and prevalence of 'feeling at peace'. *BMJ Support Palliat Care* 1:A22, 2011
41. Selman L, Higginson IJ, Agupio G, et al: Meeting information needs of patients with incurable progressive disease and their families in South Africa and Uganda: multicentre qualitative study. *BMJ* 338:b1326, 2009
42. Stjernsward J, Foley KM, Ferris FD: The public health strategy for palliative care. *J Pain Symptom Manage* 33:486-93, 2007
43. Harding R, Selman L, Powell RA, et al: Research into palliative care in sub-Saharan Africa. *Lancet Oncol* 14:e183-8, 2013
44. Harding R, Higginson IJ: Inclusion of end-of-life care in the global health agenda. *Lancet Glob Health* 2:e375-6, 2014
45. Justo Roll I, Simms V, Harding R: Multidimensional problems among advanced cancer patients in Cuba: awareness of diagnosis is associated with better patient status. *J Pain Symptom Manage* 37:325-30, 2009
46. Hastie BA, Gilson AM, Maurer MA, et al: An examination of global and regional opioid consumption trends 1980-2011. *J Pain Palliat Care Pharmacother* 28:259-75, 2014
47. Dehghan R, Ramakrishnan J, Uddin-Ahmed N, et al: 'They patiently heard what we had to say... this felt different to me': the palliative care needs and care experiences of advanced cancer patients and their families in Bangladesh. *BMJ Supportive & Palliative Care* 2:145-149, 2012
48. Dehghan R, Ramakrishnan J, Ahmed N, et al: The use of morphine to control pain in advanced cancer: an investigation of clinical usage in Bangladesh. *Palliat.Med* 24:707-714, 2010
49. Singh T, Harding R: Palliative care in South Asia: a systematic review of the evidence for care models, interventions, and outcomes. *BMC Res Notes* 8:172, 2015
50. Harding R, Higginson IJ: Palliative care in sub-Saharan Africa. *Lancet* 365:1971-7, 2005
51. Harding R: Caregivers of people at the end of life: emerging evidence. *Curr Opin Support Palliat Care*, 2014
52. Sleeman KE, Gomes B, Higginson IJ: Research into end-of-life cancer care--investment is needed. *Lancet* 379:519, 2012
53. Sleeman KE, Collis E: The Liverpool care pathway: a cautionary tale. *Bmj* 347:f4779, 2013
54. National Population Projections, 2012-based Statistical Bulletin. Office for National Statistics, 2013