King's Research Portal

DOI:

[10.1016/j.brainresbull.2016.06.015](https://doi.org/10.1016/j.brainresbull.2016.06.015)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Gordon-Weeks, P. R. (2016). The role of the drebrin/EB3/Cdk5 pathway in dendritic spine plasticity, implications for Alzheimer's disease. *Brain Research Bulletin*, -. <https://doi.org/10.1016/j.brainresbull.2016.06.015>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Title: The role of the drebrin/EB3/Cdk5 pathway in dendritic spine plasticity, implications for Alzheimer's disease

Author: Phillip R. Gordon-Weeks

PII: S0361-9230(16)30139-3

DOI: <http://dx.doi.org/doi:10.1016/j.brainresbull.2016.06.015>

Reference: BRB 9047

To appear in: *Brain Research Bulletin*

Received date: 15-5-2016

Revised date: 23-6-2016

Accepted date: 25-6-2016

Please cite this article as: Phillip R.Gordon-Weeks, The role of the drebrin/EB3/Cdk5 pathway in dendritic spine plasticity, implications for Alzheimer's disease, Brain Research Bulletin <http://dx.doi.org/10.1016/j.brainresbull.2016.06.015>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**The role of the drebrin/EB3/Cdk5 pathway in dendritic spine plasticity,
implications for Alzheimer's disease.**

Phillip R. Gordon-Weeks

Centre for Developmental Neurobiology,

New Hunt's House, Guy's Campus,

King's College London,

London SE1 1UL,

United Kingdom.

Address correspondence to:

Professor Phillip R. Gordon-Weeks,
Centre for Developmental Neurobiology,
New Hunt's House, Guy's Campus,
King's College London,
London SE1 1UL,
United Kingdom.

Phone: 020 7848 6467

Email: phillip.gordon-weeks@kcl.ac.uk

Highlights

- The cytoskeleton underlies cell shape changes in response to guidance cues
- The drebrin/EB3/Cdk5 pathway couples dynamic microtubules to actin filaments
- Microtubule insertion into dendritic spines facilitates spine plasticity
- Microtubule insertion into dendritic spines requires drebrin and EB3
- Loss of drebrin contributes to cognitive decline In Alzheimer's disease

Abstract

The drebrin/EB3/Cdk5 intracellular signaling pathway couples actin filaments to dynamic microtubules in cellular settings where cells are changing shape. The pathway has been most intensively studied in neuronal development, particularly neuritogenesis and neuronal migration, and in synaptic plasticity at dendritic spines in mature neurons. Drebrin is an actin filament side-binding and bundling protein that stabilizes actin filaments. The end-binding (EB) proteins are microtubule plus-end tracking proteins (+TIPs) that localize to the growing plus-ends of dynamic microtubules and regulate their behavior and the binding of other +TIP proteins. EB3 binds specifically to drebrin when drebrin is bound to actin filaments, for example at the base of a growth cone filopodium, and EB3 is located at the plus-end of a growing microtubule inserting into the filopodium. This interaction therefore forms the basis for coupling dynamic microtubules to actin filaments in growth cones of developing neurons. Appropriate responses to growth cone guidance cues depend on actin filament/microtubule co-ordination in the growth cone, although the role of the drebrin/EB3/Cdk5 pathway in this context has not been directly tested. A similar cytoskeleton coupling pathway operates in dendritic spines in mature neurons where the activity-dependent insertion of dynamic microtubules into dendritic spines is facilitated by drebrin binding to EB3. Microtubule insertion into dendritic spines drives spine maturation during long-term potentiation and therefore has a role in synaptic plasticity and memory formation. In Alzheimer's disease and related chronic neurodegenerative diseases, there is an early and dramatic loss of drebrin from dendritic spines that precedes synapse loss and neurodegeneration and might contribute to a failure of synaptic plasticity and hence to cognitive decline.

Keywords: dendritic spines, synaptic plasticity, actin, microtubules, drebrin, EB3, ADF/cofilin, myosin II, Alzheimer's disease

Acknowledgements: This work was supported by the BBSRC (Grant no: BB/M006336/1).

1. Introduction

Drebrin is a filamentous (F)-actin side-binding and bundling protein with roles in neuronal development and synaptic plasticity (reviewed in: Dun and Chilton, 2010; Shirao and González-Billault, 2013; Gordon-Weeks and Fournier, 2014; Gordon-Weeks, 2016). Drebrin is highly expressed in neurons, where it has been intensively studied, but there are also high levels in other cell types including mesangial cells, muscle, lymphocytes, podocytes, seminiferous epithelium, endothelial cells and some metastatic tumours (Dun and Chilton, 2010; Dart and Gordon-Weeks, 2016). In the mammalian nervous system, an embryonic form, drebrin E, is replaced by an alternatively spliced adult form, drebrin A, distinguished by a 46 amino acid insert of unknown function (Kojima et al., 2016). In embryonic rat hippocampus, for example, drebrin E predominates whereas in early postnatal hippocampus, during the time of maximum synaptogenesis, both forms are present and by adolescence drebrin A predominates (Kojima et al., 2016). Drebrin E is distributed throughout the developing neuron, although highly concentrated in growth cones, whereas drebrin A is particularly concentrated in the dendritic spines of excitatory (glutamatergic) synapses in the cerebral cortex and hippocampus. The sub-cellular distribution of drebrin in growth cones is well documented whereas its distribution in dendritic spines is less clear because of their very small size. In growth cones, drebrin E is mainly localised to the transition zone (Tzone), which forms at the junction between the central and peripheral domains, where anti-parallel bundles of F-actin are found, and the base of some filopodia, where parallel actin-filament bundles occur (Geraldo et al., 2008; Mizui et al., 2009; Worth et al., 2013). Although wide-field and confocal immunofluorescence microscopy suggest that drebrin A is present in both the neck and the head of dendritic spines (Ivanov et al., 2009a) the precise localisation of the protein, particularly in relation to F-actin in spines, is unclear. Although super-resolution microscopy is increasingly being used to determine the distribution of F-actin in dendritic spines it has not yet been used to determine the distribution of drebrin (MacGillavry and Hoogenraad, 2015).

The organisation of F-actin in spines is not completely understood. In spine heads, F-actin is branched with the barbed ends of the filaments directed distally, whereas in the spine neck, F-actin can also be bundled, although whether these are parallel or anti-parallel bundles is unresolved (Fifkova and Delay, 1982; Korobova and Svitkina, 2010). Dendritic spines are thought

to develop from dendritic filopodia but these have several structural and compositional features that are different from the filopodia of growth cones and migrating cells. Although the major cytoskeletal component of dendritic filopodia is F-actin, as it is in non-dendritic filopodia, the organisation of the filaments is different. F-actin in dendritic filopodia is mainly branched and the occasional bundles are not always parallel whereas in non-dendritic filopodia the F-actin is highly bundled and parallel (Korobova and Svitkina, 2010). Consistent with these differences in F-actin organisation, dendritic filopodia contain Arp2/3, capping proteins and myosin II, and no fascin, unlike non-dendritic filopodia (Korobova and Svitkina, 2010). Arp2/3 and capping protein are necessary for F-actin branching and myosin II forms anti-parallel bundles of F-actin whereas fascin forms parallel bundles. Drebrin clearly binds to parallel F-actin bundles, which are the only ones found in growth cone filopodia, but to what extent it binds to anti-parallel bundles is unclear, although these are found in the Tzone of the growth cone, where drebrin localises (Geraldo et al., 2008; Mizui et al., 2009) and possibly in dendritic filopodia where drebrin is also present. Myosin IIB is also concentrated in the Tzone where it organises anti-parallel F-actin bundles (Rochlin et al., 1995; Medeiros et al., 2006) and, since drebrin probably binds myosin IIB (Table 1), this interaction might contribute to its Tzone localisation, as well as direct F-actin binding.

Drebrin is a multi-domain protein of which the best characterised is the N-terminal actin-depolymerising factor homology (ADFH) domain (Fig. 1, Larbolette et al., 1999; Kessels et al., 2000; Xu and Stamnes, 2006). The ADFH domain is structurally conserved among five families of actin-binding proteins including actin depolymerising factor (ADF)/cofilin, Abp1 and twinfilin (Poukkula et al., 2011). In these other families, the ADFH domain binds globular (G)- or F-actin but there is no evidence of actin binding by the drebrin ADFH domain (Hayashi et al., 1999; Chew et al., 2005; Xu and Stamnes, 2006; Grintsevich et al., 2010; Worth et al., 2013). Instead, F-actin-binding activity in drebrin resides in two adjacent domains downstream of the ADFH domain: a coiled-coil (CC) and a helical (Hel) domain (Fig. 1) that independently or co-operatively bind or bundle F-actin (Worth et al., 2013). There then follows a polyproline-rich region (PP) and a large C-terminal region called the blue box (BB, Worth et al., 2013) that harbours binding sites for the PSD scaffolding protein cupid/homer 2 (Shiraishi-Yamaguchi et al., 2009) and motifs for Siah-2 binding (Trivedi et al., 2016) (Table 1). Since profilin is a candidate drebrin interactor (Table 1) and

interacts with polyproline regions of many proteins, it is tempting to suggest that the PP domain of drebrin might bind profilin. The 46 amino acid insert that distinguishes the A form of drebrin from drebrin E is located at the C-terminal end of the Hel domain and has no known function (Fig. 1). The insert in drebrin A might confer a novel function on the molecule or modulate an existing one, such as F-actin binding. Although there has been no comparative characterization of the F-actin-binding properties of drebrin E and A, heterologous expression of either isoform in cells has similar effects on F-actin organization and stability (Dun and Chilton, 2010).

There is cell biological and *in vitro* structural evidence that drebrin stabilises F-actin. Over-expression of drebrin E or A in heterologous cells and neurons increases the levels of F-actin (Ikeda et al., 1996; Geraldo et al., 2008; Mizui et al., 2009; Worth et al., 2013) whereas, conversely, siRNA-mediated knockdown of drebrin decreases F-actin levels in growth cones of cultured cortical neurons (Mizui et al., 2009). In vascular smooth muscle cells isolated from the heterozygotes of drebrin knockout mice, where drebrin is reduced by a half, there is a reduction of F-actin (Stiber et al., 2016). Over-expressing high levels of drebrin in heterologous cells induces the formation of prominent cell processes containing thick bundles of F-actin (Shirao et al., 1992, 1994; Keon et al., 2000; Chew et al., 2005; Geraldo et al., 2008; Worth et al., 2013). Drebrin binding to F-actin *in vitro* produces a twist in the filament that leads to filament stiffening and stabilisation (Sharma et al., 2011; Mikati et al., 2013). Drebrin protects F-actin against depolymerising agents such as cytochalasins (Ikeda et al., 1996; Ivanov et al., 2009b). There is biochemical evidence that drebrin competes with many other actin-binding proteins for actin filaments including ADF/cofilin, α -actinin, tropomyosins and fascin and this might indirectly affect F-actin stability (Ishikawa et al., 1994; Sasaki et al., 1996; Zhao et al., 2006; Grintsevich et al., 2010, 2014; Sharma et al., 2011; Mikati et al., 2013). Whether these are examples of direct competition for a common binding site or indirect competition resulting from allosteric effects is not clear. There is *in vitro* evidence that drebrin, binding to F-actin, can inhibit the F-actin severing activity of ADF/cofilin (Grintsevich et al., 2014). The competition between drebrin and ADF/cofilin for F-actin binding might be an example of an allosteric effect because ADF/cofilin produces an opposing twist in F-actin to that produced by drebrin (Sharma et al., 2011). As pointed out by Cammarata et al

(2016), the potential inhibition of the F-actin severing activity of ADF/cofilin by drebrin competition might impact on the progression of microtubules into filopodia during drebrin/EB3 coupling.

Several binding partners for drebrin have been identified including: afadin, EB3, cupidin-2/homer, CXCR4, myosin II, profilin and Shia-2 (Table 1). The microtubule-binding protein EB3 is a member of a family of so-called +TIP proteins that bind to the plus-ends of microtubules and regulate their dynamics (reviewed in: van de Willige et al., 2016). Mammalian cells express three EB proteins: EB1 and EB3 are microtubule plus-end trackers whereas EB2 is at best a weak plus-end tracker and is the least studied member of the family. The EB proteins are positioned at the core of a cluster of proteins that localise to microtubule plus-ends by virtue of the fact that they harbour binding sites for other +TIP proteins including APC and CLASP (Cammarata et al., 2016). EB1 and EB3 have both common and distinct binding partners and drebrin is one of a growing number of proteins that bind preferentially to EB3 (Table 2). In neurons undergoing neuritogenesis and in growth cones, drebrin, bound to parallel F-actin at the base of filopodia, interacts with EB3 located at the plus-ends of dynamic microtubules entering filopodia (Geraldo et al., 2008; Worth et al., 2013). The evidence for this includes the demonstration of direct binding between drebrin and EB3 in filopodia using FRET/FLIM (Geraldo et al., 2008) and the observation that a dominant negative construct, EB3M, containing the drebrin binding domain of EB3 blocks neuritogenesis (Geraldo et al., 2008) and neuronal cell migration (Trivedi et al., 2016). Thus, the drebrin/EB3 interaction cross-links dynamic microtubules to F-actin in filopodia and this likely forms part of the cellular mechanism underlying the response to axon growth and growth cone guidance cues (Cammarata et al., 2016). The drebrin/EB3/Cdk5 pathway has been independently confirmed in other cellular contexts where dynamic microtubules interacting with F-actin underlies cell shape changes, suggesting that the pathway is canonical (Bazellières et al., 2012; Merriam et al., 2013; Ketschek et al., 2016).

The functions of drebrin and EB3 are regulated by direct phosphorylation (Gordon-Weeks, 2016). Drebrin exists in two structural conformations, a closed one in which the BB auto-inhibits the F-actin-binding activity in the CC domain by binding to a region centred on S142 near the junction between the ADFH domain and the CC domain (Fig. 1) and an open conformation in which binding of the BB is relieved by cyclin-dependent kinase 5 (Cdk5) phosphorylation of S142

(Worth et al., 2013). In the closed conformation, drebrin can only bind to single actin filaments whereas in the open conformation both F-actin-binding activities are available and drebrin can bundle F-actin or straddle pre-existing F-actin filaments (Fig. 1). As well as regulating drebrin F-actin binding, Cdk5 phosphorylation, by converting drebrin to the open conformation, exposes the binding site for EB3 (Worth et al., 2013). EB3 is a substrate of the mitotic kinases Aurora A and B (Ban et al., 2009) and the phosphatase calcineurin and dephosphorylation activates EB3 (Komarova et al., 2012).

2. Drebrin function in dendritic spines

The dendritic spines of excitatory (glutamatergic) synapses on cortical and hippocampal neurons undergo a series of morphological and molecular changes associated with changes in synaptic strength and memory formation (reviewed in: Rochefort and Konnerth, 2012; Hlushchenko et al., 2016). Synapse strength can either be enhanced, by short bursts of high-frequency input, leading to long-term potentiation (LTP), or reduced, by prolonged low frequency input, causing long-term depression (LTD). LTP is induced by activation of the N-methyl-D-aspartate receptor (NMDA) which leads to long-lasting incorporation of α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptors in the post-synaptic density (PSD) of dendritic spines and to changes in spine morphology. Despite spine shape being a continuum, three main spine shape categories have been recognized: thin, stubby and mushroom, in increasing order of size (reviewed in: Rochefort & Konnerth 2012; Hlushchenko et al., 2016). An increase in synaptic strength accompanying LTP is associated with spine enlargement, whereas LTD, where synaptic strength declines, is associated with a decrease in spine size and morphologically simpler spines. Changes in dendritic spine morphology are achieved by the rapid and reversible re-organization of the cytoskeleton within the spine. The dynamic behaviour of two cytoskeletal filaments, F-actin and microtubules, dominate dendritic spine changes in response to synaptic activity. Increased synaptic activity and strength is accompanied by an increase in spine F-actin and F-actin stability, which is responsible for spine enlargement, whereas a reduction in synaptic activity reduces F-actin in spines and increases F-actin dynamics (Okamoto et al., 2004; Honkura et al., 2008). Spine head volume correlates closely with the area of the PSD (Harris and Stevens, 1989) and the number of AMPA receptors (Matsuzaki et al., 2001).

2.1. Structural role of drebrin A in spine morphogenesis

Drebrin A influences dendritic spine morphology by regulating F-actin, plays a role in activity-dependent accumulation of PSD-95, the major scaffolding protein at the PSD, and in redistribution of the NMDA type of glutamate receptor (Ivanov et al., 2009b). These effects of drebrin on spine activity are dependent on F-actin binding (Ivanov et al., 2009b). Over-expression of drebrin in mature neurons in culture produces dendritic spines with abnormally long necks (Hayashi and Shirao, 1999; Ivanov et al., 2009b; Merriam et al., 2013) consistent with the idea that drebrin stabilises F-actin in the spine neck.

2.2. Activity-driven microtubule capture and insertion into dendritic spines

Dynamic microtubules in dendrite shafts can transiently enter dendritic spines by polymerizing from their plus-ends, the end at which tubulin assembly occurs preferentially (Gu et al. 2008; Hu et al. 2008; Jaworski et al. 2009; Wagner et al., 2011; Merriam et al., 2013). The frequency of invasion is low, with approximately 1% of dendritic protrusions containing a microtubule at any one time (Hu et al. 2008) and the dwell time is in the order of minutes but, importantly, dwell time and the frequency of invasion are enhanced by pharmacological interventions that mimic LTP (Gu et al. 2008, Hu et al. 2008; Mitsuyama et al. 2008; Jaworski et al. 2009; Merriam et al. 2011) and suppressed by eliciting LTD (Kapitein et al. 2011). Depolymerisation of microtubules in dendrites using drugs such as nocodazole reduces spine density along the dendrite without affecting dendrite numbers (Jaworski et al., 2009). Dendritic microtubules are oriented with their plus-ends directed either towards or away from the cell body and can therefore polymerise antero- or retrogradely, and can be nucleated acentrosomally within the dendrite (Delandre et al., 2016). Microtubules invading spines are oriented with their plus-ends distal, i.e. directed toward the spine head. This behavior is reminiscent of microtubules invading the filopodia of growth cones (Geraldo & Gordon-Weeks 2009; Gordon-Weeks & Fournier, 2013) and axonal branches (Ketschek et al., 2016). In all of these cases, the microtubule-binding +TIP protein EB3, is located at the tip of the growing microtubule. There is now considerable evidence that microtubule invasion of dendritic spines underlies morphological and physiological changes

associated with memory formation including F-actin stabilization, spine maturation and enlargement of the PSD (Dent et al. 2011). Consistent with these findings, shRNA-mediated knock-down of EB3 reduces spine enlargement (Gu et al. 2008; Jaworski et al. 2009). An early event in dendritic spine enlargement associated with increased synaptic strength is an enlargement of the PSD and a concomitant increase in PSD-95, the major scaffolding protein of the PSD. Brain-derived neurotrophic factor (BDNF) induces spine formation and maturation, an effect that is blocked by microtubule depolymerisation using nocodazole and enhanced by stabilizing microtubules with substoichiometric concentrations of taxol (Gu et al., 2008). Exposure of dendritic spines to BDNF induces an increase of the post-synaptic density protein PSD-95 in the PSD along with spine maturation and this has also been shown to depend on invasion of dendritic spines by microtubules (Hu et al. 2011). Importantly, the frequency of microtubule invasion into dendritic spines and the number of spines invaded is enhanced by over-expression of drebrin and, conversely, reduced by siRNA-mediated knock-down of drebrin (Merriam et al., 2013). Whether drebrin and EB3 are acting independently of each other or are interacting in the drebrin/EB3 pathway in spine maturation has yet to be determined.

3. Synaptotoxicity and dendritic spine loss in Alzheimer's disease

Alzheimer's disease is primarily a disease of synapse dysfunction caused by Amyloid β ($A\beta$) oligomers derived from Alzheimer's precursor protein and loss of synapses is the tightest pathological correlate of cognitive decline in Alzheimer's disease (Klein, 2013).

3.1. Drebrin loss from dendritic spines precedes synapse loss

One of the earliest molecular changes that precede synapse loss and neurodegeneration in Alzheimer's disease and related cognitive disorders is a dramatic loss of drebrin from dendritic spines (Ivanov et al., 2009a; Dun and Chilton, 2010). This occurs in mild cognitive impairment (Counts *et al.*, 2012), Alzheimer's disease (Harigaya et al., 1996; Hatanpaa et al., 1999; Shim and Lubec, 2002; Calon et al., 2004; Zhao et al., 2006) and Down's syndrome, where individuals inevitably develop Alzheimer's (Weitzdoerfer et al., 2001; Shim and Lubec, 2002). Whether or not drebrin loss from dendritic spines is causal to cognitive decline in Alzheimer's disease and related cognitive disorders is not fully resolved. There is a correlation between the degree of drebrin

reduction in dendritic spines and the severity of cognitive impairment (Kojima and Shirao, 2007). Transgenic mouse models of familial Alzheimer's disease also show a loss of drebrin from dendritic spines (Lee and Aoki 2012; Calon et al., 2004; Calon et al., 2005; Mahadomrongkul et al., 2005; Aoki et al., 2007; Lacor et al., 2007). Drebrin loss occurs before loss of pre-synaptic markers, such as synaptophysin, suggesting it precedes synapse loss (Counts et al., 2012).

Reduction of drebrin A by intracerebral administration of antisense oligonucleotides in rats causes cognitive deficits (Kobayashi et al., 2004) and transgenic mice in which the exon coding for the spliced insert of drebrin A has been deleted by homologous recombination, so that only the E form can be expressed, show deficits in homeostatic dendritic spine plasticity (Aoki et al., 2009) and subtle deficits in cognition (Kojima et al., 2010). However, there is a compensatory up-regulation of drebrin E in these mice which probably limits the severity of the synaptic impairment (Kojima et al., 2010). Nonetheless, in the drebrin A knockout mouse, LTP is impaired in the adult, but not, curiously, in the adolescent, suggesting that drebrin A, and therefore presumably the insert, has a unique functional role in dendritic spine plasticity (Kojima et al., 2016).

Drebrin loss from dendritic spines can be replicated *in vitro* in mature embryonic hippocampal cultures within hours by exposure to A β oligomers (Zhao et al., 2006; Lacor et al., 2007). Recently, proteolysis of drebrin has been proposed as the mechanism underlying the pathological loss of drebrin from dendritic spines (Chimura et al., 2015). The activity of several proteases, including calpain and caspases, are increased in Alzheimer's disease. Drebrin is a caspase-6 target (LeBlanc, 2013), but it is not known if it is cleaved by caspase-6 in Alzheimer's disease. Prolonged over-activation of NMDA-type glutamate receptors in cultured hippocampal neurons, which mimics some features of the effects of A β oligomers on neurons, causes an increase in intracellular calcium which activates calpain, a calcium-dependent protease, resulting in drebrin proteolysis and, at the same time, a decrease in F-actin (Chimura et al., 2015). Calpain activity is also responsible for the proteolytic conversion of p35, a regulatory subunit of Cdk5, to p25, a truncated and constitutively active form which enhances Cdk5 activity in Alzheimer's disease (reviewed in: Shah and Lahiri, 2014). This might abnormally increase the levels of S142 phosphorylated drebrin impacting F-actin organization. If the open conformation of drebrin, i.e. the S142 phosphorylated form, is more susceptible to proteolysis than the closed form this might lead

to a positive feed-forward loop with consequent catastrophic drebrin degradation. Although proteolysis at synapses increases in Alzheimer's disease, drebrin breakdown products have not been seen. Proteolytic fragments might have been missed because only one antibody had been used previously to detect drebrin by immunoblotting in Alzheimer's disease brain tissue and transgenic mouse models of Alzheimer's disease and the epitope that it recognises, which is in the C-terminal BB, might be lost from proteolytic fragments (Hatanpaa et al., 1999; Calon et al., 2004; Zhao et al., 2006; Counts et al. 2012). It is unclear whether the loss of drebrin through proteolysis causes a loss of drebrin function or a dominant negative effect through one of the proteolytic fragments. Future studies will also have to address the question of whether or not A β can also induce drebrin proteolysis. In contrast, brief and transient activation of NMDA receptors *in vitro* has been reported to accompany a transient shift of drebrin A from dendritic spines to dendritic shafts followed by a re-entry into spines without changes in overall drebrin levels (Sekino et al., 2006; Bosch and Hayashi, 2012; Merriam et al., 2013; Mizui et al., 2014). LTP *in vivo* is also associated with an influx of drebrin into the spine (Fukazawa et al., 2003; Bosch et al., 2014).

Changes in other actin-binding proteins that occur in neurodegenerative disease might displace drebrin from F-actin in the dendritic spine into the dendritic shaft where proteolytic degradation could occur. This could occur through direct competition with drebrin binding to F-actin or indirectly through changes to F-actin such as depolymerisation brought about by severing or structural changes in the filament. In Alzheimer's disease there is an inverse relationship between the loss of drebrin and an increase in ADF/cofilin (Zhao et al., 2006). ADF/cofilin activity, including F-actin severing, is negatively regulated by LIM kinase phosphorylation and the activity and levels of the kinase that activates LIM kinase, PAK, is reduced in Alzheimer's disease (Zhao et al., 2006). Significantly, PAK can protect against A β oligomer-induced drebrin loss (Zhao et al., 2006).

4. Conclusions and future perspectives

The drebrin/EB3/Cdk5 molecular pathway co-ordinates dynamic microtubules and F-actin interactions that underlie various cell shape changes, such as neuritogenesis, neuronal migration, growth cone turning and dendritic spine plasticity, in response to guidance cues. It is unlikely that this pathway is the only one that has evolved to couple dynamic microtubules to F-actin and indeed

other candidate pathways have been identified (Coles and Bradke, 2015). How cells utilise different pathways coupling dynamic microtubules to F-actin and under what circumstances are areas of future research. How this pathway is regulated is only just beginning to be understood. There is evidence for regulation of drebrin and EB3 by phosphorylation and additional interactors of both proteins have also been identified, although the mechanistic details of these regulatory processes have yet to be worked out. Unravelling the distinct roles played by drebrin E and A, and in particular the function of the A insert, are also important goals.

References

- Aoki, C., N. Kojima, N. Sabaliauskas, L. Shah, T.H. Ahmed, J. Oakford, T. Ahmed, H. Yamazaki, K. Hanamura, and T. Shirao. 2009. Drebrin a knockout eliminates the rapid form of homeostatic synaptic plasticity at excitatory synapses of intact adult cerebral cortex. *J Comp Neurol.* 517:105-21.
- Aoki, C., V. Mahadomrongkul, S. Fujisawa, R. Habersat, and T. Shirao. 2007. Chemical and morphological alterations of spines within the hippocampus and entorhinal cortex precede the onset of Alzheimer's disease pathology in double knock-in mice. *J Comp Neurol.* 505:352-62.
- Ban, R., et al., 2009. Mitotic regulation of the stability of microtubule plus-end tracking protein EB3 by ubiquitin ligase SIAH-1 and Aurora mitotic kinases. *J Biol Chem.* 284, 28367-81.
- Bazellieres, E., D. Massey-Harroche, M. Barthelemy-Requin, F. Richard, J.P. Arsanto, and A. Le Bivic. 2012. Apico-basal elongation requires a drebrin-E-EB3 complex in columnar human epithelial cells. *J Cell Sci.* 125:919-31.
- Bosch, M., J. Castro, T. Saneyoshi, H. Matsuno, M. Sur, and Y. Hayashi. 2014. Structural and molecular remodeling of dendritic spine substructures during long-term potentiation. *Neuron.* 82:444-59.
- Butkevich, E., S. Hulsmann, D. Wenzel, T. Shirao, R. Duden, and I. Majoul. 2004. Drebrin is a novel connexin-43 binding partner that links gap junctions to the submembrane cytoskeleton. *Curr Biol.* 14:650-8.
- Calon, F., G.P. Lim, T. Morihara, F. Yang, O. Ubeda, N. Salem, Jr., S.A. Frautschy, and G.M. Cole. 2005. Dietary n-3 polyunsaturated fatty acid depletion activates caspases and decreases NMDA receptors in the brain of a transgenic mouse model of Alzheimer's disease. *Eur J Neurosci.* 22:617-26.
- Calon, F., G.P. Lim, F. Yang, T. Morihara, B. Teter, O. Ubeda, P. Rostaing, A. Triller, N. Salem, Jr., K.H. Ashe, S.A. Frautschy, and G.M. Cole. 2004. Docosahexaenoic acid protects from dendritic pathology in an Alzheimer's disease mouse model. *Neuron.* 43:633-45.
- Cammarata, G.M., E.A. Bearce, and L.A. Lowery. 2016. Cytoskeletal social networking in the growth cone: How +TIPs mediate microtubule-actin cross-linking to drive axon outgrowth and guidance. *Cytoskeleton (Hoboken)*. DOI: 10.1002/cm.21272.
- Chew, C.S., C.T. Okamoto, X. Chen, and R. Thomas. 2005. Drebrin E2 is differentially expressed and phosphorylated in parietal cells in the gastric mucosa. *Am J Physiol Gastrointest Liver Physiol.* 289:G320-31.
- Chimura, T., T. Launey, and N. Yoshida. 2015. Calpain-Mediated Degradation of Drebrin by Excitotoxicity In vitro and In vivo. *PLoS One.* 10:e0125119.
- Counts, S.E., B. He, M. Nadeem, J. Wu, S.W. Scheff, and E.J. Mufson. 2012. Hippocampal drebrin loss in mild cognitive impairment. *Neurodegener Dis.* 10:216-9.
- Coles, C.H. and Bradke, F. 2015. Coordinating Neuronal Actin-Microtubule Dynamics, *Curr Biol*, 25 R677-691.
- Dart, A. E. and Gordon-Weeks, P. R. 2016. Drebrin and cancer. Drebrin, (Shirao, T. ed.), Springer.
- Delandre, C., R. Amikura, and A.W. Moore. 2016. Microtubule Nucleation and Organization in Dendrites. *Cell Cycle*: DOI: 10.1080/15384101.2016.1172158.
- Dent, E.W., E.B. Merriam, and X. Hu. 2011. The dynamic cytoskeleton: backbone of dendritic spine plasticity. *Curr Opin*

Neurobiol. 21:175-81.

Dun, X.P., and J.K. Chilton. 2010. Control of cell shape and plasticity during development and disease by the actin-binding protein Drebrin. *Histol Histopathol.* 25:533-40.

Fifkova, E., and R.J. Delay. 1982. Cytoplasmic actin in neuronal processes as a possible mediator of synaptic plasticity. *J Cell Biol.* 95:345-50.

Fukazawa, Y., Y. Saitoh, F. Ozawa, Y. Ohta, K. Mizuno, and K. Inokuchi. 2003. Hippocampal LTP is accompanied by enhanced F-actin content within the dendritic spine that is essential for late LTP maintenance in vivo. *Neuron.* 38:447-60.

Geraldo, S., U.K. Khanzada, M. Parsons, J.K. Chilton, and P.R. Gordon-Weeks. 2008. Targeting of the F-actin-binding protein drebrin by the microtubule plus-tip protein EB3 is required for neuritogenesis. *Nat Cell Biol.* 10:1181-9.

Geyer, M., et al., 2015. Microtubule-Associated Protein EB3 Regulates IP3 Receptor Clustering and Ca²⁺ Signaling in Endothelial Cells. *Cell Rep.* 12, 79-89.

Gill, M.B., A.S. Kato, H. Wang, and D.S. Bredt. 2012. AMPA receptor modulation by cornichon-2 dictated by transmembrane AMPA receptor regulatory protein isoform. *Eur J Neurosci.* 35:182-94.

Gordón-Alonso, M., V. Rocha-Perugini, S. Alvarez, A. Ursa, N. Izquierdo-Useros, J. Martinez-Picado, M.A. Munoz-Fernandez, and F. Sanchez-Madrid. 2013. Actin-binding protein drebrin regulates HIV-1-triggered actin polymerization and viral infection. *J Biol Chem.* 288:28382-97.

Gordon-Weeks, P. R. (2016). Phosphorylation of Drebrin and its Role in Neuritogenesis. Drebrin, (Shirao, T. ed.), Springer.

Gordon-Weeks, P.R., and A.E. Fournier. 2014. Neuronal cytoskeleton in synaptic plasticity and regeneration. *J Neurochem.* 129:206-12.

Grintsevich, E.E., V.E. Galkin, A. Orlova, A.J. Ytterberg, M.M. Mikati, D.S. Kudryashov, J.A. Loo, E.H. Egelman, and E. Reisler. 2010. Mapping of drebrin binding site on F-actin. *J Mol Biol.* 398:542-54.

Gu, J., B.L. Firestein, and J.Q. Zheng. 2008. Microtubules in dendritic spine development. *J Neurosci.* 28:12120-4.

Harigaya, Y., M. Shoji, T. Shirao, and S. Hirai. 1996. Disappearance of actin-binding protein, drebrin, from hippocampal synapses in Alzheimer's disease. *J Neurosci Res.* 43:87-92.

Harris, K.M., and J.K. Stevens. 1989. Dendritic spines of CA 1 pyramidal cells in the rat hippocampus: serial electron microscopy with reference to their biophysical characteristics. *J Neurosci.* 9:2982-97.

Hatanpaa, K., K.R. Isaacs, T. Shirao, D.R. Brady, and S.I. Rapoport. 1999. Loss of proteins regulating synaptic plasticity in normal aging of the human brain and in Alzheimer disease. *J Neuropathol Exp Neurol.* 58:637-43.

Hayashi, K., R. Ishikawa, R. Kawai-Hirai, T. Takagi, A. Taketomi, and T. Shirao. 1999. Domain analysis of the actin-binding and actin-remodeling activities of drebrin. *Exp Cell Res.* 253:673-80.

Hayashi, K., and T. Shirao. 1999. Change in the shape of dendritic spines caused by overexpression of drebrin in cultured cortical neurons. *J Neurosci.* 19:3918-25.

Hsieh, P.C., et al., 2007. p53 downstream target DDA3 is a novel microtubule-associated protein that interacts with end-

- binding protein EB3 and activates beta-catenin pathway. *Oncogene*. 26, 4928-40.
- Hlushchenko, I., M. Koskinen, and P. Hotulainen. 2016. Dendritic spine actin dynamics in neuronal maturation and synaptic plasticity. *Cytoskeleton (Hoboken)*. DOI: 10.1002/cm.21280.
- Honkura N, Matsuzaki M, Noguchi J, Ellis-Davies GC, Kasai H. 2008. The subspine organization of actin fibers regulates the structure and plasticity of dendritic spines. *Neuron*. 57:719-29.
- Hu, X., C. Viesselmann, S. Nam, E. Merriam, and E.W. Dent. 2008. Activity-dependent dynamic microtubule invasion of dendritic spines. *J Neurosci*. 28:13094-105.
- Ikeda, K., P.A. Kaub, H. Asada, K. Uyemura, S. Toya, T. Shirao, 1996. Stabilization of adhesion plaques by the expression of drebrin A in fibroblasts, *Brain Res Dev Brain Res*, 91:227-236.
- Ishikawa, R., K. Hayashi, T. Shirao, Y. Xue, T. Takagi, Y. Sasaki, and K. Kohama. 1994. Drebrin, a development-associated brain protein from rat embryo, causes the dissociation of tropomyosin from actin filaments. *J Biol Chem*. 269:29928-33.
- Ivanov, A., M. Esclapez, and L. Ferhat. 2009a. Role of drebrin A in dendritic spine plasticity and synaptic function: Implications in neurological disorders. *Commun Integr Biol*. 2:268-70.
- Ivanov, A., M. Esclapez, C. Pellegrino, T. Shirao, and L. Ferhat. 2009b. Drebrin A regulates dendritic spine plasticity and synaptic function in mature cultured hippocampal neurons. *J Cell Sci*. 122:524-34.
- Jaworski, J., L.C. Kapitein, S.M. Gouveia, B.R. Dortland, P.S. Wulf, I. Grigoriev, P. Camera, S.A. Spangler, P. Di Stefano, J. Demmers, H. Krugers, P. Defilippi, A. Akhmanova, and C.C. Hoogenraad. 2009. Dynamic microtubules regulate dendritic spine morphology and synaptic plasticity. *Neuron*. 61:85-100.
- Kapitein, L.C., K.W. Yau, S.M. Gouveia, W.A. van der Zwan, P.S. Wulf, N. Keijzer, J. Demmers, J. Jaworski, A. Akhmanova, and C.C. Hoogenraad. 2011. NMDA receptor activation suppresses microtubule growth and spine entry. *J Neurosci*. 31:8194-209.
- Keon, B. H., P. T. Jedrzejewski, D. L. Paul and D. A. Goodenough. 2000. Isoform specific expression of the neuronal F-actin binding protein, drebrin, in specialized cells of stomach and kidney epithelia. *J. Cell Sci*. 113 Pt 2: 325-336.
- Kessels, M.M., A.E. Engqvist-Goldstein, and D.G. Drubin. 2000. Association of mouse actin-binding protein 1 (mAbp1/SH3P7), an Src kinase target, with dynamic regions of the cortical actin cytoskeleton in response to Rac1 activation. *Mol Biol Cell*. 11:393-412.
- Ketschek, A., M. Spillane, X.P. Dun, H. Hardy, J. Chilton, and G. Gallo. 2016. Drebrin Coordinates the Actin and Microtubule Cytoskeleton During the Initiation of Axon Collateral Branches. *Dev Neurobiol*. DOI: 10.1002/dneu.22377.
- Klein, W.L. 2013. Synaptotoxic amyloid-beta oligomers: a molecular basis for the cause, diagnosis, and treatment of Alzheimer's disease? *J Alzheimers Dis*. 33 Suppl 1:S49-65.
- Kobayashi, R., Y. Sekino, T. Shirao, S. Tanaka, T. Ogura, K. Inada, and M. Saji. 2004. Antisense knockdown of drebrin A, a dendritic spine protein, causes stronger preference, impaired pre-pulse inhibition, and an increased sensitivity to psychostimulant. *Neurosci Res*. 49:205-17.

- Kojima, N., K. Hanamura, H. Yamazaki, T. Ikeda, S. Itohara, and T. Shirao. 2010. Genetic disruption of the alternative splicing of drebrin gene impairs context-dependent fear learning in adulthood. *Neuroscience*. 165:138-50.
- Kojima, N., and T. Shirao. 2007. Synaptic dysfunction and disruption of postsynaptic drebrin-actin complex: a study of neurological disorders accompanied by cognitive deficits. *Neurosci Res*. 58:1-5.
- Kojima, N., H. Yasuda, K. Hanamura, Y. Ishizuka, Y. Sekino, and T. Shirao. 2016. Drebrin A regulates hippocampal LTP and hippocampus-dependent fear learning in adult mice. *Neuroscience*. 324:218-26.
- Komarova, Y.A., et al., 2012. VE-cadherin signaling induces EB3 phosphorylation to suppress microtubule growth and assemble adherens junctions. *Mol Cell*. 48, 914-25.
- Korobova, F., and T. Svitkina. 2010. Molecular architecture of synaptic actin cytoskeleton in hippocampal neurons reveals a mechanism of dendritic spine morphogenesis. *Mol Biol Cell*. 21:165-76.
- Kreis, P., R. Hendricusdottir, L. Kay, I.E. Papageorgiou, M. van Diepen, T. Mack, J. Ryves, A. Harwood, N.R. Leslie, O. Kann, M. Parsons, and B.J. Eickholt. 2013. Phosphorylation of the actin binding protein Drebrin at S647 is regulated by neuronal activity and PTEN. *PLoS One*. 8:e71957.
- Lacor, P.N., M.C. Buniel, P.W. Furlow, A.S. Clemente, P.T. Velasco, M. Wood, K.L. Viola, and W.L. Klein. 2007. Abeta oligomer-induced aberrations in synapse composition, shape, and density provide a molecular basis for loss of connectivity in Alzheimer's disease. *J Neurosci*. 27:796-807.
- Larbolette, O., B. Wollscheid, J. Schweikert, P.J. Nielsen, and J. Wienands. 1999. SH3P7 is a cytoskeleton adapter protein and is coupled to signal transduction from lymphocyte antigen receptors. *Mol Cell Biol*. 19:1539-46.
- LeBlanc, A.C. 2013. Caspase-6 as a novel early target in the treatment of Alzheimer's disease. *Eur J Neurosci*. 37:2005-18.
- Lee, D., and C. Aoki. 2012. Presenilin conditional double knockout mice exhibit decreases in drebrin a at hippocampal CA1 synapses. *Synapse*. 66:870-9.
- Li, M.W., X. Xiao, D.D. Mruk, Y.L. Lam, W.M. Lee, W.Y. Lui, M. Bonanomi, B. Silvestrini, and C.Y. Cheng. 2011. Actin-binding protein drebrin E is involved in junction dynamics during spermatogenesis. *Spermatogenesis*. 1:123-136.
- MacGillavry, H.D., and C.C. Hoogenraad. 2015. The internal architecture of dendritic spines revealed by super-resolution imaging: What did we learn so far? *Exp Cell Res*. 335:180-6.
- Mahadomrongkul, V., P.T. Huerta, T. Shirao, and C. Aoki. 2005. Stability of the distribution of spines containing drebrin A in the sensory cortex layer I of mice expressing mutated APP and PS1 genes. *Brain Res*. 1064:66-74.
- Mammoto, A., T. Sasaki, T. Asakura, I. Hotta, H. Imamura, K. Takahashi, Y. Matsuura, T. Shirao, and Y. Takai. 1998. Interactions of drebrin and gephyrin with profilin. *Biochem Biophys Res Commun*. 243:86-9.
- Matsuzaki, M., G.C. Ellis-Davies, T. Nemoto, Y. Miyashita, M. Iino, and H. Kasai. 2001. Dendritic spine geometry is critical for AMPA receptor expression in hippocampal CA1 pyramidal neurons. *Nat Neurosci*. 4:1086-92.
- Medeiros, N.A., D.T. Burnette, and P. Forscher. 2006. Myosin II functions in actin-bundle turnover in neuronal growth cones. *Nat Cell Biol*. 8:215-26.

- Merriam, E.B., D.C. Lumbard, C. Viesselmann, J. Ballweg, M. Stevenson, L. Pietila, X. Hu, and E.W. Dent. 2011. Dynamic microtubules promote synaptic NMDA receptor-dependent spine enlargement. *PLoS One*. 6:e27688.
- Merriam, E.B., M. Millette, D.C. Lumbard, W. Saengsawang, T. Fothergill, X. Hu, L. Ferhat, and E.W. Dent. 2013. Synaptic regulation of microtubule dynamics in dendritic spines by calcium, F-actin, and drebrin. *J Neurosci*. 33:16471-82.
- Mikati, M.A., E.E. Grintsevich, and E. Reisler. 2013. Drebrin-induced stabilization of actin filaments. *J Biol Chem*. 288:19926-38.
- Mitsuyama, F., G. Niimi, K. Kato, K. Hirose, K. Mikoshiba, M. Okuya, K. Karagiozov, Y. Kato, T. Kanno, H. Sanoe, and T. Koide. 2008. Redistribution of microtubules in dendrites of hippocampal CA1 neurons after tetanic stimulation during long-term potentiation. *Ital J Anat Embryol*. 113:17-27.
- Mizui, T., N. Kojima, H. Yamazaki, M. Katayama, K. Hanamura, and T. Shirao. 2009. Drebrin E is involved in the regulation of axonal growth through actin-myosin interactions. *J Neurochem*. 109:611-22.
- Mizui, T., Y. Sekino, H. Yamazaki, Y. Ishizuka, H. Takahashi, N. Kojima, M. Kojima, and T. Shirao. 2014. Myosin II ATPase activity mediates the long-term potentiation-induced exodus of stable F-actin bound by drebrin A from dendritic spines. *PLoS One*. 9:e85367.
- Okamoto K, Nagai T, Miyawaki A, Hayashi Y. 2004. Rapid and persistent modulation of actin dynamics regulates postsynaptic reorganization underlying bidirectional plasticity. *Nature neuroscience*. 7:1104-12.
- Pérez-Martínez, M., M. Gordon-Alonso, J.R. Cabrero, M. Barrero-Villar, M. Rey, M. Mittelbrunn, A. Lamana, G. Morlino, C. Calabia, H. Yamazaki, T. Shirao, J. Vazquez, R. Gonzalez-Amaro, E. Veiga, and F. Sanchez-Madrid. 2010. F-actin-binding protein drebrin regulates CXCR4 recruitment to the immune synapse. *J Cell Sci*. 123:1160-70.
- Poukkula, M., E. Kremneva, M. Serlachius, and P. Lappalainen. 2011. Actin-depolymerizing factor homology domain: a conserved fold performing diverse roles in cytoskeletal dynamics. *Cytoskeleton (Hoboken)*. 68:471-90.
- Rehm, K., L. Panzer, V. van Vliet, E. Genot, and S. Linder. 2013. Drebrin preserves endothelial integrity by stabilizing nectin at adherens junctions. *J Cell Sci*. 126:3756-69.
- Rochefort, N. L. and Konnerth, A. 2012 Dendritic spines: from structure to in vivo function. *EMBO reports*, **13**, 699-708.
- Rochlin, M.W., K. Itoh, R.S. Adelstein, and P.C. Bridgman. 1995. Localization of myosin II A and B isoforms in cultured neurons. *J Cell Sci*. 108:3661-70.
- Sasaki, Y., K. Hayashi, T. Shirao, R. Ishikawa, and K. Kohama. 1996. Inhibition by drebrin of the actin-bundling activity of brain fascin, a protein localized in filopodia of growth cones. *J Neurochem*. 66:980-8.
- Sekino, Y., S. Tanaka, K. Hanamura, H. Yamazaki, Y. Sasagawa, Y. Xue, K. Hayashi, and T. Shirao. 2006. Activation of N-methyl-D-aspartate receptor induces a shift of drebrin distribution: disappearance from dendritic spines and appearance in dendritic shafts. *Mol Cell Neurosci*. 31:493-504.
- Shah, K., Lahiri, D.K., 2014. Cdk5 activity in the brain - multiple paths of regulation. *J Cell Sci*. 127, 2391-400.
- Sharma, S., E.E. Grintsevich, M.L. Phillips, E. Reisler, and J.K. Gimzewski. 2011. Atomic force microscopy reveals drebrin induced remodeling of f-actin with subnanometer resolution. *Nano Lett*. 11:825-7.

- Shim, K.S., and G. Lubec. 2002. Drebrin, a dendritic spine protein, is manifold decreased in brains of patients with Alzheimer's disease and Down syndrome. *Neurosci Lett.* 324:209-12.
- Shiraishi-Yamaguchi, Y., Y. Sato, R. Sakai, A. Mizutani, T. Knopfel, N. Mori, K. Mikoshiba, and T. Furuichi. 2009. Interaction of Cupidin/Homer2 with two actin cytoskeletal regulators, Cdc42 small GTPase and Drebrin, in dendritic spines. *BMC Neurosci.* 10:25.
- Shirao, T., and C. Gonzalez-Billault. 2013. Actin filaments and microtubules in dendritic spines. *J Neurochem.* 126:155-64.
- Shirao, T., K. Hayashi, R. Ishikawa, K. Isa, H. Asada, K. Ikeda and K. Uyemura. 1994. Formation of thick, curving bundles of actin by drebrin A expressed in fibroblasts. *Exp. Cell Res.* 215(1): 145-153.
- Shirao, T., N. Kojima and K. Obata. 1992. Cloning of drebrin A and induction of neurite-like processes in drebrin-transfected cells. *Neuroreport* 3(1): 109-112.
- Stiber, J.A., J.H. Wu, L. Zhang, I. Nepliouev, Z.S. Zhang, V.G. Bryson, L. Brian, R.C. Bentley, P.R. Gordon-Weeks, P.B. Rosenberg, and N.J. Freedman. 2016. The Actin-Binding Protein Drebrin Inhibits Neointimal Hyperplasia. *Arterioscler Thromb Vasc Biol.* 36:984-93.
- Sweet, E.S., M.L. Previtera, J.R. Fernandez, E.I. Charych, C.Y. Tseng, M. Kwon, V. Starovoytov, J.Q. Zheng, B.L. Firestein. 2011. PSD-95 alters microtubule dynamics via an association with EB3, *J Neurosci.* 31:1038-1047.
- Takahashi, H., H. Yamazaki, K. Hanamura, Y. Sekino, and T. Shirao. 2009. Activity of the AMPA receptor regulates drebrin stabilization in dendritic spine morphogenesis. *J Cell Sci.* 122:1211-9.
- Trivedi, N., Cain, B., Stabley, D., Ramahi, J., Wong, D., Temirov, J., Kerekes, R. A., Gordon-Weeks, P.R. & Solecki, D. J. 2016. Drebrin mediated microtubule-actomyosin coupling steers two-stroke nucleokinesis and migration pathway selection in cerebellar granule neurons. *Nat. Commun.*, (in press).
- van de Willige, D., C.C. Hoogenraad, and A. Akhmanova. 2016. Microtubule plus-end tracking proteins in neuronal development. *Cell Mol Life Sci.* 73:2053-77.
- Weitzdoerfer, R., M. Dierssen, M. Fountoulakis, and G. Lubec. 2001. Fetal life in Down syndrome starts with normal neuronal density but impaired dendritic spines and synaptosomal structure. *J Neural Transm Suppl.* 59-70.
- Worth, D.C., C.N. Daly, S. Geraldo, F. Oozeer, and P.R. Gordon-Weeks. 2013. Drebrin contains a cryptic F-actin-bundling activity regulated by Cdk5 phosphorylation. *J Cell Biol.* 202:793-806.
- Xu, W., and M. Stamnes. 2006. The actin-depolymerizing factor homology and charged/helical domains of drebrin and mAbp1 direct membrane binding and localization via distinct interactions with actin. *J Biol Chem.* 281:11826-33.
- Yamazaki, H., N. Kojima, K. Kato, E. Hirose, T. Iwasaki, T. Mizui, H. Takahashi, K. Hanamura, R.T. Roppongi, N. Koibuchi, Y. Sekino, N. Mori, and T. Shirao. 2014. Spikar, a novel drebrin-binding protein, regulates the formation and stabilization of dendritic spines. *J Neurochem.* 128:507-22.
- Zhao, L., Q.L. Ma, F. Calon, M.E. Harris-White, F. Yang, G.P. Lim, T. Morihara, O.J. Ubeda, S. Ambegaokar, J.E. Hansen, R.H. Weisbart, B. Teter, S.A. Frautschy, and G.M. Cole. 2006. Role of p21-activated kinase pathway defects in the cognitive deficits of Alzheimer disease. *Nat Neurosci.* 9:234-42.

Figure legends

Figure 1. **Model of a molecular mechanism for coupling of F-actin to microtubules by the drebrin/EB3/Cdk5 pathway in filopodia.**

The diagram shows the domain structure of drebrin in the “closed” conformation when the BB domain binds at the border between the ADFH and CC domains where S142 is located (1). In this conformation drebrin can bind to actin filaments but does not bundle or straddle pre-existing actin filament bundles. One filopodial actin filament, represented by a black line, is also shown. The location of the 46 amino acid insert that distinguishes the drebrin A isoform from drebrin E is also shown (A insert). Activation of Cdk5 by intracellular signalling pathways driven by extrinsic guidance cues leads to S142 phosphorylation (2) and the consequent relief of the intramolecular occlusion of one actin filament binding domain by the BB domain. This produces an “open” conformation and drebrin now bundles actin filaments, or straddles pre-existing actin filament bundles, by co-operative binding of the CC and Hel domains. Drebrin in the “open” conformation and bound to actin filaments at the base of filopodia is now in an appropriate state to interact with the +TIP protein EB3 located at the plus end of a microtubule (purple line) entering the filopodium, and thereby couple actin filaments to a microtubule (3). ©Worth *et al.* 2013. Modified from and originally published in *J. Cell Biol.*, doi: 10.1083/jcb.201303005.

Table 1
Drebrin interactors

Interactor	Function	Reference
afadin	Adherens junction protein	Rehm et al., 2013
Arp3	Junction dynamics during spermatogenesis. Component of the F-actin nucleating and branching Arp2/3 complex	Li <i>et al.</i> , 2011
cupidin/homer 2	Post-synaptic density scaffolding protein	Shiraishi-Yamaguchi et al., 2009
connexin-43	Gap junction protein	Butkevich et al., 2004
CXCR4	Chemokine receptor for the chemokine CXCL12	Pérez-Martínez et al., 2010; Gordón-Alonso et al., 2013
EB3	+TIP protein regulating microtubule dynamics	Geraldo et al., 2008; Bazellières et al., 2012
myosin IIB	Actin-binding motor protein	Bazellières et al., 2012
profilin	Actin monomer-binding protein	Mammoto et al., 1998
PTEN	Lipid phosphatase, negative regulator of PI3K signalling pathway, tumour suppressor	Kreis et al., 2013
Siah-2	E3 ubiquitin ligase	Trivedi et al., 2016
spikar	transcriptional co-activator, dendritic spine density	Yamazaki et al., 2013

Table 2

EB3 interactors not shared with EB1

Interactor	Function	Reference
drebrin	Stabilises F-actin, couples F-actin to microtubules	Geraldo et al., 2008
Aurora A and B	Mitotic kinases, phosphorylate EB2 and EB3 but not EB1	Ban et al., 2009
calcineurin	Ca ²⁺ -dependent phosphatase	Komarova et al., 2012
DDA3	microtubule-associated protein, activator of β -catenin pathway	Hsieh et al., 2007
inositol 1,4,5-trisphosphate receptor	inositol trisphosphate activated Ca ²⁺ channel in endoplasmic reticulum	Geyer et al., 2015
PSD-95	Post-synaptic density scaffolding protein	Sweet et al., 2011
Siah-1	E3 ubiquitin ligase	Ban et al., 2009