King's Research Portal

DOI:

[10.1007/s10803-015-2361-5](https://doi.org/10.1007/s10803-015-2361-5)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Salazar, F., Baird, G., Chandler, S., Tseng, E., O'sullivan, T., Howlin, P., Pickles, A., & Simonoff, E. (2015). Co-occurring Psychiatric Disorders in Preschool and Elementary School-Aged Children with Autism Spectrum Disorder. *Journal of Autism and Developmental Disorders*, 45(8), 2283-2294. <https://doi.org/10.1007/s10803-015-2361-5>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Journal of Autism and Developmental Disorders

Co-occurring Psychiatric Disorders in Preschool and Elementary School-aged Children with Autism Spectrum Disorder

--Manuscript Draft--

Manuscript Number:	JADD-D-14-00406R2
Full Title:	Co-occurring Psychiatric Disorders in Preschool and Elementary School-aged Children with Autism Spectrum Disorder
Article Type:	Article
Keywords:	autism, autism spectrum disorder, psychopathology, child behaviour problems, prevalence
Corresponding Author:	Emily Simonoff Institute of Psychiatry London, UNITED KINGDOM
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Institute of Psychiatry
Corresponding Author's Secondary Institution:	
First Author:	Fernando Salazar, MBBS
First Author Secondary Information:	
Order of Authors:	Fernando Salazar, MBBS
	Gillian Baird, MB BChir, FRCPCH
	Susie Chandler, PhD
	Evelin Tseng, FRCPCH
	Tony O'sullivan, F.R.C.Paed., FRCPCH, DCH
	Patricia Howlin, BA, MSc, PhD, FBPS
	Andrew Pickles, M.A., M.Sc., Ph.D. FMedSci
	Emily Simonoff
Order of Authors Secondary Information:	
Abstract:	We employed a clinical sample of young children with ASD, with and without intellectual disability, to determine the rate and type of psychiatric disorders and possible association with risk factors. We assessed 101 children (57 males, 44 females) aged 4.5 to 9.8 years. 90.5% of the sample met the criteria. Most common diagnoses were: generalized anxiety disorder (66.5%), specific phobias (52.7%) and attention deficit hyperactivity disorder (59.1%). Boys were more likely to have oppositional defiant disorder (OR 3.9). Higher IQ was associated with anxiety disorders (OR 2.9) and older age with agoraphobia (OR 5.8). Night terrors was associated with parental psychological distress (OR 14.2). Most young ASD children met the criteria for additional psychopathology.
Response to Reviewers:	<p>Dear Dr Volkmar</p> <p>We appreciate the opportunity to respond to these additional comments from our reviewers. These are summarized below.</p> <p>Reviewer #1:</p> <p>The authors have made substantial and specific improvements to the manuscript based on the reviewers' comments, and it is much improved. With a few minor corrections, it will be ready for publication in this reviewer's opinion. Specifically:</p>

1. Abstract: the abstract does not mention the instrument under study, please add the PAPA to line 4.
This has been done.

2. the 90.5% sentence on line 5 can be dropped, it has no meaning since we don't know the criteria.
This has been done.

3. the final sentence should use the conventional term, "children with ASD" rather than "ASD children".
This has been done.

4. There remain multiple grammatical errors/missing words in the manuscript. use of a professional copy-editor is advised prior to resubmission.
The manuscript has been carefully re-read for grammatical and other errors and these have been modified.

5. final point: in the discussion section, where the rate of ODD is mentioned, as the authors have thought deeply about these issues, could you add a sentence giving your view on how one might think about ODD in an individual with ASD? many specialists in asd do not consider it a valid diagnosis in many instances, as individuals with asd often lack the intentionality and understanding of their actions seen in neurotypical individuals with ODD.
We have added a further discussion of this on page 16, lines 3-7. In many ways, this gets to the core of a number of issues as they pertain to additional psychopathology in people with ASD. This is an important issue. Psychiatric diagnoses are based on behavioural observations and symptom reports, and there may be many different ways of getting to the same final common pathway. However, the ODD diagnosis as implemented in PAPA/CAPA does not require intentionality, e.g., to harm or distress others, and there is no doubt that the behaviour of children with ASD may have this effect. Similarly, children without ASD may show symptoms of DDD without intending to distress others.

The previous revision substantially extended elements of this discussion, including, for example, the first few paragraphs of the Discussion.

Reviewer #2: Thanks for your revisions.

I still feel the authors haven't addressed my point that DSM-IV specifically stated that ADHD could not be diagnosed if PDD (including autism and Aspergers) was. Yet they are using cases diagnosed with DSM-IV ADHD as evidence of comorbidity (i.e. they are living in a DSM-5 world made out of DSM-IV components). That's fine, but I think that they need to state that is the case - i.e. explain the controversy around this and the continuing controversy despite DSM-5 allowing for comorbidity. A simple point by way of explanation - e.g. the ADHD rating scale gives as one of its symptoms 'blurts out answers before questions have been completed'. If a child with ASD does this excessively is that a symptom of ASD or ADHD? DSM-IV says its the former (presumably) - does DSM-5 make it any clearer in terms of what is actually going on in the child? No. It is even less conclusive. It could be either due to ASD or ADHD or both in DSM-5 world. Its the lack of specificity around symptoms that concerns me and the likelihood (especially in USA) that a child with ASD diagnosed with ADHD will automatically be put on stimulant medication. This study suggests a rate of 59% of ASD children having comorbid ADHD. I'm concerned that these findings will be used as evidence for more children with ASD being diagnosed with a condition (ADHD) which is so simplistic in concept that almost any child with ASD will fit its diagnostic criteria, especially seeing as ADHD is diagnosed mainly by parent / teacher report. Is that really telling us what ASD is? I don't think so. I'd appreciate it if the authors could place a bit more emphasis on the complexity and continuing nature of this debate.

This point overlaps with the final point raised by Reviewer 1, of whether (and how) diagnostic criteria can be applied to the symptoms expressed in children with ASD. As indicated above, we agree this is a critical question that will require other study designs (clinical trials, biomarker studies) to address in more depth. The implementation of ADHD criteria outside the usual DSM-IV rules has been specified on page 9, lines 18-20.
We agree about the complexity of distinguishing true additional psychopathology from

autistic behaviours (see response to reviewer 1) and indeed we are writing a Review on the topic. This applies not just to ADHD but many diagnoses. In our previous revision, we made a specific reference to this on page 13, lines 17-19. We have added a further sentence to the end of the paragraph, "DSM-5 has removed this exclusion, taking an agnostic approach to the reasons for overlap, consistent with our approach in this study. However, it should be noted that ADHD medication may be less effective in children with ASD." We hope this addresses the concern about inappropriate use of medication.

We hope these additional revisions address all the reviewers' comments and we once again thank them for their careful consideration of this manuscript.

Yours sincerely

Co-occurring Psychiatric Disorders in Preschool and Elementary School-aged Children with Autism Spectrum Disorder

Fernando Salazar, M.R.C.Psych.

South London and Maudsley NHS Foundation Trust, Assessment, Liaison & Outreach Team, Woodland House, Cranbrook Road, Staplehurst, Kent TN12 0ER, United Kingdom

Gillian Baird, MB BChir, FRCPCH

Guy's & St Thomas' NHS Foundation Trust, St Thomas' Hospital, Westminster Bridge Road, London SE1 7EH, United Kingdom

Susie Chandler, PhD

Department of Psychology, King's College London, Institute of Psychiatry, De Crespigny Park, Denmark Hill, London SE5 8AF, United Kingdom

Evelin Tseng, FRCPCH

Consultant Community Paediatrician, Bromley Healthcare CIC, 1st Floor, Beckenham Beacon, 379 Croydon Road, Beckenham, Kent, BR3 3QL, United Kingdom

Tony O'sullivan, F.R.C.Paed., FRCPCH, DCH

Consultant community paediatrician Lewisham Healthcare, Children First Lewisham c/o Diane Bishop at Kaleidoscope, 32 Rushey Green, Catford, London SE64JF

Patricia Howlin, BA, MSc, PhD, FBPS

Department of Psychology, King's College London, Institute of Psychiatry, De Crespigny Park, Denmark Hill, London SE5 8AF, United Kingdom

Title page with all author contact information; Abstract with key words and corresponding author email

Andrew Pickles, M.A., M.Sc., Ph.D. FMedSci

Department of Biostatistics and NIHR Biomedical Research Centre for Mental Health, Institute of Psychiatry, De Crespigny Park, Denmark Hill, London SE5 8AF, United Kingdom

Emily Simonoff, M.D., F.R.C.Psych.

Department of Child and Adolescent Psychiatry and NIHR Biomedical Research Centre for Mental Health, Institute of Psychiatry, De Crespigny Park, Denmark Hill, London SE5 8AF,

Running head: Psychiatric Disorders in Elementary School Children with ASD

United Kingdom

ABSTRACT (word count: 114/120)

We employed a clinical sample of young children with ASD, with and without intellectual disability, to determine the rate and type of psychiatric disorders and possible association with risk factors. We assessed 101 children (57 males, 44 females) aged 4.5 to 9.8 years. 90.5% of the sample met the criteria. Most common diagnoses were: generalized anxiety disorder (66.5%), specific phobias (52.7%) and attention deficit hyperactivity disorder (59.1%). Boys were more likely to have oppositional defiant disorder (OR 3.9). Higher IQ was associated with anxiety disorders (OR 2.9) and older age with agoraphobia (OR 5.8). Night terrors was associated with parental psychological distress (OR 14.2). Most young ASD children met the criteria for additional psychopathology.

Keywords: autism, autism spectrum disorder, psychopathology, child behaviour problems, prevalence

Correspondence concerning this article should be addressed to:

Emily Simonoff

Professor of Child & Adolescent Psychiatry and Head of Department

Email: Emily.simonoff@kcl.ac.uk

Tel (00 44) (0) 20 7848 5312

Fax (00 44) (0) 20 7708 5600

Running head: Psychiatric Disorders in Elementary School Children with ASD

1 Co-occurring Psychiatric Disorders in Preschool and Elementary School-aged Children with

2 Autism Spectrum Disorder

3

4

ABSTRACT (word count: 127/120)

We employed a clinical sample of young children with ASD, with and without intellectual disability, to determine the rate and type of psychiatric disorders and their association with a range of possible risk factors. We assessed 101 children (57 males, 44 females) aged 4.5 to 9.8 years with the Preschool-age Psychiatric Assessment. The overwhelming majority of the sample had one or more DSM-IV psychiatric disorders. The most common diagnoses were: generalized anxiety disorder (66.5%), specific phobias (52.7%) and attention deficit hyperactivity disorder (59.1%). Boys were more likely to have oppositional defiant disorder (OR 3.9). Higher IQ was associated with anxiety disorders (OR 2.9) and older age with agoraphobia (OR 5.8). Night terrors were associated with parental psychological distress (OR 14.2). These findings have implications for the management of young children with ASD.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2 **Keywords:** autism, autism spectrum disorder, psychopathology, child behaviour problems,
3 prevalence

1 INTRODUCTION

2 Autism spectrum disorder (ASD) is now known to occur in more than 1% of the
3 population (Baio, 2012; Baird et al., 2006). Coupled with life-course persistence and
4 significant psychosocial impairment, ASD causes a high levels of social and economic
5 burden (Knapp, Romeo, & Beecham, 2009). Recent studies have demonstrated that additional
6 psychiatric disorders commonly occur in children with ASD with many studies reporting
7 aggregated rates as high as 70-90% (de Bruin, Ferdinand, Meester, de Nijs, & Verheij, 2007;
8 Leyfer et al., 2006; Mattila et al., 2010; Mukaddes & Fateh, 2010; Simonoff et al., 2008).
9 These rates are four to six times higher than those reported in the general population
10 (Costello, Mustillo, Erkanli, Keeler, & Angold, 2003). They are also higher than in those
11 found in many studies of children with intellectual disability (ID), e.g., (Dekker & Koot,
12 2003). In ASD, common co-occurring disorders in late childhood include attention deficit
13 hyperactivity disorder (ADHD), oppositional defiant disorder (ODD) and anxiety disorders,
14 with emergence of depression and obsessive compulsive disorder in adolescence/adult life
15 (Mazefsky et al., 2008; Simonoff et al., 2008; Skokauskas, Gallagher, Skokauskas, &
16 Gallagher, 2010; van Steensel, Bogels, & de Bruin, 2013).

17 While there is a substantial literature in older children, adolescents and adults
18 describing the prevalence and correlates of co-occurring psychiatric disorders in ASD, the
19 topic has received less attention among preschool and elementary schoolchildren. However,
20 diagnostic tools are now available for this age group, including questionnaires such as the 1.5
21 to 5 years old version of the Child Behavior Checklist (Ivanova et al., 2010), the Infant
22 Toddler Social-Emotional Assessment (Carter, Briggs-Gowan, Jones, & Little, 2003), and
23 diagnostic interviews such as the Preschool Age Psychiatric Assessment (PAPA) (Helen Link
24 Egger et al., 2006). Furthermore, recent studies support the use of current diagnostic concepts

1 of psychopathology in younger children. Egger and Angold found similar architecture for
2 common psychopathology when comparing preschool and older children and adolescents
3 (Angold & Egger, 2007; S. Sterba, Egger, & Angold, 2007). Lecavalier used factor analysis
4 and demonstrated the validity of DSM-IV syndromes in three to five year-old children with
5 ASD (Lecavalier, Gadow, Devincent, Houts, & Edwards, 2011).

6 There have been two approaches to evaluating psychopathology in ASD: first,
7 standardized measures designed for and evaluated in the general population and, second,
8 bespoke instruments in which the mode of inquiry and/or criteria for symptom endorsement
9 have been specifically modified for use in ASD populations. An advantage of the former
10 approach is the ability to make direct comparisons with the general population, both about
11 prevalence rates and risk/protective factors. However, researchers have raised a number of
12 concerns about the suitability of standardized measures. When used directly with people with
13 ASD rather than informants, their understanding and ability to provide accurate responses
14 should be considered; typically developing younger children will give inaccurate responses
15 rather than volunteer they do not understand questions (Breton et al., 1995). Furthermore,
16 subtle distinctions between core symptoms of autism and those related to other
17 psychopathology may not be well-differentiated in instruments designed for the general
18 population. (Leyfer et al., 2006) highlight that their Autism Comorbidity Interview (ACI-PL),
19 based on the Kiddie-SADS, includes probes and definitions designed to make such
20 distinctions, e.g., between lack of social interest and social avoidance. Similarly, (Mazefsky
21 et al., 2008) showed that using a diagnostic interview with modified criteria reduced the rate
22 of additional psychopathology in ASD.

23 We chose the PAPA in the present study because we thought it provided the best
24 compromise between the two methods. As only parents are interviewed in this age group,

1 there is no need to consider whether probes are understood by participants. There remains a
2 broader concern about access to the mental state of younger children with poorer
3 communication skills, but this applies to all interview measures. The PAPA was designed for
4 use with parents of younger children with limited verbal skills and therefore relies more
5 heavily on behavioral examples. We are unaware of any published studies using the PAPA in
6 populations with autism but have applied its sister instrument, the Child and Adolescent
7 Psychiatric Assessment (CAPA) in parent interviews of older children with ASD (Simonoff
8 et al., 2008). The availability of general population norms for the PAPA allows direct
9 comparison to findings in the general population, which is not possible with autism-specific
10 measures such as the ACI-PL or BISCUIT (Matson, Boisjoli, Hess, & Wilkins, 2011).

11 In this study we describe the prevalence of parent-reported DSM-IV disorders in a
12 clinical sample of children with ASD aged five to nine years. We examine the role of several
13 well-recognized risk factors for psychopathology in the general population, including sex, ID,
14 parental psychological distress, family socioeconomic position and ethnic minority status.
15 We also explore the relationship of psychopathology to the child's age and autism severity.

16 17 **METHODS**

18 *Participants*

19 The eligible population for the study was all children born between September 2000
20 and August 2004 (four to eight years at recruitment) with an ASD diagnosis and living in
21 either of two London boroughs (Bromley and Lewisham, outer and inner London
22 respectively). The Social Communication Questionnaire Lifetime version [SCQ; (Rutter,
23 Bailey, & Lord, 2003)] was used to measure ASD symptoms. Clinical diagnosis was

1 established following a multidisciplinary neurodevelopmental and social communication
2 assessment led by a single community pediatrician for each borough. The teams used multi-
3 source information (parents, teachers, social workers), observation of the child (at clinic or at
4 home and or school) and structured assessments such as the Autism Diagnostic Interview-
5 Revised (Le Couteur et al., 1989), the Developmental, Dimensional and Diagnostic Interview
6 (Skuse et al., 2004), the Diagnostic Interview for Social and Communication Disorders
7 (Wing, Leekam, Libby, Gould, & Larcombe, 2002) and the Autism Diagnostic Observation
8 Schedule (Lord et al., 2000). All cases who, in the subsequent research assessment, had an
9 SCQ score below 10 (n=7) were considered by the lead pediatrician who reviewed the
10 clinical notes to confirm the ASD diagnosis.

11 The primary care services and local autism support groups identified 447 eligible
12 children fulfilling the criteria; this number represents 1.38% of the estimated population in
13 2010 for this age and geographical area (Large & Ghosh, 2006).

14 *Procedure*

15 In the first phase of the study all eligible families were mailed information about the
16 study, including an invitation to participate, consent form and a return envelope. Non-
17 responders were subsequently contacted by phone. Responses were received from 362 (81%)
18 (see Figure 1). Of these, 277 parents (62%) completed the Developmental Behavior Checklist
19 (Einfeld & Tonge, 2002), the Profile of Neuropsychiatric Symptoms (Santosh, 2006) and the
20 Social Communication Questionnaire, lifetime version, as well as questionnaire on family
21 demographic characteristics, described below. A cognitive assessment was undertaken,
22 usually in the child's school, and teachers were asked to complete parallel behavior
23 questionnaires.

24 *Insert Figure 1 about here*

The second phase of the study aimed to assess psychopathology in more detail. A sample of 131 children was selected from the 277 participants. All participating girls (n=50) were invited, in order to provide a sufficiently large female sample for comparisons according to sex. A stratified random sample of boys (N=81) was selected to include equal numbers of boys on each of the following characteristics (1) IQ ($<70/\geq 70$); (2) residing in the two London boroughs; (3) child's age (4.5 to 6.7/6.8 to 9.9 years) and (4) SCQ ($<21/\geq 22$). The mean time gap between the two phases was 373 days (SD 135, range 64-697).

Insert figure 1 about here

Measures

Child characteristics

Autism symptom severity was determined using the parent-reported SCQ, lifetime version; the published cut-off of ≥ 22 was used to split the sample into two groups.

IQ was measured in 258 cases by two study psychologists trained in the cognitive assessment of children with ASD and using one or more of the following tests, depending on the child's age and developmental level: the Mullen Scales of Early Learning [MSEL; (Mullen, 1997)], the Wechsler Preschool and Primary Scale of Intelligence [WPPSI-III-UK; (Wechsler, 2004)], and the Wechsler Intelligence Scale for Children [WISC-IV-UK (Wechsler, Rust, & Golombok, 2004)]. The British Picture vocabulary Scales were used to assess receptive language in 240 children (Dunn, Dunn, Whetton, & Burley, 1997). Parents were also asked to provide an estimate of their child's 'functional age' (see Supplementary Appendix for details of questions). In 197 cases where both direct measure and estimate were available, the mean test-based IQ was 72.8 (SD = 27.4), parent-derived estimate 68.9 (SD=23.2) with a correlation of 0.71 ($p<0.001$). Therefore, in 15 cases where tested IQ was

not available, the parent-derived estimates were substituted; in 4 cases neither measure was available.

DSM-IV symptoms and disorders present in the last three months were assessed using the electronic version of the Preschool Age Psychiatric Assessment (PAPA), a semi-structured, parent-reported interview for preschool children aged 2 to 5 years (H.L. Egger & Angold, 2004). Interviews with the main caregiver were conducted in person, in most cases in the family home, and audio-taped. The majority of interviews were conducted by a specialist trainee (equivalent to clinical fellow) in Child and Adolescent Psychiatry (FS, N=88 interviews) and the rest by two research psychologists with specialist autism experience. All interviewers attended a two day training course on the PAPA. Validation of interview administration and coding was performed by the trainer (from Duke University) at an early stage in the study. Subsequently, specific coding issues were discussed on an *ad hoc* basis both with the US trainer and ES (who is author of the sister instrument, the CAPA). Weekly meetings with the principal investigator (GB) reviewed any specific questions. These included decisions about whether items met PAPA criteria for additional psychopathology or were more appropriately considered symptoms of ASD. The detailed PAPA symptom criteria were used rigorously but in an agnostic fashion to endorse individual symptoms. Standardized algorithms previously developed and reported were used to determine diagnoses (Helen Link Egger et al., 2006). In the case of ADHD, the criterion excluding the diagnosis in the presence of ASD was waived.

Parental characteristics

The Kessler Psychological Distress Scale (K-10) was used to measure parental psychological distress (Kessler et al., 2002). The K-10 includes questions about cognitive, behavioral, emotional and psychophysiological symptoms that are elevated among people

with a wide range of different mental disorders. The 10-item questionnaire has good internal consistency; a score ≥ 24 indicate nonspecific psychological distress and was the cut-off for a binary variable. Parental education was dichotomized according to whether either parent had attained GCSEs (equivalent to a high school diploma) or higher. Families were characterized according to whether there was at least one parent who was employed; ethnicity was classified as white or other.

The study was approved by Guy's Hospital Research Ethics Committee (approval number 08/H0804/37).

Data Analysis

Data reduction and analysis used Stata 11 (StataCorp, 2009). Inverse probability weights were generated to account for the study design, in which all females were included (and therefore assigned a probability weight of 1) while boys were selected according to IQ, residence, age and autism severity. All the prevalence rates and other statistics presented in this paper reflect the overall ASD population from which the sample was drawn. Wald test statistics (adjusted t and F tests) and p values were calculated using the linearization version of the robust parameter covariance matrix as implemented by the *svy* procedures of Stata 11. Logistic regression was used for binary outcomes.

RESULTS

A total of 101 families (77% of the invited group) took part in the interview stage. This included 57 boys and 44 girls with a mean age of 6.7 years. Table 1 summarizes the sample characteristics. Comparison between the participating and non-participating children showed that participating children had lower SCQ scores but otherwise were similar in terms

of child and family characteristics. On the basis of measured IQ (and not including adaptive function criteria), 56% had IQs in the normal range, 13% mild, 10% moderate ID, 15% severe and 6% profound ID.

Insert Table 1 about here

Prevalence

Table 2 shows the weighted three-month prevalence rates for DSM-IV psychiatric disorders overall and stratified by child characteristics. It reveals that 90.5% [95% confidence intervals (CIs) 84.2, 96.7%] of the sample had at least one psychiatric disorder. The prevalence of emotional disorders was 80.0% (CIs 72.0, 88.1%) and of behavioral disorders 28.7% (95% CIs 18.2, 39.2%). The most common emotional disorders were GAD [66.5% (95% CIs 57.0, 76.0%)], followed by specific phobias [52.7% (95% CIs 40.2, 65.2%)]. Social anxiety occurred in 15.1% (95% CIs 6.2, 24.0%) and somewhat surprisingly, agoraphobia in 18.0% (95% CIs 9.2, 26.7%). Major depressive disorder was present in 14.6% (95% CIs 6.0, 23.2%). ADHD occurred in 59.1% (95% CIs 47.3, 70.9%). ODD was present in 28.7% (95% CIs 18.2, 39.2%) while conduct disorder was present in only 2.0% (95% CIs 0.6, 1%), all of whom also met criteria for ODD.

Insert Table 2 about here

Aggregation among psychiatric disorders

The pattern of overlap amongst ADHD, ODD/CD and emotional disorders is shown in Figure 2. Both ODD and ADHD rarely occurred in isolation from other disorders, in contrast to emotional disorders which presented without ADHD or ODD in 26.5% of cases (weighted). In terms of the *number* of co-occurring disorders, 11.5% of the sample had one, 28.5% had two and 51.4% three or more disorders. Multiple anxiety disorders frequently co-

1 occurred; 75.6% of cases that had any anxiety disorder met criteria for at least two diagnoses,
2 with GAD and specific phobia the most common combination, occurring in 24.3%. In
3 evaluating the number of co-occurring disorders overall, when anxiety disorders are collapsed
4 to a single diagnostic entity, 24.9% of the sample had one diagnosis, 31.8% two and 34.2%
5 three or more diagnoses.

6 *Insert Figure 2 about here*

7 *Effects of child characteristics*

8 Male sex was a risk factor for ADHD (OR 2.9, 95% CIs 1.2, 6.9), ODD (OR 3.9,
9 95% CIs 1.3, 11.8) and tic disorder (OR 5.5, 95% CIs 1.1, 27.6). Higher IQ was associated
10 with anxiety disorders as an aggregate group (referred to as *any anxiety disorder*) (OR 2.9,
11 95% CIs 1.0, 8.1) but not with any specific one and not with other disorders. Older children
12 were more likely to have anxiety disorders including GAD (OR 4.5, 95% CIs 1.7, 11.8),
13 separation anxiety (OR 5.5, 95% CIs 1.5, 20.8) and agoraphobia (OR 5.8, CIs 1.6, 21.3). As a
14 *post hoc* analysis, we explored the relationship of any anxiety disorder to the presence of
15 language, defined by the first question on the SCQ (the child's ability to string a few words
16 in a meaningful sentence) and found a non-significant trend for the presence of language to
17 increase the probability of any anxiety disorder (OR 2.7, 95% CIs 0.9, 8.5, $p=.08$).

18 Participants with more autism symptoms were more likely to have diagnoses of
19 agoraphobia (OR 5.4, 95% CIs 1.3, 22.7), ODD (OR 3.5, 95% CIs 1.1, 10.6) and night terrors
20 (OR 7.5, 95% CIs 1.3, 42.1).

21 *Effects of parental characteristics*

22 Table 3 shows weighted rates of disorders by family characteristics. Night terrors
23 were associated with both parental psychological distress (OR 14.2, 95% CIs 2.4, 84.6) and

parental unemployment (OR 8.9, 95% CIs 1.4, 56.0). Lower parental educational level was associated with higher rates of any emotional disorder (OR 8.7, 95% CIs 1.7, 43.0) and ODD (OR 4.5, CIs 1.3, 15.3) but with a lower rate of enuresis (OR 0.2, CIs 0, 1.0).

Insert Table 3 about here

DISCUSSION

This study is currently amongst the few to describe the prevalence and associated risk factors for psychiatric disorders among younger children with ASD. Virtually all (90.5%) met criteria for at least one DSM-IV disorder. The aggregate rate is substantially higher than that for a similar age group in the general population, where rates of 12-16% have been reported (H. L. Egger & Angold, 2006; Wichstrom et al., 2012). Furthermore, rates among children with ID, albeit somewhat older, are reported to be 20-40%, substantially lower than those in our present study (Dekker & Koot, 2003). However the present findings are broadly in line with those reported in children and adolescents with ASD (de Bruin et al., 2007; Leyfer et al., 2006; Mattila et al., 2010; Mukaddes & Fateh, 2010; Simonoff et al., 2008). These high rates raise questions about diagnostic classification, its application in people with ASD, and the extent to which core autism symptoms may overlap with other psychopathology. Broadly there are three alternative explanations. First, the autism spectrum phenotype may include symptoms that are usually considered part of another disorder, such as difficulties with sustained attention or social anxiety. It was on this basis that DSM-IV excluded the diagnosis of ADHD in the presence of ASD; DSM-5 has removed this exclusion, taking an agnostic approach to the reasons for overlap, consistent with our approach in this study. However, it should be noted that ADHD medication may be less effective in children with ASD (Harfterkamp et al., 2012; RUPP Autism Network, 2005; Simonoff, et al., 2013).

Second, ASD, or characteristics associated with it, may increase the risk of other symptoms/disorders. Hence, we have shown that cognitive rigidity in ASD is associated with anxiety (Hollocks et al., 2014). People with ASD experience higher rates of bullying (Montes, Halterman, Montes, & Halterman, 2007; Rowley et al., 2012) and other adverse life events (H. Green, McGinity, Meltzer, Ford, & Goodman, 2005) which may also increase their susceptibility to additional psychopathology. Finally, structured instruments for the general population may inadvertently miscode autistic symptoms to other domains of psychopathology. Thus, (Mazefsky et al., 2012) demonstrated that using an interview in which symptom definitions are modified to account for ASD presentation led to a very substantial reduction in rates. In the present study, PAPA interviews were undertaken by researchers with extensive experience of ASD who were trained not to misattribute ASD symptoms to other psychopathology, but nevertheless high rates of psychopathology are reported. Other methods will be required to test these alternative models, including risk factor studies and clinical trials.

With respect to emotional disorders, we found anxiety disorders were very common both individually and in association with other disorders, with rates higher than those reported in older age ASD groups (Simonoff et al., 2008; Witwer & Lecavalier, 2010). For the overall anxiety disorder category, as well as for some individual disorders, we found an association with higher IQ and older age. Studies in older ASD populations vary, with some reporting a similar relationship with higher IQ, others finding no association or one to lower IQ (Amr, Raddad, El-Mehesh, Mahmoud, & El-Gilany, 2011; K. D. Gadow, Devincent, Pomeroy, & Azizian, 2005; Simonoff et al., 2008; Sukhodolsky et al., 2008). In this study with younger children we speculate that more developed language, partly indexed by IQ, supports the communication of worries, so that parents are more aware of these emotions.

1 Alternatively, higher IQ, as well as older age, may expose children to more anxiety-
2 provoking situations including greater interaction with peers. Finally, higher IQ/
3 developmental level may allow them to engage in higher-order cognitions, such as worries
4 about past events, the future, self-efficacy, which are the hallmarks of anxiety disorders such
5 as GAD. We note that separation anxiety, which largely involves observable behaviors, was
6 associated with older age but not higher IQ. In this case, older children may have had more
7 experiences of separation, including school attendance, which may have led to greater
8 anxiety.

9 The rate of social phobia (15.1%) is similar to that reported in late childhood by
10 (Simonoff et al., 2008) but higher than that of others (Leyfer et al., 2006; Mattila et al., 2010;
11 Mukaddes & Fateh, 2010). It is uncertain at present whether this represents variation among
12 populations, differences between instruments or chance variation. In our study interviewers
13 were trained to distinguish social anxiety and avoidance from lack of social interest without
14 evidence of anxiety.

15 The high rate of agoraphobia (18.0%) is also noteworthy. Parental accounts indicated
16 that children exhibited anticipatory anxiety and/or avoidance behavior in settings such as
17 public transport or noisy shops. One explanation is that abnormal responses to high levels of
18 sensory stimuli may be causative but this needs further exploration (Goldsmith, Van Hulle,
19 Arneson, Schreiber, & Gernsbacher, 2006; S. A. Green, Ben-Sasson, Green, & Ben-Sasson,
20 2010).

21 Finally the rate of depression in our study is higher than in the general population of
22 the same age, with general population studies reporting rates of 2.1% and 2.0% respectively
23 (Egger & Angold, 2006; Wichstrom et al., 2012).

1 Our rate of ADHD, 59.1%, and its association with male sex, is consistent with other
2 studies (de Bruin et al., 2007; Mukaddes et al., 2010; Witwer, Lecavalier, & Norris, 2012)
3 although slightly higher than others (Leyfer, et al., 2006; Simonoff et al., 2008). The rate of
4 ODD (28.7%) is strikingly similar to what we reported using the CAPA with older children
5 as well as that in other studies (Leyfer et al., 2006; Simonoff et al., 2008; Witwer et al.,
6 2012). The low rate of CD is expected and consistent with reports in older children
7 (Simonoff, Jones, et al., 2013; Simonoff et al., 2008). We made diagnoses of ODD and CD
8 irrespective of the child's intention to cause distress or harm to others. While this is in line
9 with the diagnostic algorithms, many clinicians distinguish children on the basis of whether
10 there is 'intentionality' in their behaviour. However, a lack of intention to cause distress may
11 not be limited to those with ASD, e.g., children with ADHD. This further highlights that
12 'behavior that challenges' or 'challenging behaviour,' which is common in ASD, does not
13 conform to the structure of CD and should be considered separately (McClintock, Hall, &
14 Oliver, 2003).

15 With respect to tic disorders, the literature reports highly variable rates and ours lie in
16 the middle; the present rate of 17.4% is strikingly similar to the rate of 13.8% that we
17 reported in older children using the CAPA (Simonoff, et al., 2008). Interview reports are not
18 the optimal method for identification as, on the one hand, informants may not have noticed
19 tics and, on the other, may confuse them with stereotypies and mannerisms in ASD. Finally,
20 despite inherent methodological difficulties in the study of sleep disorders, we report rates
21 for night terrors that are similar to previous studies for this age group of children with ASD
22 (Krakowiak, Goodlin-Jones, Hertz-Picciotto, Croen, & Hansen, 2008; Richdale & Schreck,
23 2009; Williams, Sears, & Allard, 2004). The reported association between ASD severity and

parental psychological distress has been reported elsewhere (Doo & Wing, 2006; Mayes & Calhoun, 2009) .

Co-occurrence amongst additional psychiatric disorders

The finding that most children with ASD meet criteria for multiple disorders fits with previous studies (de Bruin et al., 2007; Gjevik, Eldevik, Fjaeran-Granum, & Sponheim, 2011; Leyfer et al., 2006; Mattila et al., 2010; Mukaddes & Fateh, 2010; Mukaddes et al., 2010; Simonoff et al., 2008). The extent of co-occurrence is higher than that reported in the general population. This may in part reflect issues about classification (Rutter, 2011; S. K. Sterba et al., 2010). We also found high levels of co-occurrence among the three main domains of ADHD, ODD/CD and emotional disorders.

Risk factors and correlates of psychiatric disorders

This is one of the first studies to include a sufficient number of girls with ASD to evaluate sex differences. The only differences found were higher rates of ADHD and ODD in boys, findings mirroring those among typically developing children (H. L. Egger & Angold, 2006). There is a relative absence of associations with age, bar those already described with respect to anxiety, but we note the limited age range of the present sample.

As with our previous studies we found that ID, a well-known risk factor for most psychiatric disorders in the general child population, was *not* associated with most additional disorders in this sample (Charman et al., 2011; Eisenhower, Baker, & Blacher, 2005; Simonoff et al., 2008). With respect to family characteristics, a similar lack of association to that reported here has been found in previous studies indicating that these factors may be less important in ASD than in typically developing children, possibly due to different causal pathways (Kenneth D. Gadow, Devincent, & Schneider, 2008; Gjevik et al., 2011; Simonoff

et al., 2008). We failed to find associations between parental psychological distress and psychiatric disorders; this finding is at odds with several other studies (Davis & Carter, 2008; Hastings, 2003; Tehee, Honan, & Hevey, 2009) but in line with our previous report in later childhood, where psychopathology was unrelated to parenting stress or maternal psychological distress (Simonoff et al., 2008). Many of the previous studies are of small sample sizes. Further, measures of ‘parenting stress’ often include items indexing characteristics of the parent-child relationship, which may be influenced by ASD symptoms and other psychopathology. In the present study, we measured parental psychological distress, rather than parenting stress, which may partially account for the differences. One large study demonstrated a link between parental psychopathology/psychological distress and the presence of ASD in contrast to typical development (Totsika, Hastings, Emerson, Lancaster, & Berridge, 2011) and an earlier meta-analysis demonstrated considerable inconsistency among studies but an overall relationship with parental psychopathology that was moderated by the child’s level of functioning (Yirmiya & Shaked, 2005). More research is needed to disentangle these relationships and understand the link to additional psychopathology as opposed to ASD characteristics.

Strengths and limitations

Strengths of this study include its careful sampling method and the large proportion of females. To our knowledge, this is the first study of psychopathology in ASD to over-sample females allowing a better comparison across the sexes. This study uses community-based clinics providing universal diagnostic services and hence includes all children with an ASD diagnosis in the sampling frame. There are also several limitations. Children had clinical rather than research diagnoses of ASD. Although there were few differences between those participating and declining, the non-participating children had higher SCQ scores. Because

1 SCQ scores were associated with social disadvantage, the sample may under-represent
2 families with higher levels of psychosocial adversity as well as those with most severe autism
3 symptoms. Another limitation is the use of an instrument, the PAPA, that has not been
4 validated in populations with either ASD or ID. However, we believe that the careful
5 methodology allows our findings to be compared to general population samples. As with a
6 number of other studies, the use of parent-only assessments, such as the PAPA, limits the
7 way in which ‘pervasiveness’ is defined for diagnoses such as ADHD. However, because the
8 ADHD diagnostic algorithm enquires parents to indicate knowledge of the child’s behaviour
9 in other settings (such as school), it is not clear whether this will increase or decrease the
10 reported prevalence. We note that parents of children in this age group are usually closely
11 involved in their children’s activities outside the home and are likely to be able to report on
12 their behaviour in these other settings. This is a moderately sized sample, but because of the
13 exploration of moderating factors, some of the associations we report have wide confidence
14 intervals and need to be interpreted with caution.

15 In summary, our study shows that the high rates of psychiatric disorders in ASD
16 previously reported for older children and adolescents also occur in younger children. Many
17 of these disorders have evidence-based interventions, at least for typically developing
18 children. Therefore, clinical implications include the need to systematically assess for
19 additional psychopathology at an early stage when children are diagnosed with ASD, as
20 recommended in the UK (NICE, 2011). There is also a need for validated screening and
21 diagnostic tools for psychopathology for this population and this should be a high priority for
22 future research. Finally, in the absence of effective treatment for the core symptoms of ASD,
23 and with the knowledge that additional psychopathology persists in ASD, it is essential to

1 1 develop an evidence base for treatment of co-occurring psychopathology (Simonoff, Jones, et
2
3 2 al., 2013) .
4
5 3
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

- Amr, M., Raddad, D., El-Mehesh, F., Mahmoud, E.-H., & El-Gilany, A.-H. (2011). Sex differences in Arab children with autism spectrum disorders. *Research in Autism Spectrum Disorders*, 5(4), 1343-1350.
- Angold, A., & Egger, H. L. (2007). Preschool psychopathology: lessons for the lifespan. *Journal of Child Psychology & Psychiatry & Allied Disciplines*, 48(10), 961-966.
- Baio, J. (2012). Prevalence of autism spectrum disorders: autism and developmental disabilities monitoring network, 14 sites, United States, 2008. Morbidity and Mortality Weekly Report, Surveillance Summaries. *Centers for Disease Control and Prevention*, 61(3), 1-19.
- Baird, G., Simonoff, E., Pickles, A., Chandler, S., Loucas, T., Meldrum, D., et al. (2006). Prevalence of disorders of the autism spectrum in a population cohort of children in South East Thames - The Special Needs and Autism Project. *The Lancet* 368 (9531), 210-215.
- Breton, J. J., Bergeron, L., Valla, J. P., Lepine, S., Houde, L., & Gaudet, N. (1995). Do children aged 9 through 11 years understand the DISC Version 2.25 questions? *Journal of the American Academy of Child and Adolescent Psychiatry*, 34(7), 946-954.
- Carter, A. S., Briggs-Gowan, M. J., Jones, S. M., & Little, T. D. (2003). The infant-toddler social and emotional assessment (ITSEA): Factor structure, reliability, and validity. *Journal of Abnormal Child Psychology*, 31(5), 495-514.
- Charman, T., Pickles, A., Simonoff, E., Chandler, S., Loucas, T., & Baird, G. (2011). IQ in children with autism spectrum disorders: Data from the SNAP project. *Psychological Medicine* 41(3), 619-627.

- 1 Costello, E. J., Mustillo, S., Erkanli, A., Keeler, G., & Angold, A. (2003). Prevalence and
2 development of psychiatric disorders in childhood and adolescence. *Archives of*
3 *General Psychiatry*, 60(8), 837-844.
- 4 Davis, N. O., & Carter, A. S. (2008). Parenting stress in mothers and fathers of toddlers with
5 autism spectrum disorders: associations with child characteristics. *Journal of Autism*
6 *& Developmental Disorders*, 38(7), 1278-1291.
- 7 de Bruin, E., Ferdinand, R. F., Meester, S., de Nijs, P. F., & Verheij, F. (2007). High rates of
8 psychiatric co-morbidity in PDD-NOS. *Journal of autism and developmental*
9 *disorders*, 37(5), 877-886.
- 10 Dekker, M. C., & Koot, H. M. (2003). DSM-IV disorders in children with borderline to
11 moderate intellectual disability I: Prevalence and impact. *Journal of American*
12 *Academy of Child and Adolescent Psychiatry*, 42(8), 915-922.
- 13 Doo, S., & Wing, Y. K. (2006). Sleep problems of children with pervasive developmental
14 disorders: Correlation with parental stress. *Developmental Medicine and Child*
15 *Neurology*, 48(8), 650-655.
- 16 Dunn, L. M., Dunn, L. M., Whetton, C., & Burley, J. (1997). *British Picture Vocabulary*
17 *Scales* (Second ed.). Windsor: NFER-Nelson Publishing Company.
- 18 Egger, H. L., & Angold, A. (2004). The Preschool Age Psychiatric Assessment (PAPA): A
19 structured parent interview for diagnosing psychiatric disorders in preschool children.
20 *Handbook of infant, toddler, and preschool mental health assessment*, 223-243.
- 21 Egger, H. L., & Angold, A. (2006). Common emotional and behavioral disorders in preschool
22 children: presentation, nosology, and epidemiology. *Journal of Child Psychology and*
23 *Psychiatry, and Allied Disciplines*, 47(3-4), 313-337.

- 1 Egger, H. L., Erkanli, A., Keeler, G., Potts, E., Walter, B. K., & Angold, A. (2006). Test-
2 retest reliability of the Preschool Age Psychiatric Assessment (PAPA). *Journal of the*
3 *American Academy of Child & Adolescent Psychiatry*, 45(5), 538-549.
- 4 Einfeld, S. L., & Tonge, B. J. (2002). *Manual for the Developmental Behaviour Checklist*
5 (Second ed.). Sidney: University of New South Wales and Monash University.
- 6 Eisenhower, A. S., Baker, B. L., & Blacher, J. (2005). Preschool children with intellectual
7 disability: syndrome specificity, behaviour problems, and maternal well-being.
8 *Journal of Intellectual Disability Research.* , 49(9), 657-671.
- 9 Gadow, K. D., Devincent, C., & Schneider, J. (2008). Predictors of psychiatric symptoms in
10 children with an autism spectrum disorder. *Journal of Autism & Developmental*
11 *Disorders*, 38(9), 1710-1720.
- 12 Gadow, K. D., Devincent, C. J., Pomeroy, J., & Azizian, A. (2005). Comparison of DSM-IV
13 symptoms in elementary school-age children with PDD versus clinic and community
14 samples. *Autism*, 9, 392-415.
- 15 Gjevik, E., Eldevik, S., Fjaeran-Granum, T., & Sponheim, E. (2011). Kiddie-SADS reveals
16 high rates of DSM-IV disorders in children and adolescents with autism spectrum
17 disorders. *Journal of Autism & Developmental Disorders*, 41(6), 761-769.
- 18 Goldsmith, H., Van Hulle, C. A., Arneson, C., Schreiber, J., & Gernsbacher, M. (2006). A
19 population-based twin study of parentally reported tactile and auditory defensiveness
20 in young children. *Journal of Abnormal Child Psychology*, 34(3), 378-392.
- 21 Green, H., McGinley, A., Meltzer, H., Ford, T., & Goodman, R. (2005). *Mental health of*
22 *children and young people in Great Britain, 2004*. Basingstoke: Palgrave Macmillan.
- 23 Green, S. A., Ben-Sasson, A., Green, S. A., & Ben-Sasson, A. (2010). Anxiety disorders and
24 sensory over-responsivity in children with autism spectrum disorders: is there a causal
25 relationship? *Journal of Autism & Developmental Disorders*, 40(12), 1495-1504.

- 1 Harfterkamp, M., van de Loo-Neus, G., Minderaa, R. B., van der Gaag, R. J., Escobar, R.,
2 Schacht, A., et al. (2012). A randomized double-blind study of atomoxetine versus
3 placebo for attention-deficit/hyperactivity disorder symptoms in children with autism
4 spectrum disorder. *Journal of the American Academy of Child & Adolescent*
5 *Psychiatry*, 51(7), 733-741.
- 6 Hastings, R. P. (2003). Child behaviour problems and partner mental health as correlates of
7 stress in mothers and fathers of children with autism. *Journal of Intellectual Disability*
8 *Research* 47(pt 4-5), 231-237.
- 9 Hollocks, M. J., Jones, C. R. G., Pickles, A., Baird, G., Happé, F., Charman, T., et al. (2014).
10 The association between social cognition and executive functioning and symptoms of
11 anxiety and depression in adolescents with autism spectrum disorders. *Autism*
12 *Research*, 7(2), 216-228.
- 13 Ivanova, M. Y., Achenbach, T. M., Rescorla, L. A., Harder, V. S., Ang, R. P., Bilenberg, N.,
14 et al. (2010). Preschool psychopathology reported by parents in 23 societies: testing
15 the seven-syndrome model of the child behavior checklist for ages 1.5-5.
16 [Comparative Study]. *Journal of the American Academy of Child & Adolescent*
17 *Psychiatry*, 49(12), 1215-1224.
- 18 Kessler, R. C., Andrews, G., Colpe, L. J., Hiripi, E., Mroczek, D. K., Normand, S.-L., et al.
19 (2002). Short screening scales to monitor population prevalences and trends in non-
20 specific psychological distress. *Psychological Medicine*, 32, 959-976.
- 21 Knapp, M., Romeo, R., & Beecham, J. (2009). Economic cost of autism in the UK. *Autism*,
22 13(3), 317-336.
- 23 Krakowiak, P., Goodlin-Jones, B., Hertz-Picciotto, I., Croen, L. A., & Hansen, R. L. (2008).
24 Sleep problems in children with autism spectrum disorders, developmental delays,

- and typical development: a population-based study. *Journal of Sleep Research*, 17(2), 197-206.
- Large, P., & Ghosh, K. (2006). A methodology for estimating the population by ethnic group for areas within England. *Population Trends*, 123, 21.
- Le Couteur, A., Rutter, M., Lord, C., Rios, P., Robertson, S., Holdgrafer, M., et al. (1989). Autism Diagnostic Interview: A standardized investigator-based instrument. *Journal of Autism and Developmental Disorders*, 19(3), 363-387.
- Lecavalier, L., Gadow, K. D., Devincent, C. J., Houts, C. R., & Edwards, M. C. (2011). Validity of DSM-IV syndromes in preschoolers with autism spectrum disorders. *Autism*, 15(3), 1-17.
- Leyfer, O. T., Folstein, S. E., Bacalman, S., Davis, N. O., Dinh, E., Morgan, J., et al. (2006). Comorbid psychiatric disorders in children with autism: Interview development and rates of disorders. *Journal of Autism and Developmental Disorders*, 36(7), 849-861.
- Lord, C., Risi, S., Lambrecht, L., Cook, E. H., Leventhal, B. L., DiLavore, P. C., et al. (2000). The Autism Diagnostic Observation Schedule-Generic: A standard measure of social and communication deficits associated with the spectrum of autism. *Journal of Autism and Developmental Disorders*, 30, 205-223.
- Matson, J. L., Boisjoli, J. A., Hess, J. A., & Wilkins, J. (2011). Comorbid psychopathology factor structure on the Baby and Infant Screen for Children with aUtism Traits-Part 2 (BISCUIT-Part 2). *Research in Autism Spectrum Disorders*, 5(1), 426-432.
- Mattila, M. L., Hurtig, T., Haapsamo, H., Jussila, K., Kuusikko-Gauffin, S., Kielinen, M., et al. (2010). Comorbid psychiatric disorders associated with Asperger syndrome/high-functioning autism: a community- and clinic-based study. *Journal of Autism & Developmental Disorders*, 40(9), 1080-1093.

- 1 Mayes, S. D., & Calhoun, S. L. (2009). Variables related to sleep problems in children with
2 autism. *Research in Autism Spectrum Disorders*, 3(4), 931-941.
- 3 Mazefsky, C. A., Folstein, S. E., Lainhart, J. E., Mazefsky, C. A., Folstein, S. E., & Lainhart,
4 J. E. (2008). Overrepresentation of mood and anxiety disorders in adults with autism
5 and their first-degree relatives: what does it mean? *Autism research : Official Journal*
6 *of the International Society for Autism Research*, 1(3), 193-197.
- 7 Mazefsky, C. A., Oswald, D. P., Day, T. N., Eack, S. M., Minshew, N. J., & Lainhart, J. E.
8 (2012). ASD, a psychiatric disorder, or both? Psychiatric diagnoses in adolescents
9 with high-functioning ASD. *Journal of Clinical Child & Adolescent Psychology*,
10 41(4), 516-523.
- 11 McClintock, K., Hall, S., & Oliver, C. (2003). Risk markers associated with challenging
12 behaviours in people with intellectual disabilities: a meta-analytic study. *Journal of*
13 *Intellectual Disability Research*, 47(6), 405-416.
- 14 Montes, G., Halterman, J. S., Montes, G., & Halterman, J. S. (2007). Bullying among
15 children with autism and the influence of comorbidity with ADHD: a population-
16 based study. *Ambulatory Pediatrics*, 7(3), 253-257.
- 17 Mukaddes, N. M., & Fateh, R. (2010). High rates of psychiatric co-morbidity in individuals
18 with Asperger's disorder. *World Journal of Biological Psychiatry*, 11(2_2), 486-492.
- 19 Mukaddes, N. M., Herguner, S., Tanidir, C., Mukaddes, N. M., Herguner, S., & Tanidir, C.
20 (2010). Psychiatric disorders in individuals with high-functioning autism and
21 Asperger's disorder: similarities and differences. *World Journal of Biological*
22 *Psychiatry*, 11(8), 964-971.
- 23 Mullen, E. M. (1997). *Mullen Scales of Early Learning*. Los Angeles, CA: Western
24 Psychological Services.

- 1 National Institute for Clinical Excellence (2011). *Recognition, referral and diagnosis of*
2 *children and young people on the autism spectrum.*
- 3 Research Units on Pediatric Psychopharmacology (RUPP) Autism Network (2005).
4 Randomized, controlled, crossover trial of methylphenidate in pervasive
5 developmental disorders with hyperactivity. *Archives of General Psychiatry*, 62(11),
6 1266-1274.
- 7 Richdale, A. L., & Schreck, K. A. (2009). Sleep problems in autism spectrum disorders:
8 Prevalence, nature, & possible biopsychosocial aetiologies. *Sleep Medicine Reviews*,
9 13(6), 403-411.
- 10 Rowley, E., Chandler, S., Bair, G., Simonoff, E., Pickles, A., Loucas, T., et al. (2012). The
11 experience of friendship, victimisation and bullying in children with an autism
12 spectrum disorder: Associations with child characteristics and school placement.
13 *Research in Autism Spectrum Disorders*, 6(3), 1126-1134.
- 14 Rutter, M. (2011). Research review: Child psychiatric diagnosis and classification: concepts,
15 findings, challenges and potential. *Journal of Child Psychology & Psychiatry &*
16 *Allied Disciplines*, 52(6), 647-660.
- 17 Rutter, M., Bailey, A., & Lord, C. (2003). *The Social Communication Questionnaire* (1st
18 ed.). Los Angeles: Western Psychological Services.
- 19 Santosh, P. J. (2006). Profile of Neuropsychiatric Symptoms.
- 20 Simonoff, E., Jones, C., Baird, G., Pickles, A., Happe, F., & Charman, T. (2013). The
21 persistence and stability of psychiatric problems in adolescents with autism spectrum
22 disorders. *Journal of Child Psychology & Psychiatry*, 54(2), 186-194.
- 23 Simonoff, E., Pickles, A., Charman, T., Chandler, S., Loucas, T., & Baird, G. (2008).
24 Psychiatric disorders in children with autism spectrum disorders: prevalence,

comorbidity and associated factors. *Journal of the American Academy of Child and Adolescent Psychiatry*, 47(8), 921-929.

Simonoff, E., Taylor, E., Baird, G., Bernard, S., Chadwick, O., Liang, H., et al. (2013). Randomized controlled double-blind trial of optimal dose methylphenidate in children and adolescents with severe attention deficit hyperactivity disorder and intellectual disability. *Journal of Child Psychology and Psychiatry*, 54(5), 527-535.

Skokauskas, N., Gallagher, L., Skokauskas, N., & Gallagher, L. (2010). Psychosis, affective disorders and anxiety in autistic spectrum disorder: prevalence and nosological considerations. *Psychopathology*, 43(1), 8-16.

Skuse, D., Warrington, R., Bishop, D., Chowdhury, U., Lau, J., Mandy, W., et al. (2004). The developmental, dimensional and diagnostic interview (3di): a novel computerized assessment for autism spectrum disorders. *Journal of the American Academy of Child and Adolescent Psychiatry*, 43(5), 548-558.

StataCorp. (2009). *Stata Statistical Software Release 11.1*. College Station, TX: Stata Corporation.

Sterba, S., Egger, H. L., & Angold, A. (2007). Diagnostic specificity and nonspecificity in the dimensions of preschool psychopathology. *Journal of Child Psychology & Psychiatry & Allied Disciplines*, 48(10), 1005-1013.

Sterba, S. K., Copeland, W., Egger, H. L., Jane Costello, E., Erkanli, A., & Angold, A. (2010). Longitudinal dimensionality of adolescent psychopathology: testing the differentiation hypothesis. *Journal of Child Psychology & Psychiatry & Allied Disciplines*, 51(8), 871-884.

Sukhodolsky, D. G., Scahill, L., Gadow, K. D., Arnold, L. E., Aman, M. G., McDougle, C. J., et al. (2008). Parent-rated anxiety symptoms in children with pervasive developmental

- disorders: frequency and association with core autism symptoms and cognitive functioning. *Journal of Abnormal Child Psychology*, 36(1), 117-128.
- Tehee, E., Honan, R., & Hevey, D. (2009). Factors contributing to stress in parents of individuals with autistic spectrum disorders. *Journal of Applied Research in Intellectual Disabilities* 22, 34-42.
- Totsika, V., Hastings, R. P., Emerson, E., Lancaster, G. A., & Berridge, D. M. (2011). A population-based investigation of behavioural and emotional problems and maternal mental health: associations with autism spectrum disorder and intellectual disability. *Journal of Child Psychology & Psychiatry*, 52(1), 91-99.
- van Steensel, F. J., Bogels, S. M., & de Bruin, E. I. (2013). Psychiatric comorbidity in children with autism spectrum disorders: a comparison with children with ADHD. *Journal of Child and Family Studies*, 22, 368-376.
- Wechsler, D. (2004). *Wechsler Preschool and Primary School Intelligence Scale* (Third ed.). London: Harcourt Assessment.
- Wechsler, D., Rust, J., & Golombok, S. (2004). *Wechsler Intelligence Scale for Children* ® - Fourth UK Edition (WISC-IV UK). London: Harcourt Assessment.
- Wichstrom, L., Berg-Nielsen, T. S., Angold, A., Egger, H. L., Solheim, E., & Sveen, T. H. (2012). Prevalence of psychiatric disorders in preschoolers. *Journal of Child Psychology & Psychiatry*, 53(6), 695-705.
- Williams, P. G., Sears, L. L., & Allard, A. (2004). Sleep problems in children with autism. *Journal of Sleep Research*, 13(3), 265-268.
- Wing, L., Leekam, S. R., Libby, S. J., Gould, J., & Larcombe, M. (2002). The Diagnostic Interview for Social and Communication Disorders: Background, inter-rater reliability and clinical use. *Journal of Child Psychology and Psychiatry*, 43(3), 307-325.

Witwer, A. N., & Lecavalier, L. (2010). Validity of comorbid psychiatric disorders in youngsters with autism spectrum disorders. *Journal of Developmental and Physical Disabilities*, 22(4), 367-380.

Witwer, A. N., Lecavalier, L., & Norris, M. (2012). Reliability and validity of the children's interview for psychiatric syndromes--parent version in autism spectrum disorders. *Journal of Autism & Developmental Disorders*, 42(9), 1949-1958.

Yirmiya, N., & Shaked, M. (2005). Psychiatric disorders in parents of children with autism: a meta-analysis. *Journal of Child Psychology and Psychiatry*, 46(1), 69-83.

Figure Caption Sheet

Figure 1. Sample selection and response rates

Figure 2. Pattern of overlap co-morbidities amongst children^a

Table 1. Comparison between characteristics of children and parents invited to in-depth stage

Table 2. Weighted Rates (95% confidence intervals) of Disorders by Child Characteristics

Table 3. Weighted Rates (95% CIs) of Disorders by Family Characteristics

Figure 1 Sample selection and response rates

Figure 2: Pattern of overlap co-morbidities amongst children^a

^aEmotional disorder= Depression + anxiety disorders

Table 1
Comparison between characteristics of children and parents invited to in-depth stage

	<u>Geographical area</u>			<u>Comparison of participated vs declined</u>		
	Lewisham	Bromley	p	Declined	Participated	p
Total Number (females)	87 (36)	44 (14)		30 (6)	101 (44)	
Mean Age at stage 1	6.7	6.8	0.59	6.7	6.7	0.81
(SD, range)	(1.2, 4.5-9.3)	(1.1, 4.7-9.1)		(1.1, 4.9-9.0)	(1.1, 4.5-9.3)	
Mean IQ	61.5	77.8	0.001	69.1	66.4	0.65
(SD, range)	(28.0, 19.0-114.0)	(25.1, 19.0-120.0)		(30.4, 19.0-114)	(28.0, 19.0-120)	
BPVS ^b	82.0 (15.1, 40-112)	93.8 (18.4, 40-120)	0.004	85.8 (16.6, 41-106)	86.1 (17.2, 40 -120)	.95
(SD, range)						
SCQ	19.6	23.7	0.002	24.3	20.0	0.006
(SD, range)	(7.2, 3-42)	(6.9, 6-36)		(7.3, 9.0-42.0)	(7.0, 3.0-34)	
Parental ethnicity white N	27 (31)	37 (84)	0.000	12 (40)	52 (51.5)	0.27

(%)

Parent Employed N (%)	53 (64.6) ^a	34 (77.2)	0.14	15 (60) ^a	73 (73.3)	0.27
Parent GCSEs or higher N	45 (54.9) ^a	24 (54.5)	0.97	13 (52) ^a	56 (55.4)	0.76

(%)

SCQ=Social Communication Questionnaire; BPVS=British Picture Vocabulary Scales; GCSEs are equivalent to a high school diploma

^a 5 missing values ^b 27 missing values, 7 because of inability to access test

Table 2
Weighted Rates (95% confidence intervals) of Disorders by Child Characteristics

DSM-IV	Total	<u>Sex</u>			<u>IQ</u>			<u>Age at assessment</u>			<u>Autism Severity</u>		
Disorders	Sample	Females	Males	Odds	IQ<70	IQ>= 70	Odds	<=7.5y	>7.5	Odds	SCQ<=21	SCQ>21	Odds
	N=101	N=44	N=57	Ratio	N=57	N=44	Ratio	N=47	N=54	Ratio	N=62	N=39	Ratio
Any DSM	90.5%	81.8%	92.7%	2.8	87.3%	92.1%	1.7	89.5%	91.3%	1.2	88.0%	94.5%	2.4
Disorder	(84.2-96.7)	(70.1-93.5)	(85.0-100.0)	(0.7-10.7)	(77.3-97.3)	(84.1-100.0)	(0.4-7.0)	(77.8-100.0)	(85.3-97.3)	(0.3-5.3)	(78.9-97.1)	(87.5-100.0)	(0.5-11.7)
Any DSM	80.0%	68.2%	83.1%	2.3	68.6%	85.9%	2.8	70.1%	88.7%	3.3*	78.8%	82.0%	1.2
Emo Dis. ^a	(72.0-88.1)	(54.1-82.3)	(73.6-92.6)	(0.9-5.8)	(55.1-82.0)	(75.8-96.1)	(1.0-8.0)	(54.2-86.0)	(82.2-95.1)	(1.2-9.3)	(66.9-90.8)	(73.5-90.5)	(0.5-3.0)
Any Anxiety	78.9%	65.9%	82.2%	2.4	66.5%	85.2%	2.9*	69.2%	87.4%	3.0*	78.8%	79.0%	1.0
Disorder	(70.7-87.0)	(51.5-80.3)	(72.6-91.8)	(0.9-6.0)	(52.6-80.4)	(75.0-95.4)	(1.0-8.1)	(53.2-85.1)	(80.5-94.3)	(1.1-8.5)	(66.9-90.8)	(69.7-88.2)	(0.4-2.5)
Agoraphobia	18.0%	13.6%	19.1%	1.5	10.3%	21.9%	2.5	6.4%	28.1%	5.8**	8.4%	33.4%	5.4*
	(9.2-26.7)	(3.2-10.0)	(8.4-29.8)	(0.5-4.6)	(3.0-17.5)	(9.2-40.1)	(0.8-7.1)	(1.7-12.6)	(12.5-43.7)	(1.6-21.3)	(0.0-17.7)	(16.1-50.7)	(1.3-22.7)

		24.0)	29.8)			34.6)							
Specific	52.7%	50.0%	53.4%	1.1	45.6	56.4	1.6	62.0%	44.6%	0.5	53.4%	51.5%	0.9
Phobia	(40.2-65.2)	(34.8-65.1)	(38.2-68.6)	(0.5-2.7)	(29.4-61.5)	(39.4-73.4)	(0.6-4.0)	(44.3-80.0)	(27.3-61.9)	(0.2-1.4)	(37.0-69.8)	(32.5-70.6)	(0.3-2.5)
Panic	3.1%	2.3%	3.4%	1.5	0%	4.7%	-	0%	5.9%	-	1.8%	5.3%	3.1
Disorder	(0.0-7.5)	(0-6.78.0)	(1.2-8.7)	(0.1-20.4)		(0.0-11.3)			(0.0-14.0)		(0.0-4.3)	(0.0-16.0)	(0.2-39.8)
Social	15.1%	11.4%	16.0%	1.5	5.3%	20.1%	4.5	8.0%	21.2%	3.0	12.9%	18.7%	1.6
Phobia	(6.2-24.0)	(1.7-21.0)	(5.1-26.9)	(0.4-5.2)	(0.0-13.7)	(7.4-32.7)	(0.7-28.0)	(0.1-16.0)	(6.0-36.0)	(0.7-12.7)	(2.4-23.3)	(2.7-34.7)	(0.4-6.3)
Generalized	66.5%	59.1%	68.4%	1.5	53.5%	73.2%	2.4	49.3%	81.5%	4.5**	66.2%	67.0%	1.0
Anxiety Dis.	(57.0-76.0)	(44.2-74.0)	(57.1-79.7)	(0.7-3.4)	(38.4-68.5)	(60.8-85.5)	(1.0-5.8)	(33.4-65.2)	(71.0-92.0)	(1.7-11.8)	(53.7-78.8)	(52.6-81.4)	(0.4-2.5)
Separation	18.6%	15.9%	19.3%	1.3	9.9%	23.0%	2.7	6.8%	28.8%	5.5**	20.0%	16.2%	0.8
Anxiety	(9.4-27.8)	(4.8-27.0)	(8.1-30.4)	(0.4-3.8)	(0.1-19.8)	(10.4-35.6)	(0.7-9.9)	(0.0-13.7)	(12.8-44.8)	(1.5-20.8)	(8.3-31.9)	(1.5-30.8)	(0.2-2.8)

Major	14.6%	4.5%	17.2%	4.3	6.3%	18.8%	3.4	7.8%	20.6%	3.0	15.4%	13.3%	0.8
Depression	(6.0-23.2)	(0.0-10.9)	(6.4-27.9)	(0.8-22.4)	(0.0-18.1)	(6.2-31.5)	(0.4-32.7)	(0.0-17.5)	(5.3-35.8)	(0.5-17.3)	(4.8-26.0)	(0.0-28.1)	(0.2-3.9)
Oppositional	28.7%	11.4%	33.1%	3.9*	19.5%	33.3%	2.0	26.3%	30.7%	1.2	18.8%	44.7%	3.5*
Defiant Dis.	(18.2-39.2)	(1.8-21.0)	(20.2-46.1)	(1.3-11.8)	(3.9-35.1)	(19.7-47.1)	(0.6-6.5)	(10.7-42.0)	(16.6-44.8)	(0.4-3.5)	(5.2-32.4)	(28.3-61.1)	(1.1-10.6)
Conduct	2.0%	0%	2.6%	-	0%	3.1%	-	0%	3.8%	-	0%	5.3%	-
Disorder	(0.0-6.1)		(0.0-7.6)			(0.0-9.2)			(0.0-11.4)			(0.0-16.0)	
ADHD	59.1%	38.6%	64.4%	2.9*	60.7%	58.3%	0.9	56.6%	61.3%	1.2	53.5%	68.2%	1.9
	(47.3-70.9)	(23.9-53.4)	(50.0-78.7)	(1.2-6.9)	(44.1-77.4)	(42.6-74.0)	(0.3-2.3)	(39.1-74.1)	(45.6-76.9)	(0.5-3.2)	(37.0-70.0)	(52.6-83.9)	(0.7-5.0)
Motor &	17.4%	4.5%	20.7%	5.5*	16.6%	17.8%	1.1	17.3%	17.5%	1.0	10.9%	28.0%	3.1
Voc Tic Dis.	(8.1-26.7)	(0.0-10.9)	(9.2-32.3)	(1.1-27.6)	(1.4-31.8)	(6.2-29.5)	(0.3-4.2)	(5.0-29.6)	(3.8-31.3)	(0.3-3.7)	(0.6-21.2)	(10.3-45.7)	(0.8-12.7)
Encopresis	1.9%	4.5%	1.3%	0.3	2.7%	1.5%	0.6	4.2%	0%	-	0.8%	3.9%	5.3
	(0.4-4.3)	(0.0-)	(0.0-3.8)	(0.0-3.3)	(0.0-6.5)	(0.0-4.6)	(0.0-6.6)	(0.0-9.3)			(0.0-2.2)	(0.0-9.7)	(0.4-67.4)

		10.8)											
Enuresis	13.5%	16.0%	13.0%	0.8	22.2%	9.0%	0.3	20.7%	7.2%	0.3	11.8%	16.2%	1.4
	(5.5-21.5)	(0.0-	(0.0-0.2)	(0.2-2.6)	(6.5-38.0)	(1.2-	(0.0-1.2)	(7.8-33.7)	(0.0-15.8)	(0.0-1.3)	(2.3-21.4)	(2.2-30.3)	(0.4-5.7)
		0.27)				16.9)							
Night	4.9%	6.8%	4.4%	0.6	1.4%	6.7%	5.1	7.9%	2.2%	0.3	1.5%	10.3%	7.5*
Terrors	(1.1-8.6)	(0.0-	(0.0-8.7)	(0.1-3.0	(0.0-4.1)	(1.3-	(0.6-	(0.6-15.3)	(0.0-5.3)	(0.0-1.6)	(0.0-3.6)	(1.1-19.5)	(1.3-42.1)
		14.5)				12.0)	45.0)						

^(a)Any Emotional Disorder= Mood Disorder +/- Any Anxiety Disorder; * p<0.05; ** p<0.01

Table 3

Weighted Rates (95% CIs) of Disorders by Family Characteristics

Disorders	K-10			Education			Employment			Ethnicity		
	Low	high	Odds	GCSEs	Below	Odds	Parent	No	Odds	Other	White	Odds
	N= 71	N= 26	Ratio	or above	GCSEs	Ratio	employed	parent	Ratio	N= 49	N= 52	Ratio
				N= 76	N= 25		N= 72	employed				
								N=29				
Any DSM	91.4%	86.5%	0.6	88%	98.1%	7.2	87.3%	98.4%	8.8	90.8%	94.7%	1.8
Disorder	(86.2-	(68.6-	(0.1-	(79.6-	(94.4-	(0.8-	(78.7-	(95.2-	(1.0-	(84.1-	(89.1-	(0.4-
	96.6)	100.0)	3.0)	96.3)	100.0)	65.3)	95.9)	100)	79.0)	97.4)	100.0)	7.6)
Any DSM	81.4%	73.4%	0.6	74.7%	96.3%	8.7**	78.9%	82.8%	1.2	79.4%	84.7%	1.4
Emo. Dis. ^a	(71.9-	(53.4-	(0.2-	(63.9-	(91.0-	(1.7-	(68.8-	(68.7-	(0.4-	(69.0-	(72.7-	(0.4-
	90.9)	93.5)	2.2)	85.5)	100.0)	43.0)	89.1)	96.9)	4.2)	89.6)	96.8)	4.8)
Any Anx	81.4%	70.7%	0.6	73.8%	94.4%	6.0*	78%	81.1%	1.2	77.9%	83.8%	1.5
Disorder	(71.9-	(50.4-	(0.2-	(62.9-	(88.0-	(1.5-	(67.7-	(66.7-	(0.4-	(67.4-	(71.6-	(0.5-
	90.9)	90.9)	1.9)	84.7)	100.0)	23.5)	88.3)	95.6)	3.9)	88.4)	96.0)	4.6)

Formatted Table

Formatted: Left

Agoraphobia	15.1%	26.9%	2.0	16.6%	22.1%	1.4	16.5%	21.5%	1.4	23.7%	13.2%	0.5
	(4.3-	(6.0-	(0.5-	(6.4-	(2.7-	(0.4-	(5.4-	(3.6-	(0.3-	(8.7-	(1.8-24.6)	(0.1-
	25.8)	47.8)	9.1)	26.8)	41.5)	5.6)	27.7)	39.3)	5.7)	38.7)		1.9)
Specific	54.4%	43.1%	0.6	47.2%	69.4%	1.4	50.1%	59.1%	1.4	54.2%	53.8%	1.0
Phobia	(39.4-	(19.9-	(0.2-	(32.5-	(48.6-	(0.4-	(35.3-	(37.6-	(0.5-	(37.3-	(35.5-	(0.4-
	69.4)	66.3)	1.9)	61.9)	90.2)	5.6)	64.9)	80.5)	4.1)	71.2)	72.0)	2.7)
Panic	4.4%	0%	-	4.2%	0.0%	-	3.5%	2.2%	0.6	1.3%	5.1%	4.0
Disorder	(0.0-			(0.0-			(0.0-9.3)	(0.0-6.6)	(0.0-	(0.0-	(0.0-13.4)	(0.3-
	10.3)			9.9)					8.4)	3.9)		56.7)
Social	11.4%	17.6%	1.7	12.5%	22.9%	2.0	15.0%	15.3%	1.0	14.3%	16.6%	1.2
Phobia	(1.6-	(0.0-	(0.3-	(3.0-	(0.2-	(0.4-	(4.3-	(0.0-	(0.2-	(10.9-	(3.1-30.2)	(0.3-
	21.3)	37.0)	8.8)	22.0)	45.6)	10.0)	25.7)	32.6)	5.1)	27.5)		5.4)
GAD	73.0%	47.0%	0.3	63.4%	75.9%	1.8	68.1%	62.5%	0.8	70.8%	65.6%	0.8
	(62.1-	(23.7-	(0.1-	(51.9-	(55.9-	(0.5-	(57.8-	(41.5-	(0.3-	(58.4-	(50.2-	(0.3-
	84.0)	70.3)	1.0)	75.0)	95.8)	6.2)	78.4)	83.5)	2.1)	83.1)	81.0)	2.0)
Separation	18.0%	19.0%	1.0	15.5%	28.0%	2.1	19.0%	17.4%	0.9	19.0%	19.0%	1.0

Anxiety	(7.5-28.4)	(0.0-38.5)	(0.3-4.5)	(5.5-25.5)	(5.0-51.0)	(0.5-8.7)	(8.8-29.3)	(0.2-34.7)	(0.2-3.3)	(4.2-33.9)	(5.6-32.5)	(0.3-3.9)
Major Depression	16.9%	10.1%	0.5	15.3%	12.4%	0.8	17.6%	7.1%	0.4	17.5%	12.5%	0.7
	(6.0-27.8)	(0.0-26.0)	(0.0-4.0)	(5.7-25.0)	(0.0-30.0)	(0.1-4.5)	(6.7-28.5)	(0.0-20.8)	(0.0-3.3)	(3.3-31.6)	(0.2-24.8)	(0.1-3.3)
Oppositional Defiant Dis.	23.0%	40.8%	2.3	20.5%	53.7%	4.5*	25.2%	37.4%	1.7	22.1%	36.6%	2.0
	(9.8-36.2)	(20.8-60.7)	(0.7-7.3)	(8.5-32.4)	(31.0-76.3)	(1.3-15.3)	(11.7-38.6)	(17.7-57.1)	(0.5-5.7)	(7.0-37.2)	(19.9-53.3)	(0.6-6.7)
Conduct Disorder	0%	8.2%	-	0%	8.2%	-	0%	7.1%	-	4.2%	0%	-
		(0.0-23.9)			(0.0-24.0)			(0.0-20.8)		(0.0-12.4)		
ADHD	56.9%	61.6%	1.2	56.2%	68.0%	1.6	56.9%	64.6%	1.4	66.4%	54.9%	0.6
	(42.1-71.8)	(40.7-82.5)	(0.4-3.5)	(41.8-71.0)	(47.4-88.4)	(0.5-5.0)	(42.4-71.5)	(45.3-84.0)	(0.5-3.9)	(50.9-81.9)	(37.0-72.6)	(0.2-1.7)
Motor & Vocal Tic	18.6%	8.3%	0.4	14.0%	28.0%	2.4	21.5%	7.2%	0.3	18.0%	17.6%	1.0
	(7.1-	(0-19.8)	(0.0-	(3.8-	(6.3-	(0.6-	(9.8-	(0.0-	(0.0-	(3.3-	(5.3-29.9)	(0.3-

Dis.	30.0)		2.1)	24.0)	49.8)	9.3)	33.3)	17.1)	1.4)	33.0)		3.6)
Encopresis	0%	5.9%	-	2.6%	0%	-	2.0%	1.6%	0.8	0%	4.0%	-
		(0.0-14.8)		(0.0-5.7)			(0.0-5.2)	(0.0-4.8)	(0.0-10.0)		(0.0-8.9)	
Enuresis	10.2%	21.1%	2.3	16.7%	3.8%	0.2*	10.8%	20.3%	2.1	13.7%	14.0%	1.0
	(2.0-18.5)	(1.1-41.0)	(0.5-10.4)	(6.7-26.7)	(0.0-9.0)	(0.0-10.0)	(2.5-19.1)	(2.1-38.4)	(0.5-8.6)	(1.6-25.7)	(3.2-24.9)	(0.3-3.9)
Night	1.3%	15.7%	14.2**	4.5%	6.0%	1.4	1.6%	12.9%	8.9*	2.4%	7.5%	3.3
Terrors	(0.0-3.1)	(1.5-29.9)	(2.4-84.6)	(0.5-8.5)	(0.0-15.1)	(0.2-8.8)	(0.0-4.0)	(1.1-24.7)	(1.4-56.0)	(0.0-5.8)	(0.5-14.6)	(0.5-20.7)

^(a)Any Emotional Disorder= Mood Disorder +/- Any Anxiety Disorder; * p<0.05; ** p<0.01

Fernando Salazar, South London and Maudsley NHS Foundation Trust, London, UK

Gillian Baird, Guy's & St Thomas' NHS Foundation Trust, London, UK

Susie Chandler, Department of Psychology, King's College London, London, UK

Evelin Tseng, Bromley Healthcare CIC, Kent, UK

Tony O'sullivan, Lewisham Healthcare, Children First Lewisham, London, UK

Patricia Howlin, Department of Psychology, King's College London, London, UK

Andrew Pickles, Department of Biostatistics, King's College London, London, UK

Emily Simonoff, Department of Child and Adolescent Psychiatry, King's College London,
London, UK

No changes in author affiliation subsequent to the time of study

No acknowledgments to report

Correspondence concerning this article should be addressed to:

Emily Simonoff

Professor of Child & Adolescent Psychiatry and Head of Department

Email: Emily.simonoff@kcl.ac.uk

Tel (00 44) (0) 20 7848 5312

Fax (00 44) (0) 20 7708 5600

Supplementary Material

[Click here to download Supplementary Material: Supplementary Appendix.dot](#)