King's Research Portal

DOI:

[10.1016/j.tem.2016.07.002](https://doi.org/10.1016/j.tem.2016.07.002)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Gnudi, L., Coward, R., & Long, D. (2016). Diabetic nephropathy: perspective on novel molecular mechanisms. *TRENDS IN ENDOCRINOLOGY AND METABOLISM*. <https://doi.org/10.1016/j.tem.2016.07.002>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Diabetic nephropathy: perspective on novel molecular mechanisms

Luigi Gnudi,¹ Richard JM Coward,² David A Long.³

¹Cardiovascular Division, King's College London, SE1 9NH, UK;

²Academic Renal Unit, Dorothy Hodgkin Building, University of Bristol, Whitson Street, Bristol BS1 3NY, UK;

³Developmental Biology and Cancer Programme, Institute of Child Health, University College London, WC1N 1EH, UK.

Corresponding Authors:

Luigi Gnudi, e-mail : luigi.gnudi@kcl.ac.uk

David Long, PhD, e-mail : d.long@ucl.ac.uk

Word count: 3617

Keywords: diabetes, glomerulus, oxidative stress, podocytes, endothelium, SGLT2

Abstract

Diabetes is the major cause of end stage renal disease globally, and novel treatments are urgently needed. Current therapeutic approaches for diabetic nephropathy are focusing on the inhibition of the renin angiotensin aldosterone system, on glycaemic and lipid control, and life style changes. In this review we will highlight new molecular insights in our understanding of the initiation and progression of diabetic nephropathy including glomerular insulin resistance, dysregulation of cellular substrate utilisation, podocyte-endothelial communication and inhibition of tubular sodium coupled glucose reabsorption. We believe these mechanisms offer new therapeutic targets that can be exploited to develop important renoprotective treatments for diabetic nephropathy over the next decade.

INTRODUCTION

Diabetes mellitus is a metabolic disorder associated with chronic micro- and macrovascular complications. One of the most feared chronic microvascular complications is diabetic nephropathy (DN), currently the leading cause of end-stage renal disease (ESRD) in the Western world. Strikingly, 40-45% of patients with type-1 diabetes (T1D) develop DN and reach ESRD or die before its onset. Moreover, clinicians face a ~30% prevalence of patients with type-2 diabetes (T2D) and DN, with 45% of patients currently on dialysis having a primary diagnosis of diabetes, a population also at high risk of developing cardiovascular disease [1].

An early sign of DN is an increased amount of urinary protein, manifested by “albuminuria”, which correlates with, and can predict, the progression of renal damage. Albuminuria arises from defects in the permeability of the glomerular filtration barrier consisting of glomerular endothelial cells (GECs) separated from specialized epithelia, called podocytes, by the glomerular basement membrane (GBM)[1]. Podocytes have extensive inter-digitating foot processes connected together by a slit diaphragm composed of proteins including nephrin and neph1, which interact with cytoplasmic adaptor and signalling proteins (PI3-Kinase, CD2AP, AKT, podocin). Nephrin is also linked with the podocyte actin cytoskeleton; the protein tyrosine kinase Fyn promotes nephrin phosphorylation which enhances its interaction with PI3-Kinase and PI3K-dependent phosphorylation of AKT and subsequently increases Rac1 activity, leading to modification of the actin cytoskeleton with maintenance of a normal podocyte anatomical structure and function [2, 3]. The structure and integrity of the glomerulus is also maintained by a complex local autocrine/paracrine network between the podocyte and the GECs consisting of vascular growth factors and vasoactive peptides

which is disrupted in DN [4]. The GECs are highly fenestrated with a unique ultrastructure lacking fenestrae diaphragms which facilitate water and small solutes permeability [5]. GECs are covered by a glycocalyx consisting mainly of proteoglycans which include core proteins such as syndecan and attached glycosaminoglycan side chains which appear to be important in regulating the permeability of the glomerulus [6].

Animal and human studies have established that the metabolic and haemodynamic changes that occur in diabetes lead to ultrastructural alterations of the glomerular filtration barrier, including podocyte foot process fusion and detachment, GBM thickening, a reduced endothelial cell glycocalyx, mesangial extracellular matrix accumulation and glomerulosclerosis (**Figure 1**). These structural glomerular changes correlate with increasing albuminuria which has been proposed to be a marker of generalised systemic vascular dysfunction by the “Steno hypothesis” [7] and could represent a common pathogenetic mechanism for renal and extra-renal chronic vascular complications in diabetes [1].

Over the last 5-10 years our understanding of the molecular and cellular pathways by which diabetic kidney disease results in damage to the glomerular filtration barrier has increased. In this review, we will outline recent advances in glomerular insulin signalling, oxidative and endoplasmic reticulum (ER) stress and podocyte-endothelial communication that have revealed new exciting therapeutic directions for DN.

Insulin resistance as a mechanism for the predisposition of DN

Insulin is a metabolic hormone which not only regulates glucose and the metabolism of other substrates but also directly modulates the biology of specific cells in a variety of tissues. In both T1D and T2D patients, the ability of insulin to elicit cellular responses is impaired, a concept termed “cellular insulin resistance”, and is associated with DN [8]. Insulin resistance correlates with the development of microalbuminuria both in T1D and T2D patients and patients with T1D are more likely to have a strong family history of insulin resistance when compared with those without kidney disease [9]. Insulin resistance has been implicated in the development of glomerular hypertension and hyperfiltration [10], seen in the initial phase of diabetic kidney disease [11]. Furthermore, in both T1D and T2D patients, insulin resistance *per se* contributes to higher salt sensitivity, which closely associates with increases in blood pressure, albuminuria, and a decline in renal function [12, 13].

Within the kidney many different cell types are insulin sensitive and express functional insulin receptors [14-17]. Furthermore, transgenic mouse models have revealed that inducing insulin resistance in different nephron compartments results in a variety of unfavourable renal phenotypes. In the glomerulus, approximately a decade ago it was discovered that human podocytes respond to insulin [14], and express the hallmark components of insulin sensitive cells including the insulin receptor, and key glucose transporters including GLUT4 and GLUT1. To elucidate the biological significance of insulin signalling in these cells, podocyte specific insulin receptor knockout mice were generated [18]. These animals developed albuminuria and a number of features of DN, including increased matrix production, glomerulosclerosis, and GBM thickening, but all in normoglycaemic conditions, suggesting that insulin resistance of this cell *per se* may be an important driver in glomerular diseases. Insulin signaling is important in

other parts of the nephron. Deletion of the insulin receptor in tubular epithelial cells widespread led to reduced natriuresis and hypertension [16]. Recent studies have begun to dissect out the precise function of the insulin receptor in specific tubular segments. These experiments revealed that loss of the insulin receptor in proximal tubules results in gluconeogenesis [19] while deletion in collecting ducts increased natriuresis and lowered blood pressure [20].

Diabetes provides an ideal environment consisting of increased adiposity, hyperglycaemia, and inflammation which are all important players in promoting podocyte insulin resistance and glomerular dysfunction [17](**Table 1**). A recent study has identified SMAD3 within the inflammation/ fibrosis pathway as an important modulator of podocyte insulin sensitivity in a model of obesity related DN [21]. In this work, mice fed a high fat diet exhibited an increase in kidney and podocyte SMAD3 expression levels which resulted in a severely fibrotic kidney; in these conditions SMAD3 knockout animals were protected from kidney damage and fibrosis. In parallel, fatty acid palmitate induces a SMAD3-mediated podocyte insulin resistance paralleled by mitochondrial dysfunction *in vitro*. These responses were exaggerated when animals became albuminuric, and could be rescued by SMAD3 blockade and restoration of podocyte insulin signalling [21]. Other studies have demonstrated that both Nucleotide-binding oligomerization domain containing protein 2 (NOD2)[22] and Toll-like receptor (TLR)[23] mediated-inflammation have an adverse effect on podocyte survival, insulin action, and glomerular permeability to protein. Decreased circulating adiponectin [24, 25], increased free fatty acids (FFA) levels [26], and defects in insulin action promote glomerular cells and podocyte dysfunction, and

albuminuria [27, 28]. Epigenetic mechanisms may also be important in determining insulin resistance [29]. This concept has not been studied in great detail to-date, but Kumar and colleagues have shown that insulin resistance induced by palmitate in human urinary podocyte cell lines is associated with an increase in histone H3K36me2 and reduced H3K27me3 on the promoter region of FOXO1, a regulator of gluconeogenic genes. This effect was long-lasting and persisted even after the normalisation of palmitate levels [30].

Glomerular insulin resistance, endoplasmic reticulum (ER) stress and autophagy in diabetic glomerulopathy

There are many consequences of insulin resistance within glomeruli, which are likely to contribute to the progression of DN. One key mechanism is changes to the mitochondria and the closely connected ER [31]. Mitochondrial metabolic overload results in increased cellular oxidative stress and ER-stress which leads to the activation of unfolded protein response (UPR)[32]. UPR is a positive cellular response that in its early phase either refolds accumulated unfolded proteins, or degrades unfolded protein by the ubiquitin-proteasome pathway. Misfolded proteins are detected by the ER membrane stress sensors protein kinase RNA-like ER kinase (PERK), inositol-requiring protein 1 α (IRE1 α) and activating transcription factor 6 α (ATF6 α) and its activator X-box binding protein-1 (XBP-1), which, in turn, activates several signalling events and trigger a compensatory response to prevent further accumulation of misfolded protein. However, when the unfolded protein and cellular damage exceeds a threshold, chronic and unresolved stress results in a change from an adaptive to pro-apoptotic responses [32].

164

165 There is some evidence that glucose/oxidative stress-mediated ER stress plays a role
166 in chronic vascular complications in DN [33]. Hyperglycemia, or increased glycation of
167 proteins have been shown to mediate apoptosis partly through increases in ER stress
168 in cultured murine podocytes [34, 35]. Activation of the UPR has also been observed
169 in mouse glomerular mesangial cells exposed to glucose and glucosamine [36], and
170 in kidneys from diabetic rats administered streptozotocin for 16 weeks [37]. Microarray
171 analysis of human biopsies from patients with established DN showed that UPR genes
172 were upregulated proportionally to the severity of diabetic renal lesions [38]. Finally,
173 recent experimental evidence has demonstrated that pharmacological inhibition of ER
174 stress and stabilization of the UPR is beneficial in diabetic glomerulopathy [39].

175

176 Two studies have used transgenic mice to link podocyte insulin resistance with
177 mitochondrial function and ER stress. Ising and colleagues generated a mouse model
178 of podocyte mitochondrial dysfunction by specifically knocking out a key molecule in
179 this cell involved in mitochondrial fusion called prohibitin-2 [40]. This caused a severe
180 phenotype including glomerulosclerosis, renal failure and death at approximately a
181 month of age. They then went on to inhibit both the insulin receptor and IGF-1 receptor
182 (IGF1R) contemporaneously with podocyte-specific knockdown of prohibitin-2.
183 Inhibiting the insulin receptor alone, or in combination with the IGF1R was partially
184 protective and resulted in a significantly longer life span of the mice [40]. This suggests
185 that insulin resistance could reflect a “protective” resetting of cellular substrate
186 utilisation to shield from excess substrate flow to mitochondria with “impaired”
187 respiratory capabilities. In another study, Madhusudhan *et al.* have elegantly shown
188 that under diabetic conditions ER adaptive mechanisms are impaired in the podocyte

and that this is exacerbated when the cell is rendered more insulin resistant. Studying human and murine DN they discovered that nephropathy was associated with alterations in the UPR with impairment of the nuclear translocation of XBP-1. Genetic ablation of the transcription factor XBP-1 or activation of ATF6 (downstream of XBP-1) in the podocyte of diabetic mice aggravates DN. Of interest, mice with genetically impaired podocyte insulin signalling exhibited impaired UPR (XBP-1 activation) that was associated with more severe diabetic kidney disease when compared with diabetic controls [41].

Autophagy, regulated by the mammalian target of rapamycin complex 1 (mTORC1) is, with the UPR, essential to maintain cellular homeostasis and in the context of ER stress contributes towards the elimination of toxic and damaged cellular components [42]. Haploinsufficiency of mTORC1 in podocytes or administration of rapamycin (a mTORC1 inhibitor) resulting in activation of autophagy [43], has been shown to prevent progressive DN [44, 45]. In contrast, mTORC1 activation in podocytes, resulting in inhibition of autophagy, leads to accelerated DN [46]. Loss of insulin sensitivity in cultured podocytes results in suppression of autophagy and addition of rapamycin in these cells attenuates insulin resistance [47].

Insulin resistance, the glomerular cell cytoskeleton and other mechanisms

Experiments using podocyte cell lines have begun to reveal other downstream targets of insulin resistance which may play a role in DN. Addition of exogenous insulin to human podocytes in culture led to cytoskeletal rearrangement [18], a process which has been pharmacologically targeted using small molecules as a novel therapy for DN

[48]. Other studies [49] have identified the cytoskeleton protein septin-7 as playing an important role in the regulation of insulin-mediated translocation of GLUT4 vesicles to the plasma membrane and the control of podocyte glucose transport. Insulin may also modulate calcium signalling in podocytes which has been shown to be important in maintaining cytoskeletal dynamics by altering the expression of canonical transient receptor potential-6 channel-TRPC6 [50] and large-conductance Ca^{2+} -activated K^{+} channels [51].

Insulin stimulates the Phosphoinositide 3-kinase (PI3K) pathway and causes AKT activation. In normal physiology, insulin stimulation of podocytes results in AKT phosphorylation (activation), while, in insulin-resistant disease settings such as diabetes, a number of reports have shown an early loss of glomerular AKT phosphorylation whilst AKT signaling is maintained in the tubular compartment of the kidney [28]. AKT exists in three isoforms with AKT2 being located specifically in the podocyte within the kidney [52]. A loss of podocyte AKT2 activation is detrimental when there is chronic kidney disease associated with nephron loss [52]. AKT2 is the major isoform through which insulin signals [53]. It is currently not completely clear if the loss of renal AKT activation is detrimental in the setting of diabetes as a number of studies have shown an increase in AKT phosphorylation in the vasculature in experimental animal models of diabetes [54-58], and pharmacological inhibition of the AKT activation by AS101, may confer renoprotection in diabetes [59]. More work will have to be performed to dissect the exact role of AKT in diabetic kidney disease.

Insulin can also modulate the renin-angiotensin-aldosterone system, critical for regulating glomerular haemodynamics in DN, by increasing the expression and activity

of angiotensin converting enzyme-2 (ACE2)[60]. Further work is required to identify other downstream targets of podocyte insulin signalling ideally using systems biology genomic and proteomic approaches. Candidate molecules altered by insulin signalling might include recently identified genes found to be associated with the early stages of albuminuria in in-bred strains of mice [61].

Reactive oxygen species and diabetic nephropathy

Over the last decade, an attractive unifying hypothesis has been put forward to explain diabetic microvascular complications; specifically it was postulated that an excess in cellular substrate availability leads to an increase in reactive oxygen species (ROS) which in turn drives vascular complications in DN [62]. However, this unifying hypothesis has been challenged by the negative results of antioxidant-based clinical trials [63], and a new theory of “mitochondrial hormesis” has been proposed [64], whereby the increased mitochondrial superoxide production is considered an indicator of healthy mitochondria and physiologic oxidative phosphorylation.

Recent research has found a reduction of superoxide in the kidneys of streptozotocin (STZ)-induced diabetic Akita-mice, as assessed by a combination of *in vivo* real-time transcutaneous fluorescence, confocal microscopy, and electron paramagnetic resonance analysis [65]. The authors of this study found that chronic exposure to high glucose levels (as occurs in diabetes) results in disrupted mitochondria, which was associated with a reduced respiration and a lowering in mitochondrial superoxide. Interestingly, genetic or pharmacological correction of mitochondrial dysfunction by

improving substrate utilisation was recently found to be renoprotective in a mouse model of tubulointerstitial fibrosis [66].

From experimental animal studies it appears that increased cytosolic superoxide and other non-mitochondrial sources of ROS generation play a prominent role in diabetic kidney disease and that strategies involving a more targeted (towards specific cellular compartments such as the cytosol) antioxidant approach, may be important to optimize renoprotection in diabetes [67]. Along these lines, human studies have shown that leukocytes obtained from patients with diabetes and DN (when compared with non-diabetics or patients with diabetes without DN) have a reduced maximal respiration and reserve capacity [68, 69] suggesting that chronic metabolic stress in the presence of a reduced mitochondrial function (being this primary or secondary) will manifest with low ATP-linked respiration, low reserve capacity and reduced mitochondrial ROS generation.

It could be speculated that metabolic stress could initially (early phase) promote an excess production of mitochondrial superoxide [62] that will lead, in a subsequent chronic phase (late phase), towards mitochondrial damage, progressive deterioration in bioenergetic cellular function, reduced ATP synthesis and cell death. Future work will address these questions, and need to evaluate whether cells are able to maintain adequate number of healthy mitochondria which can then burn sufficient substrates for energy production and maintain a “balanced” level of ROS (**Figure 2**).

AMP-activated protein kinase (AMPK) is a stress-activated kinase that is activated in response to depleting ATP to preserve cell survival under conditions of reduced

substrate utilisation. AMPK activation has been involved in mitochondrial biogenesis by leading to increased mitochondrial substrate utilisation and ATP generation, in parallel with stimulation of antioxidant gene expression to ensure an optimal redox balance [70]. Reduced AMPK, as seen in the diabetic kidney of both rodents and humans [65], is associated with reduced catabolic activity (mitochondrial function) [65], and reduced AMPK-mediated inhibition of NADPH oxidase (Nox2) resulting in increased ROS production [71, 72]. Taken together, these results suggest that, in the diabetic kidney, upregulation of AMPK could be therapeutically beneficial in DN [73] to regulate nutrient utilisation and mitochondrial function towards maintenance of an optimum redox balance.

Overall, more work is required in the DN field to dissect between these opposing theories specifically by examining mitochondrial function and specific ROS moieties both *in vitro* and in tissues.

Vascular endothelial growth factor-A (VEGFA) and the glycocalyx

Recent studies have shown a connection between insulin resistance and the subsequent production of VEGFA in podocytes [74]. This finding is likely to be important in the setting of DN with many elegant studies using transgenic mice highlighting the importance of podocyte VEGFA levels in the progression of this condition [4]. A new aspect of VEGFA signalling in the glomerulus is potential cross talk between VEGFA secreted from podocytes and the GECs glycocalyx in the setting of diabetes. There is clear evidence that the GECs glycocalyx is lost both systemically and within the diabetic glomerulus, and that this contributes to both cardiovascular and

renal complications [6]. Mechanistically there are a number of pathways which led to loss of the glomerular glycocalyx including hyperglycaemia [75], and ROS [76].

During the early phases of diabetes an increase in VEGFA causes glycocalyx shedding from the GECs. Furthermore, the inhibitory isoform of VEGFA called VEGF-A_{165b} also plays a role in maintaining the GECs glycocalyx in diabetes. Oltean et al. [77] have shown that in diabetic patients with progressive nephropathy, the renal expression of VEGF-A_{165b} is lost. They went on to develop a number of murine models of DN and have shown that genetic overexpression or pharmacological administration of VEGF-A_{165b} to the mouse, acting through VEGF receptor 2 in the GECs, restores damaged glomerular endothelial glycocalyx and improves renal function. VEGF-A_{165b} also improved the permeability of isolated human diabetic glomeruli suggesting the response is conserved across murine and human species [77].

VEGFA signalling is only one component of a complex system of molecular cross talk between the podocyte and glycocalyx. New insights have revealed that molecules produced by the endothelium can signal to the podocyte and then back to the glomerular glycocalyx. Using transgenic murine models and conditionally immortalised murine podocytes and GECs, Garsen et al have shown that endothelin-1 (ET-1), an endothelial derived vasoconstrictor, is released by the GECs in diabetic conditions and leads to shedding of the glycocalyx [78]. This is prevented by deleting the ET-1 receptor specifically in the podocyte. This is therapeutically intriguing as there are ET-1 receptor antagonists that have been shown to ameliorate early microalbuminuric diabetic kidney disease [79]. In the future, therapeutic approaches to maintain the GECs glycocalyx should be explored in more detail (**Figure 3**).

SGLT2 and kidney disease

Poor glycaemic control and hyperinsulinaemia (at least in the early phase of diabetes) lead to upregulation of SGLT2 expression and proximal tubular SGLT2-mediated sodium-glucose reabsorption [80], which in turns is believed to also contribute to higher blood pressure levels [81]. SGLT2 inhibitors have recently been developed as oral hypoglycaemic agents [82]. The SGLT2 antagonists block the sodium-coupled energy dependent glucose proximal tubular reabsorption resulting in improvement in diabetes control, weight loss and blood pressure lowering. Recent clinical trials have demonstrated a dramatic cardiovascular [83] and renoprotective [84] effect of the SGLT2 inhibitor empaglifozin.

The mechanism by which SGLT2 inhibitors exert their renoprotective effects is currently unknown. One possibility is that the improvement in renal disease is secondary to activation of tubuloglomerular feedback, a prime mechanism that determines a reduction in glomerular capillary pressure [85]. A complementary mechanism may be that the inhibition of enhanced tubuli sodium-coupled glucose transport seen in diabetes would result in diminished tubulointerstitial injury and progression of DN [81]. The use of SGLT2 inhibitors in combination with inhibitors of the renin-angiotensin-aldosterone system in patients with diabetes may confer some renoprotection via upregulation of ACE2 and angiotensin 1-7/ 1-9 [86] which retains a vasodilatory, anti-proliferative, anti-inflammatory and anti-oxidative stress effect [87]. Inhibition of SGLT2 increased expression and activity in diabetes [85] could lead to restriction of chronic excess in substrates (glucose) availability, known to result in AMPK inhibition [88], and promote a favourable renal outcome. Studies will have to

dissect whether inhibition of SGLT2 could result in AMPK activation and secondary renoprotection.

In parallel to these “renal mechanisms”, natriuresis and plasma volume contraction paralleled by blood pressure reduction has been also been proposed as a “systemic” renoprotective mechanisms for SGLT2 inhibitors [89](**Figure 4**).

Concluding Remarks and Future Perspectives

Glomerular cellular insulin resistance plays an important role in mitochondrial dysfunction-ER stress and the UPR, which contribute to glomerular cell dysfunction and progressive kidney disease. AMPK, with its important role in mitochondrial function, could represent a potential target for treatment in DN; more studies are required to assess the role of AMPK on podocyte biology and the regulation of the glomerular filtration barrier. A link between the tubular compartment and the glomerulus is evident in the pathophysiology of tubular SGLT2-mediated Na-coupled glucose reabsorption in diabetes, however it is not yet completely clear what are the mechanisms underlying these beneficial effects. Future studies will need to better dissect the cellular mechanisms underlying the proposed pathways outlined in this article, specifically focusing on the physiology of the nephron as a whole entity, and by identifying potentially targetable molecules for future treatment.

Acknowledgements

LG is supported by Diabetes UK, European Foundation for the Study of Diabetes, Heart Research UK, Diabetes Research Wellness Foundation, and National Institute for Health and Research, Biomedical Research Centre at Guy’s and St Thomas Foundation Trust in partnership with King’s College London.

388 DAL is supported by project grants from Diabetes UK (13/0004763 and 15/0005283),
389 Kidney Research UK (RP36/2015), a Medical Research Council New Investigator
390 Award (MR/J003638/1) and by the National Institute for Health Research Biomedical
391 Research Centre at Great Ormond Street Hospital for Children NHS Foundation Trust
392 and University College London.

393 RJMC is supported by an MRC Senior Clinical Fellowship MR/K010492/1, Kidney
394 Research UK, Wellcome Trust, and the European Union.

395

396

TABLE 1

Diabetes results in inflammation, increased adiposity and chronic hyperglycaemia which drive podocyte insulin resistance resulting in disruption to podocytes and the glomerulus. (NOD2: Nucleotide-binding oligomerization domain containing protein 2, TLR: Toll-like receptor, SMAD: vertebrate homologues of *Sma* and *Mad*; FFAs: free fatty acids; SHP-1: Src homology-2 domain-containing phosphatase-1)

Diabetes related phenotypes	Molecular mechanisms	Glomerular phenotype	REF #
Chronic inflammation	Increased NOD2	Increased pro-inflammatory responses and impaired insulin signaling	[22]
	Increased TLR	Podocyte inflammation and insulin resistance	[23]
	Increased SMAD	Mitochondrial dysfunction and insulin resistance in podocytes	[21]
Obesity	Decreased adiponectin	Albuminuria and increased oxidative stress in podocytes	[24]
	Increased FFAs	Inhibition of insulin-stimulated glucose uptake in human podocytes	[26]
Elevated blood sugar	Increased SHP-1	Podocyte insulin resistance and detachment	[27]
	Ubiquitination and degradation of components of insulin signaling pathway	Diabetic glomerulopathy and albuminuria	[28]

Figure legends:

Figure 1: Schematic structure of a normal and diabetic glomeruli.

(A) Schematic representation of a normal glomerular structure. (B) The major glomerular structural changes occurring in diabetic glomerulopathy. Note the extensive mesangial expansion, the thickening of the glomerular basement membrane (GBM), the detachment of podocyte, and the impairment in the glycocalyx and glomerular endothelial cells (GECs).

419 **Figure 2: Hypothetical shift of superoxide level imbalance in diabetic**
 420 **complications.**

421 Acute (early phase) exposure of cells to elevated glucose levels results in upregulation
 422 of glucose oxidation with pyruvate-mediated stimulation of the tricarboxylic acid (TCA)
 423 cycle with increased production of electron donors (NADH, FADH₂) that, via the
 424 electron transport chain, will results in an excess generation of superoxide (O₂⁻). Cells
 425 chronically exposed to elevated glucose levels (late phase) will result in reduction in
 426 the availability of acetyl-CoA (secondary to inhibition of pyruvate dehydrogenase
 427 activity) for the mitochondria resulting in reduced electron transport chain activity, a
 428 fall in mitochondrial ATP production, less mitochondria superoxide production and
 429 cellular dysfunction.

432

433

434 **Figure 3: Glomerular cell cross-talk and glycocalyx.**

435 Transmission electron microscopy image of the glomerular filtration barrier (podocyte
 436 glomerular basement membrane (GBM), glomerular endothelial cells (GEC), and
 437 glycocalyx) highlighting how molecules produced by the podocytes and endothelium
 438 (via the podocyte) can signal to the glomerular glycocalyx. Recently identified key
 439 molecules such as VEGF-A_{165b}, VEGF-C, and angiopoietin-1 (ANGPT1) confer a
 440 beneficial effect (green arrows) towards glycocalyx maintenance. Conversely VEGF-
 441 A₁₆₅ and Endothelin (secreted by GECs and signals to the Endothelin-1 receptor in the
 442 podocyte causing this cell to release heparanase, which then acts on the glomerular
 443 glycocalyx to cleave heparin sulphate) promote shedding of the GECs glycocalyx (red
 444 arrows).

449 **Figure 4: Proposed SGLT2 inhibition-mediated renoprotective mechanisms.**

450 SGLT2 inhibition blocks sodium-glucose coupled glucose reabsorption at the S1 S2
 451 segment of the proximal tubule. The net result is loss of glucose and sodium (the latter
 452 especially in patients on renin angiotensin aldosterone blockade) in the urine, with
 453 secondary weight loss, improvement in glycaemic control, blood pressure fall, and
 454 plasma volume contraction. These effects confer cardiac and renal protection in
 455 patients with diabetes. (ACE2: angiotensin converting enzyme 2)

References:

- 1 Gnudi, L., Gentile, G., Ruggenenti, P. (2016) The patient with diabetes mellitus. In *Oxford Textbook of Clinical Nephrology* (Turner, N., Lamiere, N., Goldsmith, D.J., Wineearls, C.G., Himmelfarb, J., Remuzzi, G., ed), pp. 1199-1247, Oxford University Press
- 2 Benzing, T. (2004) Signaling at the slit diaphragm. *J Am Soc Nephrol* 15, 1382-1391
- 3 Zhu, J., et al. (2008) Nephrin mediates actin reorganization via phosphoinositide 3-kinase in podocytes. *Kidney Int* 73, 556-566
- 4 Gnudi, L., et al. (2015) Vascular growth factors play critical roles in kidney glomeruli. *Clin Sci (Lond)* 129, 1225-1236
- 5 Haraldsson, B., et al. (2008) Properties of the glomerular barrier and mechanisms of proteinuria. *Physiol Rev.* 88, 451-487
- 6 Salmon, A.H., et al. (2012) Loss of the endothelial glycocalyx links albuminuria and vascular dysfunction. *J Am Soc Nephrol* 23, 1339-1350
- 7 Deckert, T., et al. (1989) Albuminuria reflects widespread vascular damage. The Steno hypothesis. *Diabetologia* 32, 219-226
- 8 Orchard, T.J., et al. (2002) Nephropathy in type 1 diabetes: a manifestation of insulin resistance and multiple genetic susceptibilities? Further evidence from the Pittsburgh Epidemiology of Diabetes Complication Study. *Kidney Int* 62, 963-970
- 9 Karalliedde, J. and Gnudi, L. (2016) Diabetes mellitus, a complex and heterogeneous disease, and the role of insulin resistance as a determinant of diabetic kidney disease. *Nephrol Dial Transplant* 31, 206-213

479 10 Sasson, A.N. and Cherney, D.Z. (2012) Renal hyperfiltration related to diabetes
480 mellitus and obesity in human disease. *World J Diabetes* 3, 1-6

481 11 Gnudi, L., *et al.* (2007) Mechanical forces in diabetic kidney disease: a trigger for
482 impaired glucose metabolism. *J.Am.Soc.Nephrol.* 18, 2226-2232

483 12 Trevisan, R., *et al.* (1998) Enhanced responsiveness of blood pressure to sodium
484 intake and to angiotensin II is associated with insulin resistance in IDDM patients with
485 microalbuminuria. *Diabetes* 47, 1347-1353

486 13 Vedovato, M., *et al.* (2004) Effect of sodium intake on blood pressure and
487 albuminuria in Type 2 diabetic patients: the role of insulin resistance. *Diabetologia* 47,
488 300-303

489 14 Coward, R.J., *et al.* (2005) The human glomerular podocyte is a novel target for
490 insulin action. *Diabetes* 54, 3095-3102

491 15 Butlen, D., *et al.* (1988) Insulin receptors along the rat nephron: [125I] insulin
492 binding in microdissected glomeruli and tubules. *Pflugers Arch* 412, 604-612

493 16 Tiwari, S., *et al.* (2008) Impaired sodium excretion and increased blood pressure in
494 mice with targeted deletion of renal epithelial insulin receptor. *Proc Natl Acad Sci U S*
495 *A* 105, 6469-6474

496 17 Lay, A. and Coward, R.J. (2014) Recent advances in our understanding of insulin
497 signalling to the podocyte. *Nephrol Dial Transplant* 29, 1127-1133

498 18 Welsh, G.I., *et al.* (2010) Insulin signaling to the glomerular podocyte is critical for
499 normal kidney function. *Cell Metab* 12, 329-340

500 19 Tiwari, S., *et al.* (2013) Deletion of the insulin receptor in the proximal tubule
501 promotes hyperglycemia. *J Am Soc Nephrol* 24, 1209-1214

502 20 Li, L., *et al.* (2013) Reduced ENaC activity and blood pressure in mice with genetic
 503 knockout of the insulin receptor in the renal collecting duct. *Am J Physiol Renal Physiol*
 504 304, F279-288

505 21 Sun, Y.B., *et al.* (2015) Smad3 deficiency protects mice from obesity-induced
 506 podocyte injury that precedes insulin resistance. *Kidney Int* 88, 286-298

507 22 Du, P., *et al.* (2013) NOD2 promotes renal injury by exacerbating inflammation and
 508 podocyte insulin resistance in diabetic nephropathy. *Kidney Int* 84, 265-276

509 23 Cha, J.J., *et al.* (2013) Renal protective effects of toll-like receptor 4 signaling
 510 blockade in type 2 diabetic mice. *Endocrinology* 154, 2144-2155

511 24 Sharma, K., *et al.* (2008) Adiponectin regulates albuminuria and podocyte function
 512 in mice. *J Clin Invest* 118, 1645-1656

513 25 Sweiss, N. and Sharma, K. (2014) Adiponectin effects on the kidney. *Best Pract*
 514 *Res Clin Endocrinol Metab* 28, 71-79

515 26 Lennon, R., *et al.* (2009) Saturated fatty acids induce insulin resistance in human
 516 podocytes: implications for diabetic nephropathy. *Nephrol Dial Transplant* 24, 3288-
 517 3296

518 27 Drapeau, N., *et al.* (2013) Expression of SHP-1 induced by hyperglycemia prevents
 519 insulin actions in podocytes. *Am J Physiol Endocrinol Metab* 304, E1188-1198

520 28 Mima, A., *et al.* (2011) Glomerular-specific protein kinase C-beta-induced insulin
 521 receptor substrate-1 dysfunction and insulin resistance in rat models of diabetes and
 522 obesity. *Kidney Int* 79, 883-896

523 29 Duque-Guimaraes, D.E. and Ozanne, S.E. (2013) Nutritional programming of
 524 insulin resistance: causes and consequences. *Trends Endocrinol Metab* 24, 525-535

525 30 Kumar, S., *et al.* (2016) Fatty acid induced metabolic memory involves alterations
526 in renal histone H3K36me2 and H3K27me3. *Mol Cell Endocrinol* 422, 233-242

527 31 Kornmann, B., *et al.* (2009) An ER-mitochondria tethering complex revealed by a
528 synthetic biology screen. *Science* 325, 477-481

529 32 Hetz, C. (2012) The unfolded protein response: controlling cell fate decisions under
530 ER stress and beyond. *Nature reviews. Molecular cell biology* 13, 89-102

531 33 Zhuang, A. and Forbes, J.M. (2014) Stress in the kidney is the road to pERdition:
532 is endoplasmic reticulum stress a pathogenic mediator of diabetic nephropathy? *The*
533 *Journal of endocrinology* 222, R97-111

534 34 Cao, Y., *et al.* (2014) Role of endoplasmic reticulum stress in apoptosis of
535 differentiated mouse podocytes induced by high glucose. *Int J Mol Med* 33, 809-816

536 35 Chen, Y., *et al.* (2008) Effect of taurine-conjugated ursodeoxycholic acid on
537 endoplasmic reticulum stress and apoptosis induced by advanced glycation end
538 products in cultured mouse podocytes. *Am J Nephrol* 28, 1014-1022

539 36 Cheng, D.W., *et al.* (2006) An analysis of high glucose and glucosamine-induced
540 gene expression and oxidative stress in renal mesangial cells. *Arch Physiol Biochem*
541 112, 189-218

542 37 Liu, G., *et al.* (2008) Apoptosis induced by endoplasmic reticulum stress involved
543 in diabetic kidney disease. *Biochem Biophys Res Commun* 370, 651-656

544 38 Lindenmeyer, M.T., *et al.* (2008) Proteinuria and hyperglycemia induce
545 endoplasmic reticulum stress. *J Am Soc Nephrol* 19, 2225-2236

546 39 Cao, A.L., *et al.* (2016) Ursodeoxycholic acid and 4-phenylbutyrate prevent
547 endoplasmic reticulum stress-induced podocyte apoptosis in diabetic nephropathy.
548 *Lab Invest*

549 40 Ising, C., *et al.* (2015) Inhibition of insulin/IGF-1 receptor signaling protects from
550 mitochondria-mediated kidney failure. *EMBO Mol Med* 7, 275-287

551 41 Madhusudhan, T., *et al.* (2015) Defective podocyte insulin signalling through p85-
552 XBP1 promotes ATF6-dependent maladaptive ER-stress response in diabetic
553 nephropathy. *Nature communications* 6, 6496

554 42 Kroemer, G., *et al.* (2010) Autophagy and the integrated stress response. *Mol Cell*
555 40, 280-293

556 43 Kim, J., *et al.* (2011) AMPK and mTOR regulate autophagy through direct
557 phosphorylation of Ulk1. *Nat Cell Biol* 13, 132-141

558 44 Godel, M., *et al.* (2011) Role of mTOR in podocyte function and diabetic
559 nephropathy in humans and mice. *J Clin Invest* 121, 2197-2209

560 45 Xiao, T., *et al.* (2014) Rapamycin promotes podocyte autophagy and ameliorates
561 renal injury in diabetic mice. *Mol Cell Biochem* 394, 145-154

562 46 Inoki, K., *et al.* (2011) mTORC1 activation in podocytes is a critical step in the
563 development of diabetic nephropathy in mice. *J Clin Invest* 121, 2181-2196

564 47 Xu, Y., *et al.* (2016) Autophagy downregulation contributes to insulin resistance
565 mediated injury in insulin receptor knockout podocytes in vitro. *PeerJ* 4, e1888

566 48 Schiffer, M., *et al.* (2015) Pharmacological targeting of actin-dependent dynamin
567 oligomerization ameliorates chronic kidney disease in diverse animal models. *Nat Med*
568 21, 601-609

569 49 Wasik, A.A., *et al.* (2012) Septin 7 forms a complex with CD2AP and nephrin and
570 regulates glucose transporter trafficking. *Mol Biol Cell* 23, 3370-3379

571 50 Kim, E.Y., *et al.* (2012) Insulin increases surface expression of TRPC6 channels in
572 podocytes: role of NADPH oxidases and reactive oxygen species. *Am J Physiol Renal*
573 *Physiol* 302, F298-307

574 51 Kim, E.Y. and Dryer, S.E. (2011) Effects of insulin and high glucose on mobilization
575 of slo1 BKCa channels in podocytes. *J Cell Physiol* 226, 2307-2315

576 52 Canaud, G., *et al.* (2013) AKT2 is essential to maintain podocyte viability and
577 function during chronic kidney disease. *Nat Med* 19, 1288-1296

578 53 George, S., *et al.* (2004) A family with severe insulin resistance and diabetes due
579 to a mutation in AKT2. *Science* 304, 1325-1328

580 54 Zdychova, J. and Komers, R. (2005) Emerging role of Akt kinase/protein kinase B
581 signaling in pathophysiology of diabetes and its complications. *Physiol Res* 54, 1-16

582 55 Wu, D., *et al.* (2009) PKC-beta1 mediates glucose-induced Akt activation and TGF-
583 beta1 upregulation in mesangial cells. *J Am Soc Nephrol* 20, 554-566

584 56 Kato, M., *et al.* (2006) Role of the Akt/FoxO3a pathway in TGF-beta1-mediated
585 mesangial cell dysfunction: a novel mechanism related to diabetic kidney disease. *J*
586 *Am Soc Nephrol* 17, 3325-3335

587 57 Xin, X., *et al.* (2005) Glucose-induced Akt1 activation mediates fibronectin
588 synthesis in endothelial cells. *Diabetologia* 48, 2428-2436

589 58 Kim, S.Y., *et al.* (2009) Role of kidney ADP-ribosyl cyclase in diabetic nephropathy.
590 *Am J Physiol Renal Physiol* 296, F291-297

591 59 Shemesh, Il, *et al.* (2014) AS101 prevents diabetic nephropathy progression and
592 mesangial cell dysfunction: regulation of the AKT downstream pathway. *PLoS One* 9,
593 e114287

594 60 Riera, M., *et al.* (2014) Effect of insulin on ACE2 activity and kidney function in the
595 non-obese diabetic mouse. *PLoS One* 9, e84683

596 61 Long, D.A., *et al.* (2013) Albuminuria is associated with too few glomeruli and too
597 much testosterone. *Kidney Int*

598 62 Brownlee, M. (2001) Biochemistry and molecular cell biology of diabetic
599 complications. *Nature* 414, 813-820

600 63 Forbes, J.M., *et al.* (2008) Oxidative stress as a major culprit in kidney disease in
601 diabetes. *Diabetes* 57, 1446-1454

602 64 Sharma, K. (2015) Mitochondrial hormesis and diabetic complications. *Diabetes*
603 64, 663-672

604 65 Dugan, L.L., *et al.* (2013) AMPK dysregulation promotes diabetes-related reduction
605 of superoxide and mitochondrial function. *J Clin Invest* 123, 4888-4899

606 66 Kang, H.M., *et al.* (2015) Defective fatty acid oxidation in renal tubular epithelial
607 cells has a key role in kidney fibrosis development. *Nat Med* 21, 37-46

608 67 Forbes, J.M. and Cooper, M.E. (2013) Mechanisms of diabetic complications.
609 *Physiological reviews* 93, 137-188

610 68 Czajka, A., *et al.* (2015) Altered Mitochondrial Function, Mitochondrial DNA and
611 Reduced Metabolic Flexibility in Patients With Diabetic Nephropathy. *EBioMedicine* 2,
612 499-512

613 69 Chacko, B.K., *et al.* (2015) The Bioenergetic Health Index is a sensitive measure
614 of oxidative stress in human monocytes. *Redox biology* 8, 43-50

615 70 Hardie, D.G., *et al.* (2012) AMPK: a nutrient and energy sensor that maintains
616 energy homeostasis. *Nature reviews. Molecular cell biology* 13, 251-262

617 71 Sedeek, M., *et al.* (2010) Critical role of Nox4-based NADPH oxidase in glucose-
618 induced oxidative stress in the kidney: implications in type 2 diabetic nephropathy. *Am*
619 *J Physiol Renal Physiol* 299, F1348-1358

620 72 Eid, A.A., *et al.* (2010) AMP-activated protein kinase (AMPK) negatively regulates
621 Nox4-dependent activation of p53 and epithelial cell apoptosis in diabetes. *J Biol*
622 *Chem* 285, 37503-37512

623 73 Al-Rasheed, N.M., *et al.* (2015) Renoprotective Effects of Fenofibrate via
624 Modulation of LKB1/AMPK mRNA Expression and Endothelial Dysfunction in a Rat
625 Model of Diabetic Nephropathy. *Pharmacology* 95, 229-239

626 74 Hale, L.J., *et al.* (2013) Insulin directly stimulates VEGF-A production in the
627 glomerular podocyte. *Am J Physiol Renal Physiol* 305, F182-188

628 75 Singh, A., *et al.* (2011) High glucose causes dysfunction of the human glomerular
629 endothelial glycocalyx. *Am J Physiol Renal Physiol* 300, F40-48

630 76 Singh, A., *et al.* (2013) Reactive oxygen species modulate the barrier function of
631 the human glomerular endothelial glycocalyx. *PLoS One* 8, e55852

632 77 Oltean, S., *et al.* (2014) Vascular Endothelial Growth Factor-A165b Is Protective
633 and Restores Endothelial Glycocalyx in Diabetic Nephropathy. *J Am Soc Nephrol*

634 78 Garsen, M., *et al.* (2016) Endothelin-1 Induces Proteinuria by Heparanase-
635 Mediated Disruption of the Glomerular Glycocalyx. *J Am Soc Nephrol*

636 79 de Zeeuw, D., *et al.* (2014) The endothelin antagonist atrasentan lowers residual
637 albuminuria in patients with type 2 diabetic nephropathy. *J Am Soc Nephrol* 25, 1083-
638 1093

639 80 Rahmoune, H., *et al.* (2005) Glucose transporters in human renal proximal tubular
640 cells isolated from the urine of patients with non-insulin-dependent diabetes. *Diabetes*
641 54, 3427-3434

642 81 Vallon, V. and Thomson, S.C. (2012) Renal function in diabetic disease models:
643 the tubular system in the pathophysiology of the diabetic kidney. *Annual review of*
644 *physiology* 74, 351-375

645 82 Ferrannini, E. and Solini, A. (2012) SGLT2 inhibition in diabetes mellitus: rationale
646 and clinical prospects. *Nat Rev Endocrinol* 8, 495-502

647 83 Zinman, B., *et al.* (2015) Empagliflozin, Cardiovascular Outcomes, and Mortality in
648 Type 2 Diabetes. *N Engl J Med* 373, 2117-2128

649 84 Wanner, C., *et al.* (2016) Empagliflozin and Progression of Kidney Disease in Type
650 2 Diabetes. *N Engl J Med*

651 85 Vallon, V. (2015) The mechanisms and therapeutic potential of SGLT2 inhibitors in
652 diabetes mellitus. *Annu Rev Med* 66, 255-270

653 86 Tikellis, C., *et al.* (2008) ACE2 deficiency modifies renoprotection afforded by ACE
654 inhibition in experimental diabetes. *Diabetes* 57, 1018-1025

655 87 Burrell, L.M., *et al.* (2004) ACE2, a new regulator of the renin-angiotensin system.
656 *Trends Endocrinol Metab* 15, 166-169

657 88 Viollet, B., *et al.* (2010) AMPK inhibition in health and disease. *Crit Rev Biochem*
658 *Mol Biol* 45, 276-295

659 89 Rajasekeran, H., *et al.* (2016) Sodium-glucose cotransporter 2 inhibition and
660 cardiovascular risk reduction in patients with type 2 diabetes: the emerging role of
661 natriuresis. *Kidney Int* 89, 524-526

662

663

664

665

OUTSTANDING QUESTIONS BOX

- What are the main pathway/s that link insulin action, mitochondrial function and UPR?
- What are the mitochondrial-driven mechanisms that predispose towards faster kidney disease progression in diabetes? Is there an alteration in the mitochondria driven UPR-mediated response towards cell survival or is it a primary alteration in UPR response?
- Does AMPK, with its important role as a regulator of nutrient utilisation and mitochondrial function, represent a real answer to diabetes mediated ER-stress mitochondria dysfunction, UPR, and, if so, what are the mechanisms?
- Could targeting the glycocalyx be a new therapeutic approach for diabetic nephropathy? What are the important molecular signals from the podocyte and endothelium which regulate the glycocalyx?
- How does SGLT2 inhibition confer reno-protection? Is it about SGLT2-driven sodium and volume loss (systemic effect) or tubuloglomerular feedback and inflammation (intrarenal effect)? Are these mechanisms behind the renoprotective effects of these drugs?

TRENDS BOX:

- Insulin resistance is a key mechanism for diabetic glomerulopathy.
- Disruption in the molecular communication between glomerular podocytes and endothelia is critical in the progression of diabetic nephropathy.
- Raised (but not too elevated) mitochondrial superoxide cellular levels in parallel with healthy mitochondria are protective against progression of diabetic kidney disease.
- A reduction in maximal mitochondrial respiration and reserve capacity could represent an important driving force for kidney disease progression in diabetes.
- Inhibition of SGLT2-mediated sodium-coupled glucose transport confers renoprotection of similar magnitude of inhibitors of the renin-angiotensin-aldosterone system.