King's Research Portal

DOI:

[10.3109/15412555.2012.707258](https://doi.org/10.3109/15412555.2012.707258)

Document Version

Early version, also known as pre-print

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Hogg, L., Garrod, R., Thornton, H., McDonnell, L., Bellas, H., & White, P. (2012). Effectiveness, Attendance, and Completion of an Integrated, System-Wide Pulmonary Rehabilitation Service for COPD: Prospective Observational Study. *Copd-Journal of chronic obstructive pulmonary disease*, 9(5), 546-554.
<https://doi.org/10.3109/15412555.2012.707258>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

**Open Access document
downloaded from King's Research Portal
<https://kclpure.kcl.ac.uk/portal>**

Citation to published version:

Hogg, L., Garrod, R., Thornton, H., McDonnell, L., Bellas, H., & White, P. (2012). Effectiveness, Attendance, and Completion of an Integrated, System-Wide Pulmonary Rehabilitation Service for COPD: Prospective Observational Study. *Copd-Journal of chronic obstructive pulmonary disease*, 9(5), 546-554.

The original publication is available at:

<http://dx.doi.org/10.3109/15412555.2012.707258>

This version:

<https://kclpure.kcl.ac.uk/portal/en/publications/effectiveness-attendance-and-completion-of-an-integrated-systemwide-pulmonary-rehabilitation-service-for-copd-prospective-observational-study%283d1a366d-6a9d-4779-9967-546a3239f487%29.html>

The copyright in the published version resides with the publisher.

When referring to this paper, please check the page numbers in the published version and cite these.

General rights

Copyright and moral rights for the publications made accessible in King's Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications in King's Research Portal that users recognise and abide by the legal requirements associated with these rights.'

- Users may download and print one copy of any publication from King's Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the King's Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Effectiveness, attendance, and completion of an integrated, system-wide pulmonary rehabilitation service for COPD: prospective observational study.

Journal:	<i>COPD: Journal Of Chronic Obstructive Pulmonary Disease</i>
Manuscript ID:	Draft
Manuscript Type:	Original Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	Hogg, Lauren; Guy's and St Thomas' NHS Foundation Trust, Department of Physiotherapy Garrod, Rachel ; King's College London, King's Health Partners, King's College Hospital NHS Foundation Trust, Department of Physiotherapy Thornton, Hannah; Kings College London, King's Health Partners, Department of Primary Care and Public Health Sciences McDonnell, Lynn; Guy's and St Thomas' NHS Foundation Trust, Department of Physiotherapy Bellas, Helene; King's College London, King's Health Partners, King's College Hospital NHS Foundation Trust, Department of Physiotherapy White, Patrick; Kings College London, King's Health Partners, Department of Primary Care and Public Health Sciences
Keywords:	pulmonary , rehabilitation, utilisation, integrated care, community, referral

SCHOLARONE™
Manuscripts

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

Effectiveness, attendance, and completion of an integrated, system-wide pulmonary rehabilitation service for COPD: prospective observational study.

¹Lauren Hogg, ²Rachel Garrod, ³Hannah Thornton, ¹Lynn McDonnell, ⁴Helene Bellas, ³Patrick White*

¹Department of Physiotherapy, Guy's and St Thomas' NHS Foundation Trust, St Thomas' Hospital, London, United Kingdom

²King's College London, King's Health Partners, King's College Hospital NHS Foundation Trust, Department of Physiotherapy, London, United Kingdom

³King's College London, King's Health Partners, Department of Primary Care and Public Health Sciences, London, United Kingdom

⁴ King's College Hospital NHS Foundation Trust, Department of Physiotherapy, London, United Kingdom

*Correspondence: Dr Patrick White, King's College London, King's Health Partners, Department of Primary Care and Public Health Sciences, 42 Weston Street, London SE1 3QD, UK. Tel +44 207 848 8679; Fax +44 207 848 6620. Email: patrick.white@kcl.ac.uk

Keywords: Pulmonary; rehabilitation; utilisation; integrated care; community; referral

Abstract

Pulmonary rehabilitation (PR) is one of the most effective treatments for COPD but not widely available. Uptake is poor and completion rates are low. In this integrated PR service we report on effectiveness, attendance, and completion of twice weekly rolling recruitment and once weekly cohort recruitment programmes in two hospital and five community PR sites. The hospital and two of the community programmes were ‘rolling’ recruitment twice weekly for 8 weeks. Three community programmes ran in once weekly cohorts for 8 weeks. Predictors of attendance, completion and effectiveness were sought. 1114 eligible COPD patients were referred. 812(73%) attended assessment, 656(59%) started and 441(40%) completed. Significant improvements were seen in incremental shuttle walk test (ISWT) (mean 68.3m; 95%CI 59.3-77.4), Chronic Respiratory Questionnaire self-report dyspnoea scale (CRQ-SR) (0.94; 0.80-1.07), Hospital Anxiety and Depression Scale anxiety (0.9; 0.5-1.2) and depression (1.1; 0.8-1.4) components, exceeding the minimum clinically important difference for ISWT and CRQ-SR. Twice weekly compared with once weekly programmes showed similar improvement. Patients were less likely to complete if they were deprived (4th quintile of deprivation 0.56; 0.33-0.94, 5th quintile 0.57; 0.34-0.85), reported MRC dyspnoea scale 4 (0.61; 0.37-0.97) or 5 (0.39; 0.16-0.93), or had been referred by their general practitioner (0.42; 0.24-0.74) (pseudo R² 0.103). PR is effective for COPD in real-world practice achieving results comparable to trials. Only a small proportion of the variance in attendance and completion of PR was explained by demographic characteristics, disease severity, psychological morbidity and source of referral despite the large number of participants.

Introduction

Evidence for the benefits of pulmonary rehabilitation (PR) in COPD is broadly based and widely accepted.[1,2] Its prescription is recommended in national and international guidelines for patients with symptomatic disease.[3,4] A number of randomised controlled trials and meta-analyses have reported on the beneficial effects of PR on exercise capacity, dyspnoea, quality of life and improvements in health care utility.[5-8] Low rates of referral, uptake and completion have been widely reported and have implications for service delivery. Although the availability of PR has improved in the United Kingdom, and many programmes are now available to patients referred from primary care as well as secondary care, few studies have examined the effectiveness and applicability of PR in conventional health care.[9,10] In a retrospective analysis of the trials included in a Cochrane Airways meta-analysis of rehabilitation, Bjoernshave et al highlighted selective inclusion criteria and significant drop-out rates or non-completion in several studies.[11] Seventy five percent of participants in 26 trials were non-completers due to ineligibility for study inclusion or drop-out. In the USA, Cote et al found that 53% of participants either declined to take part in a trial or dropped out.[12]

Trials investigating the predictors of drop-out from PR have highlighted a number of associated phenotypical features. In a recent systematic review Keating et al identified obstacles to initial PR uptake: disruption to valued routine, uncertainty of the referrer in its effectiveness, inconvenient timing, travel issues, and low perceived benefit.[13] Most of the studies included were small and several were qualitative. Obstacles to PR completion were illness and co-morbidities, travel, current smoking, lack of social support, COPD exacerbations, and low perceived benefit.[13] Severe disease including severe dyspnoea was associated with drop-out in a retrospective study of 239 patients by Sabit et al in the UK, and in the study by Cote et al in the USA, and Garrod and colleagues found that quadriceps weakness and depression were also predictors of drop out in the UK.[12,14,15] Whilst these studies suggest that severity of symptoms and co-morbidities may be associated with higher risk of drop-out, prior identification of participants at risk of non-attendance remains difficult and no robust predictive models are available.

There is also a paucity of information concerning the effectiveness of different models of rehabilitation, optimal duration, frequency of supervision and method of recruiting.[16] In a randomised controlled trial evaluating the frequency of supervised sessions O'Neill et al have suggested that once weekly programmes may lead to similar outcomes as twice weekly programmes. However since this small study was not powered for equivalence, results should be

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

interpreted with caution. Within the current economic climate the frequency of supervised sessions remains an important clinical consideration. The impact of the method of recruitment, for example cohort recruitment (all patients start and finish together) or rolling recruitment (continuous programme, new patients start every week), has not been adequately tested. Community programmes are likely to be as effective as hospital programmes.[17] Increasingly, data show benefits of programmes integrated between primary and secondary care for COPD.[18,19]

We report outcomes from a prospective observational study on the effectiveness of an integrated system-wide service of pulmonary rehabilitation, where rehabilitation is provided in hospital and community settings, with rolling recruitment to twice weekly supervised programmes and cohort recruitment to once weekly supervised programmes. We have evaluated attendance at assessment and rates of completion, and have sought to define predictors of effectiveness, attendance at assessment and completion.

Methods

We have analysed data from an integrated PR service in two inner London boroughs across community and acute hospital settings between April 2008 and March 2010. Ethical approval was obtained from the Proportional Research Ethics Review Committee, St Thomas Hospital, London, (Research Ethics Committee reference number 09/H0701/90).

Participants

Patients with a diagnosis of COPD were eligible. Patients were excluded if they were not appropriate for rehabilitation due to cardiovascular instability or significant musculoskeletal limitations at referral or assessment. Referrals were received from primary and secondary care including general practitioners, practice nurses, community COPD teams, in-patient COPD teams, in-patient physiotherapists, and respiratory physicians in outpatient clinics. Within two weeks, patients were sent written confirmation of receipt of referral, together with a PR information leaflet produced by the British Lung Foundation. Patients were telephoned to offer an appointment for assessment at the most suitable location followed by written confirmation. Patients requiring transport used hospital transport services to attend hospital sites. All patients received a reminder telephone call prior to the appointment. Those who did not attend were offered one further assessment appointment.

Pulmonary rehabilitation programmes

Programmes took place in seven centres: two hospital physiotherapy gyms and five community settings (two local authority gyms, one health centre and two community halls). The two hospital and two of the community programmes were delivered on a rolling recruitment basis. Patients on rolling programmes attended twice weekly supervised sessions for eight weeks (16 sessions) and were encouraged to exercise at home for at least one additional session.[20] Three community programmes used cohort recruitment whereby patients attended once weekly supervision over 8 weeks (8 sessions) and were encouraged to exercise at home for at least two additional sessions, using a locally developed home exercise video.[21] Patients attended rehabilitation at a site of their choice. Once weekly cohort programmes were offered to enable attendance at a local venue where a rolling programme may not have been available.

Standardised exercise and education was delivered across all sites in accordance with guidelines.[20] Exercise consisted of cardiovascular and limb strengthening activities in line with individual baseline function. Walking was a component in all programmes and intensity was determined at 85% peak oxygen consumption from baseline walk testing.[22] Patients with a current prescription for ambulatory oxygen used it during exercise. Where evidence of desaturation to less than 85% was evident on exercise, supplementary oxygen was provided and those patients were offered further assessment of ambulatory oxygen need.[20]

Measures

Data collected included age, gender, ethnicity, postcode (to obtain the Index of Multiple Deprivation Score - IMD), MRC dyspnoea scale, referrer, forced expiratory volume in the first second (FEV₁).[23,24] The IMD score is based on national census and local authority data and reflects deprivation specific to a geographical area. IMD scores in 2007 ranged nationally from 0 (the least deprived) to 86 (the most deprived). For analysis the IMD scores were categorised into quintiles based on the scores of all patients at referral: 6.86-27.1; 27.2-34.4; 34.5-39.01; 39.02-43.4; 43.41-60.41. Age was categorised in four groups: 0-54; 55-64; 65- 74; 75 and over. The following measures were completed before and after a course of PR: Incremental Shuttle Walk Test (ISWT), Self-Reported Chronic Respiratory Disease Questionnaire (CRQ-SR) and the Hospital Anxiety and Depression Scale (HADS).[25-27]

For the purpose of analysis the ISWT was categorised into quartiles based on the scores recorded at baseline in the patients attending assessment: 0-130 metres; 131-220 metres ; 221-340 metres ;

341-1020 metres. The anxiety and depression components of the HAD scale were also categorised into three categories based on previous study in a general population. The first category, 0-7 is normal, the second category 8-10 represents “risk” of anxiety or “risk” of depression, and the third category 11 or more represents “caseness” for anxiety or depression.[27] Patients were categorised as “completers” if they had attended at least 8 sessions (50%) on a rolling recruitment programme or had attended at least 6 sessions (75%) on a cohort recruitment programme, irrespective of attendance at final assessment.

Analysis

Analysis was carried out using SPSS (SPSS Inc, Chicago). Predictors of attendance at assessment were investigated using univariate and multivariate logistic regression with the independent variables age, gender, referrer and IMD score. Course completion was investigated using univariate and multivariate logistic regression with the independent variables age, gender, referrer, IMD score, MRC score, Global Initiative for Chronic Obstructive Lung Disease (GOLD) stage (based on FEV1), and ISWT, CRQ and HADS at baseline. Because of the number of comparisons in the univariate logistic regression of completion, we set the threshold for statistical significance at p=0.01 to allow for the increased possibility of finding a significant association by chance. Effectiveness of PR was assessed by comparing mean scores in ISWT, CRQ and HADS before and after PR using paired t-tests and 95% confidence intervals. We assessed the outcome not only in the significance of the difference but also in whether the lower end of the 95% confidence interval for the difference was greater than the minimal clinically important difference (MCID) for the ISWT (47 metres) and each domain of the CRQ-SR (0.5) and for the HADS (anxiety 1.32 and depression 1.4).[28-30] We composed a binary outcome variable based on the MCID for the outcome measure ISWT and for each domain of the CRQ-SR to seek, using multiple logistic regression, predictors of effectiveness of PR among demographic variables and baseline measures of severity.

Results

1266 people with COPD were referred to pulmonary rehabilitation between April 2008 and March 2010 (mean age (SD) 68.1 (11.0) yrs; male 52%; IMD 35.8 (9.2)). Figure 1 shows the recruitment pathway for all referrals. 812 (73%) eligible patients attended for assessment. The course was completed by 441 patients (40% of those referred, 54% of those who attended assessment and 67% of those who started the course). 635 (57%) patients were referred from primary care (13% general practitioner (GP), 21% practice nurse, 23% community COPD clinic), 41% from secondary care (19%

respiratory physician outpatient clinics, 13% in-patient multidisciplinary COPD team, 9% in-patient physiotherapist), and 2% from other referrers.

Attendance at assessment

The characteristics of the 812 patients who attended assessment are shown in Table 1 together with the adjusted odds ratios (multivariate analysis) for their attendance compared to referred patients who did not attend. Patients were less likely to attend assessment if they were under 55 years or over 74 yrs, or were referred by the in-patient COPD team, hospital physiotherapist or specialist COPD community clinic. Gender and deprivation score did influence attendance at assessment.

Completion of Pulmonary Rehabilitation

The characteristics of patients who attended PR are shown in Table 2. Unadjusted odds ratios (univariate analysis) show differences in characteristics between those who completed and those who dropped out. Factors associated with lower rates of completion were: GP referral, second to lowest quintile of deprivation (not the lowest), MRC score 4 or 5, baseline ISWT distance of less than 220m, lower baseline CRQ score in the domains of fatigue, emotion and disease mastery, and a HAD anxiety or depression score of 11 and above. Type of programme (once weekly cohort or twice weekly rolling recruitment) was not associated with completion.

In multivariate analysis GP referral, depression score of 11 and above, MRC score 4 or 5, and higher deprivation remained independently associated with lower rates of completion (Table 3).

Effectiveness

Statistically significant improvements were evident overall and in both the twice weekly rolling and the once weekly cohort recruitment groups for ISWT, all domains of the CRQ, and HADS anxiety and depression scores (Table 4). The mean change and the lower limit of its 95% confidence interval in ISWT and all domains of the CRQ exceeded the minimal clinically important differences (MCID) in patients completing PR overall and in those in twice weekly rolling recruitment groups. The MCID of the HADS anxiety element was not reached but participants in the twice weekly rolling recruitment group reached the MCID for depression. In the once weekly cohort recruitment programme, the mean change in ISWT and CRQ also exceeded the MCIDs but the lower limit of the 95% confidence interval was less than the MCID with respect to the ISWT and the CRQ emotion, fatigue and mastery domains and higher than the MCID for CRQ dyspnoea.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

No consistent predictors of effectiveness were found across the outcome measures of the ISWT and the four domains of the CRQ-SR. For example, patients with MRC dyspnoea score of 5 at baseline were less likely to improve in ISWT more than 47 metres (OR 0.054, 95% CI 0.004-0.669) or in CRQ-SR Dyspnoea domain by more than 0.5 (OR 0.054, 95% CI 0.004-0.669) when adjusting for age, sex, referrer, IMD score, and GOLD stage. Yet these patients were no less likely to improve in CRQ-SR emotion, fatigue and mastery domains. Similar sporadic associations were observed with other demographic variables and baseline severity scores. Overall pulmonary rehabilitation was equally effective in patients with mild disease, in patients over seventy five years, and in patients from more deprived backgrounds.

Discussion

In this large observational evaluation of an integrated PR service for COPD across primary and secondary care, PR was effective in improving exercise capacity, reducing dyspnoea, improving quality of life, and reducing anxiety and depression. Seventy three percent of referred patients attended for assessment and 40% completed the course of treatment. This research confirms the findings of clinical trials in a real world setting with no prior selection of participants. It answers the criticism that has been made of some trials in which many patients suitable for PR programmes were excluded.[11] It demonstrates the effectiveness of PR in everyday clinical practice, but it also shows that the key obstacles to its delivery are to be found in the take-up and completion of the treatment by those referred.

Its strength comes from its large sample size, the absence of restrictive patient selection criteria, and its application over two years. The capacity of this PR programme has been in excess of 600 annual places for at least five years for a population of about 5000 COPD patients. Access is open to clinicians from primary and secondary care and the programme has been promoted widely. PR is shown here to be effective in all patients with COPD irrespective of age and socio-economic deprivation, in patients with moderate, severe, and very severe disease.[28,29] The conclusions that can be drawn are limited by the absence of a control group. Nonetheless the size of the improvement achieved does match that observed in trials.[1] The baseline ISWT was not preceded by a training walk but the improvement seen in the ISWT matched the improvement in dyspnoea and quality of life.

As with twice weekly rehabilitation, changes in outcomes in the once weekly cohort were statistically significant. However, the lower limits of the outcomes' 95% confidence intervals in the once weekly cohort only exceeded the MCID with respect to the dyspnoea element of the CRQ-SR. The analysis was not powered to look at differences between once and twice weekly provision of rehabilitation, and patients were not randomised to once only or twice only, but this finding gives more confidence in the advantage of twice weekly over once weekly attendance.

Attendance at assessment

Eligible referrals to PR came from primary and secondary care, with slightly more from the former. This reflects an important shift in primary care awareness of PR. We were limited in comparing attenders and non-attenders before assessment by the information provided by the referrer, a problem highlighted by Keating et al in their review.[13] We cannot explain why people below 55 or above 75 years were less likely to attend assessment. This may have reflected a greater likelihood of employment in the younger group, and poor mobility and co-morbidities in the older group. Poor attendance at assessment by patients referred from a specialist COPD community clinic or while they were in hospital may reflect more complex needs and more severe disease. Recent data shows an important benefit of rehabilitation after exacerbation leading to hospital admission, but recruitment during this acute phase is more difficult.[8,31] Rehabilitation providers may have to consider whether the needs of these patients are different.

Completion of rehabilitation

The rates of completion are similar to those observed in trial settings from the time of contact or screening. Completion rates after randomisation in trials of PR are usually about 75%, compared to 54% after assessment in participants in this study.[11] This is not surprising because participants in trials are generally less affected by co-morbidities and have usually signalled their commitment to participation by completing consent forms. Patients referred by general practitioners were as likely to attend assessment as patients referred by respiratory physicians in outpatient clinics, but they were less likely to complete rehabilitation. Further exploration of the methods of preparation of patients for PR may be justified. As with Fan et al and Garrod et al, we found that depression was a predictor of drop out. [15,32,33] While these factors together with deprivation and more severe disease were significantly associated with reduced completion, they only explained 10% of the variance (pseudo $r^2 = 0.103$) in completion rate after assessment.[34] This was despite the large

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

numbers in our multivariate analysis (n=657) and the extensive data on deprivation, source of referral, severity, quality of life, exercise capacity and mental health.

A new understanding is required of the reasons why nearly half of patients assessed for PR and 60% of those referred fail to complete the treatment. We know that the determinants of behaviour with respect to exercise are complex.[35] A new approach is required which seeks to understand from the patients' perspective why they fail to complete PR. Qualitative research may lead to the development of new hypotheses which could be tested in less reductionist fashion, perhaps adopting Bayesian statistical methods to allow for more complex relationships.

Conclusions

Despite our concerns about high levels of failure to attend assessment and to complete pulmonary rehabilitation, it is clear from this study that pulmonary rehabilitation is effective in the routine clinical care of COPD in those who complete the course. Clinicians from all sectors of health services should be able to refer their patients. They should be aware that rates of attendance at assessment may be 75% or less and the course of treatment may be completed by only 40% of all those referred. Providers should plan services that make allowance for these low rates of attendance and completion. There is no reason to suspect that age or socio-economic deprivation are factors that will prevent patients from taking advantage of what is one of the most useful treatments in the management of COPD.

Acknowledgements

Funder: Guy's and St Thomas' Charity, Reference no: G060703.

We are grateful to Paul Seed of King's College London for statistical advice, and to Dr Jack Barker of King's College Hospital NHS Foundation Trust who developed and manages the database on which all clinical data relating to pulmonary rehabilitation in King's College Hospital and in community services in Lambeth Southwark is collected

Declaration of Interest

The authors report no conflict of interest

References

Reference List

1. Lacasse Y, Goldstein R, Lasserson TJ, Martin S: **Pulmonary rehabilitation for chronic obstructive pulmonary disease**. *Cochrane Database Syst Rev* 2006, CD003793.
2. Ries AL, Bauldoff GS, Carlin BW, Casaburi R, Emery CF, Mahler DA *et al.*: **Pulmonary Rehabilitation: Joint ACCP/AACVPR Evidence-Based Clinical Practice Guidelines**. *Chest* 2007, **131**: 4S-42S.
3. National Clinical Guideline Centre. Chronic obstructive pulmonary disease: management of chronic obstructive pulmonary disease in adults in primary and secondary care. 1-673. 2010. London, National Clinical Guideline Centre.

Ref Type: Report

4. Qaseem A, Wilt TJ, Weinberger SE, Hanania NA, Criner G, Van Der Molen T *et al.*: **Diagnosis and Management of Stable Chronic Obstructive Pulmonary Disease: A Clinical Practice Guideline Update from the American College of Physicians, American College of Chest Physicians, American Thoracic Society, and European Respiratory Society**. *Ann Intern Med* 2011, **155**: 179-191.
5. Nici L, Donner C, Wouters E, Zuwallack R, Ambrosino N, Bourbeau J *et al.*: **American Thoracic Society/European Respiratory Society statement on pulmonary rehabilitation**. *Am J Respir Crit Care Med* 2006, **173**: 1390-1413.
6. Griffiths TL, Phillips CJ, Davies S, Burr ML, Campbell IA: **Cost effectiveness of an outpatient multidisciplinary pulmonary rehabilitation programme**. *Thorax* 2001, **56**: 779-784.
7. Griffiths TL, Burr ML, Campbell IA, Lewis-Jenkins V, Mullins J, Shiels K *et al.*: **Results at 1 year of outpatient multidisciplinary pulmonary rehabilitation: a randomised controlled trial**. *Lancet* 2000, **355**: 362-368.

8. Seymour JM, Moore L, Jolley CJ, Ward K, Creasey J, Steier JS *et al.*: **Outpatient pulmonary rehabilitation following acute exacerbations of COPD.** *Thorax* 2010, **65**: 423-428.

9. Yohannes A, Stone R, Lowe D, Pursey N, Buckingham R, Roberts C: **Pulmonary rehabilitation in the United Kingdom.** *Chron Respir Dis* 2011, **8**: 193-199.

10. Garrod R, Ford K, Daly C, Hoareau C, Howard M, Simmonds C: **Pulmonary rehabilitation: analysis of a clinical service.** *Physiother Res Int* 2004, **9**: 111-120.

11. Bjoernshave B, Korsgaard J, Nielsen CV: **Does pulmonary rehabilitation work in clinical practice? A review on selection and dropout in randomized controlled trials on pulmonary rehabilitation.** *Clin Epidemiol* 2010, **2**: 73-83.

12. Cote CG, Celli BR: **Pulmonary rehabilitation and the BODE index in COPD.** *Eur Respir J* 2005, **26**: 630-636.

13. Keating A, Lee A, Holland AE: **What prevents people with chronic obstructive pulmonary disease from attending pulmonary rehabilitation? A systematic review.** *Chron Respir Dis* 2011, **8**: 89-99.

14. Sabit R, Griffiths TL, Watkins AJ, Evans W, Bolton CE, Shale DJ *et al.*: **Predictors of poor attendance at an outpatient pulmonary rehabilitation programme.** *Respir Med* 2008, **102**: 819-824.

15. Garrod R, Marshall J, Barley E, Jones PW: **Predictors of success and failure in pulmonary rehabilitation.** *Eur Respir J* 2006, **27**: 788-794.

16. Nici L, Raskin J, Rochester CL, Bourbeau JC, Carlin BW, Casaburi R *et al.*: **Pulmonary rehabilitation: WHAT WE KNOW AND WHAT WE NEED TO KNOW.** *J Cardiopulm Rehabil Prev* 2009, **29**: 141-151.

17. Waterhouse JC, Walters SJ, Oluboyede Y, Lawson RA: **A randomised 2 x 2 trial of community versus hospital pulmonary rehabilitation, followed by telephone or conventional follow-up.** *Health Technol Assess* 2010, **14**: i-xi, 1.

18. Casas A, Troosters T, Garcia-Aymerich J, Roca J, Hernandez C, Alonso A *et al.*: **Integrated care prevents hospitalisations for exacerbations in COPD patients.** *Eur Respir J* 2006, **28**: 123-130.

19. Roberts JA, Maslin TK, Bakerly ND: **Development of an integrated chronic obstructive pulmonary disease service model in an inner-city region in the UK: initial findings and 12-month results.** *Prim Care Respir J* 2010, **19**: 390-397.

20. British Thoracic Society Standards of Care Sub-committee on Pulmonary Rehabilitation: **Pulmonary rehabilitation.** *Thorax* 2001, **56**: 827-834.

21. Moore J, Fiddler H, Seymour J, Grant A, Jolley C, Johnson L *et al.*: **Effect of a home exercise video programme in patients with chronic obstructive pulmonary disease.** *J Rehabil Med* 2009, **41**: 195-200.

22. Singh SJ, Morgan MD, Hardman AE, Rowe C, Bardsley PA: **Comparison of oxygen uptake during a conventional treadmill test and the shuttle walking test in chronic airflow limitation.** *Eur Respir J* 1994, **7**: 2016-2020.

23. Noble M, McLennon D, Wilkinson K, Whitworth A, Barnes H, Dibben C: **The English Indices of Deprivation 2007**. London, UK: Department for Communities and Local Government; 2008.
24. Bestall JC, Paul EA, Garrod R, Garnham R, Jones PW, Wedzicha JA: **Usefulness of the Medical Research Council (MRC) dyspnoea scale as a measure of disability in patients with chronic obstructive pulmonary disease**. *Thorax* 1999, **54**: 581-586.
25. Singh SJ, Morgan MD, Scott S, Walters D, Hardman AE: **Development of a shuttle walking test of disability in patients with chronic airways obstruction**. *Thorax* 1992, **47**: 1019-1024.
26. Williams JE, Singh SJ, Sewell L, Guyatt GH, Morgan MD: **Development of a self-reported Chronic Respiratory Questionnaire (CRQ-SR)**. *Thorax* 2001, **56**: 954-959.
27. Bjelland I, Dahl AA, Haug TT, Neckelmann D: **The validity of the Hospital Anxiety and Depression Scale. An updated literature review**. *J Psychosom Res* 2002, **52**: 69-77.
28. Singh SJ, Jones PW, Evans R, Morgan MD: **Minimum clinically important improvement for the incremental shuttle walking test**. *Thorax* 2008, **63**: 775-777.
29. Schunemann HJ, Puhan M, Goldstein R, Jaeschke R, Guyatt GH: **Measurement properties and interpretability of the Chronic respiratory disease questionnaire (CRQ)**. *COPD* 2005, **2**: 81-89.
30. Puhan MA, Frey M, Buchi S, Schunemann HJ: **The minimal important difference of the hospital anxiety and depression scale in patients with chronic obstructive pulmonary disease**. *Health Qual Life Outcomes* 2008, **6**: 46.
31. Puhan MA, Scharplatz M, Troosters T, Steurer J: **Respiratory rehabilitation after acute exacerbation of COPD may reduce risk for readmission and mortality -- a systematic review**. *Respir Res* 2005, **6**: 54.
32. Fan VS, Giardino ND, Blough DK, Kaplan RM, Ramsey SD: **Costs of pulmonary rehabilitation and predictors of adherence in the national emphysema treatment trial**. *COPD* 2008, **5**: 105-116.
33. Fan VS, Giardino ND, Blough DK, Kaplan RM, Ramsey SD: **Costs of pulmonary rehabilitation and predictors of adherence in the national emphysema treatment trial**. *COPD* 2008, **5**: 105-116.
34. McFadden DL: **Conditional logit analysis of qualitative choice behavior**. In *Frontiers in Econometrics*. Edited by Zarembka P. New York: Academic Press; 2011:105-142.
35. van Sluijs EM, Griffin SJ, van Poppel MN: **A cross-sectional study of awareness of physical activity: associations with personal, behavioral and psychosocial factors**. *Int J Behav Nutr Phys Act* 2007, **4**: 53.

Table 1. Characteristics of attenders at assessment for pulmonary rehabilitation.

(Adjusted odds ratios for association with attendance)

Characteristic	Attended assessment n (% of all referred)	Adjusted odds ratios† (95% CI) n = 1084
Age (years)		
0-54	83 (60)	1
55-64	225 (77)	2.17 (1.38-3.44) ^a
65-74	276 (78)	2.24 (1.43-3.5) ^a
75+	228 (69)	1.49 (0.96-2.3)
Gender		
Female	382 (71)	1
Male	430 (74)	0.92 (0.7-1.22)
Referrer		
Consultant respiratory physician	168 (80)	1
In-patient COPD multidisciplinary team	96 (63)	0.45 (0.28-0.73) ^a
In-patient physiotherapist	60 (61)	0.43 (0.25-0.75) ^a
GP	110 (78)	0.94 (0.55-1.61)
Practice nurse	180 (76)	0.78 (0.49-1.24)
Community COPD clinic	181 (72)	0.63 (0.4-0.98) ^a
Other	17 (71)	0.67 (0.26-1.75)
Deprivation quintiles (IMD)		
6.86-27.1	137 (72)	1
27.2-34.4	174 (80)	1.3 (0.81-2.06)
34.5-39.01	153 (69)	0.8 (0.52-1.23)
39.02-43.4	164 (77)	1.18 (0.75-1.84)
43.41-60.41	162 (67)	0.76 (0.49-1.16)

^a = significant effect. † Adjusted for other variables in the table.

Pseudo R² (McFadden 1974) for multiple logistic regression = 0.064. Describes how well (6.4%) the model performs when compared to a perfect prediction model.[34]

Table 2. Characteristics of patients who completed a pulmonary rehabilitation course and of those who dropped out.

(Unadjusted odds ratios for association with completion)

	Attended Assessment n=812		Unadjusted odds ratio (95% CI)	p value^a
Characteristic	Drop Out^b n=371 (%)	Completer n=441 (%)		
Age:			1	
up to 54 years	44 (11.9)	39 (8.8)		
55 to 64 years	113 (30.5)	112 (25.4)	1.12 (0.08-1.85)	0.66
65 to 74 years	108 (29.1)	168 (38.1)	1.76 (1.07-2.88)	0.026
75 years +	106 (28.6)	122 (27.7)	1.3 (0.79-2.15)	0.309
Sex:				
Male (%)	182 (49.1)	248 (56.2)	1.33 (1.01-1.76)	0.04
Source of referral:			1	
Consultant respiratory physician	71 (19.1)	97 (22.0)		
In-patient COPD team	44 (11.9)	52 (11.8)	0.87 (0.52-1.43)	0.574
In-patient physiotherapist	32 (8.6)	28 (6.3)	0.64 (0.35-1.16)	0.14
GP	65 (17.5)	45 (10.2)	0.51 (0.31-0.83)	0.006 ^c
Practice nurse	69 (18.6)	111 (25.2)	1.17 (0.77-1.81)	0.46
Community COPD clinic	80 (21.6)	101 (22.9)	0.92 (0.61-1.41)	0.72
Other	10 (2.7)	7 (1.6)	0.51 (0.77-1.81)	0.2
IMD score of deprivation:				
Quintiles:			1	
6.86-28.1	63 (17.2)	97 (22.5)		
28.11-35.02	75 (20.4)	92 (21.3)	0.8 (0.51-1.24)	0.31
35.03-39.57	60 (16.3)	92 (21.3)	0.99 (0.63-1.57)	0.99
39.58-43.85	90 (24.5)	78 (18.1)	0.56 (0.36-0.87)	0.01 ^c
43.86-60.41	79 (21.5)	72 (16.7)	0.59 (0.38-0.93)	0.02
Ethnicity:	n=282	n=409	1	
White British & Irish	255 (90.4)	347 (84.8)		
White Other	3 (1.1)	21 (5.1)	5.14 (1.52-17.43)	0.009 ^{c,e}
Black	16 (5.7)	31 (7.6)	1.42 (0.76-2.66)	0.27
Asian sub-continent ^d	7 (2.5)	6 (1.5)	0.63 (0.21-1.9)	0.41
Other	1 (0.4)	4 (1.0)	2.94 (0.33-26.46)	0.34
MRC Dyspnoea:	n=308	n=426	1	
1 and 2	46 (14.9)	93 (21.8)		
3	102 (33.1)	168 (39.4)	0.78 (0.51-1.18)	0.24
4	128 (41.6)	140 (32.9)	0.52 (0.34-0.78)	0.002 ^c
5	30 (9.7)	16 (3.8)	0.25 (0.13-0.5)	<0.001 ^c
Gold Stage n (%)	n=222	n=285	1	
I	16 (7.2)	27 (9.5)		
II	85 (38.3)	113 (39.6)	1.02 (0.6-1.75)	0.95
III	88 (39.6)	106 (37.2)	1.13 (0.65-1.94)	0.67
IV	33 (14.9)	39 (13.7)	1.43 (0.66-3.09)	0.37
Baseline ISWT:	n=252	n=390		
Quartiles:			1	
0-130m	83 (32.9)	82 (21.0)		
131-220m	75 (29.8)	92 (23.6)	1.24 (0.8-1.91)	0.33
221-340m	44 (17.5)	108 (27.7)	2.48 (1.56-3.95)	<0.001 ^c
341-1020m	50 (19.8)	108 (27.7)	2.19 (1.39-3.44)	0.001 ^c
CRQ Dyspnoea:	n=278	n=406		
Mean (SD)	2.54 (1.14)	2.74 (1.20)	1.16 (1.02-1.33)	0.029
CRQ Emotion:	n=281	n=407		
Mean (SD)	3.69 (1.39)	4.21 (1.40)	1.31 (1.17-1.47)	<0.001 ^c
CRQ Fatigue:	n=282	n=406		
Mean (SD)	3.01 (1.29)	3.44 (1.37)	1.27 (1.13-1.43)	<0.001 ^c
CRQ Mastery:	n=283	n=407		

Integrated pulmonary rehabilitation for COPD

Mean (SD)	3.83 (1.49)	4.33 (1.41)	1.27 (1.14-1.42)	<0.001 ^c
HAD Anxiety: n (%)	n=285	n=403		
<=7	113 (36.9)	204 (50.6)	1	
8-10	66 (23.2)	93 (23.1)	0.78 (0.52-1.15)	0.21
>11	106 (37.2)	106 (26.3)	0.55 (0.39-0.79)	0.001 ^c
HAD Depression: n (%)	n=283	n=403		
<7=	131 (46.3)	246 (61.0)	1	
8-10	67 (23.7)	85 (21.1)	0.68 (0.46-0.99)	0.045
>11	85 (30.0)	72 (17.9)	0.45 (0.31-0.66)	<0.001 ^c
Programme type: n (%)	n=371	n=441		
Cohort recruitment	113 (30.5)	133 (30.2)	1	
Rolling recruitment	258 (69.5)	308 (69.8)	1.01 (0.75-1.37)	0.93

^a The large number of tests of association increases the risk of finding a significant association by chance. We have therefore increased the threshold for significance by only accepting p values of 0.01 or less.

^b Drop out = patient declined at assessment, did not start PR course or started but did not complete course.

^c = significant result

^d Asian sub-continent = Bangladesh, Bhutan, India, Nepal, Pakistani, Sri Lanka

^e The importance of the significant finding with respect to White Other subjects is doubtful because the group consisted of only 3 participants

Table 3. Characteristics of patients who completed pulmonary rehabilitation compared with those who dropped out (adjusted).

Characteristic	Adjusted odds ratio for completion of the course ^a (95% CI) n=657
Referrer	
Consultant respiratory physician	1
In-patient COPD multidisciplinary team	0.87 (0.48-1.57)
Hospital physiotherapist	0.55 (0.27-1.15)
GP	0.42 (0.24-0.74) ^b
Practice nurse	0.89 (0.54-1.48)
Community COPD clinic	0.82 (0.49-1.35)
Other	0.55 (0.15-2.07)
MRC dyspnoea score	
1 or 2	1
3	0.88 (0.55-1.41)
4	0.61 (0.37-0.97) ^b
5	0.39 (0.16-0.93) ^b
Depression score (HADS)	
(Not depressed) 0-7	1
(Risk of depression) 8-11	0.77 (0.51-1.18)
(Depressed) >11	0.56 (0.37-0.85) ^b
Deprivation quintiles (IMD)	
IMD score 6.86-28.1	1
28.11-35.02	0.72 (0.43-1.2)
35.03-39.57	1.0 (0.59-1.7)
39.58-43.85	0.56 (0.33-0.94) ^b
43.86-60.41	0.57 (0.34-0.85) ^b

^a adjusted for other variables in table and age and sex

^b significant effect .

Pseudo R² (McFadden 1974) for multiple logistic regression = 0.103. Describes how well (10.3%) the model performs when compared to a perfect prediction model.[34]

Table 4. Changes over time in exercise tolerance, health-related quality-of-life, and anxiety and depression scores.

(Rolling and cohort programmes shown separately)

	All completers		Rolling		Cohort	
	n	Mean change (CI)	n	Mean change (CI)	n	Mean change (CI)
ISWT (m)	311	68.3 (59.3-77.4)	205	74.2 (63.5-85.0)	106	56.9 (40.3-73.2)
CRQ Dyspnoea	329	0.94 (0.80-1.07)	214	0.99 (0.82-1.15)	115	0.84 (0.60-1.09)
CRQ Emotion	334	0.64 (0.52-0.76)	218	0.66 (0.50-0.82)	116	0.61 (0.41-0.81)
CRQ Fatigue	333	0.7 (0.58-0.83)	217	0.74 (0.57-0.90)	116	0.64 (0.44-0.84)
CRQ Mastery	333	0.71 (0.58-0.84)	217	0.71 (0.54-0.87)	116	0.71 (0.49-0.92)
HAD Anxiety	327	0.9 (0.5-1.2)	212	0.9 (0.5-1.3)	115	0.8 (0.2-1.3)
HAD Depression	328	1.1 (0.8-1.4)	212	1.4 (0.9-1.8)	116	0.6 (0.1-1.0)

Minimally clinically important differences: ISWT 0.47m; CRQ-SR all domains 0.5; HADS anxiety 1.32, depression 1.4.
All p values < 0.001 with the exception of change in the Anxiety (p=0.009) and Depression (p = 0.02) scores for cohort programmes.

Figure Legends

**Figure 1. Recruitment pathway of all PR referrals received April 2008 – March 2010.
Percentages based on eligible referrals (n=1114).**

For Peer Review Only

Figure 1

Figure Legends

Figure 1. Recruitment pathway of all PR referrals received April 2008 – March 2010. Percentages based on eligible referrals (n=1114).

For Peer Review Only