King's Research Portal

DOI:

[10.1136/jnnp-2016-313356](https://doi.org/10.1136/jnnp-2016-313356)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Bray, B. D., Smith, C. J., Cloud, G. C., Enderby, P., James, M., Paley, L., Tyrrell, P. J., Wolfe, C. D. A., & Rudd, A. (2016). The association between delays in screening for and assessing dysphagia after acute stroke, and the risk of stroke-associated pneumonia. *Journal of Neurology, Neurosurgery and Psychiatry*.
<https://doi.org/10.1136/jnnp-2016-313356>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

The association between delays in screening for and assessing dysphagia after acute stroke and the risk of stroke-associated pneumonia

Benjamin D Bray MD MRCP (1)

Craig J Smith MD MRCP (3)

Geoffrey C Cloud FRCP (4)

Pam Enderby PhD FRCSLT (5)

Martin James MD FRCP (6)

Lizz Paley BA (7)

Pippa J Tyrrell MD FRCP (3)

Charles DA Wolfe MD FFPHM (2,8)

Anthony G Rudd FRCP (2,8)

On behalf of the SSNAP Collaboration

(1) Farr Institute of Health Informatics, University College London, UK

(2) Division of Health and Social Care Research, King's College London, UK

(3) Stroke and Vascular Research Centre, Institute of Cardiovascular Sciences, University of Manchester, Manchester Academic Health Science Centre, Manchester, UK

(4) St George's University Hospitals NHS Foundation Trust, London, UK

(5) School of Health and Related Research, University of Sheffield, Sheffield, UK

(6) Royal Devon and Exeter NHS Foundation Trust, Exeter, UK

(7) Royal College of Physicians, London, UK

(8) National Institute for Health Research Comprehensive Biomedical Research Centre, Guy's and St Thomas' NHS Foundation Trust, London, UK

Corresponding author:

Dr Benjamin Bray

Farr Institute of Health Informatics

222 Euston Road

London

Email: benjamin.bray@kcl.ac.uk

1 Fax: +44 (0)207 848 6620

2 Phone: 0207 848 6604

3

4

Abstract

Background

There is no robust evidence that screening patients with acute stroke for dysphagia reduces the risk of stroke-associated pneumonia (SAP), or of how quickly it should be done after admission. We aimed to identify if delays in bedside dysphagia screening and comprehensive dysphagia assessments by a speech and language therapist (SALT) were associated with patients' risk of SAP.

Methods

Nationwide registry based prospective cohort study of patients admitted with acute stroke in England and Wales. Multilevel multivariable logistic regression models were fitted, adjusting for patient variables and stroke severity. The exposures were time from 1) admission to bedside dysphagia screen, and 2) admission to comprehensive dysphagia assessment.

Results

Of 63650 patients admitted with acute stroke, 55838 (88%) had a dysphagia screen and 24542 (39%) a comprehensive dysphagia assessment. Patients with the longest delays in dysphagia screening (4th quartile adjusted OR 1.14, 1.03-1.24) and SALT dysphagia assessment (4th quartile adjusted OR 2.01, 1.76-2.30) had a higher risk of SAP. The risk of SAP increased in a dose-response manner with delays in SALT dysphagia assessment, with an absolute increase of pneumonia incidence of 1% per day of delay.

Conclusion

Delays in screening for and assessing dysphagia after stroke, are associated with higher risk of stroke-associated pneumonia. Since stroke-associated pneumonia is one of the main causes of mortality after acute stroke, early dysphagia assessment may contribute to preventing deaths from

- 1 acute stroke and could be implemented even in settings without access to high technology specialist
- 2 stroke care.

3

Introduction

Stroke-associated pneumonia (SAP) is a common complication of acute stroke, affecting 6-10% of patients¹. SAP independently increases the risk of mortality and is one of the main causes of death in the first few days and weeks after stroke². It is also associated with worse functional outcomes, longer length of stay and increased healthcare costs^{3,4,5,6,7,8}. One of the main risk factors for SAP is dysphagia, which affects 37-55% of patients⁹ after stroke. Dysphagia screening using a brief bedside screening tool (such as a water swallow test), and comprehensive clinical assessments of aspiration risk by speech and language therapists (SALT), are therefore performed commonly in stroke care. Typically, all appropriate patients are screened for dysphagia and those in whom dysphagia is suspected go on to receive a comprehensive assessment. Despite being well established in clinical practice, there is however very little evidence of effectiveness for these interventions. Previous studies have generally used weak designs and provided no information to guide recommendations on how quickly dysphagia assessments after stroke should occur^{10,11}. Swallowing assessment prior to eating and drinking is recommended in European¹², United States¹³ and United Kingdom clinical guidelines but none recommend a specific approach to assessment or treatment of dysphagia.

As SAP develops most frequently within the first 7 days of stroke¹, the timing of both dysphagia screening and SALT assessment after admission are likely to be of importance. In England and Wales, dysphagia screening is recommended within 4 h of admission for acute stroke, and comprehensive assessment by a SALT (if required) within 72 h of admission¹⁴. The aim of this study was to investigate the association between delays in dysphagia screening and SALT assessment and the incidence SAP within the first 7 days after admission. The hypothesis was that delays in these assessments would be associated with an increased incidence of SAP.

Methods

Data were from the Sentinel Stroke National Audit Programme (SSNAP), the national register of stroke in England and Wales, of patients aged ≥ 16 years admitted with acute stroke (ischaemic stroke or primary intracerebral haemorrhage) between April 2013 and March 2014. SSNAP is a prospective continuous register with participation from all hospitals admitting adults with stroke in England and Wales and is estimated to include 90-95% of all stroke admissions¹⁵. Ethical approval of SSNAP was granted by the Ethics and Confidentiality Committee of the National Information Governance Board. Mortality data were obtained through data linkage with the statutory register of deaths. Data linkage was carried out by a third party and the investigators used an anonymised dataset with all patient identifiers removed.

Dysphagia screening was defined as use of a bedside swallow screening test by an appropriately trained clinician (typically a trained nurse). The exact dysphagia screening protocol was not specified by SSNAP. The times from admission to documented dysphagia screen and comprehensive assessment by SALT were recorded to the nearest minute for all patients in whom these were carried out. For patients who had a stroke as an inpatient, the time from stroke onset was used instead of time of admission. All patients without clinical exclusions (e.g. being treated palliatively only) were eligible for dysphagia screening, and eligibility for comprehensive dysphagia assessment was determined by clinical indications, such as a positive dysphagia screen or clinical suspicion of dysphagia. Patients (n=965) admitted directly to an intensive care unit (ICU) on admission were excluded from the primary analysis, since most of these patients would have been intubated.

SAP was defined as the administration of antibiotics for a new clinical diagnosis of pneumonia in the first seven days after admission and was determined by the treating physician.

Statistical analysis

1 The adjusted odds of SAP were estimated by fitting multivariable logistic regression models. Time
2 from admission to dysphagia screen and SALT dysphagia assessment were analysed both as
3 continuous variables and by division into quartiles. When included as a continuous variable they
4 were fitted as restricted cubic regression splines using the POSTRCSPLINE module¹⁶. Spline
5 coefficients cannot be interpreted directly and so the models were displayed graphically, showing
6 the modelled association (and 95% confidence interval) between time to dysphagia screen or
7 comprehensive dysphagia assessment and estimated adjusted SAP incidence. Multilevel
8 multivariable logistic regression models were also fitted using quartiles of these times to enable
9 quantification of the study results into odds ratios and also to account for clustering at the hospital
10 level. These models were specified as two level models with hospital level random intercepts.

11 All models included age, sex, stroke subtype (ischaemic, primary intracerebral haemorrhage, or
12 undetermined), pre stroke functional level using the modified Rankin Scale (mRS), place of stroke
13 (out of hospital or inpatient), vascular comorbidity (heart failure, hypertension, atrial fibrillation,
14 diabetes mellitus, previous stroke or transient ischaemic attack) and either NIH Stroke Scale (NIHSS)
15 or level of consciousness on admission.

16 Data were complete for all data items apart from the NIHSS on admission, which was available for
17 73% of patients. Models were therefore also fitted using level of consciousness on admission as a
18 proxy for severity (available for 100% of patients) and the results of the models compared to explore
19 the effect of these missing data on the results.

20 We carried out several sensitivity analyses. Firstly, competing risk from early mortality was explored
21 by excluding patients dying or starting palliative care in the first 3 days. Secondly, models were fitted
22 including a variable indicating change (increase, no change, decrease) in the level of consciousness in
23 the first seven days after stroke, to explore the possible confounding effect of changing
24 consciousness level. Thirdly, ICU patients excluded in the main analysis were included in a complete
25 data set analysis. Fourthly, time from admission to stroke unit admission was included in the models

1 as a possible confounder. Finally, we fitted models of 30 day all cause mortality, excluding patients
2 dying or starting palliative care in the first 3 days (on the grounds that death in these latter patients
3 is more likely due directly to brain injury from the stroke rather than stroke associated pneumonia).
4 These models explored whether delays in dysphagia screening and assessment were associated with
5 mortality after stroke.

1 Results

2 There were 63650 patients with acute stroke included in the cohort, admitted to 199 hospitals. Of
3 these, 55838 (87.7%) had a dysphagia screen, and 24542 (38.6%) proceeded to a comprehensive
4 assessment. The characteristics of the whole cohort, and the subgroups of patients according to
5 receipt of dysphagia screening and comprehensive dysphagia assessment are described in Table 1.
6 Patients in whom a dysphagia screen was not performed had a greater incidence of inpatient stroke
7 and primary intracerebral haemorrhage, lower level of consciousness on admission and were less
8 likely to have NIHSS on admission completed.

9

	Cohort	Dysphagia screening not performed	Dysphagia screening performed	Comprehensive dysphagia assessment
<i>n (%)</i>	63650	7812 (12.3)	55838 (87.8)	24542 (38.6)
Median age(IQR)	77 (67-85)	80 (70-86)	77 (67-85)	80 (70-87)
Female (n, %)	32054 (50.4)	4264 (54.5)	27790 (49.8)	13160 (53.6)
Stroke type (n, %)				
Ischaemic	56167 (88.2)	5948 (76.1)	50219 (91.0)	21751 (89.6)
Primary intracerebral haemorrhage	6575 (10.3)	1592 (20.4)	4983 (8.9)	2523 (10.4)
Undetermined	908 (1.4)	272 (3.5)	636 (1.1)	268 (1.1)
Inpatient stroke (n, %)	3155 (5.0)	974 (12.5)	2181 (3.9)	1599 (6.5)
Pre-Stroke mRS (n,%)				
0	36208 (57.9)	3808 (48.8)	32400 (58.0)	12174 (49.6)
1	9726 (15.3)	1109 (14.2)	8617 (15.4)	3919 (16.0)
2	6036 (9.5)	851 (10.9)	5185 (9.3)	2614 (10.7)
3	6708 (10.5)	1073 (13.7)	5635 (10.1)	3163 (12.9)
4	3734 (5.9)	670 (8.6)	3064 (5.5)	1988 (8.1)
5	1248 (2.0)	301 (3.9)	937 (1.7)	684 (2.8)
Admission NIHSS complete (n,%)	46447 (73.0)	3792 (48.4)	42655 (76.4)	17041 (69.4)
Median admission NIHSS (IQR)	4 (2-9)	5 (2-19)	4 (2-9)	7 (3-14)
Level of consciousness on admission				
0 (Alert)	53433 (84.0)	4961 (63.5)	48472 (86.8)	19156 (78.1)
1 (Not alert: Responds to voice)	6032 (9.5)	1069 (13.7)	4963 (8.9)	3697 (15.1)
2 (Not alert: Responds to pain)	2498 (3.9)	828 (10.6)	1670 (3.0)	1267 (5.2)
3 (Totally unresponsive)	1687 (2.7)	954 (12.2)	733 (1.3)	422 (1.7)
Co-morbidity (n,%)				
Heart failure	3463 (5.4)	491 (6.3)	2972 (5.3)	1625 (6.6)
Hypertension	34212 (53.9)	3930 (50.3)	30382 (54.4)	13323 (54.3)
Atrial fibrillation	13159 (20.7)	1801 (23.1)	11358 (20.3)	5929 (24.2)
Diabetes mellitus	12372 (19.4)	1482 (18.9)	10890 (19.5)	4720 (19.2)
Previous stroke/TIA	17626 (27.7)	2144 (27.4)	15482 (27.7)	7109 (29.0)
Time from onset to admission (n, %)				
Unknown (e.g. wake up stroke)	24668 (38.8)	4233 (54.2)	20435 (36.6)	9631 (39.2)
0-179 minutes	21504 (33.8)	2113 (27.1)	19391 (34.7)	9086 (37.0)
180-359 minutes	6144 (9.7)	539 (6.9)	5605 (10.0)	2289 (85.6)
360+ minutes	11334 (17.8)	927 (11.9)	10407 (18.6)	3536 (14.4)
Thrombolysis (n, %)	7087 (11.1)	417 (5.3)	6670 (12.0)	3415 (3.1)
SAP (n,%)	5533 (8.7)	1077 (13.8)	4456 (8.0)	3592 (14.6)
30 day mortality (n,%)	8397 (13.2)	2701 (34.6)	5696 (10.2)	3599 (14.7)

1

2 **Table 1. Characteristics of the study cohort**

1

2 The overall incidence of SAP was 8.7%. SAP incidence was highest in the dysphagic group referred
3 for comprehensive dysphagia assessment (14.6%). Thirty day mortality was 13.2% overall, 10.2% in
4 patients screened for dysphagia, 14.7% in patients referred for SALT assessment, and 34.6% in
5 patients in whom a dysphagia screen was not carried out.

6 The median time from admission to dysphagia screening was 2.9 hours (IQR 1.3-5.7 hours) and for
7 comprehensive dysphagia assessment was 22.9 hours (IQR 6.2-49.4 hours). In unadjusted analyses
8 there was a strong association between time from admission to dysphagia screen and incidence of
9 SAP, rising from 7-8% from 0-8 hours and increasing to 15% by 72 hours after admission. Although
10 the association was attenuated after adjusting for patient characteristics, there was still a modest
11 association (equating to approximately 1% absolute increase in the incidence of SAP) between
12 delays in dysphagia screening and incidence of SAP (Figure 1). After adjustment, patients in the
13 fourth quartile (i.e. those with the longest delays in dysphagia screening) had 36% higher odds of
14 SAP compared to those in the first quartile (aOR 1.36, 1.20-1.53) (Table 2).

15

		Time (mins)	OR	95% CI	p
Univariable (n=55838)	1st quartile	0-79	REF		
	2nd quartile	80-176	0.89	0.81-0.98	0.016
	3rd quartile	177-344	0.85	0.77-0.94	0.001
	4th quartile	≥345	1.33	1.21-1.46	<0.0001
Multivariable, including NIHSS (n=42655)	1st quartile	0-79	REF		
	2nd quartile	80-176	0.94	0.83-1.05	0.27
	3rd quartile	177-344	1.06	0.94-1.20	0.36
	4th quartile	≥345	1.36	1.20-1.53	<0.0001
Multivariable, including level of consciousness (n=55838)	1st quartile	0-79	REF		
	2nd quartile	80-176	0.92	0.83-1.01	0.08
	3rd quartile	177-344	0.89	0.81-0.99	0.03
	4th quartile	≥345	1.14	1.03-1.24	0.008

Table 2. Odds ratio for SAP in univariable and multivariable models of time from admission to dysphagia screening. All multivariable models were also adjusted for age, sex, stroke type, pre-stroke functional level, place of stroke and comorbidity, and measure of stroke severity (NIHSS or level of consciousness)

There was a strong relationship between delays in comprehensive dysphagia assessment and incidence of SAP, and delays in comprehensive dysphagia assessment were associated with an absolute increase in the risk of SAP of 3% over the first 24 hours (Figure 2). Delays in SALT dysphagia assessment beyond 24 hours were associated with an additional 4% absolute increase in the incidence of SAP (approximately 3-fold increase in the relative incidence). Patients in the slowest quartile had 1.98 (1.67-2.35) the odds of SAP compared to patients receiving the quickest SALT dysphagia assessments (Table 3). Findings were similar in the sensitivity analyses (Supplementary material). The secondary analysis of 30 day mortality broadly supported these findings: there was very weak evidence that delays in dysphagia assessment were associated with an increase in 30 day mortality (aOR 1.14, 0.99-1.30 in the slowest quarter). There was moderately strong evidence that delays in comprehensive SLT assessment were associated with an increase in mortality risk in the second (aOR 1.22, 1.02-1.47), third (aOR 1.55, 1.29-1.85) and fourth (aOR 1.35, 1.12-1.63) quarters

- 1 of time to assessment respectively. Unlike pneumonia, a dose-response relationship was not
- 2 demonstrated for the association with mortality (Supplementary material)

		Time (mins)	OR	95% CI	p
Univariable (n=24542)	1st quartile	0-369	REF		
	2nd quartile	370-1371	1.53	1.34-1.74	<0.0001
	3rd quartile	1372-2961	1.95	1.71-2.22	<0.0001
	4th quartile	≥2962	2.65	2.33-3.01	<0.0001
Multivariable, including NIHSS (n=17041)	1st quartile	0-369	REF		
	2nd quartile	370-1371	1.35	1.15-1.60	<0.0001
	3rd quartile	1372-2961	1.61	1.37-1.91	<0.0001
	4th quartile	≥2962	1.98	1.67-2.35	<0.0001
Multivariable, including level of consciousness (n=24542)	1st quartile	0-369	REF		
	2nd quartile	370-1371	1.40	1.22-1.60	<0.0001
	3rd quartile	1372-2961	1.60	1.41-1.84	<0.0001
	4th quartile	≥2962	2.01	1.76-2.30	<0.0001

Table 3. Odds ratio for SAP in univariable and multivariable models of time from admission to SALT dysphagia assessment. All multivariable models were also adjusted for age, sex, stroke type, pre-stroke functional level, place of stroke and comorbidity, and measure of stroke severity (NIHSS or level of consciousness)

Discussion

In this national cohort of unselected stroke patients, we found that there was evidence of a modest association between delays in performing dysphagia screening and the risk of SAP. There was stronger evidence for an association between the risk of SAP and delays in carrying out a comprehensive dysphagia assessment. Although limited by the risk of residual confounding, these findings provide the first evidence from a large multicentre cohort that prompt dysphagia screening and comprehensive dysphagia assessment stroke are associated with clinically significant reductions in the risk of SAP, one of the principal causes of early death after stroke.

Detecting dysphagia through the use of bedside screening assessments and comprehensive dysphagia assessments carried out by a SALT is widely recommended in clinical guidelines¹²⁻¹⁴. However, these recommendations are largely based on consensus and there is little direct evidence for dysphagia screening or assessment after stroke¹⁷. Previous studies have been limited to ecological studies demonstrating an association between site level rates of screening assessment and SAP rates after stroke^{8,10,11}. Several observational studies have described an association between dysphagia screening at any time after stroke and reduced post stroke mortality^{10,18,19,20}, and dysphagia screening was a component of a stroke care bundle found in a cluster randomised controlled trial to reduce death and dependency after stroke²¹. By contrast, an analysis of the “Get with the Guidelines – Stroke” registry data from the USA found that dysphagia screening was associated with a higher risk of SAP, although the results suggest that confounding by stroke severity contributed to the observed association²². There is no current evidence of how quickly dysphagia assessment should occur after stroke or good quality evidence of whether dysphagia assessment reduces the risk of SAP. As a result of this lack of evidence, dysphagia screening has been dropped from the list of stroke quality indicators used in the USA²³, although they remain part of quality indicators used in the UK²⁴.

1 If our findings represent causal effects, then they imply that dysphagia screening and assessment is
2 effective in reducing the risk of SAP. Since SAP is one of the main causes of death in acute stroke,
3 reducing the risk of SAP would be expected to lead to reduced mortality after stroke. The secondary
4 analyses of mortality in this study provide supporting evidence that this might be the case. We
5 would however emphasise caution in interpreting the mortality findings - there are many causes of
6 death in acute stroke and so reductions in SAP will only prevent a proportion of deaths (in keeping
7 with the reduced effect sizes for mortality we observed) and we did not observe a dose-response
8 relationship between delays in comprehensive dysphagia assessment and mortality, suggesting that
9 there might be additional confounding or bias not accounted for in the analysis.

10 There are several possible mechanisms for why delays in dysphagia assessment might lead to an
11 increased risk of SAP and further studies would be required to test these hypotheses and confirm
12 (or refute) a causal relationship: early screening may reduce the risk of inappropriate administration
13 of oral fluid or food, prompt measures to reduce aspiration risk through positioning, nursing care
14 and appropriate feeding strategies, and avoid unnecessary nasogastric tube insertions. As well as
15 exploring mechanisms, further research might usefully also explore organisational aspects of
16 dysphagia assessment, such as the use of specific assessment and treatment protocols and the
17 relationship between specialist SALT provision on stroke units and patient outcomes.

18 These data are strengthened by being drawn from a national register of unselected patients,
19 reducing the risk of selection bias. Similarly, the study used clinical rather than administrative data,
20 providing more detail than would be available from routine administrative data alone. There are
21 however a number of limitations. Firstly SAP was not defined by specific criteria but was based on
22 the judgement of the treating physician, and we did not have information on the date of diagnosis of
23 SAP and whether it occurred before or after dysphagia screening and assessment. Nonetheless, the
24 overall rate of SAP observed in this study is consistent with other studies, suggesting that differences
25 in ascertainment between centres was not a significant source of bias. In addition, although data

completeness was high, NIHSS data were not available for one quarter of patients. We used level of consciousness as a proxy for this and found that the findings were similar, but having complete data on stroke severity may have strengthened the study. The dataset lacked information on the nature of the bedside dysphagia screening tools used, the details of the comprehensive assessment (e.g. videofluoroscopy or fibre-optic evaluation of swallowing), and the results of these assessments. Further studies should aim to capture in more detail the components of these interventions. The main limitation of this study is the risk of residual confounding. The hypothesis that early dysphagia screening and SALT assessment reduce the risk of SAP could be tested in a cluster randomised controlled trial of a protocol of expedited comprehensive dysphagia assessment and this would help guide clinical practice in an important area of stroke care which currently has a poor evidence base. Implementing faster dysphagia assessments in clinical practice is principally a matter of training healthcare professionals appropriately and in most instances does not require expensive medical equipment. Dysphagia screening has been identified by the World Stroke Organisation as being achievable even in health economies with the lowest level of resources²⁵; ensuring that all stroke patients receive rapid dysphagia assessments could therefore be a part of global efforts to improve the outcomes of acute stroke, even in settings without advanced specialist stroke care.

Summary

Delays in screening for dysphagia and carrying out SALT dysphagia assessments after stroke are associated with an increased risk of SAP. This hypothesis that expedited dysphagia screening and assessments reduce the risk of SAP would be testable in an appropriately designed trial or controlled evaluation. In the meantime, these findings suggest that reducing delays in screening and assessing for dysphagia in people with acute stroke should be a focus of quality improvement in stroke care.

Acknowledgements

We would like to thank the many hundreds of individuals and organisations who have contributed to collecting data for SSNAP, without whom this study would not have been possible.

Funding

SSNAP is funded by the Healthcare Quality Information Department on behalf of the Department of Health. The research was supported by the National Institute for Health Research Biomedical Research Centre based at Guy's and St Thomas' NHS Foundation Trust and King's College London; MJ is supported by the NIHR Collaboration for Leadership in Applied Health Research and Care (CLAHRC) for the South West Peninsula. No funders were involved in the design and conduct of the study; the collection, management, analysis, and interpretation of the data; or the preparation, review, or approval of the manuscript. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR, or the Department of Health. Disclosures: none

Contributions

BDB - study design, analysis, writing

CJS - study design, writing

GCC - writing

PE - writing

MJ - writing

LP - analysis, writing

PJT - writing

CDAW - writing

AGR - writing

References

1. Westendorp WF, Nederkoorn PJ, Vermeij JD, Dijkgraaf MG, van de Beek D. Post-stroke infection: a systematic review and meta-analysis. *BMC Neurol.* 2011; 11:110
2. Vernino S, Brown RD Jr, Sejvar JJ, Sicks JD, Petty GW, O'Fallon WM. Cause-specific mortality after first cerebral infarction: a population-based study. *Stroke.* 2003 ;34:1828-32
3. Katzan IL, Cebul RD, Husak SH, Dawson NV, Baker DW. The effect of pneumonia on mortality among patients hospitalized for acute stroke. *Neurology.* 2003;60:620-625
4. Finlayson O, Kapral M, Hall R, Asllani E, Selchen D, Saposnik G; on behalf of the Investigators of the Registry of the Canadian Stroke Network. Risk factors, inpatient care, and outcomes of pneumonia after ischemic stroke. *Neurology.* 2011;77:1338-1345
5. Katzan IL, Dawson NV, Thomas CL, Votruba ME, Cebul RD. The cost of pneumonia after acute stroke. *Neurology.* 2007;68:1938-43
6. Ingeman A, Andersen G, Hundborg HH, Svendsen ML, Johnsen SP. In-hospital medical complications, length of stay, and mortality among stroke unit patients. *Stroke.* 2011;42:3214-3218
7. Koennecke HC, Belz W, Berfelde D, Endres M, Fitzek S, Hamilton F, et al. Factors influencing in-hospital mortality and morbidity in patients treated on a stroke unit. *Neurology.* 2011;77:965-972
8. Odderson IR, Keaton JC, McKenna BS. Swallow management in patients on an acute stroke pathway: quality is cost effective. *Arch Phys Med Rehabil.* 1995 ;76:1130-3
9. Martino R, Foley N, Bhogal S, Diamant N, Speechley M, Teasell R. Dysphagia after stroke. Incidence, diagnosis, and pulmonary complications. *Stroke.* 2005;36:2756-2763
10. Hinchey JA, Shephard T, Furie K, Smith D, Wang D, Tonn S; Stroke Practice Improvement Network Investigators. Formal dysphagia screening protocols prevent pneumonia. *Stroke.* 2005;36:1972-1976
11. Titsworth WL, Abram J, Fullerton A, Hester J, Guin P, Waters MF, et al. Prospective quality initiative to maximize dysphagia screening reduces hospital-acquired pneumonia prevalence in patients with stroke. *Stroke.* 2013; 44:3154-60

12. European Stroke Organisation (ESO) Executive Committee; ESO Writing Committee. Guidelines for management of ischaemic stroke and transient ischaemic attack 2008. *Cerebrovasc Dis.* 2008;25:457-507
13. Jauch EC, Saver JL, Adams HP Jr, Bruno A, Connors JJ, Demaerschalk BM, et al; American Heart Association Stroke Council; Council on Cardiovascular Nursing; Council on Peripheral Vascular Disease; Council on Clinical Cardiology. Guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke.* 2013;44:870-947
14. National Clinical Guideline for Stroke, 4th Edition. Intercollegiate Stroke Working Party 2012. Available at: www.rcplondon.ac.uk/sites/default/files/national-clinical-guidelines-for-stroke-fourth-edition.pdf URL accessed 10th June 2015
15. Sentinel Stroke National Audit Programme. Available at: www.strokeaudit.org URL accessed 10th June 2015
16. Buis M. POSTRCSPLINE: Stata module containing post-estimation commands for models using a restricted cubic spline. Available at: <http://EconPapers.repec.org/RePEc:boc:bocode:s456928> URL accessed 10th June 2015
17. Luker JA, Wall K, Bernhardt J, Edwards I, Grimmer-Somers K. Measuring the quality of dysphagia management practices following stroke: a systematic review. *Int J Stroke.* 2010;5:466-76
18. Turner M, Barber M, Dodds H, Murphy D, Dennis M, Langhorne P, et al; Scottish Stroke Care Audit. Implementing a simple care bundle is associated with improved outcomes in a national cohort of patients with ischemic stroke. *Stroke.* 2015 ;46:1065-70
19. Bravata DM, Wells CK, Lo AC, Nadeau SE, Melillo J, Chodkowski D, et al. Processes of care associated with acute stroke outcomes. *Arch Intern Med.* 2010;170:804-10
20. Bray BD, Ayis S, Campbell J, Hoffman A, Roughton M, Tyrrell PJ, et al. Associations between the organisation of stroke services, process of care, and mortality in England: prospective cohort study. *BMJ.* 2013;346:f2827

21. Middleton S, McElduff P, Ward J, Grimshaw JM, Dale S, D'Este C, et al. ; QASC Trialists Group. Implementation of evidence-based treatment protocols to manage fever, hyperglycaemia, and swallowing dysfunction in acute stroke (QASC): a cluster randomised controlled trial. Lancet. 2011;378:1699-706
22. Masrur S, Smith EE, Saver JL, Reeves MJ, Bhatt DL, Zhao X, et al. Dysphagia screening and hospital-acquired pneumonia in patients with acute ischemic stroke: findings from Get with the Guidelines--Stroke. J Stroke Cerebrovasc Dis. 2013 ;22:e301-9
23. Donovan NJ, Daniels SK, Edmiaston J, Weinhardt J, Summers D, Mitchell PH; on behalf of the American Heart Association Council on Cardiovascular Nursing and Stroke Council. Dysphagia screening: state of the art. Invitational conference proceeding from the state-of-the-art nursing symposium, international stroke conference 2012. Stroke. 2013;44:e24-e31
24. Stroke Quality Standards. National Institute for Health and Care Excellence 2010. Available at: www.nice.org.uk/guidance/qs2 URL accessed 10th June 2015
25. Lindsay P, Furie KL, Davis SM, Donnan GA, Norrving B. World Stroke Organization global stroke services guidelines and action plan. Int J Stroke. 2014 ;9 Suppl A100:4-13

- 1 **Figure 1.** *Modelled relationship between estimated incidence of SAP in the first seven days of*
2 *admission and time to dysphagia screening. (A): Multivariable model including NIHSS. (B):*
3 *Multivariable model including level on consciousness*
- 4 **Figure 2.** *Modelled relationship between estimated incidence of SAP in the first seven days of*
5 *admission and the time to SALT dysphagia assessment. (A): Multivariable model including NIHSS.*
6 *(B): Multivariable model including level on consciousness*