King's Research Portal

DOI:

[10.1016/j.ehbc.2003.12.006](https://doi.org/10.1016/j.ehbc.2003.12.006)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Patton, R. (2004). Excessive drinkers may benefit from brief interventions, but screening in general practice for case finding is not efficient. *Evidence-Based Healthcare*, 8(1), 10. <https://doi.org/10.1016/j.ehbc.2003.12.006>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

COMMENTARY

A recent government report ¹ estimated that alcohol misuse cost the NHS £1.7 billion per year. Brief interventions are effective in reducing levels of alcohol consumption and associated harm ², however successful intervention depends upon reliable and efficient identification of the target population. Screening questionnaires facilitate the detection of alcohol problems ³, and their use is recommended as a precursor to brief intervention ⁴.

This review assesses the effectiveness of screening for alcohol problems in terms of positive outcomes resulting from exposure to brief intervention. For every 1000 patients screened, 90 are identified as having potential alcohol problems, 25 receive brief intervention and of these between 2 and 3 will have reduced their alcohol consumption to below recommended levels after one year. The authors conclude that screening for alcohol problems in general practice is not feasible.

Should these findings dissuade us from undertaking screening and brief intervention as part of everyday practice? All of the studies reviewed for this meta-analysis employed exclusion criteria whereby patients who screened as positive did not qualify to receive brief interventions. In fact 72% of such patients were excluded in this way. It is likely that in clinical practice far greater numbers of patients may actually receive interventions, and thus derive benefit.

Are relatively low numbers of patients experiencing benefit from exposure to brief interventions a cause for concern? Overall 12% of such patients reduced their drinking to below *recommended* limits, we do not know how many more reduced their consumption significantly, but remained above these thresholds. The authors report a pooled NNT of 10 for the studies assessed; most clinicians would agree that this is an acceptable figure.

This paper has shown a low return in terms of patients drinking below recommended limits compared to all those who were screened. It does not however demonstrate that brief intervention itself is ineffective, rather that screening an entire population (under the conditions prescribed by research protocol) will not yield encouraging results. Is there an alternative to screening all patients? Future research that tests the effectiveness of screening procedures in a *pragmatic* ⁵ way may provide the answer.

Robert Patton, Department of Psychological Medicine, Imperial College London.

Title, name(s) and affiliation to be placed at bottom of commentary:

Literature cited:

- (1) Strategy Unit. Alcohol harm reduction project: Interim analytical report. 2003. London, Cabinet Office.
- (2) Moyer A, Finney JW, Swearingen CE, Vergun P. Brief interventions for alcohol problems: a meta-analytic review of controlled investigations in treatment-seeking and non-treatment-seeking populations. *Addiction* 2002; 97:279-292.
- (3) Fiellin DA, Reid MC, O'Connor PG. Screening for Alcohol Problems in Primary Care. *Archives of Internal Medicine* 160, 1977-1989. 2000.
- (4) Babor TF, Higgins-Biddle JC. Brief Intervention For Hazardous and Harmful Drinking: A Manual for Use in Primary Care. 2001. Geneva, World Health Organisation.
- (5) Tunis SR, Stryer DB, Clancy CM. Practical clinical trials: increasing the value of clinical research for decision making in clinical and health policy. *JAMA* 2003; 290:1624-1632.

COMMENTATOR'S CONTACT DETAILS

Name	<i>Include any qualifications that you want cited</i> Robert Patton
Title	Mr
Institution	Imperial College London
Postal Address	13.09 Claybrook Centre, 37 Claybrook Road, London, W6 8LN
Email	r.patton@imperial.ac.uk