King's Research Portal

DOI:

[10.1016/j.psychres.2016.10.033](https://doi.org/10.1016/j.psychres.2016.10.033)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Firth, J., Vancampfort, D., Schuch, F. B., Yung, A. R., & Stubbs, B. (2016). Moving beyond the weight-loss paradigm of exercise interventions for mental illness. *Psychiatry Research*, 246, 392–393.
<https://doi.org/10.1016/j.psychres.2016.10.033>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Author's Accepted Manuscript

Moving beyond the weight-loss paradigm of exercise interventions for mental illness

Joseph Firth, Davy Vancampfort, Felipe B. Schuch, Alison R Yung, Brendon Stubbs

PII: S0165-1781(16)31760-7
DOI: <http://dx.doi.org/10.1016/j.psychres.2016.10.033>
Reference: PSY10030

To appear in: *Psychiatry Research*
Accepted date: 17

Cite this article as: Joseph Firth, Davy Vancampfort, Felipe B. Schuch, Alison R Yung and Brendon Stubbs, Moving beyond the weight-loss paradigm of exercise interventions for mental illness, *Psychiatry Research*, <http://dx.doi.org/10.1016/j.psychres.2016.10.033>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting galley proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Moving beyond the weight-loss paradigm of exercise interventions for mental illness

Joseph Firth^{1*}, Davy Vancampfort^{2,3}, Felipe B. Schuch⁴, Alison R Yung^{1,5}, Brendon Stubbs^{6,7}

¹*Institute of Brain, Behaviour and Mental Health, University of Manchester, Manchester, UK;*

²*KU Leuven Department of Rehabilitation Sciences, Leuven, Belgium;*

³*KU Leuven Department of Neurosciences, UPC KU Leuven, Leuven, Belgium;*

⁴*Hospital de Clínicas de Porto Alegre, Porto Alegre, Brazil;*

⁵*Greater Manchester West NHS Mental Health Foundation Trust, Manchester, UK;*

⁶*Physiotherapy Department, South London and Maudsley NHS Foundation Trust, London, UK;*

⁷*Health Service and Population Research Department, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, UK.*

***Corresponding author:** Joseph Firth, Institute of Brain, Behaviour and Mental Health, University of Manchester, Room 3.306, Jean McFarlane Building, Oxford Road, Manchester, M13 9PL, UK. Tel: +44 (0)161 306 7914. joseph.firth@postgrad.manchester.ac.uk

We read with interest the recent correspondence by Carneiro et al. 2016, which presents a strong case for moving beyond the 'weight loss paradigm' of exercise interventions for people with depression. The authors present the argument, along with supporting evidence, that although 'weight loss' may often be advertised as the key benefit of exercise to encourage people towards physical activity, this is in fact rarely achieved from short-term exercise

programs (Malhotra et al., 2015). Furthermore, as mentioned by the authors, recent meta-analyses and clinical trials have provided strong evidence that even in the absence of weight-loss, exercise can still significantly reduce symptoms of mental disorders such as depression and schizophrenia (Carneiro et al 2015; Firth et al. 2015) - showing that weight loss is not a necessary outcome for the psychological benefits of exercise.

However, it should not be forgotten that all of the benefits of exercise for people with mental disorders are ultimately dependent on their engagement with exercise sessions – which requires motivation. A recent meta-analysis of motivations towards exercise reported by 6321 psychiatric patients found that most perceive ‘physical health’ as the primary reason to exercise (Firth et al. 2016a). More specifically, ‘weight loss’ was the most widely-endorsed physical health benefit, with 83% of the patients reported exercising for this reason. Therefore, we must confront the reality that although weight loss may not be a necessary (or even achievable) target outcome of exercise interventions for people with mental illnesses, their belief that exercise could help them to lose bodyweight is an important factor influencing their uptake and engagement in physical activity.

Nevertheless, ‘improving fitness’ is also a popular reason to exercise among people with mental illnesses – endorsed by 75% of patients in the aforementioned meta-analysis (Firth et al. 2016). Studies in younger people with first-episode psychosis have even shown that fitness is prioritized over weight-loss as an exercise goal in this population (Firth et al. 2016b). This is perhaps because obesity is less likely to be an immediate issue in the early stages of illness (Vancampfort et al. 2015), whereas improving one’s energy levels and physical capabilities can always be desirable. Additionally, cardiorespiratory fitness (unlike bodyweight) is readily

modified by exercise interventions – with meta-analytic data in psychiatric populations showing significant increases in “VO₂ Max” (the gold standard of fitness measurements) within just 12 weeks (Vancampfort et al. 2016a).

It should also be considered that whereas weight loss seems to have no significant relationship with psychological benefits of exercise (Carneiro et al. 2015, Firth et al. 2015), the opposite may be true for fitness. Specifically, clinical trials in people with mental illnesses have shown that fitness improvements from exercise interventions also predict increases in brain structure and functioning (Pajonk et al. 2010; Kimhy et al. 2015). Furthermore, a recent meta-regression of 10 controlled exercise trials in 385 patients with schizophrenia found that the fitness improvements were one of the strongest predictors of cognitive enhancement following exercise interventions (Firth et al. 2016c). Although the correlation across studies was non-significant ($p=0.08$), this could indicate that interventions which are most effective for improving fitness may also be most likely to improve cognition.

Additionally, a wealth of research in the general population has consistently found that high levels of physical fitness are more protective of cardiovascular disease morbidity and mortality than low bodyweight / body mass index (Kodama et al. 2009). Thus, we agree with Carneiro et al. (2016) that the focus of exercise interventions for people with mental illnesses should be shifted from ‘weight loss’ to ‘fitness improvement’. In all, the available evidence indicates that shifting the focus in this way would improve the delivery, engagement and outcomes of exercise interventions, since providing these attainable goals will boost a patient’s motivation towards exercise (Vancampfort et al. 2016), and targeting fitness over fat loss is more broadly

beneficial for reducing cardiometabolic risk and improving both physical and cognitive functionality.

In moving even further ‘beyond the weight loss paradigm’ of exercise interventions in mental healthcare, the role of resistance training should also be considered, as this is currently under-researched. Although weight training and resistance exercise is commonly associated only with gaining muscle, there are many other physical and psychological benefits of such training methods for people with mental illnesses. First, this may offer an alternative method for attenuating the metabolic dysfunction associated with psychotropic medications, as resistance exercise has equally positive effects as aerobic training on the metabolic features of diabetes, significantly reducing body fat and improving insulin within 16 weeks (Bacchi et al. 2012).

Second, exercise interventions for people with depression and schizophrenia which incorporate resistance training alongside or independently of aerobic exercise have found significant improvements in symptoms, cognition and real-world functioning (Firth et al. 2016d, Krogh et al. 2014, Strassnig et al. 2015).

Since the metabolic and neurocognitive benefits of resistance training may occur through independent mechanisms to aerobic exercise (Cassilhas et al. 2013, Jurca et al. 2005), it could be hypothesized that exercise interventions which incorporate both of these methods may confer maximal benefits for patients. Again however, motivation must be considered, and real-life implementation of these research findings could mean offering patients the choices of different aerobic and/or resistance exercises to add into their personalized training programs. Indeed, feasibility studies of patients with psychosis which offer customized aerobic and resistance exercise programs have already proven effective for engaging patients, and

beneficial for both physical and mental health (Firth et al. 2015, Curtis et al. 2014). Clearly, physical activity should not be reduced to a weight management strategy, but rather prioritized as a self-management strategy for anyone living with mental illness. Encouraging and supporting patients to undertake types of exercise which matches their own preferences and personal goals will help to beyond the damaging 'weight loss paradigm' of exercise interventions for people with mental illness, ensuring that future interventions are motivating, engaging and effective for improving recovery.

Disclosures

All authors declare no conflict of interests

Acknowledgements

JF is funded by an MRC Doctoral Training Scholarship.

References

- Bacchi, E., Negri, C., Zanolin, M.E., Milanese, C., Faccioli, N., Trombetta, M., Zoppini, G., Cevese, A., Bonadonna, R.C., Schena, F., Bonora, E., 2012. Metabolic effects of aerobic training and resistance training in type 2 diabetic subjects a randomized controlled trial (the RAED2 study). *Diabetes care*, 35 (4), 676-682.
- Carneiro, L.S., Mota, M.P., Vieira-Coelho, M.A., Rosenbaum, S., Fonseca, A.M., Vasconcelos-Raposo, J., 2016. Exercise for adults with depressive symptoms: beyond the weight loss paradigm. *Psychiatry Research*, in press.

Carneiro, L.S., Fonseca, A.M., Vieira-Coelho, M.A., Mota, M.P., Vasconcelos-Raposo, J., 2015.

Effects of structured exercise and pharmacotherapy vs. pharmacotherapy for adults with depressive symptoms: A randomized clinical trial. *Journal of Psychiatric Research* 71, 48-55.

Cassilhas, R.C., Lee, K.S., Fernandes, J., Oliveira, M.G., Tufik, S., Meeusen, R., de Mello, M.T.,

2012a. Spatial memory is improved by aerobic and resistance exercise through divergent molecular mechanisms. *Neuroscience* 202, 309–317.

Curtis, J., Watkins, A., Rosenbaum, S., Teasdale, S., Kalucy, M., Samaras, K. and Ward, P.B.,

2016. Evaluating an individualized lifestyle and life skills intervention to prevent antipsychotic-induced weight gain in first-episode psychosis. *Early Intervention in Psychiatry*, 10 (3), 267-276.

Firth, J., Cotter, J., Elliott, R., French, P., Yung, A.R., 2015. A systematic review and meta-analysis of exercise interventions in schizophrenia patients. *Psychological Medicine* 45 (7), 1343-1361.

Firth, J., Rosenbaum, S., Stubbs, B., Gorczynski, P., Yung, A. R., & Vancampfort, D., 2016a.

Motivating factors and barriers towards exercise in severe mental illness: a systematic review and meta-analysis. *Psychological Medicine* 46 (14), 2869-2881.

Firth, J., Rosenbaum, S., Stubbs, B., Vancampfort, D., Carney, R., Yung, A.R., 2016b. Preferences and motivations for exercise in early psychosis. *Acta Psychiatrica Scandinavica*, 134 (1), 83-84.

Firth, J., Stubbs, B., Rosenbaum, S., Vancampfort, D., Malchow, B., Schuch, F., Elliott, R.,

Nuechterlein, K.H. and Yung, A.R., 2016c. Aerobic exercise improves cognitive functioning in people with schizophrenia: a systematic review and meta-analysis. *Schizophrenia Bulletin*, in press; doi: 10.1093/schbul/sbw115.

Firth, J., Carney, R., Elliott, R., French, P., Parker, S., McIntyre, R., McPhee, J.S. and Yung, A.R., 2016. Exercise as an intervention for first-episode psychosis: a feasibility study. *Early Intervention in Psychiatry*, in press; doi: 10.1111/eip.12329.

Jurca, R.A., Lamonte, M.J., Barlow, C.E., Kampert, J.B., Church, T.S., Blair, S.N., 2005. Association of muscular strength with incidence of metabolic syndrome in men. *Medicine and Science in Sports and Exercise*, 37 (11), 1849-1855.

Kimhy, D., Vakhrusheva, J., Bartels, M.N., Armstrong, H.F., Ballon, J.S., Khan, S., Chang, R.W., Hansen, M.C., Ayanruoh, L., Lister, A., Castrén, E., 2015. The impact of aerobic exercise on brain-derived neurotrophic factor and neurocognition in individuals with schizophrenia: a single-blind, randomized clinical trial. *Schizophrenia Bulletin*, 41 (4), 859-868.

Kodama, S., Saito, K., Tanaka, S., Maki, M., Yachi, Y., Asumi, M., Sugawara, A., Totsuka, K., Shimano, H., Ohashi, Y., Yamada, N., 2009. Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. *JAMA*, 301 (19), 2024-2035.

Krogh, J., Speyer, H., Norgaard, H.C., Moltke, A., Nordentoft, M., 2014. Can exercise increase fitness and reduce weight in patients with schizophrenia and depression? *Front Psychiatry* 5, 89.

Malhotra, A., Noakes, T., Phinney, S., 2015. It is time to bust the myth of physical inactivity and obesity: you cannot outrun a bad diet. *British Journal of Sports Medicine* 49 (15), 967-U912.

Pajonk, F.G., Wobrock, T., Gruber, O., Scherk, H., Berner, D., Kaizl, I., Kierer, A., Müller, S., Oest, M., Meyer, T., Backens, M., 2010. Hippocampal plasticity in response to exercise in schizophrenia. *Archives of general psychiatry*, 67 (2), 133-143.

Strassnig, M.T., Signorile, J.F., Potiaumpai, M., Romero, M.A., Gonzalez, C., Czaja, S., Harvey, P.D., 2015. High velocity circuit resistance training improves cognition, psychiatric symptoms and neuromuscular performance in overweight outpatients with severe mental illness. *Psychiatry research*, 229 (1), 295-301.

Vancampfort, D., Stubbs, B., Mitchell, A.J., De Hert, M., Wampers, M., Ward, P.B., Rosenbaum, S., Correll, C.U., 2015. Risk of metabolic syndrome and its components in people with schizophrenia and related psychotic disorders, bipolar disorder and major depressive disorder: a systematic review and meta-analysis. *World Psychiatry*, 14(3), 339-347.

Vancampfort, D., Rosenbaum, S., Schuch, F., Ward, P.B., Richards, J., Mugisha, J., Probst, M. and Stubbs, B., 2016a. Cardiorespiratory fitness in severe mental illness: a systematic review and meta-analysis. *Sports Medicine*, 1-10.

Vancampfort, D., Moens, H., Madou, T., De Backer, T., Vallons, V., Bruyninx, P., Vanheuverzwijn, S., Mota, C.T., Soundy, A., Probst, M., 2016b. Autonomous motivation is associated with the maintenance stage of behaviour change in people with affective disorders. *Psychiatry Research* 240, 267-271.

Highlights

- Exercise interventions for people with mental illness do not consistently result in weight loss, but can significantly reduce cardiovascular risk and psychiatric symptoms
- Focusing on fitness outcomes of exercise, as opposed to weight-loss, may provide patients with more achievable and valuable goals
- The role of resistance training interventions for psychiatric populations is currently under-researched. Incorporating this alongside aerobic exercise could confer substantial benefits for cardiometabolic health and mental well-being