King's Research Portal

DOI:

[10.1111/jcpe.12630](https://doi.org/10.1111/jcpe.12630)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Sun, X., Bernabé, E., Liu, X., Gallagher, J. E., & Zheng, S. (2016). Do Contextual Factors Have a Role in Periodontal Disease? *Journal of Clinical Periodontology*, 44(1), 13-21. <https://doi.org/10.1111/jcpe.12630>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Received Date : 24-May-2016

Revised Date : 13-Sep-2016

Accepted Date : 29-Sep-2016

Article type : Other

Do Contextual Factors Have a Role in Periodontal Disease?

Running Title: Contextual factors and periodontal disease

**Xiangyu SUN^{1,2,#}, Eduardo BERNABÉ^{2,#}, Xuenan LIU¹, Jennifer E. GALLAGHER^{2,*},
Shuguo ZHENG^{1,**}**

¹ Department of Preventive Dentistry, Peking University School and Hospital of Stomatology, Department of Preventive Dentistry, National Engineering Laboratory for Digital and Material Technology of Stomatology, Beijing Key Laboratory of Digital Stomatology, Haidian District, Beijing, People's Republic of China

² Division of Population and Patient Health, King's College London Dental Institute at Guy's, King's College and St. Thomas' Hospitals, London, United Kingdom

These authors contribute equally to this work.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/jcpe.12630

This article is protected by copyright. All rights reserved.

*** Corresponding Author.**

Prof. Jennifer E. Gallagher MBE

Division of Population and Patient Health, King's College London Dental Institute, Denmark Hill Campus, Bessemer Road, London SE5 9RS, United Kingdom.
Phone: +44 (0) 20 3299 3481; Email: jenny.gallagher@kcl.ac.uk

**** Co-corresponding Author.**

Prof. Shuguo Zheng

Department of Preventive Dentistry, Peking University School of Stomatology, National Engineering Laboratory for Digital and Material Technology of Stomatology, Beijing Key Laboratory of Digital Stomatology, 22 Zhongguancun Avenue South, Beijing 100081, People's Republic of China.
Phone: +86 (0) 10 8219 5510; Email: zhengsg86@gmail.com

Conflict of Interest and Source of Funding Statement

No external funding, apart from the support of the authors' institution, was available for this study. The authors declare that there are no conflicts of interests in this study.

Key words: periodontal disease, adults, multilevel analysis, population characteristics, social environment

Abstract

Aim: To explore the association between contextual factors and periodontal disease.

Methods: We pooled individual-level data from 20,204 35-44-year-olds and 9,666 65-74-years-olds, who participated in the 3rd National Oral Health Survey in China (2005), with province-level data from different sources. The association of provincial macroeconomic factors (Gross Domestic Product [GDP] per capita and Gini coefficient), health resources (public health expenditure, dentist to population ratio and dental therapist to population ratio), demographic composition (proportion of rural population and minority ethnic groups) and tobacco industry (tobacco crops, cigarette production and proportion of smokers) with the numbers of teeth with periodontal pocket depth (PPD) ≥ 4 mm and loss of attachment (LOA) ≥ 4 mm were assessed in multilevel models, controlling for individual-level demographic, socioeconomic and behavioural factors.

Results: Only the proportion of smokers was associated with the number of teeth with PPD ≥ 4 mm and was only among senior adults. However, public health expenditure, dentist to population ratio and the proportion of minority ethnic groups were associated with the number of teeth with LOA ≥ 4 mm in both age groups. GDP per capita was also associated with the number of teeth with LOA ≥ 4 mm only among 35-44-year-olds.

Conclusion: Contextual factors may contribute to periodontal disease and help explain geographical inequalities among Chinese adults.

Clinical Relevance

Scientific rationale for study: Most research on risk factors for periodontal disease has focused on the impact of individual characteristics. The physical and social environment where people live and work has been largely overlooked.

Principal findings: Some, but not all, contextual factors were associated with periodontal disease, independent of well-known risk factors acting at individual level.

Practical implications: This study shows that the context matters for periodontal health. We may improve our understanding of periodontal disease with a broader look at the features of the environment that shape people's behaviours and oral health.

Introduction

Most studies on risk factors for periodontal disease have focused on the impact of the characteristics of individuals (Chapple et al., 2015, Genco and Borgnakke, 2013). There has been comparatively little emphasis on the impact of other factors beyond individuals, such as community or neighbourhood context. The term context is broadly used here to encompass elements of the physical (healthy food outlets, tobacco availability and advertising, and safe play areas for children) and social environment (social cohesion or violence) (Diez Roux, 2007). The physical and social environments can be hazardous (air/water pollution or crime). They can also limit the choices and resources available to individuals, repeatedly exposing people to stressful conditions, which may in turn exert a direct effect on health (chronic activation of the neuro-endocrine system) and an indirect effect through health behaviours as a coping mechanism (Macintyre and Ellaway, 2009, Diez Roux and Mair, 2010, Stafford and McCarthy, 2005). In the case of periodontal diseases, neighbourhood conditions could play a direct role through exposure to environmental tobacco smoke (Akinkugbe et al., 2016). They could also influence behaviours, such as smoking and regular dental check-ups. Furthermore, the deprivation level of a geographical area can influence the health budget and supply of health professionals and thereby increase or decrease access to care.

Although previous studies have shown geographical inequalities in periodontal disease within (Eke et al., 2016, Lopez et al., 2009) and between countries (Kassebaum et al., 2014), only a few studies have looked at contextual determinants as explanations for these differences. The

two contextual factors that have received more attention in the literature are area-level deprivation and income inequality. A national study in the US showed that individuals living in poorer neighbourhoods were more likely to have periodontitis, even after controlling for demographic, socioeconomic and behavioural characteristics of participants (Borrell et al., 2006). However, a national study in Scotland subsequently reported no association between area deprivation and periodontitis (Bower et al., 2007). The different results could be explained by the use of multilevel models in the latter study to account for the clustering of individuals within geographical areas. Failing to recognise such a hierarchical structure would underestimate standard errors, leading to an overstatement of statistical significance (Merlo et al., 2005). Similarly, preliminary ecological evidence suggested that more unequal countries have higher rates of periodontitis (Sabbah et al., 2010, Hobdell et al., 2003). A further multilevel analysis of the 2002/03 Brazilian Oral Health Survey among 35-44-year-olds showed that income inequality at municipal-level was not associated with periodontal disease (Celeste et al., 2011). Oral epidemiology should pay greater attention to the role of contextual factors in shaping oral health in general and periodontal disease in particular. The association of area composition and environments with periodontal disease highlights the possible role of area-level interventions to improve population oral health and reduce oral health inequalities. To fill this gap in knowledge, a study was set out to explore the association between contextual factors and periodontal disease.

Methods

Data source

This cross-sectional study pooled individual- and province-level data from various sources. Individual-level data were from the 3rd National Oral Health Survey of China (2005), which

covered the four World Health Organization (WHO) index ages (5-, 12-, 35- to 44-, and 65- to 74-year-olds). All 31 provinces of Mainland China participated in the survey, except for Tibet. Participants were selected using multistage stratified cluster sampling. For selection, every province was divided into urban and rural areas; urban areas were classified into three strata by population size, whereas rural areas were classified by Gross Domestic Product (GDP). One city or county was randomly selected from each stratum. Hence, three cities from urban areas and three counties from rural areas were selected from each province. For the next level, three streets or townships were randomly chosen from every city or county, respectively. Two residents' committees in these streets (or two villages in townships) were randomly chosen as survey stations. At each survey station, 20 adults and 20 senior adults were recruited randomly from the list of residents provided by each residents' committee (Figure 1). A target sample of 720 participants in each age group was initially set per province, for a total of 21,600 people nationally. For 35-44-year-olds, 23,538 participated in clinical examinations and 23,522 completed the questionnaire. For 65-74-year-olds, 23,415 were clinically examined and 12,893 completed the questionnaire (only 50% of the senior sample was invited to fill it out).

Variables selection

The number of teeth with periodontal pocket depth (PPD) ≥ 4 mm and loss of attachment (LOA) ≥ 4 mm were the two outcome measures. Clinical examinations were carried out with participants seated on a chair, and using artificial light, plane mouth mirrors and standard WHO CPI probes. All teeth, excluding third molars, were examined. PPD was assessed at six sites per tooth but only the most severe code was recorded by tooth, according to one of the following categories: 0-3, 4-5, 6+ mm. LOA was also assessed at six sites per tooth but recorded by tooth according to one of the following categories: 0-3, 4-5, 6-8, 9-11, 12+ mm

(WHO, 1997). Unified training sessions were provided to over 200 dentists in Kunming city, Yunnan, before the national survey begins. Reliability values were not calculated because of the difficulty of performing so with the large number of examiners.

Data on contextual factors were gathered from different national and international sources trying to match the survey year as close as possible. Macroeconomic factors were income inequality and average income. Income inequality was measured using the Gini coefficient, expressed as a percentage where higher values indicate greater inequality, for the period 1985-1995 (Xu and Zou, 2000). More recent income inequality data are not available due to the lack of comprehensive income surveys in China (Xie and Zhou, 2014). Average income was measured with the GDP per capita in 2005 expressed in thousand Yuan (National Bureau of Statistics of China, 2006). Health resources were measured by public health expenditure expressed as a proportion of total government spending on health for 2005 (Fang et al., 2010) and the dentist-to and dental-therapist-to-population ratios, expressed per 10 million people, in 2002 (Ministry of Health of China, 2006). Two demographic indicators were also included, namely the proportions of rural population and minority ethnic groups, extracted from the 5th National Demographic Census in 2000 (National Bureau of Statistics of China, 2001). As the largest tobacco cultivator and cigarette producer in the world (Yang et al., 2015), the strength of the Chinese tobacco industry is likely to be an important factor influencing the availability of tobacco, social norms about tobacco, smoking rates and exposure to second-hand smoke locally. Two measures relating to the level of tobacco cultivation (in ten million tons) and cigarette production (in ten trillion) in 2005 (National Bureau of Statistics of China, 2006), along with the proportion of smokers aged 15 to 69 years, taken from the 1996 national survey (Chinese Academy of Preventive Medicine, 1997), were also included.

Participants' demographic, socioeconomic and behavioural characteristics were included as covariates to separate contextual from compositional effects (when province-level differences in periodontal measures are explained by the individual composition of the provinces).

Demographic factors were sex, age, ethnicity and place of residence (urban or rural).

Participants' ethnicity was self-assigned using a list of officially recognised ethnic groups in China, and responses classified as Han or minority ethnic group. Socioeconomic factors were education and household income. Participants reported their total number of years of full-time education, and responses were regrouped in line with national cut-off points: primary school (0-9 years), junior middle school (10-12 years), senior middle school (13-15 years) and higher education (16+ years). Participants were also asked to provide an estimate of their annual household income with no pre-set categories. Income data was equivalised using the Luxembourg Income Study scale to account for family size. This was done dividing the total household income by the square root of the number of individuals in the family (Buhmann et al., 1988). After equivalisation, household income in Chinese Yuan was categorised into five groups: very low (0-4999), low (5000-9999), medium (10000-14999), high (15000-19999), and very high (20000+). Dental behaviours included toothbrushing frequency (not every day, once a day, and equal to or more than twice a day), last dental visit for any reason (never, within last year, and more than 1 year ago) and smoking status. Participants' smoking status was derived from responses to the questions 'how often do you smoke?' and "in the last month, how many cigarettes do you smoke each day?", and classified as never, former, ≤ 10 cigarettes/day, 11-20 cigarettes/day and >20 cigarettes/day.

Statistical analysis

We first present the composition of the sample of adults and senior adults according to individual and contextual factors. All contextual factors were treated as continuous variables and summarised using mean, standard deviation (SD) and minimum and maximum values.

A two-level random-intercepts and fixed-slopes model structure with individuals nested within provinces was fitted in MLwiN 2.29, using negative binomial regression as the two outcome measures were count variables with over-dispersion. Rate ratios (RR) were thus reported. The number of teeth was used as an offset variable in all models. The numbers of teeth with $PPD \geq 4\text{mm}$ and $LOA \geq 4\text{mm}$ were compared by demographic, socioeconomic, and behavioural characteristics of individuals in crude negative binomial regression models.

Our modelling strategy was first to estimate empty models (without any covariates) to evaluate the amount of unexplained variance in the numbers of teeth with $PPD \geq 4\text{mm}$ and $LOA \geq 4\text{mm}$ at province and individual levels, using the variance partition coefficient (VPC).

We then estimated the crude association between every contextual factor and periodontal measures (labelled as Model 1), and gradually adjusted them for individual-level factors (Model 2) and other province-level factors (Model 3). The reason for this strategy was to evaluate compositional and contextual effects when controlling for individual- and province-level confounders, respectively. Model 3 also allowed assessing the relative contribution of contextual factors to explaining variations in periodontal measures. All analyses were conducted using the unweighted sample as the level-2 weights, needed to compensate for the unequal probability of selection of level-2 units (Rabe-Hesketh and Skrondal, 2006, Cai, 2013), were not available with survey files. Non-weighted analyses were appropriate as our focus was on tests of association rather than deriving nationally representative estimates. More importantly, minimal differences have been observed in estimates and standard errors from weighted and unweighted multilevel regression (Carle, 2009).

RESULTS

A total of 20,204 35-44-year-olds (86%) and 9,666 65-74-year-olds (75%) had complete data in all relevant variables and were included in this analysis. The mean number of participants per province was 673 (range: 519-789) among 35-44-year-olds and 322 (range: 216-708) among 65-74-year-olds. The characteristics of the two study samples are shown in Table 1.

There were no demographic, socioeconomic or behavioural differences between the study sample and those excluded due to missing data, except for place of residence among 65-74-year-olds (i.e. urban residents were slightly overrepresented in the study sample). The mean numbers of teeth with $PPD \geq 4\text{mm}$ and $LOA \geq 4\text{mm}$ were 2.34 (SD: 4.95, 95% Confidence Interval: 2.28-2.41) and 2.04 (SD: 3.87, 95%CI: 1.98-2.09) in adults and 2.67 (SD: 4.42, 95%CI: 2.58-2.75) and 4.95 (SD: 5.04, 95%CI: 4.84-5.05) in senior adults, respectively.

Significant differences in the numbers of teeth with $PPD \geq 4\text{mm}$ and $LOA \geq 4\text{mm}$ were found by almost all demographic, socioeconomic and behavioural factors (Table 2).

The VPC from the empty models showed that between 3.3% and 4.8% of the variation in the numbers of teeth with $PPD \geq 4\text{mm}$ and $LOA \geq 4\text{mm}$ occurred at province level. Tables 3 and 4 show the individual- and contextual-level factors associated with periodontal measures in Chinese adults and senior adults, respectively. Only one contextual factor, the proportion of smokers, was associated with the number of teeth with $PPD \geq 4\text{mm}$ in the adjusted model and this was only among senior adults. A one-point increase in the proportion of smokers was associated with a 4%-increase in the mean number of teeth with $PPD \geq 4\text{mm}$. On the other hand, three contextual factors (public health expenditure, dentist to population ratio and proportion of minority groups) were significantly associated with the number of teeth with $LOA \geq 4\text{mm}$ among adults and senior adults. In both age groups, provinces with more public health expenditure had significantly more of teeth with $LOA \geq 4\text{mm}$ whereas provinces with more dentists per population and higher proportion of minority ethnic groups had

significantly fewer teeth with $\text{LOA} \geq 4\text{mm}$. GDP per capita was also positively associated with the number of teeth with $\text{LOA} \geq 4\text{mm}$, but only among adults.

Sensitivity analyses using higher thresholds (6mm) for PPD and LOA were carried out to minimise measurement error in periodontal examinations and the number of false positives.

These new analyses confirmed that GDP per capita, dentist-to-population ratio and the proportion of smokers at province level were associated with periodontal measures after adjustments. However, public health expenditure and the proportion of minority ethnic groups were no longer significantly associated with periodontal measures.

DISCUSSION

This study shows that contextual factors may contribute to periodontal disease and help explain geographical inequalities in periodontal disease. Provincial average income, dentist to population ratio and the proportion of smokers (and to a lesser extent public health expenditure and the proportion of minority ethnic groups) were associated with periodontal disease over and above the effects of established risk factors such as demographic characteristics, socioeconomic circumstances and behavioural factors.

Some limitations of this study need to be addressed. First, we used data from a relatively old survey. Despite being conducted in 2005, the 3rd National Oral Health Survey still remains the latest oral health survey available and the contemporary reference in China. No other Chinese survey provides comprehensive oral health data at province level. Second, although six periodontal sites were inspected in every tooth, only one code was recorded per tooth. Partial-mouth assessments maximise the number of people examined in the time available and encourage subjects to comply with the study protocol, but they underestimate the prevalence and severity of periodontal disease (Holtfreter et al., 2015, Susin et al., 2005).

Third, examiners were trained before data collection but examiner reliability was not assessed. This is not a unique characteristic of this survey, but rather a standard feature across national epidemiological surveys (Karimalakuzhiyil Alikutty and Bernabe, 2016), reflecting the difficulty to assess reliability when a large number of examiners is involved. That said, we found similar findings in sensitivity analyses using higher thresholds in PPD and LOA to minimise measurement error and false positives. We also found well-known risk factors of periodontal disease (smoking, toothbrushing, education and income) to be associated with our periodontal measures, suggesting that measurement error in periodontal examinations did not seriously affect the results. Fourth, although we tried to match the ecological data as close as possible to 2005, we had to go back 10 to 15 years for some indicators (income inequality and smoking rates). This should not be a big concern because contextual effects do not act instantaneously but require some time to develop (Diez Roux, 2007). Fifth, misclassification of exposure may have occurred if some participants moved between provinces between the assessment of contextual factors and the oral health survey. Interestingly, there is evidence that contextual factors were more strongly related to health measures when analysis was restricted to individuals living in their state of birth (persistent exposure group) compared to the complete sample (Gadalla and Fuller-Thomson, 2008). This means that our estimates of the association between contextual factors and periodontal disease may be somewhat conservative.

GDP per capita, dentist to population ratio and the proportion of smokers were associated with periodontal disease. Adults in more affluent provinces had more teeth with LOA.

Modernisation, economic development and increased wealth have been accompanied by a shift from diets rich in fibre and staples to diets high in fats and sugars in China (Popkin, 2014). Fibre and whole-grain diets are associated with lower risk of periodontal disease (Merchant et al., 2006, Schwartz et al., 2012). In addition, economic development could also

lead to more stressful work environments, with longer working hours and more demanding jobs. The latter argument would explain why the association was only found among adults, since most senior adults would probably be retired from work. More dentists locally could make access to dental care in general and periodontal care in particular easier, which may subsequently lead to better periodontal status. On the other hand, the proportion of smokers may be a proxy for exposure to second-hand smoke, which has been associated with periodontal disease (Akinkugbe et al., 2016). The fact that the association was only found among senior adults suggests a long-term exposure to environmental tobacco smoking.

Public health expenditure and the proportion of minority ethnic groups were also associated with periodontal disease, although these results were not confirmed in sensitivity analysis. The estimate for public health expenditure was counterintuitive since residents of provinces with higher public spending on health had more teeth with LOA. It is possible that public investment in health leads to increased life expectancy and subsequently more periodontal disease, a condition that becomes more prevalent with age (Kassebaum et al., 2014).

However, this would not explain the association found among 35-44-year-olds. Another explanation could be that public investment in health care may lead to greater provision of dental care in the form of inter-proximal restorations, which are a risk factor for periodontal disease (Broadbent et al., 2006). Moreover, individuals in provinces with larger minority ethnic groups also had fewer teeth with LOA, in spite of the fact there were no differences in LOA between ethnic groups. Having more people from the same ethnic background within the living area may generate greater cohesiveness, trust and reciprocity, attributes of a community that are related to health (Murayama et al., 2012). Contrarily, minority ethnic groups in less diverse areas may experience discrimination and marginalisation, leading to psychosocial stress and poor health (Diez Roux and Mair, 2010).

Contrary to our hypothesis, tobacco crops and cigarette production were not associated with periodontal disease. This might be attributed to the strict regulation on cigarette prices and accessibility imposed by the Chinese government across regions. Income inequality was not associated with periodontal disease either. Our Gini values for all provinces were below the 0.30 threshold effect for income inequality, above which an association with health is more likely to be found (Kondo et al., 2012). Levels of income inequality in China started to increase dramatically in the 1990s (Xie and Zhou, 2014). Therefore, it would be interesting to corroborate this finding when more recent Gini data at province level becomes available.

This study shows that the context matters for periodontal health. Our finding highlights the unique contribution of contextual factors and the role of multilevel models in separating contextual from compositional effects (the simple aggregation of individual-level factors).

Policies and interventions for prevention of periodontal disease may benefit from taking into consideration the environment in which people live and work. Further research must explore the probable mechanisms by which the physical and social environments may affect periodontal status in order to inform appropriate policy action.

In conclusion, this study provides evidence on the importance of contextual factors to periodontal disease and to explain geographical inequalities. Further research is needed on the underlying pathways linking the environment to periodontal disease.

Acknowledgements

The authors are grateful to all participants, organisers and staff who contributed to the 3rd National Oral Health Epidemiological Survey in China.

References

- Akinkugbe, A. A., Slade, G. D., Divaris, K. & Poole, C. (2016) Systematic Review and Meta-analysis of the Association Between Exposure to Environmental Tobacco Smoke and Periodontitis Endpoints Among Nonsmokers. *Nicotine & tobacco research : official journal of the Society for Research on Nicotine and Tobacco*.
- Borrell, L. N., Burt, B. A., Warren, R. C. & Neighbors, H. W. (2006) The role of individual and neighborhood social factors on periodontitis: the third National Health and Nutrition Examination Survey. *Journal of Periodontology* **77**, 444-453.
- Bower, E., Gulliford, M., Steele, J. & Newton, T. (2007) Area deprivation and oral health in Scottish adults: a multilevel study. *Community Dentistry and Oral Epidemiology* **35**, 118-129.
- Broadbent, J. M., Williams, K. B., Thomson, W. M. & Williams, S. M. (2006) Dental restorations: a risk factor for periodontal attachment loss? *Journal of clinical periodontology* **33**, 803-810.
- Buhmann, B., Rainwater, L., Schmaus, G. & Smeeding, T. M. (1988) Equivalence scales, well-being, inequality, and poverty: sensitivity estimates across ten countries using the luxembourg income study (lis) database. *Review of Income and Wealth* **34**, 115-142.
- Cai, T. (2013) Investigation of ways to handle sampling weights for multilevel model analyses. *Sociological Methodology* **43**, 178-219.
- Carle, A. C. (2009) Fitting multilevel models in complex survey data with design weights: Recommendations. *BMC Medical Research Methodology* **9**, 49.

Celeste, R. K., Fritzell, J. & Nadanovsky, P. (2011) The relationship between levels of income inequality and dental caries and periodontal diseases. *Cadernos de Saude Publica* **27**, 1111-1120.

Chapple, I. L., Van der Weijden, F., Doerfer, C., Herrera, D., Shapira, L., Polak, D., Madianos, P., Louropoulou, A., Machtei, E., Donos, N., et al. (2015) Primary prevention of periodontitis: managing gingivitis. *Journal of clinical periodontology* **42 Suppl 16**, S71-76.

Chinese Academy of Preventive Medicine (1997) *Smoking and health in China: 1996 National prevalence survey of smoking pattern*. Beijing: China Science and Technology Press.

Diez Roux, A. V. (2007) Neighborhoods and health: where are we and where do we go from here? *Revue d'Epidemiologie et de Sante Publique* **55**, 13-21.

Diez Roux, A. V. & Mair, C. (2010) Neighborhoods and health. *Annals of the New York Academy of Sciences* **1186**, 125-145.

Eke, P. I., Zhang, X., Lu, H., Wei, L., Thornton-Evans, G., Greenlund, K. J., Holt, J. B. & Croft, J. B. (2016) Predicting Periodontitis at State and Local Levels in the United States. *Journal of Dental Research* **95**, 515-522.

Fang, P., Dong, S., Xiao, J., Liu, C., Feng, X. & Wang, Y. (2010) Regional inequality in health and its determinants: evidence from China. *Health Policy* **94**, 14-25.

Gadalla, T. M. & Fuller-Thomson, E. (2008) Examining the lag time between state-level income inequality and individual disabilities: a multilevel analysis. *American Journal of Public Health* **98**, 2187-2190.

Genco, R. J. & Borgnakke, W. S. (2013) Risk factors for periodontal disease. *Periodontology* 2000 **62**, 59-94.

Hobdell, M. H., Oliveira, E. R., Bautista, R., Myburgh, N. G., Lalloo, R., Narendran, S. & Johnson, N. W. (2003) Oral diseases and socio-economic status (SES). *British Dental Journal* **194**, 91-96.

Holtfreter, B., Albandar, J. M., Dietrich, T., Dye, B. A., Eaton, K. A., Eke, P. I., Papapanou, P. N. & Kocher, T. (2015) Standards for reporting chronic periodontitis prevalence and severity in epidemiologic studies: Proposed standards from the Joint EU/USA Periodontal Epidemiology Working Group. *Journal of clinical periodontology* **42**, 407-412.

Karimalakuzhiyil Alikutty, F. & Bernabe, E. (2016) Long-term regular dental attendance and periodontal disease in the 1998 adult dental health survey. *Journal of clinical periodontology* **43**, 114-120.

Kassebaum, N. J., Bernabé, E., Dahiya, M., Bhandari, B., Murray, C. J. & Marcenes, W. (2014) Global burden of severe periodontitis in 1990-2010: a systematic review and meta-regression. *Journal of Dental Research* **93**, 1045-1053.

Kondo, N., van Dam, R. M., Sembajwe, G., Subramanian, S. V., Kawachi, I. & Yamagata, Z. (2012) Income inequality and health: the role of population size, inequality threshold, period effects and lag effects. *Journal of epidemiology and community health* **66**, e11.

Lopez, R., Frydenberg, M. & Baelum, V. (2009) Contextual effects in the occurrence of periodontal attachment loss and necrotizing gingival lesions among adolescents. *European Journal of Oral Sciences* **117**, 547-554.

Macintyre, S. & Ellaway, A. (2009) Neighbourhood and health. In: *Understanding health inequalities*, (ed.) H. Graham, pp. 84-100. Berkshire: McGraw-Hill Education.

Merchant, A. T., Pitiphat, W., Franz, M. & Joshipura, K. J. (2006) Whole-grain and fiber intakes and periodontitis risk in men. *The American Journal of Clinical Nutrition* **83**, 1395-1400.

Merlo, J., Chaix, B., Yang, M., Lynch, J. & Rastam, L. (2005) A brief conceptual tutorial of multilevel analysis in social epidemiology: linking the statistical concept of clustering to the idea of contextual phenomenon. *Journal of Epidemiology and Community Health* **59**, 443-449.

Ministry of Health of China (2006) *National health statistical yearbook 2005*. Beijing: Peking Union Medical College Press.

Murayama, H., Fujiwara, Y. & Kawachi, I. (2012) Social capital and health: a review of prospective multilevel studies. *Journal of Epidemiology* **22**, 179-187.

National Bureau of Statistics of China (2001) *Report of 5th National Demographic Census in 2000*. Beijing: China Statistics Press.

National Bureau of Statistics of China (2006) *National statistical yearbook 2005*. Beijing: China Statistics Press.

Popkin, B. M. (2014) Synthesis and implications: China's nutrition transition in the context of changes across other low- and middle-income countries. *Obesity Reviews* **15 Suppl 1**, 60-67.

Rabe-Hesketh, S. & Skrondal, A. (2006) Multilevel modelling of complex survey data. *J Roy Stat Soc a Sta* **169**, 805-827.

Sabbah, W., Sheiham, A. & Bernabe, E. (2010) Income inequality and periodontal diseases in rich countries: an ecological cross-sectional study. *International Dental Journal* **60**, 370-374.

Schwartz, N., Kaye, E. K., Nunn, M. E., Spiro, A., 3rd & Garcia, R. I. (2012) High-fiber foods reduce periodontal disease progression in men aged 65 and older: the Veterans Affairs normative aging study/Dental Longitudinal Study. *Journal of the American Geriatrics Society* **60**, 676-683.

Stafford, M. & McCarthy, M. (2005) Neighbourhoods, housing and health. In: *Social Determinants of Health* (eds.) M. Marmot & R. G. Wilkinson, pp. 318-340. Oxford: Oxford University Press.

Susin, C., Kingman, A. & Albandar, J. M. (2005) Effect of partial recording protocols on estimates of prevalence of periodontal disease. *Journal of periodontology* **76**, 262-267.

WHO (1997) *Oral Health Surveys: Basic Methods*. Geneva: World Health Organisation.

Xie, Y. & Zhou, X. (2014) Income inequality in today's China. *Proceedings of the National Academy of Sciences of the United States of America* **111**, 6928-6933.

Xu, L. C. & Zou, H. F. (2000) Explaining the changes of income distribution in China. *China Economic Review* **11**, 149-170.

Yang, G., Wang, Y., Wu, Y., Yang, J. & Wan, X. (2015) The road to effective tobacco control in China. *Lancet (London, England)* **385**, 1019-1028.

Table 1. Characteristics of the sample of Chinese adults and senior adults

Level 1: Individual	35-44-year-olds		65-74-years-olds	
<i>Sex, n^a %</i>				
Men	9886	48.9%	5027	52.0%
Women	10318	51.1%	4639	48.0%
<i>Ethnicity, n^a %</i>				
Han	18035	89.3%	8747	90.5%
Minority ethnic groups	2169	10.7%	919	9.5%
<i>Place of residence, n^a %</i>				
Rural	9965	49.3%	4487	46.4%
Urban	10239	50.7%	5179	53.6%
<i>Education level, n^a %</i>				
Up to primary school	5120	25.3%	6666	69.0%
Junior middle school	7577	37.5%	1554	16.1%
Senior middle school	4895	24.2%	886	9.2%
Higher education	2612	12.9%	560	5.8%
<i>Equivalised income, n^a %</i>				
Very low	8220	40.7%	4937	51.1%
Low	5632	27.9%	2267	23.5%
Medium	3289	16.3%	1193	12.3%
High	1246	6.2%	523	5.4%
Very high	1817	9.0%	746	7.7%
<i>Toothbrushing frequency, n^a %</i>				
Less often than daily	2146	10.6%	2179	22.5%
Once a day	11107	55.0%	4899	50.7%
Twice a day or more	6951	34.4%	2588	26.8%
<i>Last dental visit, n^a %</i>				
Never	9156	45.3%	2779	28.7%
Within last year	3293	16.3%	1981	20.5%
Over 1 year ago	7755	38.4%	4906	50.8%
<i>Smoking status, n^a %</i>				
Never smoker	12774	63.2%	5782	59.8%
Former smoker	718	3.6%	1184	12.2%
≤10 cigarettes/day	1813	9.0%	1145	11.8%
11-20 cigarettes/day	2957	14.6%	1102	11.4%
20+ cigarettes/day	1942	9.6%	453	4.7%
<i>Mean number of teeth with PPD≥4mm (SD)</i>	2.21	(4.77)	2.45	(4.15)
<i>Mean number of teeth with LOA≥4mm (SD)</i>	2.07	(3.90)	5.28	(5.28)
<i>Mean number of teeth (SD)</i>	27.06	(2.04)	20.82	(7.01)
Level 2: Province (n=30)	Mean	(SD)	Range	
<i>Gini coefficient, %</i>	20.80	(2.30)	17.31 - 27.89	
<i>GDP per capita, thousand yuan</i>	16.44	(11.05)	5.05 - 51.47	
<i>Public health expenditure, %</i>	4.09	(0.86)	2.80 - 6.20	

<i>Dentist to population ratio^b</i>	4.15	(2.80)	1.31 - 14.56
<i>Dental therapist to population ratio^b</i>	1.01	(0.34)	0.44 - 1.69
<i>Rural population, %</i>	59.94	(16.49)	11.69 - 76.62
<i>Minority ethnic groups, %</i>	12.43	(16.24)	0.31 - 59.43
<i>Cigarette production, ten trillion</i>	6.46	(6.28)	0 - 31.57
<i>Tobacco crops, ten thousand tons</i>	8.95	(16.02)	0 - 79.1
<i>Smokers, %</i>	38.04	(4.68)	28.99 - 50.45

^a All counts are unweighted

^b Dentist to and dental therapist to population ratios are expressed per ten million population

Table 2. Numbers of teeth with periodontal pocket depth (PPD)≥4mm and loss of attachment (LOA)≥4mm, by individual-level characteristics of Chinese adults and senior adults

	35-44-year-olds (n=20204)		65-74-year-olds (n=9666)	
	PPD (95% CI)	LOA (95% CI)	PPD (95% CI)	LOA (95% CI)
<i>Sex</i>				
Men	3.04 (2.93-3.15)	2.48 (2.40-2.57)	3.05 (2.92-3.18)	5.67 (5.52-5.82)
Women	1.68 (1.60-1.76)	2.11 (2.05-2.17)	2.25 (2.14-2.37)	4.16 (4.03-4.29)
<i>P value^a</i>	<0.001	<0.001	<0.001	<0.001
<i>Ethnicity</i>				
Han	2.40 (2.33-2.47)	2.11 (2.05-2.17)	2.73 (2.64-2.82)	5.04 (4.93-5.14)
Minority ethnic groups	1.87 (1.69-2.05)	1.44 (1.31-1.57)	2.06 (1.82-2.29)	4.06 (3.77-4.35)
<i>P value</i>	<0.001	0.641	<0.001	0.169
<i>Place of residence</i>				
Rural	2.49 (2.39-2.59)	2.19 (2.11-2.27)	2.66 (2.53-2.79)	5.10 (4.95-5.25)
Urban	2.20 (2.11-2.30)	1.89 (1.82-1.96)	2.67 (2.55-2.79)	4.81 (4.67-4.94)
<i>P value</i>	<0.001	<0.001	0.007	<0.001
<i>Education level</i>				
Up to primary school	2.45 (2.31-2.59)	2.20 (2.09-2.31)	2.60 (2.49-2.70)	4.95 (4.84-5.07)

Junior middle school	2.58 (2.46-2.70)	2.17 (2.07-2.26)	2.97 (2.73-3.21)	4.82 (4.57-5.07)
Senior middle school	2.15 (2.02-2.29)	1.97 (1.87-2.08)	2.63 (2.34-2.91)	5.10 (4.74-5.46)
Higher education	1.80 (1.64-1.96)	1.46 (1.34-1.59)	2.71 (2.34-3.08)	4.93 (4.50-5.36)
<i>P value</i>	0.017	<0.001	0.433	0.004
<i>Equivalised income</i>				
Very low	2.77 (2.65-2.89)	2.12 (2.03-2.20)	2.69 (2.56-2.81)	4.91 (4.77-5.04)
Low	2.26 (2.13-2.38)	1.97 (1.87-2.07)	2.61 (2.44-2.79)	4.62 (4.42-4.82)
Medium	2.06 (1.90-2.21)	1.88 (1.75-2.01)	2.56 (2.31-2.80)	5.24 (4.93-5.55)
High	1.69 (1.48-1.91)	1.89 (1.68-2.10)	3.03 (2.61-3.45)	5.29 (4.85-5.74)
Very high	1.63 (1.45-1.81)	2.27 (2.07-2.46)	2.58 (2.27-2.89)	5.47 (5.08-5.86)
<i>P value</i>	<0.001	<0.001	0.736	<0.001
<i>Toothbrushing frequency</i>				
Less often than daily	3.09 (2.84-3.34)	2.11 (1.95-2.26)	3.00 (2.80-3.20)	4.62 (4.43-4.82)
Once a day	2.50 (2.41-2.60)	2.16 (2.08-2.23)	2.67 (2.54-2.79)	5.13 (4.99-5.28)
Twice a day or more	1.86 (1.76-1.96)	1.83 (1.74-1.91)	2.38 (2.23-2.54)	4.86 (4.67-5.06)
<i>P value</i>	<0.001	<0.001	<0.001	0.004
<i>Last dental visit</i>				
Never	2.44 (2.33-2.54)	1.95 (1.87-2.03)	3.05 (2.87-3.23)	5.46 (5.27-5.66)
Within last year	2.24 (2.08-2.41)	2.13 (2.00-2.27)	2.30 (2.12-2.48)	4.77 (4.55-4.99)
Over 1 year ago	2.28 (2.17-2.38)	2.10 (2.01-2.19)	2.60 (2.48-2.72)	4.72 (4.59-4.86)
<i>P value</i>	0.336	0.001	0.003	<0.001
<i>Smoking status</i>				
Never smoker	1.79 (1.72-1.86)	1.71 (1.65-1.77)	2.26 (2.16-2.37)	4.41 (4.29-4.53)
Former smoker	2.54 (2.17-2.92)	2.51 (2.20-2.83)	2.96 (2.69-3.23)	5.73 (5.42-6.05)
≤10 cigarettes/day	2.97 (2.70-3.23)	2.23 (2.05-2.42)	3.45 (3.15-3.75)	5.49 (5.18-5.80)
11-20 cigarettes/day	3.42 (3.21-3.63)	2.68 (2.52-2.83)	3.33 (3.05-3.62)	6.10 (5.78-6.42)
20+ cigarettes/day	3.70 (3.42-3.97)	2.86 (2.65-3.06)	3.44 (3.00-3.87)	5.51 (5.01-6.00)
<i>P value</i>	<0.001	<0.001	<0.001	<0.001

^a P values were derived from crude multilevel negative binomial regression models with the number of teeth used as an offset variable.

Table 3. Models for the associations of individual- and province-level characteristics with numbers of teeth with PPD $\geq$ 4mm and LOA $\geq$ 4mm among 35-44-years-old Chinese adults

Outcome	Contextual factor	Model 1 ^a	Model 2 ^a	Model 3 ^a
		RR ^b (95% CI)	RR ^b (95% CI)	RR ^b (95% CI)
PPD	Gini coefficient, %	0.93 (0.83-1.04)	0.92 (0.82-1.02)	0.94 (0.83-1.05)
	GDP per capita, thousand yuan	1.00 (0.98-1.03)	1.01 (0.98-1.03)	1.02 (0.95-1.10)
	Public health expenditure, %	1.00 (0.73-1.36)	1.03 (0.76-1.40)	0.99 (0.72-1.36)
	Dentist to population ratio ^c	1.04 (0.94-1.14)	1.05 (0.95-1.15)	1.08 (0.91-1.29)
	Dental therapist to population ratio ^c	1.18 (0.53-2.62)	1.15 (0.52-2.52)	1.15 (0.48-2.78)
	Rural population, %	1.00 (0.98-1.02)	1.00 (0.98-1.01)	1.03 (0.99-1.08)
	Minority ethnic groups, %	0.99 (0.97-1.01)	0.99 (0.97-1.01)	0.99 (0.97-1.00)
	Cigarette production, ten trillion	0.97 (0.93-1.01)	0.97 (0.94-1.01)	0.95 (0.87-1.05)
	Tobacco crops, ten thousand tons	0.99 (0.97-1.00)	0.99 (0.97-1.00)	1.00 (0.97-1.04)
	Smokers, %	1.03 (0.97-1.09)	1.03 (0.97-1.09)	1.04 (0.99-1.10)
LOA	Gini coefficient, %	0.95 (0.84-1.07)	0.94 (0.83-1.07)	0.97 (0.87-1.08)
	GDP per capita, thousand yuan	1.02 (1.00-1.05)*	1.03 (1.00-1.05)*	1.07 (1.01-1.14)*
	Public health expenditure, %	1.19 (0.87-1.63)	1.22 (0.87-1.71)	1.38 (1.04-1.83)*
	Dentist to population ratio ^c	1.02 (0.93-1.13)	1.03 (0.93-1.15)	0.81 (0.69-0.95)**
	Dental therapist to population ratio ^c	1.17 (0.51-2.69)	1.26 (0.52-3.07)	1.49 (0.69-3.22)
	Rural population, %	0.99 (0.97-1.01)	0.99 (0.97-1.00)	1.01 (0.98-1.05)
	Minority ethnic groups, %	0.98 (0.97-1.00)*	0.98 (0.97-1.00)*	0.99 (0.97-1.00)*
	Cigarette production, ten trillion	0.96 (0.92-1.01)	0.97 (0.97-1.01)	0.95 (0.88-1.03)
	Tobacco crops, ten thousand tons	0.98 (0.96-0.99)**	0.98 (0.96-0.99)*	0.99 (0.96-1.02)
	Smokers, %	0.97 (0.92-1.03)	0.97 (0.91-1.03)	1.04 (0.99-1.09)

^a Model 1 was unadjusted, Model 2 was adjusted for all individual-level factors and Model 3 additionally adjusted for all province-level factors.

^b Two-level negative binomial regression model was fitted and rate ratios (RR) reported. The number of teeth was used as an offset variable in all models.

^c Dentist to and dental therapist to population ratios are expressed per ten million population

* p<0.05; ** p<0.01; *** p<0.001

Table 4. Models for the associations of individual- and province-level characteristics with numbers of teeth with PPD $\geq$ 4mm and LOA $\geq$ 4mm among 65-74-years-old Chinese adults

Outcome	Contextual factor	Model 1 ^a	Model 2 ^a	Model 3 ^a
		RR ^b (95% CI)	RR ^b (95% CI)	RR ^b (95% CI)
PPD	Gini coefficient, %	0.94 (0.86-1.02)	0.94 (0.86-1.02)	0.95 (0.87-1.03)
	GDP per capita, thousand yuan	1.01 (0.99-1.02)	1.01 (0.99-1.02)	1.02 (0.97-1.07)
	Public health expenditure, %	0.92 (0.73-1.15)	0.93 (0.74-1.17)	0.85 (0.68-1.07)
	Dentist to population ratio ^c	1.05 (0.98-1.13)	1.05 (0.98-1.13)	1.07 (0.95-1.21)
	Dental therapist to population ratio ^c	1.35 (0.75-2.43)	1.27 (0.71-2.28)	1.01 (0.55-1.87)
	Rural population, %	1.00 (0.98-1.01)	1.00 (0.98-1.01)	1.02 (0.99-1.06)
	Minority ethnic groups, %	0.99 (0.98-1.00)	0.99 (0.98-1.01)	1.00 (0.98-1.01)
	Cigarette production, ten trillion	0.96 (0.94-0.99)**	0.96 (0.94-0.99)*	0.95 (0.90-1.01)
	Tobacco crops, ten thousand tons	0.99 (0.97-1.00)*	0.99 (0.98-1.00)*	1.00 (0.98-1.03)
	Smokers, %	1.03 (0.99-1.07)	1.03 (0.99-1.07)	1.04 (1.00-1.08)*
LOA	Gini coefficient, %	0.96 (0.89-1.03)	0.96 (0.88-1.03)	0.98 (0.92-1.05)
	GDP per capita, thousand yuan	1.01 (1.00-1.03)	1.02 (1.00-1.03)	1.04 (1.00-1.08)
	Public health expenditure, %	1.11 (0.90-1.35)	1.11 (0.90-1.36)	1.25 (1.04-1.49)*
	Dentist to population ratio ^c	1.00 (0.94-1.07)	1.01 (0.94-1.07)	0.86 (0.78-0.95)**
	Dental therapist to population ratio ^c	1.00 (0.59-1.69)	1.02 (0.60-1.74)	1.03 (0.62-1.70)
	Rural population, %	1.00 (0.99-1.00)	0.99 (0.98-1.00)	1.01 (0.98-1.03)
	Minority ethnic groups, %	0.99 (0.98-1.00)*	0.99 (0.98-1.00)*	0.99 (0.98-1.00)**
	Cigarette production, ten trillion	0.98 (0.95-1.01)	0.98 (0.95-1.01)	0.98 (0.93-1.04)

Tobacco crops, ten thousand tons	0.98 (0.97-0.99)**	0.98 (0.97-0.99)**	0.98 (0.96-1.00)
Smokers, %	0.98 (0.94-1.02)	0.98 (0.98-0.92)	1.02 (0.99-1.06)

^a Model 1 was unadjusted, Model 2 was adjusted for all individual-level factors and Model 3 additionally adjusted for all province-level factors.

^b Two-level negative binomial regression model was fitted and rate ratios (RR) reported. The number of teeth was used as an offset variable in all models.

^c Dentist to and dental therapist to population ratios are expressed per ten million population

* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Figure Legend

Figure 1. Sampling process for the 3rd National Oral Health Survey of China (2005). Dark grey boxes represent strata and light grey boxes represent clusters. GDP, Gross Domestic Product.