King's Research Portal

DOI:

[10.1016/j.jhep.2016.11.018](https://doi.org/10.1016/j.jhep.2016.11.018)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Kyrana, E., & Dhawan, A. (2016). Minimal hepatic encephalopathy in children, uncommon or unrecognised? - Time to act. *Journal of Hepatology*. <https://doi.org/10.1016/j.jhep.2016.11.018>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Accepted Manuscript

Minimal hepatic encephalopathy in children, uncommon or unrecognised? -
Time to act

Eirini Kyrana, Anil Dhawan

PII: S0168-8278(16)30694-8
DOI: <http://dx.doi.org/10.1016/j.jhep.2016.11.018>
Reference: JHEPAT 6341

To appear in: *Journal of Hepatology*

Received Date: 20 November 2016
Accepted Date: 21 November 2016

Please cite this article as: Kyrana, E., Dhawan, A., Minimal hepatic encephalopathy in children, uncommon or unrecognised? -Time to act, *Journal of Hepatology* (2016), doi: <http://dx.doi.org/10.1016/j.jhep.2016.11.018>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title: Minimal hepatic encephalopathy in children, uncommon or unrecognised? - Time to act.

Authors: Eirini Kyrana¹ and Anil Dhawan¹

Institution: ¹Paediatric Liver, GI and Nutrition Centre and MOWATLabs King's College Hospital, London

Corresponding author: Professor Anil Dhawan, Paediatric Liver, GI and Nutrition Centre and MOWATLabs at King's College Hospital, Denmark Hill, London, SE5 9RS.

Tel: 0044-2032994408, Fax: 0044-2032993657

E-mail: anil.dhawan@nhs.net

Keywords: hepatic encephalopathy, children, chronic liver disease

Electronic word count: 1273

Both authors declare no conflict of interest. Both authors have not received any financial support for the preparation of this manuscript.

Both authors contributed equally to the preparation of this manuscript.

Title: Minimal hepatic encephalopathy in children, uncommon or unrecognised? - Time to act.

Authors: Eirini Kyrana¹ and Anil Dhawan¹

Institution: Paediatric Liver, GI and Nutrition Centre and MOWATLabs, King's College Hospital, London.

Hepatic encephalopathy has been defined as a spectrum of neuropsychiatric abnormalities seen in patients with liver dysfunction after exclusion of an intrinsic brain disease [1]. Minimal hepatic encephalopathy (MHE) has been used to describe a subset of patients with the mildest form of encephalopathy, not detected on clinical examination, but detected on psychometric testing [1, 2]. The term applies to both patients with cirrhosis and patients with portosystemic shunts without intrinsic liver disease like isolated portal vein thrombosis [1]. The assessment of children with chronic liver disease (CLD) for encephalopathy is very challenging, particularly when assessing a frightened, unwell, younger child making it even more difficult to diagnose MHE.

There are a few studies to suggest that children with CLD are at risk of developing a cognitive impairment, which is worse when the disease manifests in the first year of life [3, 4]. These children have been shown to perform worse on measures of verbal, non-verbal and overall intelligence. In addition, liver transplantation does not correct these deficiencies and children post liver transplantation have been shown to have lower overall language skills, a decrease in non-verbal intelligence and lower academic achievements [5, 6]. The precise mechanism for these deficiencies is not known. Nutritional deficiencies are a possible factor, but the presence of MHE and the effect of toxic metabolites on the developing brain could be another. The ability to identify and treat factors that may be affecting the cognitive development of these young patients could have an exponential benefit for these children's long term academic and overall life achievements. We therefore welcome in this edition of Journal of Hepatology the paper by Srivastava et al [7] which endeavours to tackle this very interesting issue of MHE in children with CLD, providing us with some insight into the mechanism of its development as well as indices on imaging that can help diagnose it.

The prevalence of MHE as reported in this study seems to be similar in children and adults at about 50% [8-10]. The ammonia hypothesis suggests that raised ammonia levels lead to an increase of glutamine in brain astrocytes causing them to swell with subsequent neuro-dysfunction with additional contributory factors being endogenous benzodiazepines and the presence of inflammation [11, 12]. MHE in adults with cirrhosis has been associated with a reduced 5 year survival rate [13] and has been associated with reduced quality of life [14]. There is no similar data available in children but some neurological abnormalities described with MHE may persist even after liver transplantation [15] making it crucial to detect MHE and offer intervention.

Various modalities have been used to diagnose MHE including EEG, critical flicker frequency, continuous reaction time, inhibitory control test, computerised test

batteries, arterial spin labelling MRI with the psychometric hepatic encephalopathy score (PHES) considered to be the gold standard [16, 17].

Srivastava et al [7] studied prospectively 67 children with CLD and compare them to 37 healthy children. The study evaluates a plethora of parameters that could potentially be associated with MHE, including blood ammonia, IL6 and TNF α levels as well as calculating PELD and MELD scores and classifying their patients according to Child class. The children undergo NPT, magnetic resonance spectroscopy (HMRS) to study brain metabolite changes (specifically glutamine, choline and myoinositol) and also have diffusion tensor imaging (DTI) derived metrics. On the basis of NPT testing the authors find a prevalence of 50.7% for MHE, which is comparable to the one reported by Razek et al [8].

The children with CLD in comparison to the healthy children had relatively higher blood ammonia, but absolute levels were still in the normal range casting some doubts towards the ammonia hypothesis. On HMRS they had higher glutamine and lower choline and myoinositol in their brain. In addition they had a higher T1 signal intensity in the globus pallidus as well as a higher mean diffusivity in all brain regions on DTI, indicating cerebral oedema. Moving on to the more interesting comparisons between the subgroup of patients with MHE and those without MHE we see that between the two groups there was no difference in age, gender or aetiology of the liver disease. The differences in Child class, PELD and MELD score did not achieve statistical significance. This observation may suggest that MHE is more of a phenomenon related to the presence of portal hypertension rather than the degree of hepatic dysfunction.

The patients with MHE had significantly higher glutamine on HMRS. As previously mentioned these patients overall had a higher mean diffusivity on all brain regions on DTI in comparison to the controls, but the patients with MHE had an even higher mean diffusivity in six brain regions including frontal white matter. NPT, which was used to diagnose the MHE, had a significant positive correlation with brain choline and brain myoinositol. NPT had a significant negative correlation with blood ammonia, IL6 and TNF α , brain glutamine and with the mean diffusivity assessed by DTI. This led to the authors analysing and concluding that in fact frontal white matter mean diffusivity was the best discriminator of MHE with a sensitivity of 73.5% and a specificity of 100%.

The study by Srivastava et al [7] provides some insight and a possible explanation for these children's poor school performance, by linking NPT testing with neuroimaging. It would have been interesting if the authors could have also correlated the presence of MHE with school performance. A large proportion of the patients studied had autoimmune liver disease, whereas a more representative cohort of patients with chronic liver disease would include more patients with cholestatic liver disease, who are affected at a younger age. Nevertheless this study brings objective neuroimaging into the evaluation of children with chronic liver disease paving the way for further studies of this kind to evaluate younger children with chronic liver disease for the presence of MHE. Although it is difficult to ascribe neurocognitive complications in children with early onset cholestatic disorders to

MHE solely, the relationship intuitively appears plausible hence an early intervention including liver transplantation could improve long term scholastic and neurocognitive outcomes.

- [1] Ferenci P, Lockwood A, Mullen K, Tarter R, Weissenborn K, Blei AT. Hepatic encephalopathy-definition, nomenclature, diagnosis, and quantification: final report of the working party at the 11th World Congresses of Gastroenterology, Vienna, 1998. *Hepatology* 2002;35:716-721.
- [2] Gitlin N, Lewis DC, Hinkley L. The diagnosis and prevalence of subclinical hepatic encephalopathy in apparently healthy, ambulant, non-shunted patients with cirrhosis. *Journal of hepatology* 1986;3:75-82.
- [3] Stewart SM, Uauy R, Kennard BD, Waller DA, Benser M, Andrews WS. Mental development and growth in children with chronic liver disease of early and late onset. *Pediatrics* 1988;82:167-172.
- [4] Stewart SM, Campbell RA, McCallon D, Waller DA, Andrews WS. Cognitive patterns in school-age children with end-stage liver disease. *Journal of developmental and behavioral pediatrics : JDBP* 1992;13:331-338.
- [5] Stewart SM, Hildebeitel C, Nici J, Waller DA, Uauy R, Andrews WS. Neuropsychological outcome of pediatric liver transplantation. *Pediatrics* 1991;87:367-376.
- [6] Krull K, Fuchs C, Yurk H, Boone P, Alonso E. Neurocognitive outcome in pediatric liver transplant recipients. *Pediatric transplantation* 2003;7:111-118.
- [7] Srivastava A, Chaturvedi S, Gupta RK, Malik R, Mathias A, Jagannathan NR, et al. Minimal hepatic encephalopathy in children with chronic liver disease: prevalence, pathogenesis and magnetic resonance-based diagnosis. *Journal of hepatology* 2016.
- [8] Razek AA, Abdalla A, Ezzat A, Megahed A, Barakat T. Minimal hepatic encephalopathy in children with liver cirrhosis: diffusion-weighted MR imaging and proton MR spectroscopy of the brain. *Neuroradiology* 2014;56:885-891.
- [9] Dhiman RK, Kurmi R, Thumburu KK, Venkataramarao SH, Agarwal R, Duseja A, et al. Diagnosis and prognostic significance of minimal hepatic encephalopathy in patients with cirrhosis of liver. *Digestive diseases and sciences* 2010;55:2381-2390.
- [10] Das A, Dhiman RK, Saraswat VA, Verma M, Naik SR. Prevalence and natural history of subclinical hepatic encephalopathy in cirrhosis. *Journal of gastroenterology and hepatology* 2001;16:531-535.
- [11] Jayakumar AR, Rama Rao KV, Norenberg MD. Neuroinflammation in hepatic encephalopathy: mechanistic aspects. *Journal of clinical and experimental hepatology* 2015;5:S21-28.
- [12] Frederick RT. Current concepts in the pathophysiology and management of hepatic encephalopathy. *Gastroenterology & hepatology* 2011;7:222-233.
- [13] Ampuero J, Simon M, Montoliu C, Jover R, Serra MA, Cordoba J, et al. Minimal hepatic encephalopathy and critical flicker frequency are associated with survival of patients with cirrhosis. *Gastroenterology* 2015;149:1483-1489.
- [14] Agrawal S, Umapathy S, Dhiman RK. Minimal hepatic encephalopathy impairs quality of life. *Journal of clinical and experimental hepatology* 2015;5:S42-48.
- [15] Lin WC, Chou KH, Chen CL, Chen HL, Lu CH, Li SH, et al. Longitudinal brain white matter alterations in minimal hepatic encephalopathy before and after liver transplantation. *PloS one* 2014;9:e105887.
- [16] Weissenborn K. Diagnosis of minimal hepatic encephalopathy. *Journal of clinical and experimental hepatology* 2015;5:S54-59.
- [17] Li Y, Liu H, Yang J, Tian X, Yang H, Geng Z. Combining arterial-spin labeling with functional MRI measurement for characterizing patients with minimal hepatic encephalopathy. *Hepatology research : the official journal of the Japan Society of Hepatology* 2016.