King's Research Portal

DOI:

[10.1103/PhysRevB.94.155104](https://doi.org/10.1103/PhysRevB.94.155104)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Caio, M. D., Cooper, N. R., & Bhaseen, M. J. (2016). Hall response and edge current dynamics in Chern insulators out of equilibrium. *Physical Review B (Condensed Matter and Materials Physics)*, 94(15), [155104]. <https://doi.org/10.1103/PhysRevB.94.155104>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Hall response and edge current dynamics in Chern insulators out of equilibrium

M. D. Caio,¹ N. R. Cooper,² and M. J. Bhaseen¹¹*Department of Physics, King's College London, Strand, London WC2R 2LS, United Kingdom*²*T. C. M. Group, Cavendish Laboratory, J. J. Thomson Avenue, Cambridge CB3 0HE, United Kingdom*

(Received 14 April 2016; published 4 October 2016)

We investigate the transport properties of Chern insulators following a quantum quench between topological and nontopological phases. Recent works have shown that this yields an excited state for which the Chern number is preserved under unitary evolution. However, this does not imply the preservation of other physical observables, as we stressed in our previous work. Here we provide an analysis of the Hall response following a quantum quench in an isolated system, with explicit results for the Haldane model. We show that the Hall conductance is no longer related to the Chern number in the postquench state, in agreement with previous work. We also examine the dynamics of the edge currents in finite-size systems with open boundary conditions along one direction. We show that the late-time behavior is captured by a Generalized Gibbs Ensemble, after multiple traversals of the sample. We discuss the effects of generic open boundary conditions and confinement potentials.

DOI: [10.1103/PhysRevB.94.155104](https://doi.org/10.1103/PhysRevB.94.155104)

I. INTRODUCTION

Topological states of matter exhibit a wealth of novel properties due to their extreme resilience to local perturbations. A striking manifestation is the quantum Hall effect [1], where the exact quantization of the Hall response is immune to sample defects [2]. This robustness is intimately linked to the topological Chern invariant, which has direct signatures in quantum transport [3,4]. Recent theory and experiments have exposed many new examples of topological phenomena, including the relativistic quantum Hall effect in graphene [5–7], topological insulators [8–11], and Majorana edge modes [12]. Work has also focused on the generation of topological Floquet states through time-dependent driving [13–15].

Out of equilibrium, much less is known about the behavior of topological states. Early works on $p + ip$ superfluids have shown that topological invariants can be preserved following quenches of the interaction strength [16,17]. In the context of Chern insulators, recent studies have shown that the Chern number is robust under unitary evolution, even following quenches between topological and nontopological phases [18,19]. However, as we stressed in Ref. [19], this does not imply the persistence of all physical observables. The presence or absence of edge states for example, depends on the final Hamiltonian and not just the initial state. In a similar way, the Hall response does not necessarily remain quantized. This is consistent with recent calculations by Wang *et al.* [20,21], who have investigated the Hall response following a quantum quench, including an external reservoir to induce decoherence and reduce to the diagonal ensemble.

In this paper we examine the Hall response following a quantum quench in a completely isolated system undergoing unitary evolution. We focus on the Haldane model [22], as recently realized using cold atomic gases [23]. We show that the Hall response is no longer quantized following a quantum quench between the topological and nontopological phases, in spite of the preservation of the Chern index in infinite-size samples. In the zero frequency limit our results agree with those of Refs. [20,21], as oscillatory off-diagonal contributions vanish. The results are in good agreement with analytical approximations based on the low-energy Dirac

Hamiltonian. We further examine the detailed properties of the edge currents following a quantum quench in finite-size systems. We show that the late-time behavior following many traversals of the sample is quantitatively described by a generalized Gibbs ensemble (GGE) [24–26]. With a view towards cold atom experiments we also consider the effects of harmonic confinement potentials.

II. MODEL

The Haldane model describes spinless fermions hopping on a honeycomb lattice with a staggered magnetic field [22]. The Hamiltonian is given by

$$\hat{H} = -t_1 \sum_{\langle i,j \rangle} (\hat{c}_i^\dagger \hat{c}_j + \text{H.c.}) - t_2 \sum_{\langle\langle i,j \rangle\rangle} (e^{i\varphi_{ij}} \hat{c}_i^\dagger \hat{c}_j + \text{H.c.}) + M \sum_{i \in A} \hat{n}_i - M \sum_{i \in B} \hat{n}_i, \quad (1)$$

where $\hat{c}_j^\dagger$ and $\hat{c}_j$ are fermionic creation and annihilation operators obeying anticommutation relations $\{\hat{c}_j, \hat{c}_j^\dagger\} = \delta_{ij}$, and $\hat{n}_i \equiv \hat{c}_i^\dagger \hat{c}_i$. Here $\langle i,j \rangle$ and $\langle\langle i,j \rangle\rangle$ indicate summation over the nearest and next-to-nearest neighbor sites, respectively, t_1 and t_2 are the associated hopping parameters, and A and B label the two sublattices. The phase $\varphi_{ij} = \pm\varphi$ corresponds to the Aharonov-Bohm phase due to the staggered magnetic field and is taken positive (negative) in the anticlockwise (clockwise) hopping direction. The associated time-reversal symmetry breaking leads to a quantum Hall effect, in the absence of a net magnetic field. The energy offset M corresponds to spatial inversion symmetry breaking, allowing both nontopological and topological phases to be explored. The phase diagram of the Haldane model is shown in Fig. 1, where we assume that $|t_2/t_1| \leq 1/3$ so that the bands may touch but not overlap [22]; see Fig. 2.

The topological and nontopological phases are distinguished by the Chern index ν [3,22,27,28], which takes the values $\nu = \pm 1$ and $\nu = 0$, respectively. This is given by

$$\nu = \frac{1}{2\pi} \int d^2k \, \Omega, \quad (2)$$

FIG. 1. Phase diagram of the Haldane model showing topological ($\nu = \pm 1$) and nontopological phases ($\nu = 0$) [22]. The triangle and the square indicate the starting points of quenches along the dashed line, as shown in Figs. 4 and 5.

where $\Omega = \partial_{k_x} A_{k_y} - \partial_{k_y} A_{k_x}$ is the Berry curvature, $A_{k_\mu} = i\langle\psi|\partial_{k_\mu}|\psi\rangle$ is the Berry connection, and the integral is performed over the first Brillouin zone. The boundaries of the topological phases correspond to $M/t_2 = \pm 3\sqrt{3} \sin \varphi$ and are independent of t_1 ; see Fig. 1. In the remainder of the paper we choose $t_2 = t_1/3$ and set $t_1 = 1$ such that M is expressed in units of t_1 . We also set the intersite spacing a to unity.

III. QUANTUM QUENCHES

As in Ref. [19], we study quantum quenches between different points of the phase diagram shown in Fig. 1. We start from the ground state of $\hat{H}$ with parameters (M_0, φ_0) and abruptly change them to new values (M, φ) . The system evolves unitarily under the new Hamiltonian, leading to a time-dependent state $|\psi(t)\rangle = \exp[-i\hat{H}(M, \varphi)t/\hbar]|\psi_0\rangle$. Here $|\psi_0\rangle$ is the initial ground state, corresponding to a band insulator of the half-filled system at (M_0, φ_0) . As shown in Refs. [18,19], the corresponding Chern number remains unchanged from its initial value, even when quenching between different phases. Nonetheless, other observables may change. In finite-size systems, the topological and nontopological phases are distinguished by the presence or absence of edge states. The repopulation of these states following a quantum quench leads to changes in the edge currents and the orbital magnetization [19]. This is accompanied by the light-cone spreading of

FIG. 2. (a) Band structure of the Haldane model [22]. (b) At low energies, the model reduces to a sum of two Dirac Hamiltonians with a gapped relativistic dispersion relation.

currents into the interior of the sample and the onset of finite-size effects. It is evident that, for a finite-size system, the concept of a Chern index must eventually break down at late times following a quantum quench, as one cannot resolve momenta on scales smaller than the inverse system size, L^{-1} [29]. The Berry phase acquired upon circulating a plaquette of size $2\pi/L$ becomes ill-defined once the Berry connection A_{k_μ} becomes as large as $|A_{k_\mu}| \sim L$, and so too does the Chern number. Since, out of equilibrium, $|\psi\rangle$ is in a superposition of ground and excited states, it oscillates at the energy difference ΔE_k . The associated Berry connection $A_{k_\mu} = \langle\psi(t)|\partial_{k_\mu}|\psi(t)\rangle$ grows in time, t , as $A_{k_\mu} \sim [\partial \Delta E_k / \partial (\hbar k_\mu)]t = vt$ where $v = \partial \Delta E_k / \partial (\hbar k_\mu)$ is a characteristic difference of band velocities, limited by $v \lesssim 2c$ with c the maximum group velocity. Thus $|A_{k_\mu}| \sim L$ when $vt \sim L$, that is, the Chern number becomes ill-defined for time scales larger than the time required for light-cone propagation across the interior of the sample, $t \gtrsim L/2c$.

IV. HALL RESPONSE

In order to further differentiate the physical characteristics of the initial and final states, we examine the Hall response following a quantum quench. We apply a time-dependent electric field $\mathbf{E}(t) = \mathbf{E}_0 \cos \omega t$ and calculate the in-phase transverse response. In the presence of periodic boundary conditions it is convenient to generate this electric field by means of an auxiliary magnetic flux threading a toroidal sample [2–4]; see Fig. 3. For recent work examining the dynamics of one-dimensional currents following “flux quenches” in ring geometries see Ref. [30].

In momentum space, the resulting Hamiltonian may be decomposed into a sum over independent modes $\hat{H} = \sum_{\mathbf{k}} \hat{H}_{\mathbf{k}}$ where $\hbar \mathbf{k} = q\mathbf{E}(t)$, or equivalently $\hbar \mathbf{k}(t) = \hbar \mathbf{k}(0) + (q\mathbf{E}_0/\omega) \sin \omega t$, where $q = -e$ is the charge of the carriers. Expanding the Hamiltonian to linear order in $\mathbf{E}_0$ yields $\hat{H}(t) = \sum_{\mathbf{k}} \hat{H}_{\mathbf{k}} + \hat{\mathbf{v}}_{\mathbf{k}} \cdot (q\mathbf{E}_0/\omega) \sin \omega t$, where $\hat{\mathbf{v}}_{\mathbf{k}} = d\hat{H}_{\mathbf{k}}/d\mathbf{k}$ is the velocity operator. Within the framework of time-dependent

FIG. 3. Setup used to evaluate the Hall response in the presence of periodic boundary conditions. An auxiliary time-dependent magnetic flux $\Phi(t)$ generates a longitudinal electric field $E_x(t) = -\frac{\partial \Phi(t)}{\partial t}$. Note that this flux does *not* correspond to the time-reversal symmetry-breaking parameter φ in the Haldane model. The Hall conductance $\sigma_{xy}(t)$ is obtained from the transverse current that is in phase with $E_x(t)$.

perturbation theory it is convenient to expand the state of the system in the basis of the unperturbed ($\mathbf{E}_0 = 0$) postquench Hamiltonian:

$$|\psi_{\mathbf{k}}(t)\rangle = \sum_{b=l,u} c_{b,\mathbf{k}}(t) e^{-iE_{b,\mathbf{k}}t/\hbar} |b,\mathbf{k}\rangle. \quad (3)$$

Here $b = l, u$ refer to the lower and upper band, respectively, and $E_{b,\mathbf{k}}$ are the energies of the single-particle states; see Fig. 2. The case where the coefficients $c_{b,\mathbf{k}}$ are constant describes the unperturbed ($\mathbf{E} = 0$) evolution of the system under the postquench Hamiltonian. In the presence of $\mathbf{E}$, first order perturbation theory yields $c_{b,\mathbf{k}}(t) \approx c_{b,\mathbf{k}} + \delta c_{b,\mathbf{k}}(t)$ where

$$\delta c_{b,\mathbf{k}}(t) = -\frac{i}{\hbar} \sum_{b'=l,u} \int_0^t dt' \mathcal{M}_{b,b'}(\mathbf{k}, t') e^{i\Delta_{b,b'}(\mathbf{k})t'} c_{b',\mathbf{k}}. \quad (4)$$

Here $\Delta_{b,b'}(\mathbf{k}) \equiv (E_{b,\mathbf{k}} - E_{b',\mathbf{k}})/\hbar$, and $\mathcal{M}_{b,b'}(\mathbf{k}, t') \equiv \langle b, \mathbf{k} | \hat{\mathbf{v}}_{\mathbf{k}} \cdot (q\mathbf{E}_0/\omega) \sin \omega t' | b', \mathbf{k} \rangle$ is the matrix element of the perturbation. In order to determine the Hall conductance, we examine the transverse current $\hat{J}_y = \sum_{\mathbf{k}} \langle \psi_{\mathbf{k}}(t) | q\hat{v}_y | \psi_{\mathbf{k}}(t) \rangle$, which flows in response to an electric field along the x direction. In this pursuit we neglect the zeroth order contribution which arises in the absence of E_0^x , due to the redistribution of carriers between the bands following the quench. At first order in E_0^x ,

$$J_y^{(1)} = 2 \text{Re} \left[\sum_{\mathbf{k}, b, b'} c_{b,\mathbf{k}}^* \delta c_{b',\mathbf{k}}(t) e^{i\Delta_{b,b'}(\mathbf{k})t'} \langle b, \mathbf{k} | e\hat{v}_y | b', \mathbf{k} \rangle \right]. \quad (5)$$

Substituting Eq. (4) into Eq. (5) and restricting attention to frequencies below the gap $\omega \ll \Delta_{b,b'}(\mathbf{k})$, one obtains

$$J_y^{(1)} = -\frac{E_x}{\hbar\omega} \sum_{\mathbf{k}, b, b'} \text{Re} \left\{ \mathcal{D} \left[\frac{e^{i\omega t}}{\omega + \Delta_{b,b'}(\mathbf{k})} + \frac{e^{-i\omega t}}{\omega - \Delta_{b,b'}(\mathbf{k})} \right] \right\}, \quad (6)$$

where $\mathcal{D} \equiv -i |c_{b,\mathbf{k}}|^2 \langle b, \mathbf{k} | q\hat{v}_y | b', \mathbf{k} \rangle \langle b', \mathbf{k} | q\hat{v}_x | b, \mathbf{k} \rangle$. In order to define the Hall conductance we extract the contribution to $J_y^{(1)}$ that is in phase with the electric field. Denoting $J_y^{(1)} = I_y \cos \omega t + \tilde{I}_y \sin \omega t$ we define $\sigma_{xy}(\omega) \equiv I_y/AE_x$, where A is the area of the sample. This yields

$$\sigma_{xy}(\omega) = \frac{q^2}{2\pi\hbar} \sum_b \int d^2k |c_{b,\mathbf{k}}|^2 \tilde{\Omega}_b(\mathbf{k}), \quad (7)$$

where

$$\tilde{\Omega}_b(\mathbf{k}) = -i \sum_{b' \neq b} \frac{\langle b, \mathbf{k} | \hat{v}_x | b', \mathbf{k} \rangle \langle b', \mathbf{k} | \hat{v}_y | b, \mathbf{k} \rangle - \text{H.c.}}{\omega^2 - \Delta_{b,b'}(\mathbf{k})^2}. \quad (8)$$

In the limit $\omega \rightarrow 0$, $\tilde{\Omega}_b(\mathbf{k})$ reduces to the Berry curvature of the b th band, and the postquench d.c. Hall conductance is given by

$$\sigma_{xy}(0) = \frac{q^2}{2\pi\hbar} \sum_b \int d^2k |c_{b,\mathbf{k}}|^2 \Omega_b(\mathbf{k}), \quad (9)$$

in agreement with Refs. [20,21]. An analogous result is also found in the context of Floquet systems [31]. Equation (9) is a generalization of the Thouless-Kohmoto-Nightingale-den Nijs (TKNN) formula [3] to handle (arbitrarily prepared) excited states. The usual TKNN formula is recovered in the limit where $|c_{l,\mathbf{k}}|^2 = 1$ and $|c_{u,\mathbf{k}}|^2 = 0$, corresponding

to the ground state with $\sigma_{xy}(0) = \nu q^2/h$. The result (9) has a straightforward physical interpretation in terms of a semiclassical Boltzmann-like approach. The center of mass velocity of a wave packet with momentum $\mathbf{k}$ and band index b is given by $\dot{\mathbf{r}}_{b,\mathbf{k}} = \partial \varepsilon_b(\mathbf{k})/\hbar \partial \mathbf{k} - (\mathbf{k} \times \hat{\mathbf{z}}) \Omega_b(\mathbf{k})$, where $\hbar \dot{\mathbf{k}} = q\mathbf{E}_0$. The second term is the anomalous velocity associated with the Berry curvature $\Omega_b(\mathbf{k})$ [32–35], where $\hat{\mathbf{z}}$ is a unit vector perpendicular to the sample. The corresponding current is given by

$$\mathbf{J} \equiv \mathbf{J}^{(0)} + \mathbf{J}^{(1)} = \sum_b \int \frac{d^2k}{(2\pi)^2} q \dot{\mathbf{r}}_{b,\mathbf{k}} |c_{b,\mathbf{k}}|^2. \quad (10)$$

This yields a contribution proportional to $\mathbf{E}_0$:

$$\mathbf{J}^{(1)} = -\frac{q^2}{2\pi\hbar} \sum_b \int d^2k |c_{b,\mathbf{k}}|^2 (\mathbf{E}_0 \times \hat{\mathbf{z}}) \Omega_b(\mathbf{k}), \quad (11)$$

in agreement with the Hall response (9). In equilibrium, this can be used to extract the Berry curvature [36] from measurements of the transverse drift [23,37].

In Fig. 4 we show numerical results for the Hall conductance in the postquench state, starting from the topological phase. It is readily seen that the values are no longer quantized in integer multiples of q^2/h . Similarly, quenches from the nontopological to the topological phase also fail to yield the quantized values found in the equilibrium ground state with $\nu = \pm 1$; see Fig. 5. Heuristically, the quench “heats up” the sample by promoting carriers to the upper band which contribute to the Hall response by the Berry curvature of that band, thus yielding a nonquantized Hall response.

FIG. 4. Hall conductance in units of q^2/h following a quantum quench from the topological phase with $\varphi = \pi/3$ and $M = 1$ (as indicated by the triangular symbol in Fig. 1) to $\varphi = \varphi'$ and $M = 1$. The gray dots are numerical results for the Haldane model obtained from Eq. (9). The black solid line is the analytical result for the corresponding quenches in the low-energy Dirac approximation, obtained by summing contributions from Eq. (13). The results are in quantitative agreement for small quenches in the vicinity of $\varphi = \pi/3$ and in qualitative agreement for larger quenches. The vertical dashed lines correspond to the boundaries of the topological phases. The Hall conductance remains numerically close to q^2/h for quenches within the same topological phase ($\nu = 1$) but does not saturate at $-q^2/h$ when quenching to the other topological phase ($\nu = -1$).

FIG. 5. Hall conductance in units of q^2/h following a quantum quench from the nontopological phase with $\varphi = \pi$ and $M = 1$ (as indicated by the square symbol in Fig. 1) to $\varphi = \varphi'$ and $M = 1$. The gray dots are numerical results for the Haldane model obtained from Eq. (9). The black solid line is the analytical result for the corresponding quenches in the low-energy Dirac approximation, obtained by summing contributions from Eq. (13). The results are in quantitative agreement for small quenches in the vicinity of $\varphi = \pi$ and in qualitative agreement for larger quenches. The vertical dashed lines correspond to the boundaries of the topological phases.

More generally, we may use Eq. (7) to determine the a.c. Hall response for frequencies smaller than the band gap, $\omega \ll \Delta_{b,b'}(\mathbf{k})$; see Fig. 6. For larger frequencies, non-diagonal contributions of the form $c_{b,\mathbf{k}}c_{b',\mathbf{k}}^*$ do not disappear under unitary evolution, and the current (5) must be employed.

Low-energy approximation

Analytical results for the Hall response (9) may be obtained within the framework of the low-energy Dirac Hamiltonian for sufficiently small quenches. At low energies, the Haldane

FIG. 6. Hall response $\sigma_{xy}(\omega)$ in units of q^2/h evaluated using Eq. (7) for frequencies below the direct band gap $\Delta \equiv \min[\Delta_{b,b'}(\mathbf{k})]$, following a quantum quench from the topological phase with $M = 1$ and $\varphi = \pi/3$, to the nontopological phase with $M = 1$ and $\varphi = \varphi'$. The three curves correspond to $\varphi' = 7\pi/8$ (circles), $\varphi' = \pi$ (triangles), and $\varphi' = 9\pi/8$ (squares), for which $\Delta \sim 0.67, 2$, and $0.67 \times t_1/\hbar$, respectively.

model may be described as the sum of two Dirac Hamiltonians $H = H_+ + H_-$ [22] where

$$H_\alpha = \begin{pmatrix} m_\alpha c^2 & -c p e^{i\alpha\theta} \\ -c p e^{-i\alpha\theta} & -m_\alpha c^2 \end{pmatrix}. \quad (12)$$

Here $\alpha = \pm$ labels the two inequivalent Dirac points, $c = 3t_1 a/2\hbar$ is the effective speed of light, $p \exp(i\theta)$ parametrizes the 2D momentum (p_x, p_y) , and $m_\alpha = (M - 3\sqrt{3}at_2 \sin \varphi)/c^2$ is the effective Dirac fermion mass [22]. In this representation, a quench of the Haldane parameters $M \rightarrow M'$ and $\varphi \rightarrow \varphi'$ corresponds to quenches of the effective masses $m_\alpha \rightarrow m'_\alpha$. Combining results for the Berry curvature [36], with the Dirac band occupation following a quench [19], one obtains

$$\sigma_{xy}^\alpha(0) = \frac{\alpha e^2}{2h} \int_0^\infty dp \frac{c p m'_\alpha (p^2 + c^2 m'_\alpha)}{\sqrt{p^2 + c^2 m_\alpha^2} (p^2 + c^2 m'_\alpha)^2}, \quad (13)$$

for the two Dirac points. This is in agreement with Refs. [20,21]. In deriving this result, the integral over the two-dimensional Brillouin zone has been replaced by an integral over the infinite two-dimensional momentum space. The solid lines in Figs. 4 and 5 correspond to the low-energy approximation $\sigma_{xy}(0) = \sigma_{xy}^+(0) + \sigma_{xy}^-(0)$. The approximation is in good agreement with the exact numerical results for the lattice model (1). For small quenches, which do not explore the nonlinear regime of the dispersion relation, the results agree quantitatively. For larger quenches, the approximation breaks down, but it still captures the qualitative features. For large quenches, with $|m - m'| \rightarrow \infty$, the contribution to the Hall conductance from a single Dirac point shows plateaus at $(\pm\pi/8)q^2/h$; see Fig. 7. These results are consistent with the notion that preservation of ν does *not* imply the preservation of the Hall response $\sigma_{xy}(0)$. In the absence of interactions, the final state is nonthermal and characterized by the occupations

FIG. 7. Postquench Hall conductance in units of q^2/h for a single Dirac point ($\alpha = +$), following a quench from $m > 0$ to m' . For the same choice of quench parameters, the other Dirac point ($\alpha = -$) displays the opposite Hall conductance. Note that for quenches in the Haldane model, the changes in mass for each Dirac point may differ, yielding a nonzero Hall response. The dashed lines correspond to the asymptotes $\sigma_{xy}(0) = \pm(\pi/8)q^2/h$, which can be determined analytically from Eq. (13).

FIG. 8. Finite-size cylindrical geometry with periodic (open) boundary conditions along the x (y) direction. When $\varphi \neq 0$ there are generically edge currents, J_1^x and J_N^x , flowing along the sample boundaries in opposite directions. After a quantum quench, the edge currents evolve as a function of time, and currents flow into the interior of the sample.

$c_{b,\mathbf{k}}$ in Eq. (9). In this excited state the Hall response need not be quantized.

V. GENERALIZED GIBBS ENSEMBLE

A quantitative description of the nonthermal postquench state can be obtained by considering the system in a cylindrical geometry with open boundary conditions along one direction; see Fig. 8. In our previous work we showed that the resulting edge currents undergo nontrivial dynamics, due to the presence or absence of edge modes in the spectrum [19].

In particular, the edge currents decay towards new values that depend on the postquench Hamiltonian and not just the initial state; see Fig. 9. This evolution is accompanied by light-cone spreading of currents into the interior of the sample, with the eventual onset of finite-size effects and resurgent oscillations. Here, we show that the long time behavior of the total edge current, after many traversals of the sample, is captured by a GGE [24–26]. The density matrix is given by

$$\hat{\rho}_{\text{GGE}} = Z^{-1} \exp \left(- \sum_{\gamma} \lambda_{\gamma} \hat{n}_{\gamma} \right), \quad (14)$$

where $\hat{n}_{\gamma}$ are the conserved occupations of a given energy state, γ labels the energy level and the momentum index, and $Z = \text{Tr} \exp(-\sum_{\gamma} \lambda_{\gamma} \hat{n}_{\gamma})$; see Appendix C. Here the λ_{γ} are generalized inverse temperatures. These are determined by the self-consistency condition that the mode occupations immediately after the quench coincide with the averages computed via the GGE, $\langle \hat{n}_{\gamma} \rangle = \text{Tr}(\hat{\rho}_{\text{GGE}} \hat{n}_{\gamma})$. This yields $\lambda_{\gamma} = \ln[(1 - \langle \hat{n}_{\gamma} \rangle)/\langle \hat{n}_{\gamma} \rangle]$ [24]. In Fig. 10 we show the comparison between the time-averaged total edge current along a single edge at late times and the predictions of the GGE. The results are in excellent agreement.

FIG. 9. Total edge current $J_1^x(t)$ in units of $qt_1 a/\hbar$ following a quantum quench within the topological phase with $\nu = -1$. The time t is measured in units of $\hbar/t_1$. We set $\varphi = \pi/3$ and quench M from 1.4 to -1.4 . The solid horizontal (green) line corresponds to the time-averaged total edge current, in the quasistationary regime before the onset of finite-size traversals. Inset: Late-time data after 29 traversals of the sample. The dashed (red) line corresponds to the prediction of the GGE. This agrees with the time average of the late-time data, as shown by the solid horizontal line (blue), to within approximately 3%.

VI. CONFINEMENT POTENTIALS

Thus far, we have explored the dynamics of the Haldane model in toroidal and cylindrical geometries, as periodic boundary conditions provide considerable simplifications for theory and simulation. In order to make clear predictions for experiment, it is instructive to consider finite-size samples with open boundaries, especially due to the use of optical traps for cold atoms. We first consider the effects of transverse confinement in the cylindrical geometry depicted in Fig. 8 before discussing the case of rotationally symmetric confinement. For earlier work exploring the effects of trapping potentials

FIG. 10. Comparison of the time-averaged total edge current in the Haldane model at late times (circles) and the prediction $\langle \hat{\rho}_{\text{GGE}} \hat{J}_1^x \rangle$ of the GGE (triangles) for quantum quenches with $\varphi = \pi/3$ held fixed and $M = 1.4 \rightarrow M'$. The time averaging is performed over a single traversal period of the finite-size system, following 29 traversals of the sample. The currents are measured in units of $qt_1 a/\hbar$.

FIG. 11. Equilibrium properties of the Haldane model in the cylindrical geometry shown in Fig. 8, with an additional harmonic confinement potential V_n given by Eq. (15). We consider the topological phase with $M = 0$ and $\varphi = \pi/3$ and set $N = 70, C = 1.2$, and $\zeta = 15$. (a) Particle density ρ_n as a function of the row index $n \in 1, \dots, 70$. At half-filling, the potential confines the particles into the region $\zeta < n < N - \zeta$, as indicated by the vertical dashed lines. This imposes an effective system size in which it is possible to observe edge effects. (b) Total longitudinal current J_n^x along the cylinder showing clearly separated edge currents, broadened due to the smooth potential. Hall currents exist in the interior of the sample due to the effective electric field generated by the trap.

on the equilibrium edge physics of topological systems see Refs. [38,39].

We consider the Haldane model in the setup shown in Fig. 8, with an additional harmonic confinement potential applied along the y direction, $V(y) \propto (y - y_0)^2$. Here y_0 is the center of the trap which we take to be located at the midpoint of the strip. It is convenient to parametrize the potential as

$$V_n = C \left(\frac{n - N/2}{\zeta - N/2} \right)^2, \quad (15)$$

where $n = 1, \dots, N$ labels the row of the strip, ζ controls the effective width of the confined sample, and C is a constant. For suitable choices of C and ζ , for a strip of a given width N , the equilibrium particle density is approximately uniform for $\zeta < n < N - \zeta$; see Fig. 11(a). The corresponding current profile in the topological phase shows clearly separated edge currents that are broadened due to the smooth confinement potential; see Fig. 11(b). In contrast to the case of hard wall boundaries [19], the bulk also contains longitudinal Hall currents if it corresponds to a topological phase with nonzero Chern number. These arise due to the effective transverse electric field generated by the harmonic potential.

Following a quantum quench, the edge currents spread towards the interior of the sample, as found in the case of hard wall boundaries [19]; see Fig. 12. The currents also spread outside the initial sample area to a lesser extent due to the harmonic confinement. The light-cone propagation is clearly visible in the presence of the trap, but the apparent speed of light differs slightly from the uniform case. This is attributed to the broadening of the edge current profiles and the presence

FIG. 12. Dynamics of the currents $|J_n^x(t)|$ following a quantum quench from the topological phase with $M = 0$ and $\varphi = \pi/3$ to the nontopological phase $M = 3$ and $\varphi = \pi/3$. We set $N = 70, C = 1.2$, and $\zeta = 15$. The spreading of the currents into the interior (and to a lesser extent the exterior) of the effective sample is visible even in the presence of the harmonic potential. However, the propagation speed departs from the effective speed of light, due to the broadening of the edge currents and the contribution of bulk Hall currents.

of the equilibrium bulk Hall currents, induced by the harmonic potential.

The case of fully open boundary conditions presents a significant increase in the numerical computations required but is not expected to lead to different results. In the presence of a rotationally symmetric harmonic trap, as shown in Fig. 13(a), broadened edge currents will flow on the effective sample boundaries; see Fig. 13(b). Likewise, equilibrium bulk Hall currents will also circulate (in the opposite sense to the edge currents) due to the effective radial electric field. Following a quantum quench, the edge currents will flow towards the interior of the sample, exhibiting light-cone propagation; this

FIG. 13. (a) Illustration of the potential landscape of a hexagonal optical lattice in the presence of a rotationally symmetric harmonic trap. (b) Circulating equilibrium currents in the resulting disk geometry. Following a quantum quench between the topological and the nontopological phases the edge currents are expected to propagate into the interior (and the exterior) of the sample at the effective speed of light. This will be smoothed out due to the broadening of the edge currents and the presence of the bulk Hall currents.

will be smoothed out by the edge current broadening and the bulk Hall currents, which are present even in equilibrium.

An estimate of the time scales for the light-cone propagation can be inferred from the experimental parameters used in Ref. [23]. Using the typical hopping parameter $t_1 \sim 10^2 \hbar$ Hz, the effective speed of light is $c \sim 10^3 a/s$, measured in intersite spacings per second, i.e., lattice site propagation occurs on millisecond time scales. The time scale for observing the onset of the current plateau in Fig. 9, and the light-cone propagation in Fig. 12, is thus of the order of several milliseconds; the unit of time in Figs. 9 and 12 is $\hbar/t_1 \sim 10^{-3}$ s. This is comparable with the time scales explored in experiment [23].

VII. CONCLUSIONS

In this paper we have explored the transport properties of Chern insulators following a quantum quench in an isolated system undergoing unitary evolution. The Hall conductance is no longer described by the ground state relation $\sigma_{xy}(0) = \nu q^2/h$, in spite of the preservation of ν in infinite-size systems; the Chern index is a property of the state, while the Hall response depends both on the state and the final Hamiltonian. In the presence of open boundary conditions, the total edge currents are described by a GGE at late times. It would be interesting to explore the quench dynamics of Chern insulators in experiment, probing their Hall response and edge current dynamics.

Note added. While this work was being finalized for publication the preprints [40,41] appeared which also investigate the Hall response out of equilibrium.

ACKNOWLEDGMENTS

We acknowledge helpful conversations with F. Essler, N. Goldman, and B. Halperin. This work was supported by EPSRC Grants EP/J017639/1 and EP/K030094/1. M.J.B. thanks the EPSRC Centre for Cross-Disciplinary Approaches to Non-Equilibrium Systems (CANES) funded under grant EP/L015854/1. M.J.B. and M.D.C. thank the Thomas Young Center. Statement of compliance with EPSRC policy framework on research data: all data accompanying this publication are directly available within the publication.

APPENDIX A: LOW-ENERGY APPROXIMATION

In Figs. 4 and 5 we compare the postquench Hall response in the Haldane model (1) with the low-energy Dirac approximation (13). For completeness we provide a derivation of the latter; see also Refs. [20,21]. For a single Dirac point, the occupation probability amplitudes of the lower and upper bands following an effective mass quench $m \rightarrow m'$ are given

by [19]

$$\begin{aligned} a(m, m', p) &= f_-(p, m) f_-(p, m') + f_+(p, m) f_+(p, m'), \\ b(m, m', p) &= f_+(p, m) f_-(p, m') - f_-(p, m) f_+(p, m'), \end{aligned} \quad (\text{A1})$$

respectively, where $f_{\pm}(p, m) \equiv \sqrt{\frac{1}{2}(1 \pm \frac{mc}{\sqrt{p^2 + m^2 c^2}})}$. Here we have parameterized the two-dimensional momentum space with (p, θ) , as in the main text. For a single Dirac point α , the Berry curvature of the upper band is given by $\alpha \Omega_D(p, m)$, where $\Omega_D(p, m) = c p m / (p^2 + m^2 c^2)^{3/2}$; the Berry curvature of the lower band has the opposite sign [36]. Substituting these results into Eq. (9) it follows that after a quench $m_{\alpha} \rightarrow m'_{\alpha}$,

$$\begin{aligned} \sigma_{xy}^{\alpha}(0) &= \frac{\alpha q^2}{2h} \int_0^{\infty} dp \Omega_D(p, m'_{\alpha}) \\ &\times (|b(m_{\alpha}, m'_{\alpha}, p)|^2 - |a(m_{\alpha}, m'_{\alpha}, p)|^2). \end{aligned} \quad (\text{A2})$$

Equation (13) follows straightforwardly [20,21].

APPENDIX B: EDGE CURRENTS

In order to study the dynamics of the edge currents, we consider the Haldane model (1) in a finite-size cylindrical geometry, with periodic (open) boundary conditions along the x (y) direction; see Fig. 8. We define the local current flowing through a site l of the lattice as [19]

$$\hat{\mathbf{J}}_l := -\frac{iq}{2} \sum_j \delta_{jl} (t_{lj} \hat{c}_l^{\dagger} \hat{c}_j - \text{H.c.}), \quad (\text{B1})$$

where t_{lj} is the hopping parameter, δ_{jl} is the vector joining site j to l , and the sum is performed over nearest and next-nearest neighbors. Each lattice index corresponds to a triplet of indices $\{m = 1, \dots, M; n = 1, \dots, N; s = A, B\}$, labeling the x and y positions of the unit cell, and the sublattice [19]. The total longitudinal current flowing along the lower edge of the cylinder in the x direction is given by $J_1^x = \langle \hat{J}_1^x \rangle = \sum_{ms} \langle \hat{J}_{m1s}^x \rangle$, as shown in Figs. 8 and 9.

APPENDIX C: GENERALIZED GIBBS ENSEMBLE

In the finite-size cylindrical geometry depicted in Fig. 8, the Hamiltonian can be diagonalized as $\hat{H} = \sum_{\Upsilon, k_x} \epsilon_{\Upsilon}(k_x) \hat{f}_{\Upsilon}^{\dagger}(k_x) \hat{f}_{\Upsilon}(k_x)$, where we have exploited the periodicity in the x direction, and $\Upsilon = 1, \dots, 2N$ labels the energy levels at each k_x point. In the noninteracting model (1) the number operators $\hat{n}_{\Upsilon}(k_x) = \hat{f}_{\Upsilon}^{\dagger}(k_x) \hat{f}_{\Upsilon}(k_x)$ are conserved. Denoting the pair of indices $\{\Upsilon, k_x\}$ by γ , the late-time values of the time-averaged edge currents are described by the GGE given in Eq. (14).

- [1] K. Klitzing, G. Dorda, and M. Pepper, *Phys. Rev. Lett.* **45**, 494 (1980).
- [2] R. B. Laughlin, *Phys. Rev. B* **23**, 5632(R) (1981).
- [3] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, *Phys. Rev. Lett.* **49**, 405 (1982).
- [4] Q. Niu, D. J. Thouless, and Y.-S. Wu, *Phys. Rev. B* **31**, 3372 (1985).

- [5] G. Murthy and R. Shankar, *Rev. Mod. Phys.* **75**, 1101 (2003).
- [6] K. S. Novoselov, A. K. Geim, S. V. Morozov, D. Jiang, M. I. Katsnelson, I. V. Grigorieva, S. V. Dubonos, and A. A. Firsov, *Nature (London)* **438**, 197 (2005).
- [7] Y. Zhang, Y.-W. Tan, H. L. Stormer, and P. Kim, *Nature (London)* **438**, 201 (2005).
- [8] C. L. Kane and E. J. Mele, *Phys. Rev. Lett.* **95**, 226801 (2005).

- [9] B. A. Bernevig, T. L. Hughes, and S.-C. Zhang, *Science* **314**, 1757 (2006).
- [10] M. König, S. Wiedmann, C. Brüne, A. Roth, H. Buhmann, L. W. Molenkamp, X.-L. Qi, and S.-C. Zhang, *Science* **318**, 766 (2007).
- [11] A. B. Khanikaev, S. H. Mousavi, W.-K. Tse, M. Kargarian, A. H. MacDonald, and G. Shvets, *Nat. Mater.* **12**, 233 (2013).
- [12] L. Fu and C. L. Kane, *Phys. Rev. Lett.* **100**, 096407 (2008).
- [13] T. Kitagawa, E. Berg, M. Rudner, and E. Demler, *Phys. Rev. B* **82**, 235114 (2010).
- [14] N. H. Lindner, G. Refael, and V. Galitski, *Nat. Phys.* **7**, 490 (2011).
- [15] M. C. Rechtsman, J. M. Zeuner, Y. Plotnik, Y. Lumer, D. Podolsky, F. Dreisow, S. Nolte, M. Segev, and A. Szameit, *Nature (London)* **496**, 196 (2013).
- [16] M. S. Foster, M. Dzero, V. Gurarie, and E. A. Yuzbashyan, *Phys. Rev. B* **88**, 104511 (2013).
- [17] M. S. Foster, V. Gurarie, M. Dzero, and E. A. Yuzbashyan, *Phys. Rev. Lett.* **113**, 076403 (2014).
- [18] L. D'Alessio and M. Rigol, *Nat. Commun.* **6**, 8336 (2015).
- [19] M. D. Caio, N. R. Cooper, and M. J. Bhaseen, *Phys. Rev. Lett.* **115**, 236403 (2015).
- [20] P. Wang and S. Kehrein, *New J. Phys.* **18**, 053003 (2016).
- [21] P. Wang, M. Schmitt, and S. Kehrein, *Phys. Rev. B* **93**, 085134 (2016).
- [22] F. D. M. Haldane, *Phys. Rev. Lett.* **61**, 2015 (1988).
- [23] G. Jotzu, M. Messer, R. Desbuquois, M. Lebrat, T. Uehlinger, D. Greif, and T. Esslinger, *Nature (London)* **515**, 237 (2014).
- [24] M. Rigol, V. Dunjko, V. Yurovsky, and M. Olshanii, *Phys. Rev. Lett.* **98**, 050405 (2007).
- [25] M. Rigol, V. Dunjko, and M. Olshanii, *Nature (London)* **452**, 854 (2008).
- [26] J.-S. Caux and R. M. Konik, *Phys. Rev. Lett.* **109**, 175301 (2012).
- [27] S.-S. Chern, *Ann. Math. Second Ser.* **47**, 85 (1946).
- [28] M. V. Berry, *Proc. R. Soc. A* **392**, 45 (1984).
- [29] We are grateful to Prof. B. I. Halperin for this argument.
- [30] Y. O. Nakagawa, G. Misguich, and M. Oshikawa, *Phys. Rev. B* **93**, 174310 (2016).
- [31] H. Dehghani, T. Oka, and A. Mitra, *Phys. Rev. B* **91**, 155422 (2015).
- [32] R. Karplus and J. M. Luttinger, *Phys. Rev.* **95**, 1154 (1954).
- [33] W. Kohn and J. M. Luttinger, *Phys. Rev.* **108**, 590 (1957).
- [34] D. Xiao, M.-C. Chang, and Q. Niu, *Rev. Mod. Phys.* **82**, 1959 (2010).
- [35] M.-C. Chang and Q. Niu, *Phys. Rev. Lett.* **75**, 1348 (1995).
- [36] H. M. Price and N. R. Cooper, *Phys. Rev. A* **85**, 033620 (2012).
- [37] M. Aidelsburger, M. Lohse, C. Schweizer, M. Atala, J. Barreiro, S. Nascimbène, N. Cooper, I. Bloch, and N. Goldman, *Nat. Phys.* **11**, 162 (2014).
- [38] T. D. Stanescu, V. Galitski, and S. Das Sarma, *Phys. Rev. A* **82**, 013608 (2010).
- [39] M. Buchhold, D. Cocks, and W. Hofstetter, *Phys. Rev. A* **85**, 063614 (2012).
- [40] Y. Hu, P. Zoller, and J. C. Budich, *arXiv:1603.00513* (2016).
- [41] J. H. Wilson, J. C. W. Song, and G. Refael, *arXiv:1603.01621* (2016).