King's Research Portal

DOI:

[10.1017/S1352465816000631](https://doi.org/10.1017/S1352465816000631)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Brooks, S. K., Chalder, T., & Rimes, K. A. (Accepted/In press). Chronic Fatigue Syndrome: Cognitive, Behavioural and Emotional processing vulnerability factors. *Behavioural and Cognitive Psychotherapy*. <https://doi.org/10.1017/S1352465816000631>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Chronic fatigue syndrome: Cognitive, behavioural and emotional processing vulnerability factors

Dr. Samantha K. Brooks¹, Professor Trudie Chalder¹ & Dr. Katharine A. Rimes²

¹Department of Psychological Medicine, King's College London, Institute of Psychiatry, Weston Education Centre, Cutcombe Road, London SE5 9RJ.

²King's College London, Institute of Psychiatry, Psychology and Neuroscience, Dept. of Psychology, Henry Wellcome Building, De Crespigny Park, London SE5 8AF.

*Requests for reprints should be addressed to Dr. Katharine Rimes, King's College London, Institute of Psychiatry, Psychology and Neuroscience, Dept. of Psychology, Henry Wellcome Building, De Crespigny Park, London SE5 8AF, UK (Email: Katharine.Rimes@kcl.ac.uk)

Abstract

Background: Cognitive-behavioural models of chronic fatigue syndrome (CFS) suggest that personality factors such as perfectionism and high moral standards may contribute to the development of CFS.

Aims: To investigate cognitive, behavioural and emotional processing risk factors for CFS.

Methods: CFS patients (n=67) at a UK specialist clinic completed questionnaires about psychological characteristics both currently and retrospectively (six months pre-CFS onset). Responses were compared to those of healthy individuals (n=73) who rated their current characteristics. Forty-four relatives retrospectively rated the premorbid psychological characteristics of the CFS participants.

Results: CFS patients showed similar levels of current perfectionism to controls, though higher premorbid perfectionism. CFS patients showed greater self-sacrificial beliefs and more unhelpful beliefs about experiencing and expressing negative emotions, both currently but more markedly prior to onset. In the six months pre-illness onset, CFS patients showed more disruption to their primary goal and greater general stress than controls. Ratings of premorbid psychological characteristics by relatives

were consistent with patients' self-reports. The extent of overinvestment in one goal was significantly associated with fatigue.

Conclusions: Perfectionism, self-sacrificial tendencies, unhelpful beliefs about emotions, and perceived stress may be present to a greater extent pre-morbidly in CFS patients compared to healthy individuals.

Keywords: chronic fatigue syndrome, aetiology, causes, risk.

Introduction

Chronic fatigue syndrome

Chronic fatigue syndrome (CFS) is a condition characterised by severe and disabling fatigue, affecting physical and mental functioning, lasting at least six months (Fukuda et al., 1994).

Various different theories have been proposed to explain the etiology and treatment of CFS, including biopsychosocial (Harvey & Wessely, 2009) and psychodynamic (Taerk & Gnam, 1994) models, and models highlighting the importance of premorbid psychological processes such as attachment and mentalisation (Luyten & van Houdenhove, 2013). CFS has frequently been associated with personality traits such as neuroticism and perfectionism (van Geelen, Sinnema, Hermans & Kuis, 2007). The present study considers CFS in the context of a cognitive-behavioural model suggesting that individuals who develop CFS tend to have a premorbid personality characterised by perfectionism, high standards for performance and personal conduct, and beliefs that negative emotions are unacceptable signs of weakness (Surawy, Hackmann, Hawton & Sharpe, 1995). It is suggested that psychosocial stress or illness may cause acute fatigue (Hatcher & House, 2003), and that premorbid psychological characteristics may contribute to this fatigue becoming chronic. Surawy et al. (1995) suggest the fatigue experienced after stress or illness may result in the individual perceiving that they are no longer meeting their goals for performance, behaviour, or independent coping, thus activating underlying unhelpful beliefs (e.g. *'If I make mistakes or show negative emotions, others will think badly of me'*). The activation of such beliefs may cause distress and greater fatigue. Typically the individual then tries harder to meet their standards, despite increasing exhaustion. Surawy et al. (1995) suggest that when this fails, the individual falls into a state of chronic exhaustion and demoralisation. More recent theories have expanded on the idea of stress vulnerability, for example suggesting that CFS involves a 'crash' in the neurobiological stress system (van Houdenhove, Van Den Eede & Luyten, 2009), a persistent elevated stress response (Wyller, Malterud & Eriksen, 2009), or dysregulated stress signal sensitivity (Sraher, Skoluda, Rohleder & Nater, 2016). Chronic stress may be associated with pathophysiological changes in CFS such as disturbances in inflammation, oxidative stress, mitochondria or energy metabolism (Tanaka et al., 2015; Sraher et al., 2016).

Perfectionism and Goal Pursuit

Luyten, van Houdenhove, Cosyns and van den Broeck (2006) investigated the relationship between premorbid and postmorbid perfectionism, fatigue and depression in individuals with CFS. Contrasting retrospective and current ratings by CFS patients with responses from healthy controls (HC), they showed that CFS patients reported higher pre and post-morbid levels of both adaptive and maladaptive perfectionism, although CFS attenuated certain aspects of perfectionism. Personal standards appeared to be lower after illness onset and participants showed less concern for mistakes. A prospective study investigating predictors of CFS following an acute episode of glandular fever (Moss-Morris, Spence & Hou, 2011) found that perfectionism predicted CFS onset. Dittner, Rimes and Thorpe (2011) used a prospective design to explore whether students with negative perfectionism (i.e. believing that failure to meet high standards was unacceptable) were more at risk for fatigue after academic stress than those without perfectionistic beliefs. Negative perfectionism was positively associated with fatigue and predicted subsequent levels of both fatigue and depression; a case-control study (Deary & Chalder, 2008) has also shown an association between unhealthy perfectionism and fatigue. Related to the concept of perfectionism, action proneness – that is, the tendency toward direct action to meet goals – has been related to CFS: a retrospective study by van Houdenhove, Neerinckx, Onghena, Lysens and Vertommen (2001) suggested that a high level of action proneness may play a perpetuating role in CFS.

Failing to meet perfectionist expectations may worsen psychological wellbeing and striving to pursue unattainable goals may contribute to fatigue. Furthermore, it has been suggested that a tendency to invest self-esteem in one goal while neglecting other goals puts one at risk of distress if something happened making it difficult to maintain progress towards the primary goal (Lam & Power, 1991). There is evidence to support this hypothesis in depression (Arieti & Bemporad, 1980; Champion & Power, 1995; Lam, Green, Power & Checkley, 1996), with research also linking this to perfectionism (Powers, Koestner & Topciu, 2005) and self-criticism (Shahar, Henrich, Blatt, Ryan & Little, 2003). The present study aimed to explore whether this tendency to invest self-esteem in one goal may also be true for CFS.

Self-sacrifice

Ware's (1993) interviews with CFS patients suggested they often tended to put others' needs before their own. This is consistent with Surawy et al.'s (1995) observations of high moral standards for behaviour pre-onset in CFS patients. The CB model suggests that believing that putting one's own needs before those of others is selfish or wrong may lead to people failing to prioritise looking after their own health despite increasing fatigue. Conversely, they may feel guilty if unable to keep up with their standards for helping others, which could also contribute to distress and fatigue. However, the concept of self-sacrifice has not been systematically investigated in this population.

Beliefs about emotions

In a cross-sectional study, White and Schweitzer (2000) found no differences between CFS patients and controls with regards to the extent to which they tried to control their reactions when they were feeling anxiety, sadness or anger. In contrast, Rimes and Chalder (2010) found that CFS patients were more likely to report beliefs about the unacceptability of experiencing or expressing negative emotions than HC. An experimental study by Rimes, Ashcroft, Bryan and Chalder (2016) found that CFS participants had lower observer-rated emotional expression than HC, despite greater distress and higher autonomic arousal. Hiding distress takes effort which could add to fatigue, and could reduce the likelihood of receiving social support.

Design of CFS studies

Many of the studies we have described thus far have been cross-sectional or prospective, with a small number of retrospective studies (e.g. Luyten et al., 2006; van Houdenhove et al., 2001) and a lack of longitudinal studies. This is reflective of CFS literature as a whole, with very few longitudinal studies and only a small number of retrospective studies, most relating to childhood trauma and experiences (e.g. Fisher & Chalder, 2003; Heim et al., 2009). The current study aimed to address this gap in the literature by using a combination of retrospective and current ratings of psychological variables.

Aims and hypotheses

This study investigated psychological characteristics of people with CFS compared to healthy individuals.

The majority of previous studies exploring psychological characteristics of CFS patients have asked about current, rather than pre-morbid, tendencies. It is important to explore pre-morbid beliefs and other psychological characteristics as, once CFS has developed, individuals may be forced to modify such tendencies due to the impact of their illness. It is also possible that some individuals may believe that previous psychological characteristics may have contributed to their difficulties and made the deliberate decision to modify them. We therefore aimed to examine both current and pre-morbid perfectionist tendencies to better understand perfectionism in CFS patients, by asking them to rate their characteristics in the six months pre-CFS onset as well as currently. As these retrospective ratings may be affected by memory or other biases, relatives of CFS patients were also asked to rate the premorbid psychological characteristics of the affected individual. The current study enables replication of the pre-morbid and post-morbid perfectionism comparison used by Luyten et al. (2006), and extends it by exploring other variables such as self-sacrifice, beliefs about emotions and goal investment.

We hypothesised that CFS patients would show greater perfectionism, greater self-sacrificial beliefs and more negative beliefs about emotions than the control group. In line with previous evidence that individuals over-invested in one area to the exclusion of others are more prone to depression, it was hypothesised that CFS patients would report more over-investment in one goal (pre-onset) than HC. We also predicted that CFS patients would report more disruption in the area of their main goal in the 6 months prior to onset than HC report over past 6 months.

Method

The study was cross-sectional and retrospective, comparing psychological factors in CFS patients versus healthy controls. CFS patients were asked to rate beliefs both currently and 6 months pre-onset. The control group were only asked to rate beliefs in the last six months.

67 patients meeting the Oxford (Sharpe et al., 1991) or Center for Disease Control (Fukuda et al., 1994) criteria for CFS who had been assessed at the specialist CFS clinic at King's College London / South London and Maudsley NHS Foundation Trust completed questionnaires after assessment but before receiving cognitive behavioural therapy. Additionally, one relative, partner or close friend of each

patient were asked to complete questionnaires. A total of 44 family/friends completed questionnaires about the CFS patient – the discrepancy between numbers is due to some participants not wanting to ask a family member to take part, or not having anyone they felt was appropriate to ask. The control group consisted of 73 individuals from the general population who self-reported that they had never received treatment for a psychological illness, matched with the CFS sample for age, gender and education level. Controls were recruited via university and social media adverts and through assistance from colleagues.

Measures

Fatigue

The Chalder Fatigue Questionnaire (Chalder et al., 1993) was used to measure fatigue. Research has reported this scale to be reliable and valid (Cella & Chalder, 2010) and the original study reported Cronbach's alpha of 0.8903 (Chalder et al., 1993).

Perfectionism

Perfectionism was assessed using Campbell and DiPaula's (2002) Perfectionistic Self-Belief Scales, including two measures of self-oriented perfectionism (Importance of Being Perfect) and two measures of socially-prescribed perfectionism (Others' High Standards and Conditional Acceptance). Support for the four subscales of perfectionism used in this measure came from Stoeber and Childs (2010) who found satisfactory reliability (Cronbach's $\alpha \geq .70$) for all scores and replicated the original findings that Perfectionist Striving appears to be the most positive form of perfectionism and Conditional Acceptance is the most negative. Relatives also completed the perfectionism measure.

Self-sacrifice

The 16-item self-sacrifice subscale of the Young Schema Questionnaire (Young & Brown, 1994), containing items such as "I put others' needs before my own, or else I feel guilty" was used to assess self-sacrificial beliefs. There is some evidence to support the reliability and validity of this scale

(Schmidt, Joiner, Young & Telch, 1994), with Cronbach's alpha between .83 - .96 for all scores.

Relatives also completed the self-sacrifice measure.

Beliefs about emotions

The 12-item Beliefs About Emotions Scale (Rimes & Chalder, 2010) was used to assess beliefs about experiencing negative thoughts and feelings, showing negative thoughts and feelings to others and beliefs about the anticipated reactions of other people, if the person were to reveal their emotions.

Good reliability and validity of this scale has been shown, with Cronbach's alpha of 0.91 (Rimes & Chalder, 2010). Relatives also completed the beliefs about emotions measure.

Goals

The Roles and Goals questionnaire (Lam & Power, 1991) asks a number of questions about an individual's main roles or goals, in different life domains: employment, hobbies and interests, most important relationship, health, and any other significant role/relationships. Investment in each role or goal was assessed by asking questions including how much this made them feel good as a person; how much energy and effort they put in; to what extent success in other areas of their life depended on them doing well in this role or goal; and to what extent they thought that life would feel meaningless or unhappy without it. Participants reported whether there had been any events causing problems in any of the five domains, and rated the extent of the problem on a scale from 0 ("no problem") to 5 ("totally disrupted this area of my life") to give a measure of goal disruption. An extra item, "How stressful did you find your work?", rated on a 4-point scale, gave a measure of work stress. Investment in each goal was calculated using the mean of the ratings on the five questions in each domain; the goal with the highest investment score was the 'primary' goal. Over-investment was calculated using investment in primary goal minus investment in remaining goals. Relatives also completed the roles and goals measure.

This measure has good reliability with Cronbach's alpha of .85 (Lam & Power, 1991). This questionnaire was only used in the premorbid version of the questionnaire for the CFS patients, due to the severe disturbance to the roles / goals for most patients' current life as a result of their condition.

Anxiety, depression, stress

The Hospital Anxiety and Depression Scale (Zigmond and Snaith, 1983) used to assess anxiety and depression has been shown to be reliable and valid, with Cronbach's alpha of .83 for the anxiety scale and .82 for the depression scale (Bjelland, Dahl, Haug & Necklemann, 2002). The Perceived Stress Scale (Cohen, Kamarck & Mermelstein, 1983) measured the degree to which situations in one's life were appraised as stressful. This scale has shown good reliability (Cronbach's alpha $\geq$.84) (Cohen et al., 1983).

Relatives completed questionnaires regarding the six months before the patient developed CFS, while the non-clinical group completed similar questionnaires concerning their feelings over the previous six months. The CFS group completed both questionnaires.

Ethics

The study was approved by the Institute of Psychiatry Ethics Committee (ref. 308/02). Full informed consent was obtained from all participants. Data was anonymised. The CFS participants were given two versions of the questionnaires to complete, one for themselves and one to pass on to a relative who returned it directly to the researchers in a stamped addressed envelope provided. Both had the same study number attached so they could be matched afterwards. There was no record of which patient was given which number questionnaire.

Analysis

Data was entered into SPSS version 21.0 (IBM Corp, 2012) for analysis. Missing items on scales were replaced with that patient's mean response in cases where less than 25% of items were missing; if more than 25% of responses had been left blank, questionnaires were excluded.

Independent sample t-tests were used to examine group differences on questionnaire scores. CFS patients' ratings of their feelings over the previous six months, and the six months pre-onset, were compared using paired-sample t-tests. Matched-pairs analysis compared CFS patients' ratings to those of their relatives. The Chi-square test was used to calculate whether there was a significant difference in the percentage of CFS patients and controls reporting disruption to the primary goal. Pearson's

correlation was used to examine the relationship between fatigue and over-investment in primary goal. Effect sizes (Cohen's *d*) are reported for significant results.

Results

Sociodemographics and clinical characteristics

CFS patients and healthy controls were of similar age, with a mean age of 41.2 years for CFS patients and 37.5 years for controls ($t(100)=-1.301, p=.196$). Approximately 79.1% of CFS patients and 69.9% of controls were female ($X^2(1, N = 122) = 0.49, p = 0.36$). There was no significant difference in the proportion of each group who were of Caucasian ethnicity (96.7% of CFS, 89.2% of HC; $X^2(1, N = 99) = 2.34, p = 0.14$). The HC were more likely to be living with a partner (85.0%, compared to 55.2% of CFS patients; $X^2(1, N = 106) = 9.43, p = 0.002, d=0.68$), and were more likely to be in paid employment (86.3% compared to 32.8% of CFS patients; $X^2(1, N = 106) = 28.22, p = 0.000, d=1.28$). CFS patients were more likely to report having had previous psychological problems than HC (53.7% versus 26.0% respectively; $X^2(1, N = 106) = 5.35, p = 0.017, d=0.59$) as well as current psychological problems (52.2% of patients versus 5.5% of controls; $X^2(1, N = 106) = 24.03, p = 0.000, d=1.20$). Mean duration of CFS was 74.6 months. As expected, CFS participants had significantly higher scores on the Chalder Fatigue Scale (Mean 18.4, SD 8.74) than the HC (Mean 12.4, SD 3.73; $t(103)=-4.067, p=.000, d=0.89$); the HADS-depression scale (Mean 7.97, SD 4.82 vs. 2.53, 3.05 ; $t(103)=-6.382, p=.000, d=1.35$) and the HADS-anxiety scale (Mean 10.4, SD 5.52 vs. 5.30, 3.38 ; $t(103)=-5.247, p=.000, d=1.11$).

Psychological characteristics: current and premorbid

The scores on the self-report measures for psychological characteristics are shown in Table 1; for the CFS participants there are 'current' ratings (ratings for the previous six months) and also 'retrospective' ratings (the six months prior to CFS onset) while the healthy controls rated the previous six months only.

TABLE 1 APPROXIMATELY HERE

For ratings of the previous six months, independent t-tests indicated that CFS patients reported significantly higher levels of self-sacrifice ($t(62)=-2.370$, $p=.020$, $d=.0.74$) and more negative beliefs about emotions ($t(94)=-2.190$, $p=.031$, $d=0.82$) than healthy controls, but did not report significantly higher levels of perfectionism ($t(94)=-.864$, $p=.390$).

Paired t-tests were used to compare ratings for CFS participants for the previous six months with the six months prior to onset. These indicated that CFS patients reported significantly higher total perfectionism ($t(56)=-3.234$, $p=.002$, $d=0.42$), as well as scored higher on the subscales Conditional Acceptance ($t(64)=-3.233$, $p=.002$, $d=0.38$), Others' High Standards ($t(58)=-4.015$, $p=.000$, $d=0.47$) and Importance of Being Perfect ($t(65)=-3.591$, $p=.001$, $d=0.37$). Patients also reported higher self-sacrifice ($t(55)=-4.367$, $p=.000$, $d=0.25$) and more negative beliefs about emotions ($t(57)=-3.857$, $p=.000$, $d=0.36$) in the six months pre-CFS onset than at the current time.

Role and goal investment

Ratings of goal involvement and stress are shown in Table 2.

TABLE 2 APPROXIMATELY HERE

There were no significant differences in the scores for investment in work, health, hobby or 'other' roles and goals for the two groups; however, the control group reported significantly more investment in relationship goals than CFS patients ($t(101)=2.882$, $p=.005$, $d=0.60$). The two groups did not differ significantly in terms of over-investing in one goal. However, as predicted, CFS patients reported significantly greater disruption to their primary goal than healthy controls ($t(101)=-4.157$, $p=.000$, $d=1.04$), indicating that they found it more difficult to modify goals after a disrupting event than controls. There were no significant differences between the two groups in hours spent on work or work

stress, but CFS patients reported significantly higher perceived general stress ($t(99)=-2.340$, $p=.021$, $d=0.47$).

As expected, there was a significant correlation between fatigue and the extent to which the primary goal was overvalued relative to other goals, for the sample as a whole: $r(99) = .220$, $p=.029$. However this relationship did not remain significant after controlling for depression: $r(95) = .193$, $p=.058$.

Relatives' ratings about the CFS patient

Consistency was found between CFS participants' scores and their relatives' ratings. Results for the 44 CFS patients whose relatives completed questionnaires are presented in Table 3 alongside the results of the relatives' questionnaires.

TABLE 3 APPROXIMATELY HERE

There were no significant differences found between patients' pre-morbid scores and relatives' ratings about the patients for pre-morbid perfectionism ($t=-.988$, $p=.330$), self-sacrifice ($t=1.115$, $p=.272$), beliefs about emotions ($t=-.656$, $p=.516$), general stress ($t=-.150$, $p=.881$), or investment in any of the goals: work ($t=.887$, $p=.380$), hobby ($t=.964$, $p=.342$), relationship ($t=.723$, $p=.474$), health ($t=.636$, $p=.528$), or other role/goal ($t=-.656$, $p=.521$).

Discussion

This study provides evidence for several different psychological vulnerability factors for CFS – perfectionist standards; self-sacrificial beliefs/behaviours; negative beliefs about experiencing/expressing negative emotions; perceived stress and primary goal disruption.

After CFS onset, perfectionism (with the exception of Perfectionist Striving, a positive form of perfectionism), self-sacrifice and beliefs about emotions were all less negative than pre-onset, although self-sacrifice and beliefs about emotions were still more negative in CFS patients than controls. It should be noted that these findings reflect mean-level differences; it should not be assumed that all patients will have shown premorbid psychological factors. Additionally, the design of the study cannot prove that these are vulnerability factors or whether they interact with virus to influence onset; however, we can explore the differences between groups to support our hypothesis that psychological vulnerability factors contribute at least in part to the onset of CFS.

As predicted, CFS patients were more likely to report disruption to their primary goal in the six months pre-CFS onset compared to healthy individuals rating the previous six months. There was no evidence of over-investment in their main goal or time spent pursuing goals, but they reported more stress than healthy individuals in the six months pre-onset.

These findings are consistent with a cognitive behavioural model which suggests that CFS onset is triggered by a virus or other stressor making it difficult for the individual to maintain their usual high standards in their valued activities (e.g. Surawy et al., 1995). This model suggests that due to premorbid psychological characteristics of perfectionism and unhelpful beliefs about the unacceptability of negative emotions and putting others' needs before their own, the CFS-prone individual becomes distressed by this disruption to their valued goal and finds it difficult to access help. However, a recent paper in a general population found that the association between unhelpful beliefs about emotions and fatigue was not mediated by lower social support-seeking, but by greater emotional avoidance and lower self-compassion (Sydenham, Beardwood & Rimes, 2016). These possible processes require further investigation in people with CFS.

Beliefs that one should hide negative emotions and put others' needs before one's own could be regarded as forms of emotional and moral perfectionism. However it should be noted that these may also relate to other psychological constructs. For example people may put others' needs before their own because of low self-esteem or fear of rejection.

Developing CFS appears to be associated with a modification of some unhelpful beliefs: this needs to be considered in research investigating the psychological characteristics of people with CFS. It is unlikely that psychological intervention explains this change as participants were recruited at the beginning of treatment. It may be that getting ill helps individuals identify and re-evaluate unhelpful beliefs and behaviours, or it may be that they are simply too tired to maintain previous high standards. Some individuals may find that being forced to seek help or reduce their high standards helps to challenge their previous beliefs about the negative consequences of these actions. Future community-based studies could investigate further how psychological characteristics may change after fatigue onset even without treatment, and how this might impact on the course of the condition. Cognitive behavioural treatment of CFS typically addresses perfectionist attitudes which may be important for preventing relapse after recovery even if the patient does not feel currently able to pursue their desired high standards.

Of course it should be noted that these findings may also apply to individuals with other chronic conditions such as multiple sclerosis. Future research should consider the role of the psychological factors investigated here.

Limitations

As this study was retrospective and based on self-report data, the findings are open to bias. Memory and other biases can affect results. The inclusion of relatives' ratings was helpful in that they closely matched the participants' own ratings and therefore suggest that bias was minimal. However a prospective study is needed to confirm that these are premorbid vulnerability factors.

A further limitation is that the CFS participants completed the same questionnaires twice in one sitting (i.e. once to rate premorbid characteristics and once to rate current characteristics) which could have affected responses.

Additionally, it is important to note that although relatives' ratings were shown to confirm patient reports and therefore gave us some insight into premorbid traits and behaviour, it is possible that a gradual (rather than sudden) onset of CFS could account for premorbid characteristics. For example, it may be that having to cope with increasing fatigue, pain and anxiety during the onset of CFS leads to perfectionism as a means of coping – therefore relatives may have perceived the patients to be perfectionistic pre-CFS but this may be associated with the gradual onset of the illness rather than an innate personality trait. It would be useful for similar future studies to collect information on whether CFS onset was gradual or sudden so that this can be taken into account.

Unfortunately we did not keep information about participants' refusal rate and so cannot make demographic comparisons between participants and non-responders, which may have been useful.

Our results apply to a group of participants attending a tertiary CFS service and may not be generalizable to other CFS populations; for example, in primary care the patients are likely to be more divergent.

The current paper considered only psychological vulnerability factors for CFS, and we acknowledge that despite our significant results there are likely to be other influences in illness development – for example, biological factors such as cortisol. Future research could explore both the vulnerability factors discussed here and other, non-personality-related variables.

Implications

Our results suggest that CFS patients have a tendency to be perfectionist and self-sacrificial prior to illness onset; hold negative beliefs about emotions; and are more likely to have experience disruption in their primary goal prior to onset. These findings, if replicated, could have important implications for understanding the aetiology and treatment of CFS. For example, it may be necessary within CBT to address perfectionism, beliefs about emotions, self-sacrifice, and addressing disrupted goals. Even if perfectionist standards about performance, emotions or self-sacrificial behaviour have been attenuated

since onset because the individual is not able to maintain their desired standards, addressing these attitudes may be important for preventing relapse after recovery. It may be necessary to ask the patient about those factors directly or use questionnaires to assess them, because patients do not typically come to treatment presenting these as the problem and may not realise that they have, for example, more negative beliefs about emotions than other people. As well as addressing people's unhelpful beliefs about emotions it may also be the case that emotional processing needs to be encouraged in treatment for CFS. Indeed, Godfrey, Chalder, Ridsdale, Seed and Ogden (2007) found that the extent of emotional processing that occurred during therapy for chronic fatigue (either CBT or counselling) was a significant predictor of outcome. One way of addressing unhelpful beliefs about emotions might be mindfulness-based cognitive therapy (MBCT), a pilot study of which has suggested that such therapy may be useful for improving fatigue, depression and unhelpful beliefs in people with CFS (Rimes & Wingrove, 2013). For any treatment approach, it may be particularly challenging to improve outcomes for patients with very strong negative beliefs about emotions or consider perfectionism as adaptive and ego-syntonic; indeed, research has suggested that strong unhelpful beliefs about emotions are associated with poorer recovery rates in CBT for CFS (Flo & Chalder, 2014). It may be beneficial to draw on CBT for perfectionism (c.f. Shafran, Egan & Wade, 2010) to enhance the treatment of CFS, as this treatment was developed for individuals who usually have positive beliefs about perfectionism. Interventions should also consider focusing on roles and goals and strategies for modifying these if they are disrupted or problematic.

Acknowledgements and Conflicts of Interest

This research received no specific grant from any funding agency, commercial or not-for-profit sectors. Professor Chalder receives research support from the Biomedical Research Centre for the South London and Maudsley NHS Foundation Trust and the Institute of Psychiatry. This organisation had no role no involvement in study design, collection of data, analysis, writing up, or decision to submit. We express thanks to the individuals who participated in the study, and to all therapists involved in

collecting the data: Suzanne Roche, Sue Wilkins, Bella Stensnas, Mary Burgess and Brendan Thomas.

We would like to thank Helen Roy for her help with recruiting healthy control participants.

T Chalder is the author of the following books: Chalder T. (1995) *Coping with Chronic Fatigue*. Sheldon Press. London; Chalder T & Hussain K. (2002) *Self help for Chronic Fatigue Syndrome. A guide for young people*. Blue Stallion Publication. Oxon; Burgess M & Chalder T. (2005) *Overcoming Chronic Fatigue*. Constable & Robinson. London. K. Rimes and S. Brooks have no conflict of interest with respect to this publication.

Ethical Standards

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, and its most recent revision.

References

- Arieti, S., & Bemporad, J. R. (1980). The psychological organization of depression. *American Journal of Psychiatry*, 137(11), 1360-1365. doi:10.1176/ajp.137.11.1360
- Bjelland, I., Dahl, A., Haug, T., & Necklemann, D. (2002). The validity of the hospital anxiety and depression scale: an updated literature review. *Journal of Psychosomatic Research*, 52(2), 69–77. doi:10.1016/S0022-3999(01)00296-3
- Campbell, J. D., & DiPaula, A. (2002). Perfectionistic self-beliefs: Their relation to personality and goal pursuit. In: G. L. Flett & P. L. Hewitt (Eds.), *Perfectionism: theory, research and treatment* (pp. 181-198). Washington DC: American Psychological Association.
- Cella, M., & Chalder, T. (2010). Measuring fatigue in clinical and community settings. *Journal of Psychosomatic Research*, 69(1), 17-22. doi: 10.1016/j.jpsychores.2009.10.007

- Chalder, T., Berelowitz, G., Hirsch, S., Pawlikowska, T., Wallace, P., & Wessely, S. (1993). Development of a fatigue scale. *Journal of Psychosomatic Research*, 37(2), 147-53. doi: 10.1016/0022-3999(93)90081-p
- Champion, L. A., & Power, M. J. (1995). Social and cognitive approaches to depression: Towards a new synthesis. *British Journal of Clinical Psychology*, 34(4), 485-503. doi:10.1111/j.2044-8260.1995.tb01484.x
- Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, 24(4), 385-396. doi:10.2307/2136404
- Deary, V., & Chalder, T. (2014). Personality and perfectionism in chronic fatigue syndrome: A closer look. *Psychology & Health*, 25(4), 465-475. doi:10.1080/08870440802403863
- Dittner, A. J., Rimes, K., & Thorpe, S. (2011). Negative perfectionism increases the risk of fatigue following a period of stress. *Psychology & Health*, 26(3), 253-268. doi:10.1080/08870440903225892
- Fisher, L., & Chalder, T. (2003). Childhood experiences of illness and parenting in adults with Chronic Fatigue Syndrome. *Journal of Psychosomatic Research*, 54, 439-443. doi:10.1016/S0022-3999(02)00458-0
- Flo, E., & Chalder, T. (2014). Prevalence and predictors of recovery from chronic fatigue syndrome in a routine clinical practice. *Behaviour Research and Therapy*, 63, 1-8. doi:10.1016/j.brat.2014.08.013
- Fukuda, K., Straus, S. E., Hickie, I., Sharpe, M., Dobbins, J. G., Komaroff, A. & the International Chronic Fatigue Syndrome Study Group. (1994). The Chronic Fatigue Syndrome: A comprehensive approach to its definition and study. *Annals of Internal Medicine*, 121(12), 953-959. doi:10.7326/0003-4819-121-12-199412150-00009

Godfrey, E., Chalder, T., Ridsdale, L., Seed, P., & Ogden, J. (2007). Investigating the 'active ingredients' of cognitive behaviour therapy and counselling for patients with chronic fatigue in primary care: Developing a new process measure to assess treatment fidelity and predict outcome. *British Journal of Clinical Psychology*, 46(3), 253-272. doi:10.1348/014466506x147420

Harvey, S. B., & Wessely, S. (2013). Chronic fatigue syndrome: identifying zebras amongst the horses. *BMC Medicine*, 7, 58. doi:10.1186/1741-7015-7-58

Hatcher, S., & House, A. (2003). Life events, difficulties and dilemmas in the onset of chronic fatigue syndrome: a case-control study. *Psychological Medicine*, 33(7), 1185-1192.
doi:10.1017/s0033291703008274

Heim, C., Nater, U. M., Maloney, E., Boneva, R., Jones, J. F., & Reeves, W. C. (2009). Childhood trauma and risk for chronic fatigue syndrome: Association with neuroendocrine dysfunction. *Archives of General Psychiatry*, 66(1), 72-80. doi:10.1001/archgenpsychiatry.2008.508

IBM Corp. Released 2012. IBM SPSS Statistics for Windows, Version 21.0. Armonk, NY: IBM Corp.

Lam, D. H., Green, B., Power, M. J., & Checkley, S. (1996). Dependency, matching adversities, length of survival and relapse in major depression. *Journal of Affective Disorders*, 37(2-3), 81-90.
doi:10.1016/0165-0327(95)00075-5

Lam, D. H., & Power, M. J. (1991). A questionnaire designed to assess roles and goals: a preliminary study. *British Journal of Medical Psychology*, 64(4), 359-373. doi:10.1111/j.2044-8341.1991.tb01673.x

Luyten, P., & van Houdenhove, B. (2013). Common and specific factors in the psychotherapeutic treatment of patients suffering from chronic fatigue and pain. *Journal of Psychotherapy Integration*, 23(1), 14-27. doi:10.1037/a0030269

- Luyten, P., van Houdenhove, B., Cosyns, N., & van den Broeck, A-L. (2006). Are patients with chronic fatigue syndrome perfectionistic - or were they? A case-control study. *Personality and Individual Differences*, 40(7), 1473-1483. doi:10.1016/j.paid.2005.10.023
- Moss-Morris, R., Spence, M. J., & Hou, R. (2011). The pathway from glandular fever to chronic fatigue syndrome: can the cognitive behavioural model provide the map? *Psychological Medicine*, 41(5), 1099-1107. doi:10.1017/s003329171000139x
- Powers, T. A., Koestner, R., & Topciu, R. A. (2005). Implementation intentions, perfectionism, and goal progress: Perhaps the road to hell is paved with good intentions. *Personality and Social Psychology Bulletin*, 31(7), 902-912. doi:10.1177/0146167204272311
- Rimes, K. A., Ashcroft, J., Bryan, L. & Chalder, T. (2016). Emotional suppression in chronic fatigue syndrome: Experimental study. *Health Psychology*, 35(9), 979-986. doi:10.1037/hea0000341
- Rimes, K. A., & Chalder, T. (2010). The Beliefs About Emotions Scale: Validity, reliability and sensitivity to change. *Journal of Psychosomatic Research*, 68(3), 285-292. doi:10.1016/j.jpsychores.2009.09.014
- Rimes, K. A., & Wingrove, J. (2013). Mindfulness-based cognitive therapy for people with chronic fatigue syndrome still experiencing excessive fatigue after cognitive behaviour therapy: a pilot randomized study. *Clinical Psychology & Psychotherapy*, 20(2), 101-117. doi:10.1002/cpp.793
- Schmidt, N. B., Joiner, T. E., Young, J. E., & Telch, M.J. (1994). The Schema questionnaire: investigation of psychometric properties and the hierarchical structure of a measure of maladaptive schemas. *Cognitive Therapy and Research*, 19(3), 295-321. doi:10.1007/bf02230402
- Shafran, R., Egan, S. & Wade, T. (2010). Overcoming perfectionism. London: Constable & Robinson.

Shahar, G., Henrich, C. C., Blatt, S. J., Ryan, R., & Little, T. (2003). Interpersonal relatedness, self-definition, and their motivational orientation during adolescence: A theoretical and empirical integration. *Developmental Psychology*, 39(3), 470-483. doi:10.1037/0012-1649.39.3.470

Sharpe, M. C., Archard, L. C., Banatvala, J. E., Borysiewicz, L. K., Clare, A. W., David, A., ... Wright, D. J. M. (1991). A report—chronic fatigue syndrome: Guidelines for research. *Journal of the Royal Society of Medicine*, 84(2), 118-121.

Strahler, J., Skoluda, N., Rohleder, N., & Nater, U. M. (2016). Dysregulated stress signal sensitivity and inflammatory disinhibition as a pathophysiological mechanism of stress-related chronic fatigue. *Neuroscience & Biobehavioural Reviews*, 68, 298-218. doi:10.1016/j.neubiorev.2016.05.008

Stoeber, J., & Childs, J. H. (2010). The assessment of self-oriented and socially prescribed perfectionism: subscales make a difference. *Journal of Personality Assessment*, 92(6), 577-585. doi:10.1080/00223891.2010.513306

Surawy, C., Hackmann, A., Hawton, K., & Sharpe, M. (1995). Chronic fatigue syndrome: A cognitive approach. *Behaviour Research and Therapy*, 33(5), 535-544. doi:10.1016/0005-7967(94)00077-w

Syndenham, M., Beardwood, J., & Rimes, K. A. (2016). Beliefs about emotions, depression, anxiety and fatigue: A mediational analysis. Published online prior to print version. doi:<http://dx.doi.org/10.1017/S1352465816000199>

Taerk, G., & Gnam, W. (1994). A psychodynamic view of the chronic fatigue syndrome. The role of object relations in etiology and treatment. *General Hospital Psychiatry*, 16(5), 319-325. doi:10.1016/0163-8343(94)90018-3

Tanaka, M., Tajima, S., Mizuno, K., Ishii, A., Konishi, Y., Miike, T., Watanabe, Y. (2015). .Frontier studies on fatigue autonomic nerve dysfunction and sleep rhythm disorder. *Journal of Physiological Sciences*, 65, 483–498. doi:10.1007/s12576-015-0399-y

Van Geelen, S. M., Sinnema, G., Hermans, H. J. M., & Kuis, W. (2007). Personality and chronic fatigue syndrome: Methodological and conceptual issues. *Clinical Psychology Review*, 27, 885-903. doi:10.1016/j.cpr.2007.01.010

Van Houdenhove, B., Neerinx, E., Onghena, P., Lysens, R., & Vertommen, H. (2001). Premorbid “overactive” lifestyle in chronic fatigue syndrome and fibromyalgia: An etiological factor or proof of good citizenship? *Journal of Psychosomatic Research*, 51(4), 571-576. doi:10.1016/s0022-3999(01)00247-1

Van Houdenhove, B., Van Den Eede, F., & Luyten, P. (2009). Does hypothalamic-pituitary-adrenal axis hypofunction in chronic fatigue syndrome reflect a ‘crash’ in the stress system? *Medical Hypotheses*, 72, 701-705. doi:10.1016/j.mehy.2008.11.044

Ware, N. (1993). Society, mind and body in chronic fatigue syndrome: an anthropological view. In: *Chronic Fatigue Syndrome* (Ciba Foundation Symposium 173) (pp.62-82). Chichester: Wiley.

White, C., & Schweitzer, R. (2000). The role of personality in the development and perpetuation of chronic fatigue syndrome. *Journal of Psychosomatic Research*, 48(6), 515-524. doi:10.1016/s0022-3999(00)00087-8

Wyller, V. B., Malterud, K., & Eriksen, H. R. (2009). Can sustained arousal explain the chronic fatigue syndrome? *Behavioural and Brain Functions*, 5, 10. doi:10.1186/1744-9081-5-10.

Young, J. E., & Brown, G. (1994). Young Schema-Questionnaire (2nd ed.). In: J. E. Young (Ed.) *Cognitive therapy for personality disorders: A schema-focused approach* (Rev. ed.) (pp. 63-76). Sarasota, FL: Professional Resource Press.

Zigmond, A. S., & Snaith, R. P. (1983). The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, 67(6), 361–370. doi:10.1037/t03589-000

Tables

Table 1. Mean (SD) of psychological characteristics for CFS patients and healthy controls

	CFS patients (retrospective ratings)	CFS patients (current)	Controls (current)
	Mean (SD)	Mean (SD)	Mean (SD)
Perfectionism	97.6 (25.41)***	87.2 (24.35)	83.3 (19.81)
Self-sacrifice	68.4 (18.54)***	64.0 (16.70)	56.6 (12.73)*
Beliefs about emotions	38.9 (18.09)***	32.8 (15.85)	26.3 (11.91)*

*Significantly different to CFS patients' current ratings * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$*

Higher scores indicate more problematic beliefs

Table 2. Mean (SD) of roles and goals, over-investment in one goal, disruption to primary goal, hours worked, work stress and perceived general stress

	CFS patients	Controls
	Mean (SD)	Mean (SD)
Work	10.9 (3.79)	9.94 (3.25)
Health	10.0 (3.16)	10.6 (2.03)
Relationship	9.93 (3.62)	11.8 (2.50)**
Other	9.35 (2.46)	10.1 (2.30)
Hobby	7.80 (3.41)	7.76 (3.25)
Over-investing in main goal	.688 (.430)	.667 (.341)
Disruption to primary goal	2.16 (1.72)	0.66 (1.11)***
Work (hours per week spent on goal)	41.2 (14.21)	39.8 (21.18)
Work stress (0-3, higher scores indicate more stress)	1.75 (1.027)	1.72 (0.88)
Perceived general stress	43.8 (5.62)	41.5 (4.03)*

*Significantly different to CFS patients' ratings * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$*

Higher scores indicate greater investment in goals, greater disruption to primary goal, and greater stress.

Table 3. Relatives' ratings of psychological characteristics of the individual with CFS

	CFS patients self-report (n=44)	Relatives' ratings about the patient (n=44)
	Mean (SD)	Mean (SD)
Perfectionism	102.8 (21.30)	96.3 (27.31)
Self-sacrifice	73.3 (16.36)	71.3 (14.28)
Beliefs about emotions	44.7 (16.10)	41.6 (15.32)
General stress	43.8 (3.96)	44.0 (3.81)
Roles & goals: Work	11.6 (3.48)	11.2 (2.80)
Roles & goals: Hobby	8.42 (3.76)	8.00 (3.27)
Roles & goals: Relationship	10.2 (3.42)	10.2 (3.27)
Roles & goals: Health	10.2 (3.00)	9.98 (3.28)
Roles & goals: Other	9.84 (2.17)	10.1 (2.46)

*Significantly different to CFS patients' ratings * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (no significant differences found)*