King's Research Portal

DOI:

[10.1016/j.mrgentox.2010.12.003](https://doi.org/10.1016/j.mrgentox.2010.12.003)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Mizerovska, J., Dracinska, H., Frei, E., Schmeiser, H. H., Arlt, V. M., & Stiborova, M. (2011). Induction of biotransformation enzymes by the carcinogenic air-pollutant 3-nitrobenzanthrone in liver, kidney and lung, after intra-tracheal instillation in rats. *Mutation Research-Genetic Toxicology And Environmental Mutagenesis*, 720(1-2), 34 - 41. <https://doi.org/10.1016/j.mrgentox.2010.12.003>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

**Open Access document
downloaded from King's Research Portal
<https://kclpure.kcl.ac.uk/portal>**

Citation to published version:

Mizerovska, J., Dracinska, H., Frei, E., Schmeiser, H. H., Arlt, V. M., & Stiborova, M. (2011). Induction of biotransformation enzymes by the carcinogenic air-pollutant 3-nitrobenzanthrone in liver, kidney and lung, after intra-tracheal instillation in rats. *Mutation Research-Genetic Toxicology And Environmental Mutagenesis*, 720(1-2), 34 - 41, doi: 10.1016/j.mrgentox.2010.12.003

The published version is available at:

DOI: 10.1016/j.mrgentox.2010.12.003

This version: Post-print

<https://kclpure.kcl.ac.uk/portal/en/publications/induction-of-biotransformation-enzymes-by-the-carcinogenic-airpollutant-3nitrobenzanthrone-in-liver-kidney-and-lung-after-intratracheal-instillation-in-rats%2844f86a65-798d-42dd-95ad-eef18fd247b3%29.html>

NOTICE: this is the author's version of a work that was accepted for publication in *Mutation Research-Genetic Toxicology And Environmental Mutagenesis*. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in *Mutation Research-Genetic Toxicology And Environmental Mutagenesis*, [Vol.720, 1-2, 2011] DOI: 10.1016/j.mrgentox.2010.12.003

The copyright in the published version resides with the publisher.

When referring to this paper, please check the page numbers in the published version and cite these.

General rights

Copyright and moral rights for the publications made accessible in King's Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications in King's Research Portal that users recognise and abide by the legal requirements associated with these rights.'

- Users may download and print one copy of any publication from King's Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the King's Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Induction of biotransformation enzymes by the carcinogenic air pollutant 3-nitrobenzanthrone in liver, kidney and lung after intratracheal instillation in rats*

**Jana Mizerovská^a, Helena Dračínská^a, Eva Frei^b, Heinz H. Schmeiser^c, Volker M. Arlt^d,
Marie Stiborová,^{a,*}**

*^aDepartment of Biochemistry, Faculty of Science, Charles University, Albertov 2030, 128 40
Prague 2, The Czech Republic*

*^bDivision of Preventive Oncology, National Center for Tumour Diseases, German Cancer Research
Center, Im Neuenheimer Feld 280,, 69120 Heidelberg, Germany*

*^cResearch Group Genetic Alteration in Carcinogenesis, German Cancer Research Center,
Im Neuenheimer Feld 280, 69120 Heidelberg, Germany*

*^dSection of Molecular Carcinogenesis, Institute of Cancer Research, Institute of Cancer Research,
Sutton, Surrey SM2 5NG, United Kingdom*

Keywords: 3-nitrobenzanthrone; enzyme induction; cytochrome P450; NAD(P)H:quinone
oxidoreductase; DNA adduct

*Corresponding author: Department of Biochemistry, Faculty of Science, Charles University,
Albertov 2030, 128 40 Prague 2, Czech Republic. Tel.: +420-221951285; fax: +420-221951283.

E-mail address: stiborov@natur.cuni.cz (M. Stiborová)

Abbreviations: 3-ABA, 3-aminobenzanthrone; Ah, aryl hydrocarbon; COX, cyclooxygenase; CYP, cytochrome P450; dA, deoxyadenosine; dA- N^6 -ABA, 2-(2'-deoxyadenosin- N^6 -yl)-3-aminobenzanthrone; dG, deoxyguanosine; dG- N^2 -ABA, 2-(2'-deoxyguanosin- N^2 -yl)-3-aminobenzanthrone; dG-C8- N -ABA, N -(2'-deoxyguanosin-8-yl)-3-aminobenzanthrone; EROD, 7-ethoxyresorufin O -deethylation; LPO, lactoperoxidase; MPO, myeloperoxidase; MROD, 7-methoxyresorufin O -demethylation; NAT, N,O -acetyltransferase; 3-NBA, 3-nitrobenzanthrone; N -OH-3-ABA, N -hydroxy-3-aminobenzanthrone; NQO1, NAD(P)H:quinone oxidoreductase; RAL, relative adduct labelling; SDS, sodium dodecyl sulphate; SULT, sulfotransferase; TLC, thin-layer chromatography;

* This work was supported in part by the Grant Agency of the Czech Republic, grants 301/09/0472 and 305/09/H008, the Ministry of Education of the Czech Republic, grants MSM0021620808 and 1M0505, the German Cancer Research Center and by Cancer Research UK.

Abstract

3-Nitrobenzanthrone (3-NBA), a carcinogenic air pollutant, was investigated for its ability to induce cytochrome P450 (CYP) 1A1/2 and NAD(P)H:quinone oxidoreductase (NQO1) in liver, kidney and lung of rats treated by intratracheal instillation. **The organs used were from a previous study performed to determine the persistence of 3-NBA-derived DNA adducts in target and non-target tissues [Bieler *et al.*, *Carcinogenesis* 28 (2007) 1117-1121].** NQO1 is the enzyme reducing 3-NBA to *N*-hydroxy-3-aminobenzanthrone (*N*-OH-3-ABA) and CYP1A enzymes oxidize human metabolite of 3-NBA, 3-aminobenzanthrone (3-ABA), to the same reactive intermediate. 3-NBA and 3-ABA are both activated to species forming DNA adducts by cytosols and/or microsomes isolated from rat lung, the target organ for 3-NBA carcinogenicity, and from liver and kidney. Each compound generated the same five DNA adducts detectable by ³²P-postlabeling. When hepatic cytosols from rats treated with 0.2 or 2 mg/kg bw of 3-NBA were incubated with 3-NBA, DNA adduct formation was 3.2- and 8.6-fold higher, respectively, than in incubations with cytosols from control animals. Likewise, cytosols isolated from lungs and kidneys of rats exposed to 3-NBA more efficiently activated 3-NBA than those of control rats. This increase corresponded to an increase in protein levels and enzymatic activities of NQO1. Incubations of hepatic, pulmonary or renal microsomes of 3-NBA-treated rats **with 3-ABA** led to an up to 9.6-fold increase in DNA adduct formation relative to controls. The highest **induction in DNA adduct levels** was **found in lung**. The stimulation of DNA adduct formation correlated with expression of CYP1A1/2 induced by the intratracheal instillation of 3-NBA. The results demonstrate that 3-NBA induces NQO1 and CYP1A1/2 in livers, lungs and kidneys of rats after intratracheal instillation, thereby enhancing its own genotoxic and carcinogenic potential.

1. Introduction

Lung cancer is the most common malignant disease worldwide and is the major cause of death from cancer. Although tobacco smoking is the overwhelming cause of lung cancer, vehicular exhaust and ambient air pollution are also implicated as causative factors [1-4]. Nitro-aromatic compounds are widely distributed environmental pollutants found in exhaust from diesel and gasoline engines and on the surface of ambient air particulate matter. The increased lung cancer risk after exposure to these environmental sources and the detection of nitro-aromatics in the lungs of non-smokers with lung cancer has led to considerable interest in assessing their potential cancer risk [1-4].

The nitroaromatic 3-nitrobenzanthrone (3-nitro-7*H*-benz[*de*]anthracen-7-one, 3-NBA) (**Fig. 1**) occurs in diesel exhaust and in airborne particulate matter [5-8]. The main metabolite of 3-NBA, 3-aminobenzanthrone (3-ABA) (**Fig. 1**) [9,10], has been found in urine samples of salt mine workers occupationally exposed to diesel emissions [6], demonstrating human exposure to 3-NBA. 3-NBA is carcinogenic in rats, causing lung tumours after intratracheal instillation [7]. It is also an exceptionally potent mutagen in the Ames *Salmonella typhimurium* assay, scoring more than 6 million revertants per nanomole in strain YG1024 overexpressing bacterial nitroreductase and *O*-acetyltransferase [5]. 3-NBA has also been shown to be genotoxic in several other short-term tests and in the transgenic Muta Mouse assay [11,12]. Its genotoxicity has been further documented by the detection of specific DNA adducts formed *in vitro* as well as *in vivo* in rodents in various tissues [11,12,14-22].

In **liver, lung and kidney** most of the metabolic activation of 3-NBA is attributable to cytosolic human and rat NAD(P)H:quinone oxidoreductase (**NQO1**). **Human** *N,O*-acetyltransferase (NAT), NAT2, followed by NAT1, sulfotransferase (SULT), SULT1A1 and, to a lesser extent, SULT1A2 are the major activating phase II enzymes [10,18,23,24]. **Human** hepatic NADPH:cytochrome P450 (CYP) reductase also effectively activates 3-NBA [17], but in mice, 3-NBA is predominantly activated by NQO1 rather than NADPH:CYP reductase [18] (**Fig. 1**).

CYP1A1 and 1A2 (CYP1A1/2) are essential for the oxidative activation of 3-ABA in human and rat livers forming the same DNA adducts that are formed *in vivo* by 3-ABA or 3-NBA [25,26]. However, other organs also have the metabolic capacity to activate 3-ABA to form DNA adducts, independent from the CYP-mediated oxidation in the liver [27]. Previous results indicate that besides CYP enzymes expressed in several extra-hepatic tissues, peroxidases may play a role in the oxidative activation of 3-ABA [27]. Mammalian prostaglandin H synthase (cyclooxygenase, COX), lactoperoxidase (LPO) and myeloperoxidase (MPO) were found to be effective in activating 3-ABA [27] (Fig. 1). In animal experiments with Wistar rats the major role of CYP1A1 in 3-ABA activation in lungs and kidneys rather than peroxidases was elucidated [24,28]. In the same rat model we treated rats with single i.p. doses of 0.4, 4 or 40 mg/kg body weight (bw) of 3-NBA and saw that 3-NBA induces the major enzymes activating itself and its metabolite 3-ABA in livers [23], lungs and kidneys [24], thereby enhancing its own genotoxic and carcinogenic potential. Since inhalation is the major route by which airborne materials gain access to the body, primary exposure of the lungs to 3-NBA would constitute a better model system. However, experimental inhalation systems, with appropriate generation and characterization of exposure atmospheres, are expensive to acquire and maintain [29]. Therefore, direct instillation of a test material into the lungs *via* the trachea has been employed in many studies as an alternative exposure to inhalation. Instillation has also certain advantages over inhalation, the foremost being that the actual dose delivered to the lungs of each animal can be defined accurately.

Therefore, the objective of the present study was to evaluate whether the major enzymes activating 3-NBA (NQO1) and 3-ABA (CYP1A) are induced after a single intratracheal instillation of 3-NBA in the lungs, livers and kidneys of rats and if so, whether this induction also leads to higher DNA adduct levels, when cytosols and microsomes from these organs are incubated with 3-NBA and 3-ABA. The organs used in this study were from rats treated in an experiment to determine the persistence of 3-NBA-derived DNA adducts in target and non-target tissues [21,22].

2. Material and Methods

2.1. Chemicals

NADPH, deoxyadenosine (dA) 3'-monophosphate, deoxyguanosine (dG) 3'-monophosphate, menadione (2-methyl-1,4-naphthoquinone) and calf thymus DNA were from Sigma Chemical Co (St Louis, MO, USA); 7-ethoxyresorufin and 7-methoxyresorufin from Fluka Chemie AG (Buchs, Switzerland). Enzymes and chemicals for the ^{32}P -postlabeling assay were obtained from sources described [30]. All these and other chemicals were reagent grade or better.

2.2. Synthesis of 3-NBA and 3-ABA

3-NBA and 3-ABA were synthesized as described [15] and their authenticity was confirmed by UV spectroscopy, electrospray mass spectra and high field proton NMR spectroscopy.

2.3. Animal experiments

The study was conducted in accordance with the German Regulations for the Care and Use of Laboratory Animals, which is in compliance with Declaration of Helsinki. **Lungs, livers and kidneys used in this study were from rats (female Sprague–Dawley rats, $n = 3$ per group) treated with a single dose of 0.2 or 2 mg/kg body weight (bw) of 3-NBA by intratracheal instillation from a previous experiment to determine the persistence of 3-NBA-derived DNA adducts in target and non-target tissues [21,22].** Animals were killed 2 days after treatment by cervical dislocation. Lungs, livers and kidneys were removed, immediately after sacrifice, frozen in liquid nitrogen and stored at -80°C until isolation of microsomal and cytosolic fractions.

2.4. Preparation of microsomal and cytosolic fractions

Microsomal and cytosolic fractions were isolated from the livers, lungs and kidneys of **rats, as** described [31,32]. Both subcellular preparations were analyzed for the presence of 3-NBA by HPLC on a CC 250/4 Nucleosil, 100-5 C18 HD column (Macherey-Nagel, 4 x 250 mm) preceded by a C-18 guard column. Eluent was 70% methanol in water, at a flow rate of 0.6 ml min^{-1} , and detection was at 254 nm. The standard of 3-NBA eluted with a retention time of 8.2 min. No 3-

NBA was detectable in microsomal and cytosolic fractions from rats that had been treated with this compound (data not shown).

2.5. Cytosolic incubations

The deaerated and argon-purged incubation mixtures, in a final volume of 750 μ l, consisted of 50 mM Tris-HCl buffer (pH 7.4), containing 0.2% Tween 20, cofactor for cytosolic NQO1 (1 mM NADPH), pooled liver, lung and kidney cytosolic sample from 3 rats, treated either with vehicle (control) or with 0.2 or 2 mg/kg bw of 3-NBA (0.5 mg of cytosolic protein), 30 μ M 3-NBA (dissolved in 7.5 μ l dimethylsulfoxide) and 0.5 mg of calf thymus DNA (2 mM dNp). The reaction was initiated by adding 3-NBA. Incubations with rat cytosols were carried out at 37°C for 3 hr; the cytosol-mediated 3-NBA-derived DNA adduct formation was found to be linear up to 4 hr [18]. Control incubations were carried out either (i) without activating system (cytosol), (ii) without NADPH, (iii) without DNA or (iv) without 3-NBA. After extraction with ethyl acetate, DNA was isolated from the residual water phase by the phenol/chloroform extraction method as described [18].

2.6. Microsomal incubations

Incubation mixtures, in a final volume of 750 μ l, consisted of 50 mM potassium phosphate buffer (pH 7.4), 1 mM NADPH, hepatic, pulmonary and renal microsomal samples from 3 rats (**pooled samples**), either control or treated with 0.2 or 2 mg/kg bw of 3-NBA (0.5 mg of microsomal protein), 100 μ M 3-ABA or 3-NBA (dissolved in 7.5 μ l dimethylsulfoxide) and 0.5 mg of calf thymus DNA. The reaction was initiated by adding 3-ABA or 3-NBA and were carried out at 37°C for 2 hr; the microsomal-mediated 3-NBA (3-ABA)-derived DNA adduct formation was found to be linear up to 3 hr [17,25]. Control incubations were carried out either (i) without activating system (microsomes), (ii) with activating system and 3-ABA or 3-NBA, but without DNA or (iii) with activating system and DNA but without 3-ABA or 3-NBA. After the incubation and extraction with ethyl acetate, DNA was isolated from the residual water phase by the phenol/chloroform extraction method as described [25].

2.7. ³²P-Postlabeling analysis

³²P-Postlabeling analysis using butanol extraction, and thin layer chromatography (TLC) and HPLC were performed as described [19]. Enrichment by butanol extraction has been shown to yield more adduct spots and a better recovery of 3-NBA (3-ABA)-derived DNA adducts than using enrichment by nuclease P1 digestion [15,19]. DNA adduct spots were numbered as reported [15,19]. As reference compounds deoxyadenosine (dAp) and deoxyguanosine (dGp) 3'-monophosphates (4 μmol/ml) (Sigma) were incubated with 3-NBA (300 μM) activated by XO (1 U/ml) (Sigma) in the presence of hypoxanthine and analyzed as **described** [13]. DNA adduct standard samples of 3-NBA, 2-(2'-deoxyadenosin-*N*⁶-yl)-3-aminobenzanthrone-3'-phosphate (dA3'p-*N*⁶-ABA), *N*-(2'-deoxyguanosin-*N*²-yl)-3-aminobenzanthrone-3'-phosphate (dG3'p-*N*²-ABA) and *N*-(2'-deoxyguanosin-8-yl)-3-aminobenzanthrone-3'-phosphate (dG3'p-C8-*N*-ABA), were prepared by reacting *N*-acetoxy-3-aminobenzanthrone with dAp or dGp and analysed as **described** [19].

2.8. Isolation of CYP1A1 and CYP1A2

Recombinant rat CYP1A1 protein was purified to homogeneity from membranes of *Escherichia coli* transfected with a modified *CYP1A1* cDNA [33,34], in the laboratory of H. W. Strobel (University of Texas, Medical School of Houston, Texas, USA) by P. Hodek (Charles University, Prague, Czech Republic). Rat CYP1A2 was purified to homogeneity from hepatic microsomes of rats treated with β-naphthoflavone **as described** [35].

2.9. Preparation of antibodies

Leghorn chicken were immunized subcutaneously three times a week with rat recombinant CYP1A1 or rat CYP1A2 antigens (0.1 mg/animal) emulsified in complete Freund's adjuvant for the first injection and in incomplete adjuvant for boosters. Immunoglobulin fraction was purified from pooled egg yolks using fractionation by polyethylene glycol 6000 [33,34].

Rabbit polyclonal antibodies against human NQO1 were from Sigma Chemical Co (St Louis, MO, USA).

2.10. Estimation of CYP1A1, CYP1A2 and NQO1 protein content in microsomes and cytosols of rat liver, lung and kidney

Immunoquantitation of rat hepatic, pulmonary and renal microsomal CYP1A1, CYP1A2 and of cytosolic NQO1 was done by sodium dodecyl sulphate (SDS)-polyacrylamide gel electrophoresis. Samples containing 70 µg microsomal or cytosolic proteins were subjected to electrophoresis on SDS/10% polyacrylamide gels [23,24]. After migration, proteins were transferred onto polyvinylidene difluoride membranes. Rat CYP1A1, CYP1A2 and NQO1 proteins were probed with the chicken polyclonal antibodies raised against CYP1A1 or CYP1A2 and rabbit antibodies against human NQO, respectively, as reported elsewhere [23,24,28,33,35]. The antibodies against rat CYP1A1 and those against rat CYP1A2 recognize CYP1A1 and CYP1A2 in liver microsomes always as two protein bands, corresponding to CYP1A1 and CYP1A2. The antibodies against human recombinant NQO1 recognize this enzyme in rat hepatic, pulmonary and renal cytosols as one protein band. Rat recombinant CYP1A1 and CYP1A2 (in SupersomesTM, Gentest Corp., Woburn, MA, USA) and human recombinant NQO1 (Sigma Chemical Co, St Louis, MO, USA) were used as positive controls to identify the bands of CYP1A1 and CYP1A2 in microsomes and NQO1 in cytosols. The antigen-antibody complex was visualized with an alkaline phosphatase-conjugated rabbit anti-chicken IgG antibody and 5-bromo-4-chloro-3-indolylphosphate/nitrobluetetrazolium as chromogenic substrate [23,24,28,33,35]. Band intensities were quantified by Elfoman software.

2.11. CYP1A1, CYP1A2 and NQO1 enzyme activity assays

The microsomal samples were characterized for CYP1A1/2 activity using 7-ethoxyresorufin *O*-deethylation (EROD) activity and for CYP1A2 using 7-methoxyresorufin *O*-demethylation (MROD) activity [33,34,36]. One unit of EROD and MROD activities is defined as the amount of enzyme catalyzing the production of 1 pmol of resorufin in 1 min. The cytosolic samples were characterised for NQO1 activity. NQO1 activity was measured essentially as described by Ernster [37], but because of low activities of this enzyme in rat cytosolic samples, the method was

improved by addition of cytochrome *c*. The standard assay system contained 25 mM Tris-HCl (pH 7.4), 0.07% bovine serum albumin, 200 μ M NADH and 10 μ M menadione (2-methyl-1,4-naphthoquinone) dissolved in methanol and 77 μ M cytochrome *c*. The enzyme activity was determined by following the conversion of cytochrome *c* at 550 nm on a Hewlett-Packard 8453 diode array spectrophotometer. One unit of activity is defined as the amount of enzyme catalyzing the conversion of 1 μ mol of cytochrome *c* in 1 min.

3. Results

3.1. The effect of intratracheal instillation of 3-NBA to rats on expression of enzymes activating 3-NBA and 3-ABA

In the present study, Western blots using polyclonal antibodies raised against NQO1, CYP1A1 and CYP1A2 showed that the expression of all these enzymes was induced after 2 days in **liver, lung and kidney of** rats that received 3-NBA by intratracheal instillation (Figs. 2 and 3).

The NQO1 protein was expressed in all organs of control (untreated) rats, being highest in the liver; 3.4- and 6.2-fold higher levels were found in liver than in lung and kidney, respectively (Fig. 2). The levels of NQO1 protein were enhanced **in liver** after intratracheal instillation of rats with 0.2 or 2 mg 3-NBA/kg bw, by 5.8- and 8.7-fold, respectively (Fig. 2). The expression of protein of this enzyme was also induced **dose-dependently** in kidney and lung, but to a lesser extent than in the **liver (Fig. 2)**. The increased expression of NQO1 in the liver was paralleled by an increased activity of this enzyme (Table 1). Because of the low activity of NQO1 in kidney and lung, the effect of treating rats with 3-NBA on NQO1 activity in these organs was not evaluated.

The expression of CYP1A1 and CYP1A2 in microsomes was also induced in rats after intratracheal instillation with 3-NBA. Chicken antibodies raised against rat CYP1A1 or CYP1A2 used in this work each recognized both CYP1A isoforms because of a high degree of homology in their proteins [38]. The efficiency of 3-NBA to induce expression of CYP1A1 was higher than that of CYP1A2 in all organs tested in this study. In lung even no induction of CYP1A2 expression was found at all (Fig. 3). The EROD and MROD activities, which were used as markers of CYP1A1/2 and CYP1A2 activities, respectively [38], were induced by 3-NBA in livers (Table 1). **Whereas** the increase in EROD activity was much lower than the increase in CYP1A1 expression, **induction of MROD closely correlated with the increase in CYP1A2 protein expression in rat liver.** Because of the low basal levels of CYP1A enzymes in kidney and lung, the effect of treating rats with 3-NBA on their activities with EROD and MROD was not determined.

3.2. The effect of intratracheal instillation of 3-NBA to rats on activation of 3-NBA by hepatic, pulmonary and renal cytosols and microsomes

Cytosolic samples isolated from liver, kidney and lung of rats 2 days **after intratracheal instillation of 3-NBA** were capable of reductively activating 3-NBA to species forming **a cluster of 5 adducts in calf thymus DNA** (see **spots 1–5 in Figure 4A** for the liver **cytosol**). Thin-layer chromatograms of ^{32}P -labeled DNA from control incubations carried out in parallel without cytosol, without DNA or without 3-NBA were devoid of adduct spots in the region of interest (data not shown). **Only in DNA from control incubations containing liver cytosols without NADPH 3-NBA-DNA adducts were found (up to 2.6 ± 0.3 adducts per 10^8 deoxynucleotides), formed by cytosolic NQO1 mediated by the basal NADPH level present in hepatic cytosols.** Cochromatographic analysis of individual spots on HPLC confirmed that adduct spots 1–5 that are formed with rat hepatic, pulmonary and renal cytosols are derived from 3-NBA by nitroreduction (data not shown). Three of these adducts were identified as 2-(2'-deoxyadenosin- N^6 -yl)-3-aminobenzanthrone (dA- N^6 -ABA; spot 1), *N*-(2'-deoxyguanosin- N^2 -yl)-3-aminobenzanthrone (dG- N^2 -ABA; spot 3) and *N*-(2'-deoxyguanosin-8-yl)-3-aminobenzanthrone (dG-C8-*N*-ABA; spots 4 and 5).

In the presence of NADPH, hepatic cytosol from control rats was 2.6- and 4.7-fold more effective to activate 3-NBA than cytosols from kidney and lung, respectively (**Fig. 5**). 3-NBA was more effectively activated by cytosols from all organs of rats treated with 3-NBA in a dose dependent manner (**Fig. 5**). Up to 8.6-fold higher levels of 3-NBA-derived DNA adducts were found in incubations with hepatic cytosols of rats treated with 2 mg 3-NBA/kg bw relative to those of control rats, whereas only 2.3- and 2-fold higher levels of adducts were found using cytosols of kidney and lung, respectively (**Fig. 5**).

Liver microsomes of control rats and those treated with 3-NBA activated 3-NBA. Only a 1.7-fold increase in 3-NBA-derived DNA adduct levels was found in liver microsomes isolated from rats treated with 2 mg 3-NBA/kg bw (Table 2). Because CYP1A1/2 were found to also activate 3-

NBA [17], this increase is probably caused by induction of these enzymes. No 3-NBA-derived DNA adducts were however detectable after incubating 3-NBA with any of the pulmonary or renal microsomes and NADPH (data not shown). Hence, it seems that the activities of microsomal NADPH:CYP reductase and/or CYP are not sufficient to activate 3-NBA to an extent that DNA adduct formation is detectable in these extrahepatic organs.

3.3. The effect of intratracheal instillation of 3-NBA to rats on activation of 3-ABA by rat hepatic, pulmonary and renal microsomes

In the presence of NADPH, rat liver, lung and kidney microsomes were capable of oxidizing 3-ABA to form DNA adducts (see Figure 4B). Microsomal activation of 3-ABA generated a pattern of DNA adducts consisting of a cluster of five adducts (spots 1-5 in Figure 4B for the liver microsomes). **No DNA adducts were generated without NADPH a cofactor of CYP-mediated system, (data not shown).** In the presence of NADPH, liver microsomes of control rats 6.4- and 12.4-fold more efficiently activated 3-ABA than microsomes isolated from kidney and lung, respectively (**Fig. 6**). Liver, lung and kidney microsomes isolated from rats after intratracheal instillation with 2 mg 3-NBA/kg bw exhibited 3.9-, 9.7- and 3.3-fold higher efficiencies to activate 3-ABA, respectively, than microsomes from control rats (**Fig. 6**). **Chromatograms** of DNA digests from control incubations carried out in parallel without microsomes, without NADPH, without DNA, or without 3-ABA were all devoid of adduct spots in the region of interest (data not shown).

4. Discussion

The present study has increased our knowledge on the potential of 3-NBA to induce several biotransformation enzymes after intratracheal instillation that are responsible for metabolic activation of this carcinogen (NQO1) and its human metabolite, 3-ABA (CYP1A1/2), in the lung, the target organ for 3-NBA carcinogenicity, in the liver and kidney of rats. The rat was used as an experimental model on the basis that the same enzymes activate 3-NBA and 3-ABA in livers of rat and human [15,17,18,23,25,27]. Therefore, the results should provide some indication of what might occur in **tissues** of humans **exposed** to this carcinogenic pollutant.

Recently, we have shown that in Wistar rats treated i.p with a single dose of 0.4, 4 or 40 mg/kg bw 3-NBA acts as an inducer of NQO1 and CYP1A1/2 **in liver**, lung and kidney [23,24]. This induction was found to lead to an increase in the activities of these enzymes and in their potential to activate 3-NBA in cytosols (NQO1) and 3-ABA in microsomes (CYP1A) to species forming DNA adducts [23,24]. 3-NBA delivered i.p. is absorbed *via* the mesenteric veins and lymphatic systems, and passes through the liver. Thus, its concentration and effect in this tissue was higher than in the distal tissues such as lung and kidney (induction of NQO1 and CYP1A1 expression in liver of these rats was up to 5- and 2-fold higher than in lungs and kidneys, respectively) [23,24]. Since **human** exposure to 3-NBA occurs primarily *via* the respiratory tract and inhaled particles, we here evaluated the potential of 3-NBA administered by intratracheal instillation on enzymes capable of reducing 3-NBA or oxidizing 3-ABA. Sprague-Dawley rats were used for this study, because this animal model was successfully utilized to evaluate the long-term persistence of 3-NBA-derived DNA adducts after treatment by instillation [22]. **The present results confirm that hepatic, renal and pulmonary cytosols and microsomes from Sprague-Dawley rats activate 3-NBA and 3-ABA to a similar extent as Wistar rats, producing the same pattern of DNA adducts [23,24,28].**

The results of this work show that 3-NBA **induces** NQO1 and CYP1A1/2 also **after intratracheal** instillation to rats. The highest increase in expression levels of the enzymes studied

was found again in the liver, followed by kidney and lung, with exception of CYP1A1 expression in the kidney. **However**, expression of this protein in this organ in control rats was hardly detectable by Western blotting, which might overestimate its **relative** increase after 3-NBA treatment. The increase in expression of CYP1A1/2 and particularly NQO1 in the liver correlated with increased enzymatic activities and the activation of 3-NBA and its metabolite, 3-ABA, by hepatic cytosols and microsomes. The induction of CYP1A1/2 and NQO1 by 3-NBA is probably caused by binding **of** this planar molecule to the aryl hydrocarbon (Ah) receptor, which is known to be responsible for induction of CYP1A [38,39] and is one of the mechanisms responsible for induction of NQO1 (see [40] for a review).

The induction of NQO1 and CYP1A1 in the livers by 3-NBA administered by intratracheal instillation was higher than in the liver of rats treated by intraperitoneal injection **[23]**; a more than one order of magnitude lower 3-NBA dose produced similar expression levels. This finding underlines the risk for humans when exposed to this carcinogenic air pollutant by inhalation. All the more so, if one considers that the influence of 3-NBA instillation on the activation of its metabolite 3-ABA was highest in lungs. Indeed, the levels of 3-ABA-derived DNA adducts generated by lung microsomes isolated from 3-NBA-treated rats were almost 10 times higher than those formed by pulmonary microsomes from controls. This is the highest value of enzyme induction determined in this study.

As it is known that both NQO1 and CYP1A1/2 are expressed in **liver, lung and kidney not only in** rats but also **in humans** [35,40,42], **these enzymes** could contribute significantly and specifically to the metabolic activation of 3-NBA and/or 3-ABA in the human lung, thereby mediating 3-NBA carcinogenicity in this organ.

Inter-individual variations in susceptibility and variations in drug-metabolizing enzyme activities in target tissues appear to be important determinants of cancer risk [41,42]. Expression levels and activities of NQO1 and CYP1A1/2 differ considerably among individuals, because the enzymes are influenced by several factors, including smoking, drugs and environmental chemicals

[38,41,43]. It should be noted that besides induction of these enzymes by several compounds including 3-NBA [23,,24,38-40,44 and present paper], different activities of NQO1 and **CYP1A1/2** are also determined by genetic polymorphisms. Thus, genetic polymorphisms in **both enzymes could** be important determinants of a possible lung cancer risk from 3-NBA. So far two polymorphisms in the human *NQO1* gene have been found in the general population, one of them being associated with an increased risk of urothelial tumours [45] and paediatric leukaemia [46]. The polymorphic expression of CYP1A1 has been attributed to altered expression of the aryl hydrocarbon (Ah) receptor, the transcription factor that modulates its regulation, or the Ah receptor nuclear translocator (Arnt) protein, its associated transcription factor [40,44,47]. Moreover, the *CYP1A1* **and CYP1A2 genes are** polymorphic [38,42,**48,49**]. So far, CYP1A1*2A, CYP1A1*2B and CYP1A1*4 polymorphisms have been found that might be associated with cancers of lung, oesophagus or breast and with acute myeloid leukaemia [50-53], **while the CYP1A2*1F polymorphism might be associated with an increased risk of colorectal cancer [48]. In addition, the combination of NAT2 plus CYP1A1*2A, CYP1A1*2C, or CYP1A2*1F genotypes, and that of CYP1A2*1F plus CYP1A2*1C may define a group of persons who are genetically susceptible to colorectal cancer [49].**

In conclusion, the results of the present study show for the first time that 3-NBA is capable of inducing NQO1 and CYP1A1/2 enzymes in rat liver, lung and kidney after intratracheal instillation. **The result of such an exposure is both an enhanced reductive activation of 3-NBA and an oxidative activation of its metabolite 3-ABA to species forming DNA adducts, because 3-NBA induces both NQO1 and CYP1A.** Because 3-NBA is a potent genotoxin and a suspected human carcinogen [17-19,22], the results found in this work support our efforts in determining the risk of 3-NBA to human health.

References

- [1] IARC monographs on the evaluation of carcinogenic risks to humans. Diesel and gasoline engine exhausts and some nitroarenes. International Agency for Research on Cancer, IARC Monogr. Eval. Carcinog. Risks Hum. 46 (1989) 1-458.
- [2] USEPA, Health assessment document for diesel engine exhaust, EPA/600/8- 90/057F, National Center for Environmental Assessment, Office of Research and Page 18 of 27 Development, U.S. Environmental Protection Agency, Washington, D.C. (2002), 669 pp.
- [3] P. Vineis, F. Forastiere, G. Hoek, M. Lipsett, Outdoor air pollution and lung cancer: recent epidemiologic evidence, *Int. J. Cancer* 111 (2004) 647-652.
- [4] E. Grashick, F. Laden, J.E. Hart, B. Rosner, T.J. Smith, D.W. Dockery, F.E. Speizer, Lung cancer in railroad workers exposed to diesel exhaust, *Environ. Health Perspect.* 112 (2004) 1539-1543.
- [5] T. Enya, H. Suzuki, T. Watanabe, T. Hirayama, Y. Hisamatsu, 3-Nitrobenzanthrone, a powerful bacterial mutagen and suspected human carcinogen found in diesel exhausts and airborne particulates, *Environ. Sci. Technol.* 31 (1997) 2772-2776.
- [6] A. Seidel, D. Dahmann, H. Krekeler, J. Jacob, Biomonitoring of polycyclic aromatic compounds in the urine of mining workers occupationally exposed to diesel exhaust, *Int. J. Hyg. Environ. Health* 204 (2002) 333-338.
- [7] E. Nagy, M. Zeisig, K. Kawamura, Y. Hisumatsu, A. Sugeta, S. Adachi, L. Möller, DNA-adduct and tumor formations in rats after intratracheal administration of the urban air pollutant 3-nitrobenzanthrone, *Carcinogenesis* 26 (2005) 1821-1828.
- [8] V. M. Arlt, 3-Nitrobenzanthrone, a potential human cancer hazard in diesel exhaust and urban air pollution: A review of the evidence, *Mutagenesis* 20 (2005) 399–410.
- [9] T. Hansen, A. Seidel, J. Borlak, The environmental carcinogen 3-nitrobenzanthrone and its main metabolite 3-aminobenzanthrone enhance formation of reactive oxygen intermediates in human A549 lung epithelial cells, *Toxicol. Appl. Pharmacol.* 221 (2007) 222-234.
- [10] M. Stiborová, V. Martínek, M. Svobodová, J. Šistková, Z. Dvořák, J. Ulrichová, V. Šimánek, E. Frei, H.H. Schmeiser, D.H. Phillips, V.M. Arlt, Mechanisms of the different DNA adduct forming potentials of the urban air pollutants 2-nitrobenzanthrone and carcinogenic 3-nitrobenzanthrone, *Chem. Res. Toxicol.* 23 (2010) 1192-120.
- [11] V.M. Arlt, K.J. Cole, D.H. Phillips, Activation of 3-nitrobenzanthrone and its metabolites to DNA-damaging species in human B-lymphoblastoid MCL-5 cells, *Mutagenesis* 19 (2004) 149-156.

- [12] V.M. Arlt, L. Zhan, H.H. Schmeiser, M. Honma, M. Hayashi, D.H. Phillips, T. Suzuki, DNA adducts and mutagenic specificity of the ubiquitous environmental pollutant 3-nitrobenzanthrone in Muta Mouse, *Environ. Mol. Mutagen.* 43 (2004) 186-195.
- [13] V.M. Arlt, C.A. Bieler, W. Mier, M. Wiessler, H.H. Schmeiser, DNA adduct formation by the ubiquitous environmental contaminant 3-nitrobenzanthrone in rats determined by ^{32}P -postlabeling, *Int. J. Cancer* 93 (2001) 450-454.
- [14] V.M. Arlt, H.R. Glatt, E. Muckel, U. Pabel, B.L. Sorg, A. H.H. Schmeiser, D.H. Phillips, Metabolic activation of the environmental contaminant 3-nitrobenzanthrone by human acetyltransferases and sulfotransferase, *Carcinogenesis* 23 (2002) 1937-1945.
- [15] V.M. Arlt, H.R. Glatt, E. Muckel, U. Pabel, B.L. Sorg, A. Seidel, H. Frank, H.H. Schmeiser, D.H. Phillips, Activation of 3-nitrobenzanthrone and its metabolites by human acetyltransferases, sulfotransferases and cytochrome P450 expressed in Chinese hamster V79 cells, *Int. J. Cancer* 105 (2003) 583-592.
- [16] V.M. Arlt, B.L. Sorg, M. Osborne, A. Hewer, A. Seidel, H.H. Schmeiser, D.H. Phillips, DNA adduct formation by the ubiquitous environmental pollutant 3-nitrobenzanthrone and its metabolites in rats, *Biochem. Biophys. Res. Commun.* 300 (2003) 107-114.
- [17] V.M. Arlt, M. Stiborová, A. Hewer, H.H. Schmeiser, D.H. Phillips, Human enzymes involved in the metabolic activation of the environmental contaminant 3-nitrobenzanthrone: evidence for reductive activation by human NADPH:cytochrome P450 reductase, *Cancer Res.* 63 (2003) 2752-2761.
- [18] V.M. Arlt, M. Stiborová, C.J. Henderson, M.R. Osborne, C.A. Bieler, E. Frei, V. Martínek, B. Sopko, C.R. Wolf, H.H. Schmeiser, D.H. Phillips, The environmental pollutant and potent mutagen 3-nitrobenzanthrone forms DNA adducts after reduction by NAD(P)H:quinone oxidoreductase and conjugation by acetyltransferases and sulfotransferases in human hepatic cytosols, *Cancer Res.* 65 (2005) 2644-2652.
- [19] V.M. Arlt, H.H. Schmeiser, M.R. Osborne, M. Kawanishi, T. Kanno, T. Yagi, D.H. Phillips, T. Takamura-Enya, Identification of three major DNA adducts formed by the carcinogenic air pollutant 3-nitrobenzanthrone in rat lung at the C8 and N² position of guanine and at the N⁶ position of adenine, *Int. J. Cancer* 118 (2006) 2139-2146.
- [20] C.A. Bieler, M. Wiessler, L. Erdinger, H. Suzuki, T. Enya, H.H. Schmeiser, DNA adduct formation from the mutagenic air pollutant 3-nitrobenzanthrone, *Mutat. Res.* 439 (1999) 307-311.
- [21] C.A. Bieler, M. Cornelius, R. Klein, V.M. Arlt, M. Wiessler, D.H. Phillips, H.H. Schmeiser, DNA adduct formation by the environmental contaminant 3-nitrobenzanthrone after intratracheal instillation in rats, *Int. J. Cancer* 116 (2005) 833-838.

- [22] C.A. Bieler, M.G. Cornelius, M. Stiborova, V.M. Arlt, M. Wiessler, D.H. Phillips, H.H. Schmeiser, Formation and persistence of DNA adducts formed by the carcinogenic air pollutant 3-nitrobenzanthrone in target and non-target organs after intratracheal instillation in rats, *Carcinogenesis* 28 (2007) 1117-1121.
- [23] M. Stiborová, H. Dračínská, J. Hájková, P. Kadeřábková, E. Frei, H.H. Schmeiser, P. Souček, D.H. Phillips, V.M. Arlt, The environmental pollutant and carcinogen 3-nitrobenzanthrone and its human metabolite 3-aminobenzanthrone are potent inducers of rat hepatic cytochromes P450 1A1 and -1A2 and NAD(P)H:quinone oxidoreductase, *Drug Metab. Dispos.* 34 (2006) 1398-1405.
- [24] M. Stiborová, H. Dračínská, J. Mizerovská, E. Frei, H.H. Schmeiser, J. Hudeček, P. Hodek D.H. Phillips, V.M. Arlt, The environmental pollutant and carcinogen 3-nitrobenzanthrone induces cytochrome P450 1A1 and NAD(P)H:quinone oxidoreductase in rat lung and kidney, thereby enhancing its own genotoxicity, *Toxicology* 247 (2008) 11-22.
- [25] V.M. Arlt, A. Hewer, B.L. Sorg, H.H. Schmeiser, D.H. Phillips, M. Stiborová, 3-Aminobenzanthrone, a human metabolite of the environmental pollutant 3-nitrobenzanthrone, forms DNA adducts after metabolic activation by human and rat liver microsomes: evidence for activation by cytochrome P450 1A1 and P450 1A2, *Chem. Res. Toxicol.* 17 (2004) 1092-1101.
- [26] J. Mizerovská, H. Dračínská V.M. Arlt, , H.H. Schmeiser, E. Frei, M. Stiborová, Oxidation of 3-aminobenzanthrone, a human metabolite of carcinogenic environmental pollutant 3-nitrobenzanthrone, by cytochromes P450 – similarity between human and rat enzymes, *Neuro Endocrinol. Lett* 30 (Suppl. 1) (2009) 52-59.
- [27] V.M. Arlt, C.J. Henderson, C.R. Wolf, H.H. Schmeiser, D.H. Phillips, M. Stiborová, Bioactivation of 3-aminobenzanthrone, a human metabolite of the environmental pollutant 3-nitrobenzanthrone: evidence for DNA adduct formation mediated by cytochrome P450 enzymes and peroxidases, *Cancer Lett.* 234 (2006) 220-231.
- [28] M. Stiborová, H. Dračínská, M. Martínková, J. Mizerovská, J. Hudeček, P. Hodek, J. Liberda, E. Frei, H.H. Schmeiser, D.H. Phillips, V.M. Arlt, 3-Aminobenzanthrone, a human metabolite of the carcinogenic environmental pollutant 3-nitrobenzanthrone, induces biotransformation enzymes in rat kidney and lung, *Mutat. Res.* 676 (2009) 93-101.
- [29] K.E. Driscoll, D.L. Costa, G. Hatch, R. Henderson, G. Oberdorster, H. Salem, R.B. Schlesinger, Intratracheal instillation as an exposure technique for the evaluation of respiratory tract toxicity: uses and limitations, *Toxicol. Sci.* 55 (2000) 24–35.
- [30] D.H. Phillips, V.M. Arlt, The ³²P-postlabelling assay for DNA adducts, *Nat. Protoc.* 2 (2007) 2772-2781.

- [31] D. Aimová, L. Svobodová, V. Kotrbová, B. Mrázová, P. Hodek, J. Hudeček, R. Václavíková, E. Frei, M. Stiborová, The anticancer drug ellipticine is a potent inducer of rat cytochromes P450 1A1 and 1A2, thereby modulating its own metabolism, *Drug Metab Dispos.* 35 (2007) 1926-1934.
- [32] M. Stiborová, E. Frei, B. Sopko, K. Sopková, V. Marková, M. Laňková, T. Kumstýřová, M. Wiessler, H.H. Schmeiser, Human cytosolic enzymes involved in the metabolic activation of carcinogenic aristolochic acid: evidence for reductive activation by human NAD(P)H:quinone oxidoreductase, *Carcinogenesis* 24 (2003) 1695-1703.
- [33] M. Stiborová, V. Martínek, H. Rýdlová, P. Hodek, E. Frei, Sudan I is a potential carcinogen for humans: evidence for its metabolic activation and detoxication by human recombinant cytochrome P450 1A1 and liver microsomes, *Cancer Res.* 62 (2002) 5678-5684.
- [34] M. Stiborová, V. Martínek, H. Rýdlová, T. Koblas, P. Hodek, Expression of cytochrome P450 1A1 and its contribution to oxidation of a potential human carcinogen 1-phenylazo-2-naphthol (Sudan I) in human livers, *Cancer Lett.* 220 (2005) 145-154.
- [35] M. Stiborová, E. Frei, M. Wiessler, H.H. Schmeiser, Human enzymes involved in the metabolic activation of carcinogenic aristolochic acids: evidence for reductive activation by cytochromes P450 1A1 and 1A2, *Chem. Res. Toxicol.* 14 (2001) 1128-1137.
- [36] M.D. Burke, S. Thompson, R.J. Weaver, C.R. Wolf, R.T. Mayer, Cytochrome P450 specificities of alkoxyresorufin O-dealkylation in human and rat liver, *Biochem. Pharmacol.* 48 (1994) 923-936.
- [37] L. Ernster, DT-Diaphorase, *Methods Enzymol.* 10 (1967) 309-317.
- [38] S. Rendic, F.J. DiCarlo, Human cytochrome P450 enzymes: A status report summarizing their reactions, substrates, inducers, and inhibitors, *Drug Metab. Rev.* 29 (1997) 413-480.
- [39] A.T. Drahashuk, B.P. McGarrigle, K.E. Larsen, J.J. Stegeman, J.R. Olson, Detection of CYP1A1 protein in human liver and induction by TCDD in precision-cut liver slices incubated in dynamic organ culture, *Carcinogenesis* 19 (1998) 1361-1368.
- [40] D. Ross, J.K. Kepa, S.L. Winski, H.D. Beall, A. Anwar, D. Siegel, NAD(P)H:quinone oxidoreductase (NQO1): chemoprotection, bioactivation, gene regulation and genetic polymorphisms, *Chem.- Biol. Interact.* 129 (2000) 77-97.
- [41] G. Smith, L.A. Stanley, E. Sim, R.C. Strange, C.R. Wolf, Metabolic polymorphism and cancer susceptibility, *Cancer Surv.* 25 (1995) 27-65.
- [42] F.P. Perera, Environment and Cancer: Who are susceptible? *Science* 278 (1997) 1068-1073.
- [43] R.M. Chen, M.W. Chou, T.H. Ueng, Induction of cytochrome P450 1A in hamster liver and lung by 6-nitrochrysene, *Arch. Toxicol.* 72 (1998) 395-401.

- [44] J. Hukkanen, O. Pelkonen, J. Hakkola, H. Raunio, Expression and regulation of xenobiotic-metabolizing cytochrome P450 (CYP) enzymes in human lung, *Crit. Rev. Toxicol.* 32 (2002) 391-411.
- [45] W.A. Schulz, A. Krummeck, I. Rosinger, P. Eickelman, C. Neuhaus, T. Ebert, B.J. Schmitz-Dräger, H. Sies, Increased frequency of a null-allele for NAD(P)H-quinone oxidoreductase in patients with urological malignancies, *Pharmacogenetics* 7 (1997) 235-239.
- [46] J. Wiemels, A. Pagnamenta, G.M. Taylor, O.B. Eden, F.E. Alexander, M.F. Greaves, A lack of functional NAD(P)H:quinone oxidoreductase allele is selectively associated with pediatric leukemias that have MLL fusions. United Kingdom Childhood Cancer Study Investigators, *Cancer Res.* 59 (1999) 4095-4099.
- [47] M. Dickins, Induction of cytochromes P450, *Curr. Top. Med. Chem.* 4 (2004) 1745-1766.
- [48] **S. Y. Bae, S.K. Choi, K.R. Kim, C.S. Park, S.K. Lee, H.K. Roh, D.W. Shin, J.E. Pie, Z.H. Woo, J.H. Kang, Effects of genetic polymorphisms of MDR1, FMO3 and CYP1A2 on susceptibility to colorectal cancer in Koreans. *Cancer Sci.* 97 (2006)774-779.**
- [49] **K. Yoshida, K. Osawa, M. Kasahara, A. Miyaishi, K. Nakanishi, S. Hayamizu, Y. Osawa, A. Tsutou, Y. Tabuchi, E. Shimada, K. Tanaka, M. Yamamoto, J. Takahashi, Association of CYP1A1, CYP1A2, GSTM1 and NAT2 gene polymorphisms with colorectal cancer and smoking. *Asian Pac. J. Cancer Prev.* 8 (2007) 438-444.**
- [50] A. D'Alo, M. Voso, F. Guidy, G. Kaseiny, S. Sica, L. Pagano, S. Hohaus, G. Leone, Polymorphisms of CYP1A1 and glutathione S-transferase and susceptibility to adult acute myeloid leukaemia, *Haematologica* 89 (2004) 664-670.
- [51] C.-X. Yang, K. Matsuo, Z. Wang, K. Tajima, Phase I/II enzyme gene polymorphisms and esophageal cancer risk: A meta-analysis of the literature, *World J Gastroenterol.* 11 (2005) 2534-2538.
- [52] Y. Li, R. Millikan, D. Bell, L. Cui, C. Tse, B. Newman, K. Conway, Cigarette smoking, cytochrome P450 1A1 polymorphisms, and breast cancer among African-American and white women, *Breast Cancer Res.* 6 (2004) R460-R473.
- [53] Y. Li, R. Millikan, D. Bell, L. Cui, C. Tse, B. Newman, K. Conway, Polychlorinated biphenyls, cytochrome P450 1A1 (CYP1A1) polymorphisms, and breast cancer risk among African American women and white women in North Carolina: a population-based case-control study, *Breast Cancer Res.* 7 (2005) R12-R18.

Table 1

Rat hepatic cytosolic NQO1 and microsomal CYP1A1/2 and CYP1A2 specific enzyme activities

Enzyme activity	Control rat	3-NBA-treated rat	
		0.2 mg/kg bw	2 mg/kg bw
NQO1 ^a [μmo/min.mg]	237.6 ± 20.1	1136.7 ± 25.6 (4.8) ^{***}	1742.5 ± 315.7 (7.3) ^{***}
CYP1A1/2 ^b (EROD) [pmo/min.mg]	10.5 ± 1.0	12.1 ± 1.2 (1.2)	20.7 ± 1.8 (2.0) ^{***}
CYP1A2 ^c (MROD) [pmo/min.mg]	10.1 ± 1.1	12.4 ± 1.3 (1.2)	18.8 ± 1.8 (1.9) ^{***}

^aNQO1 enzyme activities were assayed using menadione as a substrate as described in Materials and Methods. ^bEROD and ^cMROD activities were used to measure CYP1A1/2 and CYP1A2 enzyme activities (see Materials and Methods). **Numbers are averages and SEs of four measurements, fold increase over control is shown in parenthesis.**

^{***}Significantly different from controls: $p < 0.001$ (Student's t-test).

Table 2

DNA adduct formation by 3-NBA activated by liver microsomes of rats either control or treated with 3-NBA

	RAL ^a (mean/10 ⁸ nucleotides)					Total
	Spot 1	Spot 2	Spot 3	Spot 4	Spot 5	
control + NADPH	0.3±0.02	0.3±0.03	0.9±0.07	0.2±0.02	0.1±0.01	1.8±0.1
3-NBA (0.2 mg/kg bw) + NADPH	0.4±0.03	0.3±0.04	1.4±0.1*	0.2±0.02	0.2±0.02	2.5±0.2
3-NBA (2 mg/kg bw) + NADPH	0.4±0.03	0.4±0.03	1.7 ± 0.1*	0.5±0.05*	0.2±0.02	3.1±0.2*

^a Mean RAL (relative adduct labeling) ± SE of triplicate *in-vitro* incubations; spot 1 = dA-*N*⁶-ABA, spot 3 = dG-*N*²-ABA, spots 4/5 = dG-C8-*N*-ABA.

**p*<0.05. Values significantly different from control (Student *t*-test).

Legends to Figures

Fig. 1. Pathways of metabolic activation and DNA adduct formation of 3-nitrobenzanthrone and 3-aminobenzanthrone. See text for details. 3-NBA, 3-nitrobenzanthrone; 3-ABA, 3-aminobenzanthrone; NQO1, NAD(P)H:quinone oxidoreductase; NAT, *N,O*-acetyltransferases; SULT, sulfotransferases; COX-1, cyclooxygenase 1; CYP, cytochrome P450; LPO, lactoperoxidase; MPO, myeloperoxidase; POR, NADPH:cytochrome P450 oxidoreductase; R = -COCH₃ or -SO₃H; dA-*N*⁶-ABA, 2-(2'-deoxyadenosin-*N*⁶-yl)-3-aminobenzanthrone; dG-*N*²-ABA, *N*-(2'-deoxyguanosin-*N*²-yl)-3-aminobenzanthrone; dG-C8-*N*-ABA, *N*-(2'-deoxyguanosin-8-yl)-3-aminobenzanthrone.

Fig. 2. Induction of NQO1 in the cytosol of livers, lungs and kidneys of rats after intratracheal instillation of 0.2 or 2 mg/kg bw of 3-NBA. Mean values $\pm$ SD (*n* = 3) in percent of control values from rats given vehicle only. Inset: immunoblots of NQO1 from untreated and 3-NBA-treated rats stained with antibody against human NQO1. Bar graphs show the relative intensities of the stained bands. Hepatic, pulmonary and renal cytosols were subjected to SDS-PAGE, proteins transferred to PVDF membranes and probed with antibody as described in Material and Methods. Values significantly different from control: **p*<0.05, ****p*<0.01 (Student *t*-test).

Fig. 3. Induction of CYP1A1 (A) and CYP1A2 (B) in livers, lungs and kidneys of rats after intratracheal instillation of 0.2 or 2 mg/kg bw of 3-NBA. Mean $\pm$ SD (*n* = 3) in percent of control values from rats given vehicle only. Inset: immunoblots of microsomal CYP1A1 (A) and CYP1A2 (B) from untreated and 3-NBA-treated rats stained with anti rat CYP1A1 and CYP1A2 antibodies. Bar graphs show the relative intensities of the stained bands. Microsomes isolated from rat livers were subjected to SDS-PAGE, proteins transferred to PVDF membranes and probed with antibody as described in Material and Methods. Values significantly different from control: **p*<0.05, ****p*<0.01 (Student *t*-test).

Fig. 4: Autoradiographic profiles of DNA adducts generated (A) by 3-NBA after activation with cytosols isolated from livers of rats treated with 2 mg/kg bw of 3-NBA, and (B) by 3-ABA after activation with microsomes isolated from livers of rats treated with 2 mg/kg bw of 3-NBA using the butanol enrichment version of the ^{32}P -postlabeling assay. Spot 1 = dA- N^6 -ABA, spot 3 = dG- N^2 -ABA, spots 4/5 = dG-C8- N -ABA.

Fig. 5: DNA adduct formation by 3-NBA activated with cytosols isolated from livers, lungs and kidneys of rats, control or treated with 0.2 or 2 mg/kg bw of 3-NBA by intratracheal instillation. Mean RAL (relative adduct labeling) $\pm$ SE of triplicate *in-vitro* incubations are shown. * $p < 0.05$, *** $p < 0.01$ (Student *t*-test).

Fig. 6: DNA adduct formation by 3-ABA activated with microsomes in the presence of NADPH isolated from livers, lungs and kidneys of rats, control or treated with 0.2 or 2 mg/kg bw of 3-NBA by intratracheal instillation. Mean RAL (relative adduct labeling) $\pm$ SE of triplicate *in-vitro* incubations are shown. *** $p < 0.001$ (Student *t*-test).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6