King's Research Portal

DOI:

[10.1016/j.envres.2016.12.020](https://doi.org/10.1016/j.envres.2016.12.020)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Dimakopoulou, K., Grivas, G., Samoli, E., Rodopoulou, S., Spyrtos, D., Papakosta, D., Karakatsani, A., Chaloulakou, A., & Katsouyanni, K. (2017). Determinants of personal exposure to ozone in school children. Results from a panel study in Greece. *Environmental Research*, 154, 66-72.
<https://doi.org/10.1016/j.envres.2016.12.020>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Title: Determinants of personal exposure to ozone in school children. Results from a panel study in Greece.

Konstantina Dimakopoulou¹, Georgios Grivas², Evangelia Samoli¹, Sophia Rodopoulou¹, Dionisis Spyratos³, Despoina Papakosta³, Anna Karakatsani⁴, Archontoula Chaloulakou², Klea Katsouyanni^{1,5}.

¹Department of Hygiene, Epidemiology and Medical Statistics, National and Kapodistrian University of Athens, Medical School, 75, Mikras Asias Street, 115 27 Athens, Greece.

²School of Chemical Engineering, National Technical University of Athens, Athens, Greece.

³Pulmonary Department, G. Papanikolaou Hospital, Medical School, Aristotle University of Thessaloniki, Thessaloniki, Greece.

⁴2nd Pulmonary Department, “ATTIKON” University Hospital, Medical School, National and Kapodistrian University of Athens, Athens, Greece.

⁵Department of Primary Care & Public Health Sciences and MRC-PHE Centre for Environment and Health, King's College London, London, UK.

Corresponding author:

Klea Katsouyanni

Professor of Biostatistics and Epidemiology

Department of Hygiene, Epidemiology and Medical Statistics

University of Athens Medical School 75, Mikras Asias street

115 27 Athens, Greece

Tel: +30-210-7462086

Fax: +30-210-7462205

e-mail: kkatsouy@med.uoa.gr

<http://school.med.uoa.gr/en/faculty/faculty-details/687-2107462086.html>

and

Professor of Public Health

Department of Primary Care & Public Health Sciences and

Environmental Research Group

King's College London

Franklin-Wilkins Building, 150 Stamford Street

London SE1 9NH

e-mail: klea.katsouyanni@kcl.ac.uk

Abstract

Background: In the wider framework of the RESPOZE (ReSPiratory effects of OZone Exposure in Greek children) panel study, we investigated possible determinants of O₃ exposure of school children, measured with personal passive samplers, in Athens and Thessaloniki, Greece.

Methods: Personal exposure to O₃ was measured for five weeks spread along the academic year 2013-14, in 186 school children in Athens and Thessaloniki, Greece. At the same time, at-school outdoor measurements were performed and ambient levels of 8-hour daily maximum O₃ from fixed sites were collected. We also collected information on lifestyle and housing characteristics through an extended general questionnaire (GQ) and each participant completed daily time activity diaries (TADs) during the study period.

Results: Mean outdoor concentrations were higher during the warmer months, in the suburbs of the cities and in Athens. Personal exposure concentrations were significantly lower compared to outdoor. Daily levels of at-school outdoor and ambient levels of O₃ from fixed sites were significant determinants of personal exposure to O₃. For a 10 µg/m³ increase in at-school outdoor O₃ concentrations and PM₁₀ measurements a 20.9% (95% CI: 13% , 28%) increase in personal exposure to O₃ was found. For a half an hour more spent in transportation an average increase of 7% (95% CI: 0.3% , 14.6%) in personal exposure to O₃ was observed. Among other possible determinants, time spent in transportation (TAD variable) and duration of open windows were the ones associated with personal O₃ exposure levels.

Conclusions: Our results support the use of outdoor and ambient measurements from fixed sites in epidemiological studies as a proxy of personal exposure to O₃, but this has to be calibrated taking into account personal measurements and time-activity patterns.

Keywords: Personal exposure, Time-activity diaries, Ozone, Panel study, Air pollution

Source of Funding: The work has been co-funded by the European Commission and the Greek government by the National Strategic Reference Framework 2007-2013
Contract Ref: RESPOZE-children/2248. The authors declare no conflict of interest.

Introduction

Ozone (O₃) is one of the main components of the photochemical air pollution mixture. It is a highly reactive gas formed by the reaction with sunlight (photochemical reaction) of primary pollutants such as nitrogen oxides (NO_x) and volatile organic compounds (VOCs) originating mainly from vehicle emissions but also from solvents and industry (WHO, 2014).

Investigation of the association between health outcomes and exposure to O₃ is often conducted by using routine measurements performed at outdoor fixed sites. Time series studies in Europe, the U.S. and Asia, have associated short term exposure to outdoor O₃ with adverse health effects (Bell et al., 2004; Choi et al., 2011; Darrow et al., 2014; Gryparis et al., 2004; Ito et al., 2005; Peel et al., 2005; Peng et al., 2013; Rodopoulou et al., 2014; Ghanbari Ghazikali et al., 2016; Miri et al., 2016), while panel studies (Liu et al., 2009; Nickmilder et al., 2007) that take into account repeated observations for the same individuals at specified periods of time have reached contradictory results. Cohort studies investigating the effects of long-term exposures (Lipfert et al., 2006; Krewski et al., 2009; Jerrett et al., 2009; Smith KR et al., 2009; Tzivian, 2011; Zanobetti & Schwartz, 2011) also provided inconsistent results.

Several air quality guidelines and standards are currently in application to protect the population from harmful air pollution effects following short term exposure. The World Health Organisation (WHO) has established a guideline value of 100µg/m³ for the daily maximum 8-hours average ozone concentrations. . This recommended limit was reduced from the previous level of 120 µg/m³, based on conclusive associations between lower O₃ concentrations and daily mortality (WHO, 2014). The most recent National Ambient Air Quality Standards (NAAQS) of the US Environmental Protection Agency (EPA) for O₃ was revised downwards to 0.070 ppm (140 µg/m³)

for 8-hour exposure (EPA, 2015). The current European Union (EU) directive for O₃ defined a health protection threshold of 110 µg/m³ for 8-hour mean concentrations (E.C. Ozone Directive, 2016).

Ozone concentrations measured at fixed sites have been used as a proxy for personal exposure (Liu et al., 2009; Nickmilder et al., 2007; Sienra-Monge et al., 2004). Although those are hypothesized to be collected and thus to reflect the average population exposure, they lack information on their quantitative association with personal exposure, the range of individual exposures and the relative contribution of factors, such as housing characteristics, behavioral aspects and time activity patterns.

Personal exposure assessment of O₃ has become feasible due to passive monitors that are small and light weighted (Koutrakis et al., 1993). Studies have been carried out using passive personal O₃ devices to monitor personal exposure of both children and adults for periods up to one week each time over the study periods (Geyh et al., 1999; Lee et al., 2004; Liu et al., 1993; Ramírez-Aguilar et al., 2008). To our knowledge, similar studies have not been conducted in the Mediterranean area, even though it is considered to be at highest O₃ risk in Europe due to precursor pollutant emissions and climate. Over the time period 2000-2010, Sicard et al. (2013) have demonstrated that O₃ concentrations are still increasing in Mediterranean cities, making ozone exposure a potential major public health concern in urban areas.

In this paper, we investigate possible determinants of personal exposure assessment of O₃, using individually carried passive samplers, during 5 weeks, in a representative sample of students aged 10-11 years in the two major cities of Greece, Athens (state capital) and Thessaloniki (second largest city and one of the main ports in the Mediterranean), performed in the framework of the panel study entitled ReSPiratory effects of OZone Exposure in Greek children (RESPOZE).

Methods and Data

Study design

In the RESPOZE study, personal exposure to O₃, using passive samplers, of 186 school children was measured in Athens and Thessaloniki, Greece. Measurements were performed during a total of five weeks in Athens and four weeks in Thessaloniki per student: two weeks in Athens and one in Thessaloniki were in the fall (October-December), one in the winter (February) and the final two weeks in the spring and early summer (April-June) for both cities (calendar weeks in the two cities were close but did not coincide in order to optimize the use of the instruments) . The field work took place during the academic year 2013-14. Information on daily habits and activities was collected from two sources and on different time scales: 1. from a general questionnaire (GQ) with questions on lifestyle and housing characteristics, and 2. through a daily time activity and symptoms diary (TAD).

All children in the study were 10-11 years old, attending the 5th grade of public elementary schools. Informed consent was signed from the parents of children willing to participate prior to the initiation of the study. The study design was approved by the Ethics Committee of the University of Athens and the Ministry of Education.

Sampling

We aimed for a sample of 100 students from Athens and 100 from Thessaloniki attending public schools. Students attending public schools are required to live in a radius of about 500m around the school. 60% of the sample was drawn from areas characterized by high O₃ concentrations (typically city suburbs) and 40% from areas

with low O₃ (the city centers where ozone concentrations are lower, due to scavenging in the presence of primary pollutants), based on the school location. We sampled schools placed near a fixed monitoring site (within 2 km) in order to collect data on regulated pollutants to enable comparisons between personal exposure and fixed site measurements. We sampled 21 schools in total in Athens and 13 in Thessaloniki and our final sample consisted of 97 children in Athens and 89 in Thessaloniki. In general, the response rate of the children varied between schools and the number sampled from each school was from 1 to 19 students. More details on the sampling procedure can be found in Samoli et al. (2016)

Personal & At-School Outdoor O₃ measurements

Personal and at-school outdoor ozone measurements, resulting in weekly O₃ average concentrations, were made using the Ogawa passive ozone sampler (Ogawa & Co. USA Inc., Pompano Beach, FL), which is based on the nitrite-coated filter method (Koutrakis et al., 1993). Field blanks were collected as part of the quality assurance and control protocol. The method's limit of detection was calculated at 1.98 µg m⁻³ as three times the standard deviation of blanks. The precision was estimated by duplicate measurements, resulting in a coefficient of variation of 4.6%. The degree of agreement between passive O₃ measurements and regulatory monitoring was assessed by collocated 7-day outdoor measurements with a reference UV photometric instrument, at a site of the air quality monitoring network in Athens. Results of the comparison indicated good correlation between passive and reference methods ($r^2=0.9$). More details on the exposure assessment campaign can be found in Grivas et al.

Samplers were given to all students in Athens and Thessaloniki. A team of three field workers visited students at their schools twice for each field work week on the same weekday. At the beginning of the study week (first visit) O₃ personal samplers were distributed and were collected on the same day of the next week (second visit). The children were instructed to wear the personal O₃ samplers continuously at chest level for the 7-days over the study period of the 5 weeks. They were instructed to clip the sampler directly to outer clothing, wear the sampler continuously, except when sleeping, bathing, or engaged in an intense athletic activity, for which wearing the sampler was not allowed. During these times the samplers were placed nearby, on a surface elevated at breathing height. Approximately 85% of total personal samples were successfully collected and analyzed. Failed measurements were mainly due to children misplacing, damaging or incorrectly exposing the sampler.

In addition to the personal samplers carried by the children, for the respective 7-day measurement periods, at-school outdoor ozone measurements were made. Outdoor samplers were located on the roofs or at a high spot in the garden of 20 (out of 34) participating schools. Samplers were placed at a distance of at least 2m from the ground and were sheltered from weather elements using a PVC protective cap. Outdoor sampling locations were selected as to not be directly impacted by traffic. After completion of sampling, filters were extracted by ultrapure grade water and analyzed for nitrate ions using ion chromatography. Derived solute nitrate concentrations were blank corrected and converted to ambient ozone concentrations using data on sampling duration and rate. Samples with concentrations below the detection limit were assigned half the detection limit value.

As a result, for each participating child and over the complete study period, a dataset of 4 to 5 observations of a 7-day average personal and at-school outdoor O₃ concentration was calculated.

Data from the fixed site monitoring network

For both study areas and for the entire duration of the study period, ambient 8-hour daily maximum O₃ and daily PM₁₀ data from the fixed site monitoring networks, operated by the Ministry of Environment & Energy (www.ypeka.gr), were collected. More specifically we obtained data from network monitors that were within 2km at the most to the sampled schools. In Athens, data were used from 6 nearby stations, of which 2 are placed in the low O₃ concentration areas and 4 placed in the high O₃ concentration areas. In Thessaloniki data were used from 3 nearby stations, one in the low O₃ concentration and 2 placed in the high O₃ concentration areas. For each child a weekly average ambient O₃ and PM₁₀ concentration was calculated using data from the nearest station to the child's school and corresponding to the exact 7-days the child participated in the study.

General Questionnaire

Trained interviewers visited the families of the children and administered an extensive questionnaire before the beginning of the field work. Information was collected on demographic and socio-economic variables e.g. age (yrs), sex, father's education (yrs); life-style and residential characteristics e.g. in house cigarette smoking (yes/no), installed air-condition (yes/no), open windows (hours/day); lifestyle characteristics

e.g. way of transport to school (on foot, car/bus, motorcycle, bike), and finally information on medical and residential history.

Time Activity Diary (TAD)

Information on daily activities that can be related to O₃ personal exposure was collected through TADs for each day during the study period. The TADs were distributed during the first school visit of every study week, when the participant was given a TAD and was instructed to record (self-completed) their daily activities in 15-min intervals across a 24-hr time period structured diary form. Responses were pre-coded into four categories: "at home", "outdoors" for locations such as parks or playgrounds, "in transportation" and "indoors other than at home" for locations such as the school, or other indoor locations visited for activities or social reasons (Figure 1). Using the TAD data we calculated for each child the time (in hours) spent daily at home, outdoors, in transportation and in other indoors spaces and consequently, we calculated the corresponding weekly average of time spent in each type of location, for each of the 5 study weeks. Additionally, each TAD included questions on occurrence of daily symptoms and school absenteeism; self-performed rheometer peak expiratory flow (PEF) measurements; and information of any incident regarding the use of the ozone personal sampler. TADs were inspected for data quality, proper completion or incorrect recordings when collected by the field workers during the second school visit at the end of each study week.

Statistical methods

The main objective of this analysis is to investigate which variables affect personal ozone exposure as measured by the individual passive samplers. Therefore in our

analysis personal ozone measurements were the dependent variable. As potential determinants of the personal ozone measurements, we considered data from the TADs variables from the GQ and ambient ozone concentrations.

We investigated the distribution of O₃ personal exposure by study period and area. We investigated the association between personal exposure to O₃ and the potential determinants using linear mixed models. In order to account for the repeated measurements for each child during the 5 weeks we applied random effects models, incorporating a random intercept. Specifically, as covariates we used activity variables derived from the TADs ("Time spent indoors, at home", "Time spent outdoors", "Time spent in transportation" and "Time spent indoors, other than at home", each as continuous weekly averages in hours); the corresponding at-school outdoor O₃ (µg/m³) measurements and the nearby fixed site O₃ concentrations. Furthermore, selected variables derived from the general questionnaire were examined as possible determinants: sex (girls vs boys), in house smoking (yes vs no), air condition use (yes vs no), heating using an open fireplace (yes vs no), duration of open windows (in hours), mode of transport to school (on foot vs by car or bus; by car or bus vs on foot), out-of-school indoor and outdoor activities in the afternoon (yes vs no), mode of transport to out-of-school indoor or outdoor activities, or other activities than mentioned before during weekends (as above).

We initially investigated the associations between personal exposure to O₃ and all predictor variables used alternatively in different models. For every mixed effect model marginal R squared (R_M^2) values were derived to assess model fit (Nakagawa & Schielzeth 2013). The model with the highest R_M^2 was regarded as the start model. To this start model all predictor variables that were statistically significant in the one-determinant models were added consecutively, using a supervised forward stepwise

procedure. Only the predictor variables that remained statistically significant were included in the final model. In all models, we adjusted for the design area variables: a) study city (Thessaloniki/Athens), b) high vs low O₃ concentrations area and time period, to account for seasonality (using 4 dummy variables for the five weeks). All tests were two-sided at a significance level of 0.05.

Results

Table 1 shows the personal, at-school outdoor and ambient from fixed sites O₃ (µg/m³) concentrations. All concentrations were significantly higher in the third period (April to June) compared to the other two study periods ($p < 0.05$); higher in the suburbs (high O₃ area) compared to the center of the city (low O₃ area) ($p = 0.001$); and higher in Athens than in Thessaloniki ($p < 0.001$). Ozone personal measurements ranged from 1 to 25.4 µg/m³ in the first period (October-December 2013), from 0.8 to 17.1 9 µg/m³ in the second study period (February 2014) and from 1 to 41.9 µg/m³ in the third period (April to June 2014), while they were significantly ($p < 0.001$) lower compared to the outdoor measurements either at school or the fixed monitoring sites. As the distribution of the O₃ personal exposures deviated from normality, it was log-transformed when used as a dependent variable in the models.

Table 2 provides a description of the study population, by city and study area according to O₃ concentrations based on the school's location. Life-style and housing characteristics and time activity patterns are presented. Children residing in the two cities were similar according to age and sex (51.5% boys in Athens vs 49.4% in Thessaloniki and of the same age). In Athens, most children living in low O₃ areas went to school on foot, while children in suburbs usually went by car or bus.

Table 3 and Figure 2 present the summary statistics for daily time activity patterns as calculated from the TADs (hours) by study period, city and exposure area. Children spent on average 17 hours/day indoors at home and more time was spent outdoors during the third period (Figure 2). Children in high O₃ areas spent on average more time in transportation than those living in low O₃ areas. In Athens suburbs, children spent on average more time indoors in places other than at home compared to those in low exposed areas.

Table 4 shows the results of linear mixed models using the log transformed personal exposure to O₃ as the dependent variable. One-independent variable models denoted that the at-school outdoor O₃ measurements were strongly associated with the personal exposure to O₃. The fixed site ambient O₃ and PM₁₀ concentrations were also a significant determinants of personal exposure. Regarding the TAD variables, the strongest association was with increased time spent in transportation (p=0.049) which was associated with increased O₃ personal exposure and at home (p=0.056) associated with decreased O₃ personal exposure. The only general questionnaire variable significantly associated with personal exposure to O₃ was increased duration of open windows.

Table 5 presents the mixed models effect estimates for potential determinants of O₃ personal exposure chosen after applying a supervised forward stepwise procedure. At-school outdoor and fixed monitoring site O₃ measurements (alternatively considered), duration of open windows, PM₁₀ measurements from fixed sites and time spent in transportation were significantly associated with personal exposure to O₃. Specifically, for a half an hour more spent in transportation an average increase of 7% (95% CI: 0.3% , 14.6%) in personal exposure to O₃ is observed. For a 10 µg/m³ increase in at-school outdoor O₃ concentrations and PM₁₀ measurements a 20.9%

(95% CI: 13% , 28%) and 12.5% (95% CI: 4% , 22%) increase in personal exposure to O₃ is found, respectively. Finally, for one more hour/day with the windows open an average increase of 2% (95% CI: 0.1% , 3.7%) in personal exposure to O₃ is observed. We did not further adjust for ambient O₃ at fixed sites since the correlation between the at-school outdoor and measured from fixed sites O₃ was 0.73. The use of O₃ concentrations at fixed sites resulted in models with lower values of the marginal R squared.

Discussion

In a panel study in Greece, we measured school children's personal and at-school outdoor exposure to O₃ for 5-weeks during the academic year 2013-14. The present results indicate that ambient O₃ measurements from fixed sites and at-school outdoor concentrations are significant determinants of personal exposure to O₃, which although, is better determined by the latter. This is consistent with some of the previous findings, in which outdoor and ambient concentrations were found to be among the most important predictors of personal exposure to O₃ (Geyh et al., 1999; Liu et al., 1995; O'Neil et al., 2003; Ramirez-Aguilar et al., 2008; Xue et al., 2005). Other studies, however, conducted in the US, suggest that associations between ambient concentrations from fixed sites and personal exposure to O₃ exist only in the summer and are weak (Sarnat et al., 2005; Sarnat et al., 2006; Brown et al., 2009). The inconsistent conclusions seen in the various studies may reflect the differences in the geographic locations, life style (e.g. the use of air conditioning and more time spent indoors), the climatic conditions and the study designs.

Spending more time in transportation (TADs) was significantly associated with higher personal O₃ exposure even after adjusting for outdoor levels. These findings are also consistent with the results of similar studies conducted in children (Lee et al., 2004;

Ramirez-Aguilar et al., 2008; Xue et al., 2005) and adults (Liu et al., 1995), in which personal exposure to O₃ was determined by time activity information (i.e. time spent outdoors and time of the day children were outdoors). The same studies also found that housing characteristics (i.e. house fan usage, presence of a gas range in the house and distance between the residence and the fixed ambient monitoring stations); and lifestyle factors (i.e. having pets) are important predictors for personal exposure to O₃. Regarding housing factors, only the duration of open windows was significantly associated to personal exposure to O₃ in the present study.

Our study is based on a representative sample of school children that live in two major cities of Greece the population of which comprise about half of the Greek population. In addition, these cities provide an excellent opportunity to study O₃ effects as concentrations are high due to precursor pollutant emissions and climate conditions. The present study was conducted for a relatively long period of follow up and had a large sample of students. It obtained repeated measures of personal, at-school outdoor and ambient O₃ concentrations and time-activity information. This allowed us to fit alternate mixed models for investigating a variety of potential determinants of personal exposure to O₃. One limitation of our study is the difficulty of collecting data through the self-completion of TADs. However, exhausting data quality control procedures were followed and controls with manual inspections of the data and cross references from the field workers' progress diaries, that limited inaccurately reported data.

Based on our results a prediction model for estimating personal exposure to O₃ in children living in a Mediterranean climate can be implemented. Using our models personal exposure estimation could be applied in epidemiological studies in which the use of personal O₃ monitoring devices is not feasible. Furthermore, the validity of our

model in different populations could be easily tested by providing personal monitors to a subsample of participants before generalizing to the whole sample under investigation.

In conclusion, we found geographical and temporal differences in personal O₃ exposures. Ozone personal measurements were significantly lower compared to the outdoor measurements. At-school outdoor level of O₃ was the major predictor of personal exposure to O₃. PM₁₀ measurements from fixed sites, duration of open windows and time spent in transportation were also significantly associated with personal exposure to O₃. Our results support the use of outdoor and ambient measurements from fixed sites in epidemiological studies as a proxy of personal exposure to O₃, taking into account time-activity patterns and possibly calibration factors.

References

Bell M.L., McDermott A., Zeger S.L., Samet J.M., Dominici F. (2004). Ozone and short-term mortality in 95 US urban communities, 1987-2000. *JAMA*, 292(19):2372–8.

Brown K.W., Sarnat J.A., Suh H.H., Coull B.A., Koutrakis P. (2009). Factors influencing relationships between personal and ambient concentrations of gaseous and particulate pollutants. *Sci Total Environ.* Jun 1;407(12):3754-65.

Choi M., Curriero F.C., Johantgen M., Mills M.E., Sattler B., Lipscomb J. (2011). Association between ozone and emergency department visits: an ecological study. *Int J Environ Health Res.*, 21(3):201–21.

Darrow L.A., Klein M., Flanders W.D., Mulholland J.A., Tolbert P.E., Strickland M.J. (2014). Air pollution and acute respiratory infections among children 0-4 years of age: an 18-year time-series study. *Am J Epidemiol.*, Nov 15;180(10):968-77.

Geyh A.S., Roberts P.T., Lurmann F.W., Schoell B.M., Avol E.L. (1999) Initial field evaluation of the Harvard active ozone sampler for personal ozone monitoring. *J Expo Anal Environ Epidemiol*, 9:143-149.

Ghanbari Ghazikali M., Heibati B., Naddafi K., Kloog I., Oliveri Conti G., Polosa R., Ferrante M. (2016). Evaluation of Chronic Obstructive Pulmonary Disease (COPD)

attributed to atmospheric O₃, NO₂, and SO₂ using Air Q Model (2011-2012 year).
Environ Res, Jan;144(Pt A):99-105.

Grivas G., Dimakopoulou K., Samoli E., Papakosta D., Karakatsani A., Katsouyanni K., Chaloulakou A. Ozone exposure assessment for children in Greece: Results from the RESPOZE study. (Under Review)

Gryparis A., Forsberg B., Katsouyanni K, Analitis A, Touloumi G, Schwartz J, Samoli E, Medina S, Anderson HR, Niciu EM, Wichmann HE, Kriz B, Kosnik M, Skorkovsky J, Vonk JM, Dörtbudak Z. (2004). Acute effects of ozone on mortality from the 'air pollution and health: a European approach' project. American Journal of Respiratory and Critical Care Medicine, 170(10):1080–1087.

Ito K., De Leon S.F., Lippmann M. (2005). Associations between ozone and daily mortality: analysis and meta-analysis. Epidemiology, 16(4):446–457.

Jerrett M., Burnett R.T., Pope C.A. 3rd., Ito K., Thurston G., Krewski D., Shi Y., Calle E., Thun M. (2009). Long-term ozone exposure and mortality. New England Journal of Medicine, 360(11):1085–1095.

Koutrakis P., Wolfson J.M., Bunyaviroch A., Froehlich S.E., Hirano K., Mulik J.D. (1993). Measurement of ambient ozone using a nitrite-coated filter. Anal. Chem. 65, 209–214.

Krewski D., Jerrett M., Burnett R.T., Ma R., Hughes E., Shi Y., Turner M.C., Pope C.A. 3rd., Thurston G., Calle E., Thun M.J. (2009). Extended follow-up and spatial analysis of the American Cancer Society study linking particulate air pollution and mortality. Boston, Health Effects Institute (Research Report 140; <http://pubs.healtheffects.org/getfile.php?u=478>, accessed 17 March 2013):5–114; commentary: 115–136.

Lee K., Parkhurst W.J., Xue J., Ozkaynak H., Neuberg D., Spengler J.D. (2004). Outdoor/Indoor/Personal Ozone Exposures of Children in Nashville, Tennessee. *J. Air Waste & Manage. Assoc.*, 54:352-359.

Lipfert F.W., Wyzga R.E., Baty J.D., Miller J.P. (2006). Traffic density as a surrogate measure of environmental exposures in studies of air pollution health effects: long-term mortality in a cohort of US veterans. *Atmospheric Environment*, 40(1):154–169.

Liu L.J., Koutrakis P., Leech J., Broder I. (1995). Assessment of ozone exposures in the greater metropolitan Toronto area. *J Air Waste Manag Assoc*, 45(4):223-34.

Liu L.J., Koutrakis P., Suh H.H., Mulik J.D., Burton R.M. (1993) Use of personal measurements for ozone exposure assessment: a pilot study. *Environ Health Perspect*, 101:318-324.

Liu L., Poon R., Chen L., Frescura A.M., Montuschi P., Ciabattini G., Wheeler A., Dales R. (2009). Acute effects of air pollution on pulmonary function, airway

inflammation, and oxidative stress in asthmatic children. *Environmental Health Perspectives*, 117(4):668–674.

Miri M., Derakhshan Z., Allahabadi A., Ahmadi E., Oliveri Conti G., Ferrante M., Aval H.E. (2016) Mortality and morbidity due to exposure to outdoor air pollution in Mashhad metropolis, Iran. The AirQ model approach. *Environ Res*, Nov;151:451-457.

Nakagawa S., Schielzeth H. (2013) A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods in Ecology and Evolution* (4)2:133–142.

Nickmilder M., de Burbure C., Sylviane C., Xavier D., Alfred B., Alain D. (2007). Increase of exhaled nitric oxide in children exposed to low levels of ambient ozone. *Journal of Toxicology and Environmental Health, Part A*, 70(3–4):270–274.

O'Neill M.S., Ramirez-Aguilar M., Meneses-Gonzalez F., Hernández-Avila M., Geyh A.S., Sienra-Monge J.J., Romieu I. (2003). Ozone exposure among Mexico City outdoor workers. *J Air Waste Manag Assoc* 53:339-346.

Peel J.L., Tolbert P.E., Klein M., Metzger K.B., Flanders W.D., Todd K., Mulholland J.A., Ryan P.B., Frumkin H. (2005) Ambient air pollution and respiratory emergency department visits *Epidemiology*, 16, pp. 164–174.

Peng R.D., Samoli E., Pham L., Dominici F., Touloumi G., Ramsay T., Burnett R.T., Krewski D., Le Tertre A., Cohen A., Atkinson R.W., Anderson H.R., Katsouyanni K., Samet J.M. (2013). Acute effects of ambient ozone on mortality in Europe and North America: results from the APHENA study. *Air Qual Atmos Health*, Jun 1;6(2):445-453.

Ramírez-Aguilar M., Barraza-Villarreal A., Moreno-Macías H., Winer A.M., Cicero-Fernández P., Vélez-Márquez M.G.D., Cortez-Lugo M., Sienra-Monge J.J., Romieu I. (2008). Assessment of personal exposure to ozone in asthmatic children residing in Mexico City. *Salud Publica Mex*, 50:67-75.

Rodopoulou S., Chalbot M.C., Samoli E., Dubois D.W., San Filippo B.D., Kavouras IG. (2014) Air pollution and hospital emergency room and admissions for cardiovascular and respiratory diseases in Doña Ana County, New Mexico. *Environ Res.*, Feb;129:39-46.

Samoli E., Dimakopoulou K., Evangelopoulos D., Rodopoulou S., Karakatsani A., Veneti L., Sionidou M., Tsolakoglou I., Krasanaki I., Grivas G., Papakosta D., Katsouyanni K. (2016) Is daily exposure to ozone associated with respiratory morbidity and lung function in a representative sample of schoolchildren? Results from a panel study in Greece. *J Expo Sci Environ Epidemiol*. May 18.

Sarnat J.A., Brown K.W., Schwartz J., Coull B.A., Koutrakis P. (2005). Ambient gas concentrations and personal particulate matter exposures: implications for studying the health effects of particles. *Epidemiology*. May;16(3):385-95.

Sarnat S., Coull B., Schwartz J., Gold D, Suh H. (2006). Factors affecting the association between ambient concentrations and personal exposures to particles and gases. *Environ Health Perspect*, 114;649–654.

Sicard P., De Marco A., Troussier F., Renou C., Vas N., Paoletti E. (2013). Decrease in surface ozone concentrations at Mediterranean remote sites and increase in the cities. *Atmos Environ*. 79;705-715.

Sienra-Monge J.J., Ramírez-Aguilar M., Moreno-Macias H., Reyes-Ruiz N.I., Del Rio-Navarro B.E. (2004). Antioxidant supplementation and nasal inflammatory responses among young asthmatics exposed to high levels of ozone. *Clinical and Experimental Immunology*, 138(2):317–322.

Smith K.R., Jerrett M., Anderson H.R., Burnett R.T., Stone V., Derwent R., Atkinson R.W., Cohen A., Shonkoff S.B., Krewski D., Pope C.A. 3rd, Thun M.J., Thurston G. (2009). Public health benefits of strategies to reduce greenhouse-gas emissions: health implications of short-lived greenhouse pollutants. *Lancet*, 374(9707):2091–2103

The current Ozone Directive and other relevant legislation. (2016). Retrieved from <http://www.eea.europa.eu/publications/TOP08-98/page006.html>

Tzivian L. (2011). Outdoor air pollution and asthma in children. *Journal of Asthma*, 48(5):470–481.

US Environmental Protection Agency (EPA), 2015. 40 CFR Part 50, 51, 52, et al.
National Ambient Air Quality Standards for Ozone; Final Rule.

WHO, Regional Office for Europe, 2014. Air quality in Europe — 2014 report,
Luxembourg.

Xue J., Liu S.V., Ozkaynak H., Spengler J.D. (2005). Parameter evaluation and model
validation of ozone exposure assessment using Harvard Southern California Chronic
Ozone Exposure Study data. J Air Waste Manag Assoc. Oct;55(10):1508-15.

Zanobetti A., Schwartz J. (2011). Ozone and survival in four cohorts with potentially
predisposing diseases. American Journal of Respiratory and Critical Care Medicine,
184(7):836–841.

Table 1. Descriptive statistics of personal exposure to ozone (O₃), at-school outdoor and fixed site measurements (µg/m³), by study period, city and O₃ concentrations area.

	Athens		Thessaloniki	
Study period	Low O₃ area	High O₃ area	Low O₃ area	High O₃ area
October to December 2013				
Personal weekly O ₃ concentration (µg/m ³)				
Samples (n)	60	109	28	49
Mean (SD)	6.1 (3.51)	8.2 (5.33)	2.0 (1.37)	2.1 (1.03)
Median (25th-75th)	5.7 (4.3-7.5)	7.3 (4.3-11.0)	1 (1-2.6)	1.8 (1.2-2.8)
Min-Max	1.2 - 19	1.2 - 25.4	1 - 6.2	1 - 4.9
At-school outdoor weekly O ₃ concentration (µg/m ³)				
Mean (SD)	35.7 (5.25)	52.6 (12.62)	11.4 (2.98)	24.1 (10.49)
Median (25th-75th)	33.7 (32.4-42.2)	57.3 (42.3-60.1)	10.3 (10.3-13.4)	23.6 (16.4-24.3)
Min-Max	28.4 - 42.7	32.6 - 96.3	8.5 - 23.9	15.6 - 80.0
Fixed sites weekly average of O ₃ concentration (µg/m ³)				
Mean (SD)	32.8 (11.45)	66.7 (15.13)	28.4 (10.98)	51.8 (19.37)
Median (25th-75th)	33.9 (21.0-42.7)	65.4 (53.6-70.9)	26.7 (18.0-34.2)	49.7 (40.9-77.6)
Min-Max	17.2 - 48.9	50.6 - 105.6	17.1 - 44.9	24.2 - 100.6
February 2014				
Personal weekly O ₃ concentration (µg/m ³)				
Samples (n)	28	53	25	42
Mean (SD)	2.7 (2.24)	4.8 (3.89)	2.7 (1.72)	3.6 (1.67)
Median (25th-75th)	2.2 (0.8-3.6)	3.7 (2.6-5.2)	2.2 (1.7-3.1)	3.7 (2.6-4.4)
Min-Max	0.8 - 10.6	0.8 - 17.1	1.0 - 8.0	1.0 - 7.0
At-school outdoor weekly O ₃ concentration (µg/m ³)				
Mean (SD)	33.7 (1.58)	48.1 (12.22)	22.1 (5.88)	27.1 (4.44)
Median (25th-75th)	33.6 (32.3-34.0)	48.9 (33.4-61.9)	20.5 (17.8-23.9)	25.1 (25.1-29.0)
Min-Max	28.4 - 38.7	32.8 - 63.1	17.8 - 51.2	24.5 - 56.0
Fixed sites weekly average of O ₃ concentration (µg/m ³)				
Mean (SD)	39.7 (2.24)	75.5 (5.92)	37.9 (0.91)	50.4 (11.11)
Median (25th-75th)	39.3 (28.1-50.1)	75.0 (72.6-77.1)	38.0 (36.9-39.0)	55.6 (37.0 - 57.6)
Min-Max	17.2 - 50.3	51.4 - 85.3	36.9 - 39.0	37.0 - 69.4
April to June 2014				
Personal weekly O ₃ concentration (µg/m ³)				
Samples (n)	55	90	38	73
Mean (SD)	13.2 (7.56)	17.6 (7.10)	7.7 (5.15)	9.8 (8.34)
Median (25th-75th)	13.2 (8.3-18.7)	17.3 (12.6-21.6)	6.2 (3.4 - 11.7)	6.9 (4.5 - 10.7)
Min-Max	1.2 - 33.8	3.1 - 41.8	1 - 21.7	1.2 - 41.9
At-school outdoor weekly O ₃ concentration (µg/m ³)				

Mean (SD)	62.3 (7.93)	84.5 (10.37)	53.2 (9.60)	60.4 (9.30)
Median (25th-75th)	65.4 (60.5-66.9)	78.2 (76.4-90.4)	49.4 (44.6-62.8)	61.6 (56-61.8)
Min-Max	44.6 - 69.2	76 - 107	41.3 - 70	45.8 - 80
Fixed sites weekly average of O ₃ concentration (µg/m ³)				
Mean (SD)	35.6 (26.40)	100.0 (5.91)	72.1 (3.04)	81.3 (12.16)
Median (25th-75th)	19.7 (10.4-67.3)	101.7(99.1-104.1)	72.7 (69.3-75.6)	83.1 (69.4-92.4)
Min-Max	7.1 - 73.6	88.9 - 106.9	69.1 - 76.3	66.1 - 100.6

Table 2. Characteristics of children participating in the RESPOZE panel study during the academic year 2013-14, by city and O₃ concentrations area.

	Athens		Thessaloniki	
	Low O ₃ area (n=37)	High O ₃ area (n=60)	Low O ₃ area (n=32)	High O ₃ area (n=57)
Boys (n, %)	22 (59.5)	28 (46.7)	14 (42.4)	30 (51.7)
Age (yrs; mean, (SD))	10.3 (0.3)	10.3 (0.3)	10.4 (0.4)	10.4 (0.3)
Father's Education (yrs; mean (SD))	14.0 (2.8)	15.2 (3.7)	15.3 (3.4)	14.1 (3.3)
In house cigarette smoking (yes; n, %)	14 (38.9)	14 (23.3)	7 (21.2)	13 (22.4)
Air condition (yes; n, %)	28 (75.7)	45 (75.0)	25 (75.8)	48 (82.8)
Open windows (hours/day; mean, (SD))	10.9 (3.37)	11.8 (3.64)	12.2 (3.42)	11.2 (4.13)
Transportation to school				
• On foot (n, %)	30 (81.1)	26 (43.3)	23 (69.7)	41 (71.9)
• By car/bus (n, %)	6 (16.2)	44 (73.3)	8 (24.2)	32 (55.2)

Table 3. Mean (standard deviation) of activity variables (time spent daily; hours) derived from the time activity diaries (TADs), by study period, city and O₃ concentrations area.

	Athens		Thessaloniki	
	Low O ₃ area (n=37)	High O ₃ area (n=60)	Low O ₃ area (n=32)	High O ₃ area (n=57)
Time spent indoors, at home (hours)				
October to December 2013	18 (2.5)	17 (1.8)	17 (1.9)	17 (1.6)
February 2014	17 (2.8)	17 (1.8)	17 (1.4)	17 (1.6)
April to June 2014	17 (2.2)	17 (1.9)	17 (1.4)	17 (2.1)
Time spent outdoors (hours)				
October to December 2013	0.9 (0.9)	0.9 (0.7)	1.4 (1.1)	1.3 (0.8)
February 2014	0.7 (0.9)	0.8 (0.8)	2.0 (0.9)	1.6 (0.9)
April to June 2014	1.1 (1.1)	1.4 (1.0)	1.8 (1.2)	1.6 (1.0)
Time spent in transportation (hours)				
October to December 2013	0.8 (0.6)	0.9 (0.3)	0.5 (0.5)	0.6 (0.3)
February 2014	0.7 (0.4)	0.9 (0.4)	0.7 (0.3)	0.8 (0.4)
April to June 2014	0.7 (0.3)	0.9 (0.4)	0.4 (0.3)	0.6 (0.4)
Time spent indoors, other than at home (hours)				
October to December 2013	4.3 (1.5)	5.3 (1.7)	5.4 (1.5)	5.0 (1.1)
February 2014	4.8 (1.3)	5.0 (1.4)	4.5 (1.1)	4.6 (1.2)
April to June 2014	4.6 (1.6)	4.9 (1.6)	4.9 (1.6)	4.7 (1.2)

Table 4. Mixed models estimates (coefficients (b) per fixed increment and 95% confidence intervals (CI)) for determinants of log transformed personal exposure to O₃ (µg/m³) and percent change and 95% confidence intervals (CI) in personal exposure to O₃. Results from linear mixed models for determinants considered alternatively, adjusting for design area variables (study area: Thessaloniki vs Athens; O₃ concentration area: high vs low) and study week (4 dummy variables).

Independent variable	One-independent variable models			
TAD* variables (time spent; hours)	R_M²	b (95% C.I.)	% change (95% C.I.)	p-value
Indoors, at home (per hour/day)	0.522	-0.023 (-0.047 , 0.001)	-2.3 (-4.6 , 0.1)	0.056
Outdoors (per hour/day)	0.520	0.042 (-0.013 , 0.097)	4.3 (-1.3 , 10.2)	0.137
In transportation (per half hour/day)	0.522	0.069 (0.000 , 0.137)	7.1 (0.0 , 14.7)	0.049
Indoors, other than at home (per hour/day)	0.519	0.011 (-0.023 , 0.046)	1.1 (-2.3 , 4.7)	0.516
Fixed outdoor pollutant measurements (µg/m ³)				
At-school outdoor (per 10 µg/m ³)	0.540	0.177 (0.115 , 0.239)	19.3 (12.2 , 27.0)	<0.001
Fixed monitoring site (per 10 µg/m ³)	0.525	0.044 (0.010 , 0.079)	4.5 (1.0 , 8.2)	0.012
PM ₁₀ measurements from fixed sites (per 10 µg/m ³) (Athens & Thessaloniki)	0.523	0.083 (0.004 , 0.163)	8.7 (0.4 , 17.7)	0.039
Other potential determinants from the GQ*				
Gender (girls vs boys)	0.519	0.036 (-0.096 , 0.168)	3.7 (-9.1 , 18.3)	0.592
In house cigarette smoking (yes vs no)	0.524	-0.069 (-0.220 , 0.081)	-6.7 (-19.8 , 8.5)	0.366
Air condition (yes vs no)	0.522	-0.128 (-0.282 , 0.026)	-12.0 (-24.6 , 2.6)	0.103
Open windows (per hour/day)	0.525	0.020 (0.002 , 0.037)	2.0 (0.2 , 3.8)	0.030
Transportation to school (on foot vs all others)	0.522	0.113 (-0.024 , 0.250)	11.9 (-2.4 , 28.4)	0.107
Transportation to school (by car/bus vs all others)	0.519	-0.117 (-0.260 , 0.026)	-11.0 (-22.9 , 2.6)	0.109

R_M²: marginal R squared

*TAD: Time Activity Diary; GQ: General Questionnaire

Table 5. Mixed models estimates (coefficients (b) per fixed increment and 95% confidence intervals (CI)) for determinants of log transformed personal exposure to O₃ (µg/m³) and percent change and 95% confidence intervals (CI) in personal exposure to O₃. Results from linear mixed models for determinants mutually adjusted a. considering the at-school outdoor weekly O₃ concentrations and b. the fixed monitoring sites weekly average of O₃ concentrations and adjusted for design area variables (study area: Thessaloniki vs Athens; O₃ concentration area: high vs low) and study week (4 dummy variables).

	Independent variables	R_M^2	b (95% C.I.)	% change (95% C.I.)	p-value
Model 1					
a.	At-school outdoor O ₃ (10µg/m ³) +	0.546	0.175 (0.113,0.236)	19.1 (12.0 , 26.7)	<0.001
	Open windows (1 hour/day)		0.018 (0.001 , 0.036)	1.9 (0.1 , 3.7)	0.043
b.	Fixed monitoring site O ₃ (10µg/m ³) +	0.531	0.044 (0.009 , 0.078)	4.5 (0.9 , 8.1)	0.014
	Open windows (1hour/day)		0.019 (0.001 , 0.037)	1.9 (0.1 , 3.7)	0.033
Model 2					
a.	At-school outdoor O ₃ (10µg/m ³) +	0.551	0.187 (0.126 , 0.248)	20.6 (13.4 , 28.2)	<0.001
	Open windows (1 hour/day) +		0.019 (0.002 , 0.037)	2.0 (0.2 , 3.8)	0.028
	PM ₁₀ measurements (10µg/m ³)		0.120 (0.042 , 0.198)	12.8 (4.3 , 21.9)	0.002
b.	Fixed monitoring site O ₃ (10µg/m ³) +	0.533	0.040 (0.005 , 0.074)	4.0 (0.5 , 7.7)	0.026
	Open windows (1hour/day) +		0.020 (0.002 , 0.037)	2.0 (0.2 , 3.8)	0.026
	PM ₁₀ measurements (10µg/m ³)		0.077 (-0.002 , 0.157)	8.0 (-0.2 , 17.0)	0.056
Model 3					
a.	At-school outdoor O ₃ (10µg/m ³) +	0.552	0.186 (0.124 , 0.247)	20.9 (13.2 , 28.0)	<0.001
	Open windows (1hour/day) +		0.019 (0.001 , 0.036)	1.9 (0.1 , 3.7)	0.035
	PM ₁₀ measurements (10µg/m ³) +		0.118 (0.038 , 0.198)	12.5 (3.9 , 21.9)	0.004
	In transportation (time spent; half hour/day)		0.069 (0.003 , 0.136)	7.2 (0.3 , 14.6)	0.041
b.	Fixed monitoring site O ₃ (10µg/m ³)+	0.535	0.042 (0.007 , 0.078)	4.3 (0.7 , 8.1)	0.019
	Open windows (1hour/day)+		0.019 (0.002 , 0.037)	1.9 (0.2 , 3.7)	0.032
	PM ₁₀ measurements (10µg/m ³)+		0.072 (-0.010 , 0.153)	7.4 (-1.0 , 16.6)	0.085
	In transportation (time spent; half hour/day)		0.147 (0.012 , 0.282)	7.6 (0.6 , 15.2)	0.033

R_M^2 : marginal R squared