King's Research Portal

DOI:

[10.1001/jamapsychiatry.2017.0185](https://doi.org/10.1001/jamapsychiatry.2017.0185)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Fusar-Poli, P. (2017). Negative psychosis prevention trials. *JAMA Psychiatry*, 74(6), 651.
<https://doi.org/10.1001/jamapsychiatry.2017.0185>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

NEGATIVE TRIALS IN PSYCHOSIS PREVENTION: THE NEGLECTED IMPORTANCE OF PRETEST RISK ENRICHMENT

Dear Editor

In the wake of the recently published ω -3 polyunsaturated fatty acids multisite trial (NEURAPRO) in the journal (McGorry et al, n=304¹), the three largest studies of preventative interventions in individuals at ultra high risk (UHR) for psychosis have turned out to be negative (Morrison et al, n=288², McFarlane et al, n=292³), suggesting that it may not currently be possible to prevent psychosis. However, the actual risk of psychosis in these interventional studies was found to be extremely low, even in the control group (11.2% in McGorry et al at 1y¹, 9% at 2y in Morrison et al and 2.3% at 2y in McFarlane et al ³, implying that these negative findings are likely to be secondary to small statistical power.

On a conceptual level, these studies challenge the underlying assumption that it is acceptable to meet specific psychometric criteria at intake to be slotted into the UHR category. Actually, the predictive power of the UHR criteria is not fixed priority, but is strongly dependent on the risk enrichment of the samples to which they are applied (i.e. “pretest” risk)⁴. This is due to the psychometric characteristics of these UHR instruments: they are good to rule out psychosis, but only moderately useful to rule in psychosis⁴. Therefore, majority of the observed risk to psychosis is not gained through UHR assessment per se, but is obtained before the assessment (pretest phase) during the recruitment of these individuals. Pretest risk enrichment in samples undergoing an UHR assessment remains substantial (15% at 38 months) and highly heterogeneous (9%-24% at 38 months)⁴. Pretest risk enrichment is modulated by the type of recruitment strategies adopted to select individuals for UHR assessment⁴. For example, recruiting individuals who were already filtered by adult mental health services is associated with higher pretest risk enrichment, as compared to recruiting individuals through intensive community outreach⁵. Not surprisingly, the lowest transition risk was reported in McFarlane et al, that has adopted a strong “community outreach and education program aimed at teachers, school, and college counsellors”³. Due to the heightened pressure to recruit participants into interventional studies, it is possible that a substantial proportion of these samples had been recruited through community outreach, diluting the transition risks.

It therefore seems expedient to control the pretest risk enrichment of samples recruited into future interventional studies for psychosis prevention by using pretest risk stratification models that have been recently validated in this journal⁵.

References:

1. McGorry P, Nelson B, Markulev C, et al. Effect of ω -3 Polyunsaturated Fatty Acids in Young People at Ultrahigh Risk for Psychotic Disorders. *JAMA Psychiatry*. 2016;Nov 23, 2016.
2. Morrison AP, French P, Stewart SL, et al. Early detection and intervention evaluation for people at risk of psychosis: multisite randomised controlled trial. *Bmj*. 2012;344:e2233.
3. McFarlane WR, Levin B, Travis L, et al. Clinical and functional outcomes after 2 years in the early detection and intervention for the prevention of psychosis multisite effectiveness trial. *Schizophrenia bulletin*. 2015;41(1):30-43.
4. Fusar-Poli P, Schultze-Lutter F, Cappucciati M, et al. The Dark Side of the Moon: Meta-analytical Impact of Recruitment Strategies on Risk Enrichment in the Clinical High Risk State for Psychosis. *Schizophrenia bulletin*. 2016;42(3):732-743.
5. Fusar-Poli P, Rutigliano G, Stahl D, et al. Deconstructing pretest risk enrichment to optimize prediction of psychosis in individuals at clinical high risk. *JAMA Psychiatry*. 2016.