King's Research Portal

DOI:

[10.1016/j.conb.2016.12.009](https://doi.org/10.1016/j.conb.2016.12.009)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Dehorter, N., Marichal, N., Marín, O., & Berninger, B. (2017). Tuning neural circuits by turning the interneuron knob. *CURRENT OPINION IN NEUROBIOLOGY*, 42, 144-151. <https://doi.org/10.1016/j.conb.2016.12.009>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Tuning neural circuits by turning the interneuron knob

Nathalie Dehorter¹, Nicolás Marichal³, Oscar Marín^{1,2*} and Benedikt Berninger^{3,*}

Address

¹Centre for Developmental Neurobiology, Institute of Psychiatry, Psychology and Neuroscience, King's College London, London SE1 1UL, United Kingdom

²MRC Centre for Neurodevelopmental Disorders, King's College London, London SE1 1UL, United Kingdom

³Institute of Physiological Chemistry , University, Medical Center Johannes Gutenberg, University Mainz, Hanns-Dieter-Hüscher-Weg 19, 55128 Mainz, Germany

*Corresponding author: Oscar Marín (oscar.marin@kcl.ac.uk) or Benedikt Berninger (berningb@uni-mainz.de)

Summary

Interneurons play a critical role in sculpting neuronal circuit activity and their dysfunction can result in neurological and neuropsychiatric disorders. To temporally structure and balance neuronal activity in the adult brain interneurons display a remarkable degree of subclass-specific plasticity, of which the underlying molecular mechanisms have recently begun to be elucidated. Grafting new interneurons to pre-existing neuronal networks allows for amelioration of circuit dysfunction in rodent models of neurological disease and can reopen critical windows for circuit plasticity. The crucial contribution of specific classes of interneurons to circuit homeostasis and plasticity in health and disease underscores their generation as an important objective for emerging strategies of lineage reprogramming in vivo.

Introduction

Cortical GABAergic interneurons exert a crucial influence on the computations performed by microcircuits of the cerebral cortex by gating signal flow and sculpting activity patterns [1]. Dysfunction of inhibitory circuits can lead to neurological and neuropsychiatric disease, very typically due to abnormalities in the development and plasticity of cortical interneurons [2]. Interneurons come in many flavors and have traditionally been classified on the basis of their molecular, anatomical and physiological properties [3]. While early genetic specification seems to determine the cardinal properties of these interneurons [4], it is increasingly recognized that their final specification is subject to regulation by neuronal activity [5,6]. In this review, we discuss how interneurons contribute to neuronal plasticity of adult cortical circuits in a subclass-specific fashion and highlight new insights into the molecular mechanisms that underlie these physiological adaptations. We also review how network dysfunction can be ameliorated and plasticity reactivated by grafting specific classes of interneurons. Finally, we consider new opportunities for rebuilding inhibitory circuits by the emerging strategy of direct lineage reprogramming *in vivo*.

Modulation of neuronal plasticity by fast-spiking interneurons

The balance between synaptic excitation and inhibition is critical for cortical function, and its disruption has been associated with several developmental neuropsychiatric conditions such as autism or schizophrenia [7,8]. For instance, temporally locked excitation and inhibition is fundamental for the generation of oscillatory rhythms underlying high-order functions, and defects in these rhythms are strongly linked to neurological and psychiatric disorders [9]. Although pyramidal cells may have very different levels of activity depending on the behavioral state and their specific

engagement into particular neural assemblies, individual neurons show relatively stable ratios of excitation and inhibition due to the prominent capability of interneurons to compensate for changes in the firing of pyramidal cells [10-12].

Inhibitory neurons regulate the adaptation of neural circuits to experience by controlling the connectivity between neurons. In particular, somatostatin-expressing (SST) interneurons mediate different forms of plasticity and memory formation by providing state-dependent inhibition at multiple levels and timescales in the cortex. These interneurons are involved in feedback and feedforward inhibitory and disinhibitory circuits, thereby integrating different streams of sensory information [13,14]. Inhibitory plasticity seems to be mediated by parvalbumin-expressing (PV+), fast-spiking basket cells, which primarily target the soma of pyramidal cells. These cells are able to adjust their synapses onto pyramidal cells in response to changes in their activity ([Fig. 1a](#)). For instance, experimental suppression of the firing of individual pyramidal cells reduces their inhibition but not their excitatory drive [11], whereas enhancing the discharge of pyramidal cells potentiates perisomatic inhibition [11,15]. These synaptic changes can be induced through different mechanisms. For example, sensory deprivation experiments have revealed that weakening of feedforward inhibition in layer (L) 4 $\rightarrow$ L2/3 connections contributes to compensate for the reduction in sensory drive, thereby maintaining excitatory-inhibitory balance in L2/3 pyramidal neurons [16]. Conversely, the responses induced by the direct modification of L2/3 pyramidal cells (either increasing or decreasing their firing) suggest that feedback circuits also contribute to this form of inhibitory plasticity [11]. In addition to synaptic mechanisms, other studies have shown that fast-spiking interneurons contribute to maintain the excitatory-inhibitory balance by dynamically adjusting their intrinsic neuronal excitability in an activity-dependent manner [17,18].

The dynamic reconfiguration of PV+ fast-spiking basket cells in response to dynamic changes in network conditions has attracted much attention recently. Work from the Caroni laboratory suggests that PV+ fast-spiking basket cells may contribute to neural plasticity by dynamically oscillating between two main states in response to recent experience [19]. These cell states are characterized by specific neurochemical properties and synaptic-density ratios, and are differentially modulated by experience (Fig. 1b). For example, environmental enrichment increased the fraction of basket cells with low levels of PV and GAD67 and relatively low excitatory inputs, whereas fear conditioning increases the fraction of basket cells with high PV and GAD67 levels and that receive a relatively high number of excitatory synapses [19]. The switch from low-PV to high-PV state also involves changes in the number inhibitory synapses received by basket cells [19], which suggest that plasticity mechanisms involve adjustments in very different types of synapses. Intriguingly, recent studies have revealed that low-PV and high-PV basket cells are born at different times during neurogenesis [20], and therefore are likely engaged in different neuronal assemblies. Since the laminar location of interneurons derived from the medial ganglionic eminence (MGE) is directly linked to their neurogenesis [21], low-PV and high-PV basket cells should be largely segregated in deep and superficial layers of the neocortex [22]. This later finding makes difficult to reconcile the idea of state oscillations, as originally described in the hippocampus [19], with the organization of interneurons in the neocortex. One possibility is that early- and late-born PV+ interneurons are fated to exhibit different forms of plasticity. For example, low-PV basket cells are particularly abundant in L2/3, where they would contribute to the acquisition of new information through adult circuit plasticity [18,20].

Molecular mechanisms controlling inhibitory neuronal plasticity

It is now well established that activity does not simply promote the maturation of interneuron populations that are already pre-specified, but also directs differentiation in the adult. The effects of neuronal activity are mediated through the regulation of transcription factors that set in motion programs of gene expression controlling specific features of these cells [23]. Recent studies have focused on transcription factors that are targets of specific Ca^{2+} -dependent signaling pathways in various classes of interneurons. For instance, it has been shown that PV+ fast-spiking basket cells rely specifically on the Ca^{2+} /calmodulin-dependent kinase 1 (CaMKI) pathway to drive gene expression in response to the opening of Ca_v1 channels by excitatory inputs from pyramidal cells [24]. CaMKI promotes the expression of CREB target genes in PV+ fast-spiking basket cells, including *Pvalb* (PV) and *Gad1* (GAD67) [24]. One of the genes that might be activated downstream of CaMKI is *Er81* (*Etv1*), an activity-dependent ETS transcription factor that interacts with the transcriptional coactivator CREB-binding protein [25,26]. *Er81* regulates the intrinsic properties of PV+ fast-spiking basket cells, including the expression of Kv1.1 channel subunits, and neural activity in turn regulates the expression (and probably the nuclear translocation) of *Er81* [18]. This mechanism allows PV+ fast-spiking basket cells to dynamically adjust their excitability in response to changes in network activity. It remains to be determined how this mechanism of neural plasticity influences the function of PV+ fast-spiking basket cells [27-29], as it is clear now that they comprise a heterogeneous population of cells.

The mechanisms regulating activity-dependent changes in gene expression have also begun to unveil for other types of interneurons. For instance, the transcription factor *Npas4* is an early-response transcription factor that is induced in response to

activity in both somatostatin-expressing (SST+) interneurons and pyramidal cells. Interestingly, *Npas4* drives different gene expression programs in both cell types, reinforcing the view that activity-dependent transcriptional responses are regulated in a cell-class specific manner [30]. Another major class of interneurons is characterized by the expression of vasoactive intestinal peptide (VIP). It has been recently shown that sensory experience promotes the expression of *Igf1* specifically in VIP+ interneurons, which in turn enhances the inhibitory synaptic input onto these neurons [31].

How does the timing of different forms of activity-dependent plasticity influence their impact on neuronal circuits? Neural activity induces gene expression changes in cortical circuits, but it is unclear how these modifications impact neural circuits in the long term. A recent study has shown that activity-dependent plasticity of PV+ fast-spiking basket cells is specifically required for long-term, but not short or intermediate forms of memory consolidation [32]. This form of plasticity relies on dopamine signaling through D1/D5 receptors, and underlies the contribution of PV+ fast-spiking basket cells to fast oscillatory events during memory consolidation, including sharp-wave ripples. Similarly, transient disinhibition contributes to map stabilization following whisker deprivation, which suggest that inhibitory plasticity temporally precedes more classical forms of Hebbian plasticity during sensory map formation and stabilization [33]. The signaling pathways underlying this form of homeostatic plasticity are currently unknown, but they should be distinct from synapse-specific plasticity as they operate on different temporal and spatial scales [34]. Determining the molecular nature of this process from a temporal perspective will certainly aid to segregate interneurons based on their experience-dependent functions.

Altogether, these studies suggest that activity-induced transcriptional responses are triggered in a cell type-specific manner to achieve circuit homeostasis. In some cases, activity-dependent responses modified cardinal features in specific classes of interneurons, revealing a prominent role of neural activity in the specification of neuronal fate [18]. Future studies should aim to link global patterns of gene expression with specific neuronal fates, for example through approaches such as the recently developed ‘patch-seq’ method [35].

Cell-based tuning of cortical inhibitory circuits

Given the emerging significance of inhibitory plasticity for experience-dependent circuit remodeling, a fascinating avenue for tinkering with neuronal circuits consists in the addition of new interneurons by grafting their precursors into the brain. Nearly two decades of transplantation experiments [36,37] have unveiled that precursors from the MGE and the caudal ganglionic eminence (CGE) have a remarkable capacity to disperse within the postnatal and adult mouse cerebral cortex (as well as other CNS areas), settle within appropriate cortical layers, differentiate into lineage-specific interneurons, and functionally integrate. This has yielded interesting insights not only into the mechanisms regulating interneuron specification, migration, survival and integration, but also allowed the modification of diseased circuits and the reactivation of cortical plasticity [38].

For example, transplantation of varying numbers of interneuron precursors into the postnatal mouse cortex revealed that apoptotic cell death of interneurons is intrinsically determined and apparently not dependent on the limited supply of trophic support as posed by the neurotrophic hypothesis [39]. Another important finding of this study was that the frequency of inhibitory synaptic events observed in

surrounding pyramidal neurons did not linearly scale with interneuron density. This is suggestive of the operation of homeostatic mechanisms that adjust synaptic strength and number, and bears important implications for the interpretation of the functional consequences of interneuron addition to neural circuits. Bearing this in mind, several studies have shown that despite heterochronic transplantation into the cortex, interneuron precursors differentiate into their cognate interneuron lineages acquiring appropriate morphologies, laminar positions, neurochemical signatures and electrophysiological properties dependent on from which progenitor domain they originate. For instance, MGE-derived precursors gave rise to PV+ and SST+ interneurons [40], while VIP+ interneurons were only present in CGE transplants [37].

Based on such findings, transplantation of MGE precursors has been assessed for their ability to modify diseased neural circuits. Prime candidates for a cell-based therapy via addition of new interneurons are epilepsies [41]. Indeed, transplantation of MGE precursors into the cortex of Kv1.1 mutant mice simulating a human channelopathy prior to the appearance of seizures resulted in reduced seizure activity [42]. Interestingly, the disease-modifying effect of transplantation can be observed even after manifestation of seizures. When MGE precursors were grafted into the hippocampus of mice that had undergone pilocarpine-induced status epilepticus, there was a marked reduction in seizure occurrence and behavioral comorbidities of epilepsy compared to non-grafted controls [43]. While the earlier study showed that there was a significant increase in inhibitory input into pyramidal neurons, it would be worth repeating these studies with MGE precursors expressing optogenetic actuators to dissect the functional importance of inhibition in suppressing the epileptic activity,

as recently shown for transplantation experiments in models of Parkinson's disease [44].

Grafting of MGE-derived interneuron precursors into the adult striatum has also shown to bring about functional improvements in a rat model of Parkinson's disease [45] and reverse mechanical hypersensitivity produced by peripheral nerve injury upon grafting into the adult mouse spinal cord [46]. These effects are not restricted to rodent MGE precursors. When human embryonic stem cells were differentiated into MGE-like precursors and transplanted into the injured mouse spinal cord, a relieve of injury-related symptoms was observed [47]. These experiments highlight the remarkable capacity of interneurons derived from MGE grafts or generated from stem cells for integrating throughout the CNS and inducing functional circuit remodeling.

One of the most remarkable effects of MGE grafts is the reopening of a critical window of cortical plasticity (Fig. 2a). Cortical plasticity can be probed by closure of one eye (monocular deprivation, MD) and subsequent assessment of the resulting reorganization of connectivity between the open and closed eyes, respectively, and the primary visual cortex [48]. If MD is induced during a critical period of cortical development, then the closed eye will exhibit a reduced ability of activating neurons in the primary visual cortex compared to the open eye (a phenomenon referred to as ocular dominance plasticity, ODP). Intriguingly, the end of this critical period is closely associated with the maturation of PV+ and SST+-positive interneurons which in mice peaks at postnatal day 35 [49]. Southwell and colleagues have tested the hypothesis whether transplantation of MGE-derived precursors would result in the induction of a new window of ODP when MD was performed after the end of this critical window [50]. MGE grafts induced a new critical window the closure of which coincided with the maturation of the grafted interneurons. Remarkably, the time

course of maturation of the grafted interneurons matched the pace of endogenous interneuron differentiation suggestive of an intrinsic maturational program. More recently, the Gandhi lab has shown that the capacity of MGE grafts for reactivating ODP is not limited to early postnatal life but extends into adult [51]. Importantly, such reopened ODP allows for the reversal of visual deficits induced by visual deprivation during the earlier critical period at both circuit and behavioral levels. In keeping with the subclass-specific function of interneurons in experience-dependent function discussed above, the ODP re-opening effect is restricted to interneurons derived from MGE grafts (PV+ and SST+), while CGE grafts fail to induce a new critical window despite their successful dispersion and integration in the primary visual cortex [37,52].

The possibility of generating MGE-like progenitors from human induced pluripotent stem cells (hiPSC), which can mature into functional interneurons following grafting into the mouse cortex [53,54], allows addressing now the exciting question whether human MGE-like interneurons possess the same capacity for reactivating ODP as their rodent counterparts. Even more thrilling is the prospect that the duration of such induced critical window would be substantially prolonged along with the protracted maturation pace specific to human interneurons [55].

Prospects of inhibitory circuit remodeling by lineage reprogramming

In line with the subclass- and state-specific distribution of interneurons, it has been shown that during development the laminar distribution of cortical interneurons is under strong influence of the excitatory projection neuron subtypes they normally innervate [56-58]. Even more striking, when early postmitotic layer 2/3 callosal projection neurons were lineage-reprogrammed into layer 5-type of corticofugal

projection neurons by forced expression of the transcription factor Fezf2, the reprogrammed neurons received inhibitory synaptic input akin to endogenous layer 5 projection neurons [59] (Fig. 2b). It will be interesting to learn whether such remodeling of interneuron distribution and synaptic innervation is limited to the intrinsic window of interneuron maturation or represents a form of circuit plasticity extending into adulthood.

Direct lineage reprogramming of brain-resident cells into induced neurons has gained enormous momentum as novel means towards repairing the diseased brain [60,61]. Earlier in vitro studies discovered the possibility of reprogramming mouse astroglia into inhibitory neurons exhibiting firing properties of MGE-derived interneurons by forced expression of the transcription factors Ascl1 and Dlx2 [62]. Subsequent work revealed the feasibility of converting adult human brain pericytes into functional GABAergic neurons via two reprogramming factors Sox2 and Ascl1 [63], which also constitute the core of a reprogramming cocktail for mouse and human fibroblasts into GABAergic neurons [64]. Remarkably, these fibroblast-derived interneurons can functionally integrate when transplanted into the hippocampus. Most excitingly, direct lineage conversion of glia into neurons can be induced in vivo (Fig. 2b). However, evidence for specification of a defined subtype identity of the neurons induced via reprogramming is currently scarce. Consistent with the role of Neurog2 during the development of the cerebral cortex, neurons induced from glia in vivo by co-expression of Neurog2 and Bcl2 acquired a pyramidal neuron-like morphology and have a transcriptional profile characteristic of deep-layer projection neurons (Ctip2+, FoxP2+, Satb2- and Cux1-) [65]. Likewise, NeuroD1-induced neurons were found to be Tbr1+ and Ctip2+, again suggesting some degree of subtype specification [66]. In contrast, conversion of glia into subtype-specific

interneurons has not yet been achieved in the cerebral cortex *in vivo*, though induced GABA-immunoreactive neurons have been observed in the lesioned spinal cord following reprogramming by Sox2 [67]. Given the unique effects of different classes of interneurons on circuit function and plasticity, the development of robust strategies to induce the conversion of local glia into subtype-specific interneurons is of outstanding interest to unleash the full potential of these cells as disease-modifying agents.

Conclusions

Owing to their plasticity well into adulthood, interneurons are an emerging target for interventional circuit remodeling in the treatment of neurological and neuropsychiatric disease. To fully unleash such potential, it will be critical to uncover the molecular mechanisms that account for the remarkable flexibility of these neurons and their class-specific differences. Similarly, only a full grasp of the biology of grafted or lineage-reprogrammed interneurons will allow us to recruit these cells as means for re-tuning diseased cortical microcircuits to their healthy function.

Acknowledgements

We are grateful to members of the Marín and Berninger labs for discussions on this subject. We apologize to colleagues whose work is not cited in this review.

Regrettably, space was too limited to cite all significant original articles. Our work on this topic is supported by grants from the European Research Council (ERC-2011-AdG 293683), and the Wellcome Trust (103714MA) to O.M., and ERA-NET NEURON (01EW1604) and the German Research Foundation DFG (CRC1080) to B.B. N.D. was supported by an EMBO postdoctoral fellowship. N.M. was supported by a Human Frontier Science Program fellowship. O.M. is a Wellcome Trust Investigator.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Tremblay R, Lee S, Rudy B: **GABAergic interneurons in the neocortex: From cellular properties to circuits.** *Neuron* 2016, **91**:260-292.
2. Marín O: **Interneuron dysfunction in psychiatric disorders.** *Nat Rev Neurosci* 2012, **13**:107-120.
3. Ascoli GA, Alonso-Nanclares L, Anderson SA, Barrionuevo G, Benavides-Piccione R, Burkhalter A, Buzsaki G, Cauli B, Defelipe J, Fairén A, *et al.*: **Petilla terminology: nomenclature of features of GABAergic interneurons of the cerebral cortex.** *Nat Rev Neurosci* 2008, **9**:557-568.
4. Bartolini G, Ciceri G, Marín O: **Integration of GABAergic interneurons into cortical cell assemblies: lessons from embryos and adults.** *Neuron* 2013, **79**:849-864.
5. De Marco Garcia NV, Karayannis T, Fishell G: **Neuronal activity is required for the development of specific cortical interneuron subtypes.** *Nature* 2011, **472**:351-355.
6. Close J, Xu H, De Marco Garcia N, Batista-Brito R, Rossignol E, Rudy B, Fishell G: **Satb1 is an activity-modulated transcription factor required for the terminal differentiation and connectivity of medial ganglionic eminence-derived cortical interneurons.** *J Neurosci* 2012, **32**:17690-17705.

7. Nelson SB, Valakh V: **Excitatory/Inhibitory Balance and Circuit Homeostasis in Autism Spectrum Disorders.** *Neuron* 2015, **87**:684-698.
8. Gonzalez-Burgos G, Cho RY, Lewis DA: **Alterations in cortical network oscillations and parvalbumin neurons in schizophrenia.** *Biol Psychiatry* 2015, **77**:1031-1040.
9. Uhlhaas PJ, Singer W: **Neuronal dynamics and neuropsychiatric disorders: toward a translational paradigm for dysfunctional large-scale networks.** *Neuron* 2012, **75**:963-980.
10. Shao YR, Isett BR, Miyashita T, Chung J, Pourzia O, Gasperini RJ, Feldman DE: **Plasticity of recurrent I2/3 inhibition and gamma oscillations by whisker experience.** *Neuron* 2013, **80**:210-222.
11. Xue M, Atallah BV, Scanziani M: **Equalizing excitation-inhibition ratios across visual cortical neurons.** *Nature* 2014, **511**:596-600.
- This study provides an example of the specific regulation of E/I ratios in L2/3 pyramidal cells in the neocortex. The authors report that changes in the excitability of pyramidal neurons lead to a fine titration of inhibition by PV+ interneurons.
12. D'Amour J A, Froemke RC: **Inhibitory and excitatory spike-timing-dependent plasticity in the auditory cortex.** *Neuron* 2015, **86**:514-528.
- This study describes a form of inhibitory plasticity that requires interactions with co-activated excitatory synapses to regulate excitatory-inhibitory balance.

13. Liguz-Leczna M, Urban-Ciecko J, Kossut M: **Somatostatin and Somatostatin-Containing Neurons in Shaping Neuronal Activity and Plasticity.** *Front Neural Circuits* 2016, **10**:48.

14. Scheyltjens I, Arckens L: **The current status of somatostatin-interneurons in inhibitory control of brain function and plasticity.** *Neural Plast* 2016, **2016**:8723623.

15. Lourenco J, Pacioni S, Rebola N, van Woerden GM, Marinelli S, DiGregorio D, Bacci A: **Non-associative potentiation of perisomatic inhibition alters the temporal coding of neocortical layer 5 pyramidal neurons.** *PLoS Biol* 2014, **12**:e1001903.

16. House DR, Elstrott J, Koh E, Chung J, Feldman DE: **Parallel regulation of feedforward inhibition and excitation during whisker map plasticity.** *Neuron* 2011, **72**:819-831.

17. Campanac E, Gasselin C, Baude A, Rama S, Ankri N, Debanne D: **Enhanced intrinsic excitability in basket cells maintains excitatory-inhibitory balance in hippocampal circuits.** *Neuron* 2013, **77**:712-722.

18. Dehorter N, Ciceri G, Bartolini G, Lim L, del Pino I, Marin O: **Tuning of fast-spiking interneuron properties by an activity-dependent transcriptional switch.** *Science* 2015, **349**:1216-1220.

- This study identifies a homeostatic molecular mechanism controlling the intrinsic properties of adult PV+ fast-spiking basket cells. The activity-dependent molecular 'switch' operated by the transcription factor Er81 influences gene expression and modifies several properties in these interneurons.

19. Donato F, Rompani SB, Caroni P: **Parvalbumin-expressing basket-cell network plasticity induced by experience regulates adult learning.** *Nature* 2013, **504**:272-276.
20. Donato F, Chowdhury A, Lahr M, Caroni P: **Early- and late-born parvalbumin basket cell subpopulations exhibiting distinct regulation and roles in learning.** *Neuron* 2015, **85**:770-786.
- This study reveals that neurogenesis predicts baseline PV levels and connectivity in adult PV+ fast-spiking basket cells. PV+ interneuron subpopulations are involved in cellular networks with different functions: Early-born PV+ interneurons are required for rule consolidation, while late-born PV+ interneurons are involved in the acquisition of new information.
21. Ciceri G, Dehorter N, Sols I, Huang ZJ, Maravall M, Marín O: **Lineage-specific laminar organization of cortical GABAergic interneurons.** *Nat Neurosci* 2013, **16**:1199-1210.
22. Rymar VV, Sadikot AF: **Laminar fate of cortical GABAergic interneurons is dependent on both birthdate and phenotype.** *J Comp Neurol* 2007, **501**:369-380.
23. West AE, Greenberg ME: **Neuronal activity-regulated gene transcription in synapse development and cognitive function.** *Cold Spring Harb Perspect Biol* 2011, **3**.
24. Cohen SM, Ma H, Kuchibhotla KV, Watson BO, Buzsaki G, Froemke RC, Tsien RW: **Excitation-Transcription Coupling in Parvalbumin-Positive Interneurons Employs a Novel CaM Kinase-Dependent Pathway Distinct from Excitatory Neurons.** *Neuron* 2016, **90**:292-307.

- The authors reveal an important mechanism that regulates gene expression and long-term cellular plasticity in PV+ interneurons. They described a novel CaMK-dependent signaling pathway through Cav1 channels that triggers CREB phosphorylation, gene expression and dendritic branching.

25. Papoutsopoulou S, Janknecht R: **Phosphorylation of ETS transcription factor ER81 in a complex with its coactivators CREB-binding protein and p300.** *Mol Cell Biol* 2000, **20**:7300-7310.

26. Abe H, Okazawa M, Nakanishi S: **The Etv1/Er81 transcription factor orchestrates activity-dependent gene regulation in the terminal maturation program of cerebellar granule cells.** *Proc Natl Acad Sci USA* 2011, **108**:12497-12502.

27. Atallah BV, Bruns W, Carandini M, Scanziani M: **Parvalbumin-expressing interneurons linearly transform cortical responses to visual stimuli.** *Neuron* 2012, **73**:159-170.

28. Wilson NR, Runyan CA, Wang FL, Sur M: **Division and subtraction by distinct cortical inhibitory networks in vivo.** *Nature* 2012, **488**:343-348.

29. Lee SH, Kwan AC, Zhang S, Phoumthipphavong V, Flannery JG, Masmanidis SC, Taniguchi H, Huang ZJ, Zhang F, Boyden ES, *et al.*: **Activation of specific interneurons improves V1 feature selectivity and visual perception.** *Nature* 2012, **488**:379-383.

30. Spiegel I, Mardinly AR, Gabel HW, Bazinet JE, Couch CH, Tzeng CP, Harmin DA, Greenberg ME: **Npas4 regulates excitatory-inhibitory balance within neural circuits through cell-type-specific gene programs.** *Cell* 2014, **157**:1216-1229.

- The authors focus on transcriptional modifications regulating the function, localization, or expression of transcription factor downstream of calcium-signaling pathways during synapse development.
31. Mardinly AR, Spiegel I, Patrizi A, Centofante E, Bazinet JE, Tzeng CP, Mandel-Brehm C, Harmin DA, Adesnik H, Fagiolini M, *et al.*: **Sensory experience regulates cortical inhibition by inducing IGF1 in VIP neurons.** *Nature* 2016, **531**:371-375.
- The paper describes a molecular mechanism specific to VIP+ interneurons through which activity regulates the expression of *Igf1*, and this molecule in turn controls the formation of synaptic inputs onto these neurons.
32. Karunakaran S, Chowdhury A, Donato F, Quairiaux C, Michel CM, Caroni P: **PV plasticity sustained through D1/5 dopamine signaling required for long-term memory consolidation.** *Nat Neurosci* 2016, **19**:454-464.
- This study shows that learning-induced plasticity of PV+ interneurons is specifically required for long-term, but not short-term memory consolidation, a function that is sustained through D1/D5 dopamine receptor activation.
33. Li L, Gainey MA, Goldbeck JE, Feldman DE: **Rapid homeostasis by disinhibition during whisker map plasticity.** *Proc Natl Acad Sci U S A* 2014, **111**:1616-1621.
34. Bateup HS, Deneffio CL, Johnson CA, Saulnier JL, Sabatini BL: **Temporal dynamics of a homeostatic pathway controlling neural network activity.** *Front Mol Neurosci* 2013, **6**:28.
35. Fuzik J, Zeisel A, Mate Z, Calvigioni D, Yanagawa Y, Szabo G, Linnarsson S, Harkany T: **Integration of electrophysiological recordings with single-cell RNA-seq data identifies neuronal subtypes.** *Nat Biotechnol* 2016, **34**:175-183.

- This study underscores the subclass specific effects of interneurons on cortical plasticity. While MGE precursors give rise to interneurons that induce an additional critical period, CGE precursors fails to do so even though they disperse well within the cortex and functionally integrate as CGE lineage-specific interneurons.

36. Wichterle H, Garcia-Verdugo JM, Herrera DG, Alvarez-Buylla A: **Young neurons from medial ganglionic eminence disperse in adult and embryonic brain.** *Nat Neurosci* 1999, **2**:461-466.

37. Larimer P, Spatazza J, Espinosa JS, Tang Y, Kaneko M, Hasenstaub AR, Stryker MP, Alvarez-Buylla A: **Caudal Ganglionic Eminence Precursor Transplants Disperse and Integrate as Lineage-Specific Interneurons but Do Not Induce Cortical Plasticity.** *Cell Rep* 2016, **16**:1391-1404.

38. Southwell DG, Nicholas CR, Basbaum AI, Stryker MP, Kriegstein AR, Rubenstein JL, Alvarez-Buylla A: **Interneurons from embryonic development to cell-based therapy.** *Science* 2014, **344**:1240622.

39. Southwell DG, Paredes MF, Galvao RP, Jones DL, Froemke RC, Sebe JY, Alfaro-Cervello C, Tang Y, Garcia-Verdugo JM, Rubenstein JL, *et al.*: **Intrinsically determined cell death of developing cortical interneurons.** *Nature* 2012, **491**:109-113.

40. Du T, Xu Q, Ocbina PJ, Anderson SA: **NKX2.1 specifies cortical interneuron fate by activating Lhx6.** *Development* 2008, **135**:1559-1567.

41. Anderson SA, Baraban SC: **Cell Therapy Using GABAergic Neural Progenitors.** In *Jasper's Basic Mechanisms of the Epilepsies*, edn 4th edition. Edited by Noebels JL, Avoli M, Rogawski MA, Olsen RW, Delgado-Escueta AV: National Center for Biotechnology Information; 2012.

42. Baraban SC, Southwell DG, Estrada RC, Jones DL, Sebe JY, Alfaro-Cervello C, Garcia-Verdugo JM, Rubenstein JL, Alvarez-Buylla A: **Reduction of seizures by transplantation of cortical GABAergic interneuron precursors into Kv1.1 mutant mice.** *Proc Natl Acad Sci U S A* 2009, **106**:15472-15477.
43. Hunt RF, Girskis KM, Rubenstein JL, Alvarez-Buylla A, Baraban SC: **GABA progenitors grafted into the adult epileptic brain control seizures and abnormal behavior.** *Nat Neurosci* 2013, **16**:692-697.
44. Steinbeck JA, Choi SJ, Mrejeru A, Ganat Y, Deisseroth K, Sulzer D, Mosharov EV, Studer L: **Optogenetics enables functional analysis of human embryonic stem cell-derived grafts in a Parkinson's disease model.** *Nat Biotechnol* 2015, **33**:204-209.
45. Martinez-Cerdeno V, Noctor SC, Espinosa A, Ariza J, Parker P, Orasji S, Daadi MM, Bankiewicz K, Alvarez-Buylla A, Kriegstein AR: **Embryonic MGE precursor cells grafted into adult rat striatum integrate and ameliorate motor symptoms in 6-OHDA-lesioned rats.** *Cell Stem Cell* 2010, **6**:238-250.
46. Bràz JM, Sharif-Naeini R, Vogt D, Kriegstein A, Alvarez-Buylla A, Rubenstein JL, Basbaum AI: **Forebrain GABAergic neuron precursors integrate into adult spinal cord and reduce injury-induced neuropathic pain.** *Neuron* 2012, **74**:663-675.
47. Fandel TM, Trivedi A, Nicholas CR, Zhang H, Chen J, Martinez AF, Noble-Haeusslein LJ, Kriegstein AR: **Transplanted human stem cell-derived interneuron precursors mitigate mouse bladder dysfunction and central neuropathic pain after spinal cord injury.** *Cell Stem Cell* 2016, **19**:544-557.
48. Wiesel TN, Hubel DH: **Single-Cell Responses in Striate Cortex of Kittens Deprived of Vision in One Eye.** *J Neurophysiol* 1963, **26**:1003-1017.

49. Hensch TK: **Critical period plasticity in local cortical circuits.** *Nat Rev Neurosci* 2005, **6**:877-888.
50. Southwell DG, Froemke RC, Alvarez-Buylla A, Stryker MP, Gandhi SP: **Cortical plasticity induced by inhibitory neuron transplantation.** *Science* 2010, **327**:1145-1148.
51. Davis MF, Figueroa Velez DX, Guevarra RP, Yang MC, Habeeb M, Carathedathu MC, Gandhi SP: **Inhibitory Neuron Transplantation into Adult Visual Cortex Creates a New Critical Period that Rescues Impaired Vision.** *Neuron* 2015, **86**:1055-1066.
- This paper shows that the plasticity-inducing effect of grafted MGE-derived interneurons is not restricted to early postnatal life but can also be observed in adult animals. Reactivation of cortical plasticity allows for correction of visual impairments induced by sensory deprivation during the postnatal critical period on a circuit and behavioral level.
52. Tang Y, Stryker MP, Alvarez-Buylla A, Espinosa JS: **Cortical plasticity induced by transplantation of embryonic somatostatin or parvalbumin interneurons.** *Proc Natl Acad Sci U S A* 2014, **111**:18339-18344.
53. Nicholas CR, Chen J, Tang Y, Southwell DG, Chalmers N, Vogt D, Arnold CM, Chen YJ, Stanley EG, Elefanty AG, *et al.*: **Functional maturation of hPSC-derived forebrain interneurons requires an extended timeline and mimics human neural development.** *Cell Stem Cell* 2013, **12**:573-586.
54. Maroof AM, Keros S, Tyson JA, Ying SW, Ganat YM, Merkle FT, Liu B, Goulburn A, Stanley EG, Elefanty AG, *et al.*: **Directed differentiation and functional maturation of cortical interneurons from human embryonic stem cells.** *Cell Stem Cell* 2013, **12**:559-572.

55. Marín O: **Human cortical interneurons take their time.** *Cell Stem Cell* 2013, **12**:497-499.
56. Hevner RF, Daza RA, Englund C, Kohtz J, Fink A: **Postnatal shifts of interneuron position in the neocortex of normal and reeler mice: evidence for inward radial migration.** *Neuroscience* 2004, **124**:605-618.
57. Pla R, Borrell V, Flames N, Marín O: **Layer acquisition by cortical GABAergic interneurons is independent of Reelin signaling.** *J Neurosci* 2006, **26**:6924-6934.
58. Lodato S, Rouaux C, Quast KB, Jantrachotechatchawan C, Studer M, Hensch TK, Arlotta P: **Excitatory projection neuron subtypes control the distribution of local inhibitory interneurons in the cerebral cortex.** *Neuron* 2011, **69**:763-779.
59. Ye Z, Mostajo-Radji MA, Brown JR, Rouaux C, Tomassy GS, Hensch TK, Arlotta P: **Instructing Perisomatic Inhibition by Direct Lineage Reprogramming of Neocortical Projection Neurons.** *Neuron* 2015, **88**:475-483.
- Lineage reprogramming of early postmitotic layer 2/3 callosal projection neurons into layer 5-like corticofugal projection neurons results in a profound remodeling of inhibitory circuitry. Reprogrammed neurons in layer 2/3 exhibit an inhibitory neuron input pattern akin to endogenous layer 5 neurons.
60. Li H, Chen G: **In Vivo Reprogramming for CNS Repair: Regenerating Neurons from Endogenous Glial Cells.** *Neuron* 2016, **91**:728-738.
61. Heinrich C, Spagnoli FM, Berninger B: **In vivo reprogramming for tissue repair.** *Nat Cell Biol* 2015, **17**:204-211.
62. Heinrich C, Blum R, Gascon S, Masserdotti G, Tripathi P, Sanchez R, Tiedt S, Schroeder T, Gotz M, Berninger B: **Directing astroglia from the cerebral cortex into subtype specific functional neurons.** *PLoS Biol* 2010, **8**:e1000373.

63. Karow M, Sanchez R, Schichor C, Masserdotti G, Ortega F, Heinrich C, Gascon S, Khan MA, Lie DC, Dellavalle A, *et al.*: **Reprogramming of pericyte-derived cells of the adult human brain into induced neuronal cells.** *Cell Stem Cell* 2012, **11**:471-476.

64. Colasante G, Lignani G, Rubio A, Medrihan L, Yekhlief L, Sessa A, Massimino L, Giannelli SG, Sacchetti S, Caiazzo M, *et al.*: **Rapid Conversion of Fibroblasts into Functional Forebrain GABAergic Interneurons by Direct Genetic Reprogramming.** *Cell Stem Cell* 2015, **17**:719-734.

- This paper describes a cocktail of transcription factors around a core of Sox2 and Ascl1 [63] and complemented by Foxg1, Dlx5 and Lhx6 to derive functional GABAergic interneurons from reprogrammed mouse and human fibroblasts. These induced interneurons functionally integrate upon transplantation in vivo.

65. Gascon S, Murenu E, Masserdotti G, Ortega F, Russo GL, Petrik D, Deshpande A, Heinrich C, Karow M, Robertson SP, *et al.*: **Identification and Successful Negotiation of a Metabolic Checkpoint in Direct Neuronal Reprogramming.** *Cell Stem Cell* 2016, **18**:396-409.

- This paper demonstrates that successful reprogramming of glia into neurons depends on the ability of responding to altered metabolic demands. The authors identified Bcl2 as a key player to negotiate the transition from glycolysis to oxidative phosphorylation. Overexpression of Bcl2 markedly enhanced Neurog2-induced reprogramming of cortical glia into pyramidal-like neurons in vivo.

66. Guo Z, Zhang L, Wu Z, Chen Y, Wang F, Chen G: **In vivo direct reprogramming of reactive glial cells into functional neurons after brain injury and in an Alzheimer's disease model.** *Cell Stem Cell* 2014, **14**:188-202.
67. Su Z, Niu W, Liu ML, Zou Y, Zhang CL: **In vivo conversion of astrocytes to neurons in the injured adult spinal cord.** *Nat Commun* 2014, **5**:3338.

Figure legends

Figure 1. Activity-dependent regulation of PV+ interneuron plasticity. **(a)** Pyramidal cells received inhibitory inputs from PV cells that match their corresponding levels of activity (light to dark colors indicate low to high activity). When the activity of pyramidal cells is suppressed by expression of Kir2.1 (blue), the inhibition that these cells receive from PV interneurons is reduced and the variability of synaptic inputs in Kir2.1-expressing neurons is reduced. **(b)** Cell-specific plasticity on PV interneurons. The plasticity of early-born PV interneurons is primarily regulated through excitation whereas the plasticity of late-born PV interneurons is regulated through inhibition. PV+ fast-spiking interneurons are largely segregated in two main classes based in their birthdates, level of PV and synaptic inputs. Early-born basket cells (E9.5-E11.5; lower panel) preferentially allocate in deep layers of the cortex, express relatively high PV levels and receive strong excitatory inputs. Late-born basket cells (E13.5-E15.5; upper panel) preferentially allocate in superficial layers of the cortex, express relatively low PV levels and receive strong inhibitory inputs. In response to excitation, early-born PV basket cells specifically exhibit cell plasticity with enhanced excitatory inputs and increased PV levels. In response to inhibition, late-born PV basket cells specifically exhibit cell plasticity with enhanced inhibitory inputs and decreased PV levels. **(c)** Activity regulates CaMKI trafficking between the cytoplasm and the nucleus, thereby coupling excitation to gene expression in PV+ fast-spiking interneurons. Depolarizing stimuli induce Ca²⁺ influx via CaV1 channels, which signals CaMKI to translocate to the nucleus, induce CREB phosphorylation (P-CREB) and specific gene transcription and, ultimately, morphological changes in PV+ interneurons, including enhanced dendritic branching (red). **(d)** Activity-dependent transcriptional control of PV+ fast-spiking basket cell excitability. Expression of Er81 segregates PV+ fast-spiking basket cells in at least two major subtypes based on

their intrinsic properties, including the latency to firing at near threshold potential and synaptic inputs. PV+/Er81+ basket cells display delay to the first spike and strong excitatory inputs whereas PV+/Er81- basket cells show no delay and strong inhibitory inputs. In the absence of Er81 all PV+ interneurons show limited delay or become non-delayed, and exhibit reduced excitatory but enhanced inhibitory inputs (blue). PV, parvalbumin-expressing interneuron; PC, pyramidal Cell; ↑, increase in activity; ↓, Decrease in activity; P, phosphorylation. Modified from Refs. 11 (a); 19 and 20 (b), 24 (c); and 18 (d).

Figure 2. Cell-based remodeling of inhibitory circuits. **(a)** Grafting of MGE-derived interneuron precursors reactivates cortical plasticity. Monocular deprivation during a critical period of visual cortical plasticity (P19-35) results in the weakening of retinotopic inputs from the deprived eye (illustrated by the weak color grading) by P100. Grafting of interneuron precursors from the MGE into the visual cortex at P60 induces a new critical period that allows for cortical remapping of the input originating from the formerly closed and then reopened eye (illustrated by the enhanced color grading). **(b)** Circuit remodeling induced by lineage reprogramming. (1→2) Lineage reprogramming of layer 2/3 projection neurons into layer 5-like neurons by the transcription factor Fezf2 induces functional redistribution of perisomatic inhibitory inputs arising from PV+ interneurons to these cells. (1→3) Forced expression of Neurog2 and Bcl2 following brain injury in vivo induces conversion of local glia into Ctip2+ pyramidal-like neurons. (1→4) Selection of appropriate reprogramming factors may allow for lineage reprogramming of local glia into induced interneurons.

Figure 1

Figure 2
[Click here to download high resolution image](#)