King's Research Portal

DOI:

[10.1002/msc.1179](https://doi.org/10.1002/msc.1179)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

McIlroy, S., Sayliss, L., Browning, P., & Bearne, L. M. (2017). Aquatic Therapy for people with Persistent Knee Pain: A feasibility study. *Musculoskeletal Care*, 15(4), 350-355. <https://doi.org/10.1002/msc.1179>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Aquatic therapy for persistent knee pain

Title Page

Full title: Aquatic Therapy for people with Persistent Knee Pain: A feasibility study

Short title: Aquatic therapy for persistent knee pain

Key words: Aquatic therapy, knee, pain, exercise

Word Count: 1676

Authors:

Suzanne McIlroy, MSc BSc MCSP*

King's College Hospital

Lesley Sayliss, MSc MCSP

University of Hertfordshire

Paul Browning, MSc MCSP

University of Hertfordshire

Lindsay M. Bearne, PhD MSc MCSP*

King's College London

Corresponding Authors:

Suzanne McIlroy MSc BSc MCSP, King's College Hospital, Therapy Suite, 1st Floor Golden Jubilee Wing, Denmark Hill, London, SE5 9RS. Suzanne.McIlroy@nhs.net 0203 299 8220

Lindsay M. Bearne PhD MSc MCSP, Kings College London, Division of Health and Social Care Research, 3.26 Shepherd's House, Guy's Campus, London SE1 1UL. lindsay.bearne@kcl.ac.uk 0207 848 6283

Aquatic therapy for persistent knee pain

Annually, approximately 25% of people ≥ 55 years old in the United Kingdom (UK) experience persistent knee pain (PKP) which causes disability and reduced quality of life (Peat et al. 2001). Key management strategies comprise self-management advice, weight reduction and exercise (American Academy of Orthopedic Surgeons, National Institute for Health and Care Excellence, 2014; Fransen et al. 2015). Aquatic therapy (AT, exercise in warm water) improves pain and function in long-term musculoskeletal conditions (Barker et al. 2014; Waller et al. 2014). It utilises the properties of water including buoyancy (Harrison & Bulstrode, 1987), resistance (Petrick et al. 2001; Poyhonon et al. 2001) and heat (optimal water temperature 34-45 degrees Celcius (Pool Water Treatment Advisory Group, 2013) (Hall et al. 1990; Becker, 2009) to aid exercise performance.

Evidence from a recent meta-analysis, including eight high and three moderate quality studies (1092 participants), suggest that AT has a short term, small-medium beneficial effect on self-reported pain, joint mobility and function compared to a control group in people with lower limb osteoarthritis (OA) (Waller et al. 2014). When compared to land based exercise, evidence from meta-analyses of six studies (five with a risk of selection bias due to poor of reporting of allocation of concealment; 398 participants) suggests that AT produces similar improvements in self-reported pain and function (Lu et al. 2015). However, the longer term effects of AT are not clear (Cochrane et al, 2005; Lund et al, 2008).

There is no consensus on the most clinical or cost effective dosage or location of delivery for AT for people with PKP (Bartels et al. 2007; Holden et al. 2008). Published AT programmes range from 3x30 minutes weekly sessions for 6 weeks (Foley et al. 2003) to 2x60 minutes weekly sessions for 52 weeks (Lin et al. 2004; Cochrane et al. 2005) with some delivered in public pools at sub-optimal water temperature (Cochrane et al 2005; Wang et al. 2007, 2011). From a societal perspective, a one year AT intervention for people with lower limb OA showed a saving of £123—175/patient/ annum and incremental cost-effectiveness ratios ranged from £3,838 -£5,951 per quality-adjusted life-year in the AT group compared to usual care (Cochrane et al. 2005). However, a survey of 538 practising UK based physiotherapists (77% working in the NHS), suggests that 90% of Physiotherapists provide ≤ 6 sessions of AT for PKP patients (Holden et al. 2008). Therefore, the clinical and cost effectiveness of AT delivered at a dosage and location that reflects contemporary clinical practice needs to be established.

Aquatic therapy for persistent knee pain

To inform a definitive trial, this study evaluated the feasibility (participant recruitment, retention and outcome completion rate, adherence and adverse effects) of a randomised controlled trial (RCT) comparing six weekly sessions of AT compared to usual care on self-reported pain, function and general health in adults with PKP.

Methods:

This prospective two arm feasibility RCT received approval from the Joint University College London/University College London Hospital (UCLH) Committees on the Ethics of Human Research Committee Alpha (08/H0715/31) and the UCLH Research and Development Department (08/0103).

Participants and recruitment:

Potentially eligible patients were identified from their physiotherapy referral by a physiotherapist at one inner-city NHS hospital over a 10 week period. The lead researcher screened potential participants for eligibility via the telephone. Eligibility criteria included: adults ≥ 50 years old with self-reported PKP of >3 months duration; knee pain over the past 7 days of $>3/10$ on a Numerical Rating Scale (0=no pain, 10=worst pain imaginable) (Ferreira-Valente et al. 2011) and able and willing to provide informed consent.

Exclusion criteria comprised: self-reported early morning stiffness ≥ 30 minutes; contraindications to AT (Aquatic Therapy Association of Chartered Physiotherapists, 2015); upcoming knee surgery in the next 3 months; physiotherapy, AT or surgery for their PKP in the previous six months; the presence of another condition primarily limiting their mobility. Eligible participants attended one appointment where written informed consent was obtained prior to baseline assessment. The study target sample was 20 participants (Lancaster et al. 2004; Julious 2005; Hertzog, 2008).

Outcome Measures

Feasibility outcomes: the proportion of eligible people willing to enter the study, participant retention rate and outcome completion was recorded by the researcher. The reasons for withdrawal from the

Aquatic therapy for persistent knee pain

study (e.g. resentful demoralisation or other factors), adverse effects and attendance at AT sessions was recorded by the treating physiotherapist.

Sociodemographic and clinical outcomes: All sociodemographic characteristics (age (years), gender, Body Mass Index (NHS choices, 2015)) and clinical outcomes were collected at baseline and clinical outcomes were collected at 6 weeks by a researcher who was unaware of participant group allocation.

Self-reported disability was measured using the valid, reliable and responsive Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC) (Bellamy et al. 1988; Gentelle-Bonnassies et al. 2000; Angst et al. 2001). This 24-item questionnaire has three subscales; pain (5 items), stiffness (2 items) and function (17 items). Each item is measured on a 5-point Likert scale, with higher scores representing greater symptoms or disability.

Self-reported health status was assessed using the validated Short Form 12 (Jenkinson & Layte, 1997). The physical and mental subscale are each measured by 3 and 5-point Likert scales. Higher scores indicate better health (Gandhi et al, 2001).

Knee pain was assessed using a reliable and valid Visual Analogue Scale (0 -10cm anchors, 0 = no pain, 10 = worst pain imaginable) (McCormack et al. 1988). Participants marked a point on the 10cm line that represented their knee pain at that moment (Bellamy, 1997).

Walking ability was assessed by the 6 minute walk test (American Thoracic Society, 2002). Participants were asked to walk as far as possible around two cones, placed 10 metres apart in a straight corridor, in 6 minutes. The total distance walked (metres) in 6 minutes (6 Minute Walk Distance is reliable (Kennedy et al. 2005) and responsive to change (French et al. 2011) in people with long-term knee conditions.

Protocol

Following baseline assessment, all participants attended one 30 minute individual, self-management education session with a physiotherapist. This comprised: information on the causes of PKP, physical activity/aerobic exercise and knee exercise (e.g. quadriceps strengthening exercises); footwear advice and the use of shock absorbing insoles; activity pacing; pain (e.g. thermotherapy) and weight

Aquatic therapy for persistent knee pain

management (Arthritis Research UK, 2005; National Institute for Health and Care Excellence (NICE) (CG177), 2014).

Block randomization with computer generated random block sizes, was conducted and held by a third party unconnected with the study. Following baseline assessment and attendance at the self-management session, participants were randomly assigned to receive either usual care or AT in addition to usual care. The Physiotherapist not involved with the outcome assessment contacted the randomization administrator and informed the participant of their treatment allocation.

Aquatic Therapy Intervention Group

Participants randomised to receive AT, completed 6x30 minutes weekly group sessions of AT delivered by one of two Senior Physiotherapists who had undertaken postgraduate AT training. AT was conducted in a purpose built pool (3x5.6 metres, water temperature 33-35°C). All participants completed a circuit of exercises aimed to increase function (Table 1) (American College of Sports Medicine (2006), The American Geriatrics Society Panel on Exercise and Osteoarthritis (2001)). Participants continued to receive usual medical care as directed by their referring Physician (NICE, 2014).

Usual Care Comparison Group

Participants randomised to the comparison group continued to receive usual medical care e.g. medication, adjunctive therapies (NICE, 2014), as directed by their referring Physician.

Analysis:

Feasibility, sociodemographic and clinical outcomes were summarized using descriptive statistics as appropriate. Effect sizes were calculated using Hedges' *g* (standardized mean difference between outcome scores of AT and usual care group at 6 weeks, adjusted for different sample sizes) (Lenhard & Lenhard, 2016) and categorized as small (0.01–0.19), medium (0.2–0.79) or large (>0.79) (Cohen, 1998). Statistical analysis was performed on SPSS (version 12.0.1, SPSS Inc. Chicago).

Results:

Thirty nine patients were screened for eligibility during the 10 week recruitment period. Eighteen people met the eligibility criteria but four declined to participate due to other commitments (insufficient time) (2 males and 2 females) (Figure 1)). 14 participants ((mean (SD) age 63 (7.5) years, all female) were randomised (n=7 AT, n=7 comparison group, 78% recruitment rate). 13 participants completed the study. One participant withdrew from the study after randomisation due to their allocation to the comparison group (resentful demoralization) (Figure 1).

At baseline there were no substantial differences between the groups in any sociodemographic or clinical outcomes (Table 2) or between the participant that withdrew and those who completed the study.

Attendance at the AT was 98%. One participant did not attend one session (unrelated illness). No adverse effects were reported. All clinical measures were fully completed by the participants who completed the study.

At 6 weeks, small to medium effects were found in all clinical outcomes, favouring AT (Table 2).

Discussion:

This study demonstrated that a two arm RCT evaluating six, weekly sessions of AT compared to usual care was feasible and that our AT programme was well tolerated by people with PKP. The improvement in outcomes following AT is promising and suggests that a definitive RCT is warranted.

Our recruitment rate was higher than most previous studies of AT (Foley et al. 2003, Fransen et al. 2007) and implies that recruiting patients with PKP from NHS physiotherapy departments is feasible. Our study retention rates are good (93%), concurring with other work (Waller et al. 2014). High study retention rates are important because attrition of participants may threaten validity of RCTs, produce bias and limit statistical validity by reducing the power to detect true between group differences (Zweben et al.

Aquatic therapy for persistent knee pain

2009). Only one of our participants withdrew following randomisation to the comparison group but strategies, such as incentives, to minimise resentful demoralisation, should be considered (Adamson et al. 2006).

Our participants had high attendance at AT (98%,) which compares favourably with other studies (Foley et al. 2003; Fransen et al. 2007; Hinman et al. 2007; Lund et al. 2008; Wang et al. 2011) suggesting participants found the duration and intensity of our programme acceptable and tolerable, as there were no reported adverse effects.

Whilst our study was not powered to detect a change in any clinical variables, our findings are promising and reflect the findings of a meta-analysis of AT for other long-term musculoskeletal conditions (Barker et al. 2014) and lower limb OA (Waller et al. 2014) suggesting our programme warrants further investigation.

Our study has several strengths. Our programme is based on clinical guidelines and reflects contemporary UK practice, unlike other studies (Cochrane et al. 2005; Fransen et al. 2007; Silva et al. 2008; Wang et al. 2011). Key objective and self-reported clinical variables for PKP were included and were fully completed by our female participants, who had sociodemographic and clinical characteristics which are typical for participants with PKP entered into exercise studies (Jordan et al. 1996; Hurley et al. 2007), suggesting they did not find the assessment burdensome. Anecdotal feedback from clinicians and participants suggests that the protocol was feasible and that AT was acceptable to participants, although, formal qualitative exploration in a mixed method study is required.

This study has several limitations. We did not meet our target sample (n=20), although our recruitment rate was typical for studies of AT (5-6 participants /month (Foley et al. 2003; Lund et al. 2008). Only women were enrolled onto our study reflecting evidence from a meta-analysis evaluating the effectiveness of AT for lower limb OA, which reported that 73% of participants were female (Waller et al. 2014). Relatively few key measures were included in our study, similar to previous research (Foley et al. 2003; Cochrane et al. 2005; Silva et al. 2007; Lund et al. 2008; Gill et al. 2009; Wang et al. 2011) and no assessments were conducted after 6 weeks, so the impact of a longer study duration could not be

Aquatic therapy for persistent knee pain

captured. No formal qualitative data exploring the acceptability and experience of AT and our study protocol was collected.

This study will inform our definitive trial design. Dedicated research staff will be employed to screen referrals which may increase the identification and recruitment of potential participants. Other outcomes (e.g. quality of life, cost utility) and assessment time points (e.g. 6 and 30 months) will be included to more completely capture the effect of AT in the longer term, similar to other studies in PKP (Hurley 2007). Our AT programme will be offered at a range of times (e.g. evening) to increase accessibility and acceptability to males and others with life commitments. Records of other therapies prescribed during the trial and mixed research method will be used to formally explore the acceptability and experience of AT and the study to participants and clinicians.

Conclusion:

Our study showed that an RCT of a clinically practicable AT programme for people with PKP was feasible. It had typical recruitment, retention and outcome completion rate and was well tolerated by women with PKP. This suggests that our programme warrants further investigation and, if effective, could inform the delivery of AT for people with PKP in practice.

References

Adamson, J., Cockayne, S., Puffer, S., & Torgerson, D. J. (2006). Review of randomised trials using the post-randomised consent (Zelen's) design. *Contemporary Clinical Trials*, 27(4), 305-319.

American College of Sports Medicine (2006). ACSM's guidelines for exercise testing and prescription (7th ed.). Philadelphia: Lippincott Williams and Wilkins.

American Geriatrics Society Panel on Exercise and Osteoarthritis (2001). Exercise prescription for older adults with osteoarthritis pain: consensus practice recommendations. *Journal of the American Geriatrics Society*, 49, 808-823.

American Thoracic Society Committee on Proficiency Standards for Clinical Pulmonary Function Laboratories (2002). American Thoracic Society statement: guidelines for the six-minute walk test.

American Journal of Respiratory and Critical Care Medicine, 166,111.

Aquatic therapy for persistent knee pain

Angst, F., Aeschlimann, A., Steiner, W., & Stucki, G. (2001). Responsiveness of the WOMAC osteoarthritis index as compared with the SF-36 in patients with osteoarthritis of the legs undergoing a comprehensive rehabilitation intervention. *Annals of the Rheumatic Diseases*, 60(9), 834-840.

Aquatic Therapy Association of Chartered Physiotherapists (ATACP). (2015). *Guidance on good practice in Aquatic Physiotherapy*. London: Chartered Society of Physiotherapy.

Arthritis Research UK (2005). *Osteoarthritis of the knee. An information booklet*. Retrieved 1st March 2008 from <http://www.arc.org.uk/arthritis/patpubs/6027/6027.asp> [1 March 2008]

Barker, A. L., Talevski, J., Morello, R. T., Brand, C. A., Rahmann, A. E., & Urquhart, D. M. (2014). Effectiveness of aquatic exercise for musculoskeletal conditions: a meta-analysis. *Archives of Physical Medicine and Rehabilitation*, 95(9), 1776-1786.

Becker, B. E. (2009). Aquatic therapy: scientific foundations and clinical rehabilitation applications. *PM&R*, 1(9), 859-872. DOI: <http://dx.doi.org/10.1016/j.pmrj.2009.05.017>

Bellamy, N. (1997). Osteoarthritis clinical trials: candidate variables and clinimetric properties. *The Journal of Rheumatology*, 24(4), 768-78.

Bellamy, N., Buchanan, W. W., Goldsmith, C. H., Campbell, J. & Stitt, L. W. (1988). Validation study of WOMAC: a health status instrument for measuring clinically important patient relevant outcomes to Antirheumatic drug therapy in patients with osteoarthritis of the hip or knee. *The Journal of Rheumatology*, 15(2), 1833-1840.

Cochrane, T., Davey, R. C. & Matthes Edwards, S. M. (2005). Randomised controlled trial of the cost-effectiveness of water-based therapy for lower limb osteoarthritis. *Health Technology Assessment*, 9. DOI: <http://dx.doi.org/10.3310/hta9310>

Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*. 2nd edition. New Jersey: Lawrence Erlbaum Associates.

Aquatic therapy for persistent knee pain

Ferreira-Valente, M. A., Pais-Ribeiro, J. L. & Jensen, M. P. (2011). Validity of four pain intensity rating scales. *PAIN*, 152(10), 2399-404. DOI: 10.1016/j.pain.2011.07.005

Foley, A., Halbert, J., Hewitt, T. & Crotty, M. (2003). Does hydrotherapy improve strength and physical function in patients with osteoarthritis - a randomised controlled trial comparing a gym based and a hydrotherapy based strengthening programme. *Annals of the Rheumatic Diseases*, 62(12), 1162-1167. DOI: 10.1136/ard.2002.005272

Fransen, M., McConnell, S., Harmer, A. R., Van der Esch, M., Simic, M. & Bennell, K. L. (2015). Exercise for osteoarthritis of the knee: a Cochrane systematic review. *British Journal of Sports Medicine*, 49(24), 1554-7. DOI: 10.1136/bjsports-2015-095424

Fransen, M., Nairn, L., Winstanley, J., Lam, P. & Edmonds, J. (2007). Physical activity for osteoarthritis management: a randomized controlled clinical trial evaluating hydrotherapy or Tai Chi classes. *Arthritis Care & Research*, 57(3), 407-14. DOI: 10.1002/art.22621

French, H. P., Fitzpatrick, M. & FitzGerald, O. (2011). Responsiveness of physical function outcomes following physiotherapy intervention for osteoarthritis of the knee: an outcome comparison study. *Physiotherapy*, 97(4), 302-8. [doi:10.1016/j.physio.2010.03.002](https://doi.org/10.1016/j.physio.2010.03.002)

Gandhi, S.K., Salmon, J.W., Zhao, S.Z., Lambert, B.L., Gore, P.R. & Conrad, K. (2001). Psychometric evaluation of the 12-item short form health survey (SF-12) in osteoarthritis and rheumatoid arthritis clinical trials. *Clinical Therapeutics*, 23(7), 1080-1098.

Gentelle-Bonnassies, S., Le Claire, P., Mezieres, M., Ayral, X. & Dougados, M. (2000). Comparison of the responsiveness of symptomatic outcome measures in knee osteoarthritis. *Arthritis Care and Research*, 13(5), 280-285. DOI: 10.1002/1529-0131(200010)13:5<280::AID-ANR6>3.0.CO;2-6

Gill, S.D., McBurney, H. & Schulz, D. L. (2009). Land-based versus pool-based exercise for people awaiting joint replacement surgery of the hip or knee: results of a randomized controlled trial. *Archives of Physical Medicine and Rehabilitation*, 90(3), 388-394. DOI: [10.1016/j.apmr.2008.09.561](https://doi.org/10.1016/j.apmr.2008.09.561)

Hall, J., Bisson, D. & O'Hare, P. (1990). The physiology of immersion. *Physiotherapy*, 76(9), 517-521. DOI: [10.1016/S0031-9406\(10\)63019-2](https://doi.org/10.1016/S0031-9406(10)63019-2)

Aquatic therapy for persistent knee pain

- Harrison, R. & Bulstrode, S. (1987). Percentage weight-bearing during partial immersion in the hydrotherapy pool. *Physiotherapy Practice*, 3(2), 60-63. DOI: 10.3109/09593988709087741
- Hertzog, M. A. (2008). Considerations in determining sample size for pilot studies. *Research in Nursing & Health*, 31(2), 180-91.
- Hinman, R. S., Heywood, S. E. & Day, A. R. (2007). Aquatic physical therapy for hip and knee osteoarthritis: results of a single-blind randomised controlled trial. *Physical Therapy*, 87(1), 32-43.
- Holden, M. A., Nicholls, E. E., Hay, E. M. & Foster, N.E. (2008). Physical therapists' use of therapeutic exercise for patients with clinical knee osteoarthritis in the United Kingdom: in line with current recommendations? *Physical Therapy*, 88(10), 1109-1121. DOI: 10.2522/ptj.20080077
- Hurley, M. V., Walsh, N. E., Mitchell, H. L., Pimm, T. J., Patel, A., Williamson, E., ... & Reeves, B. C. (2007). Clinical effectiveness of a rehabilitation program integrating exercise, self-management, and active coping strategies for chronic knee pain: A cluster randomized trial. *Arthritis Care & Research*, 57(7), 1211-1219. DOI: 10.1002/art.22995
- Jenkinson, C., & Layte, R. (1997). Development and testing of the UK SF-12. *Journal of Health Services Research*, 2(1), 14-18.
- Jordan, J. M., Luta, G., Renner, J. B., Linder, G. F., Dragomir, A., Hochberg, M. C., & Fryer, J. G. (1996). Self-reported functional status in osteoarthritis of the knee in a rural southern community: the role of sociodemographic factors, obesity, and knee pain. *Arthritis & Rheumatism*, 9(4), 273-278.
- Julious, S. A. (2005). Sample size of 12 per group rule of thumb for a pilot study. *Pharmaceuticals Statistics*, 4(4), 287-91
- Kennedy, D. M., Stratford, P. W., Wessel, J., Gollish, J.D. & Penney, D. (2005). Assessing stability and change of four performance measures: a longitudinal study evaluating outcome following total hip and knee arthroplasty. *BMC Musculoskeletal Disorders*, 6(1), 1. DOI: 10.1186/1471-2474-6-3

Aquatic therapy for persistent knee pain

Lancaster, G.A., Dodd, S. & Williamson, P. R. (2004). Design and analysis of pilot studies: recommendations for good practice. *Journal of Evaluation in Clinical Practice*, 10(2), 307-12. DOI: 10.1111/j..2002.384.doc.x

Lenhard, W., Lenhard, A. (2016). Calculation of effect sizes. http://www.psychometrica.de/effect_size.html. Bibergau (Germany): Psychometrica. DOI: 10.13140/RG.2.1.3478.4245

Lin, S. Y., Davey, R. C., & Cochrane, T. (2004). Community rehabilitation for older adults with osteoarthritis of the lower limb: a controlled clinical trial. *Clinical Rehabilitation*, 18(1), 92-101. DOI: 10.1191/0269215504cr706oa

Lu, M., Su, Y., Zhang, Y., Zhang, Z., Wang, W., He, Z. ... Sheng, L. (2015). Effectiveness of aquatic exercise for treatment of knee osteoarthritis. *Zeitschrift für Rheumatologie*, 74(6), 543-52. DOI: 10.1007/s00393-014-1559-9

Lund, H., Weile, U., Christensen, R., Rostock, B., Downey, A., Bartel, E. M., ... Bliddal, H. (2008). A randomized controlled trial of aquatic and land-based exercise in patients with knee osteoarthritis. *Journal of Rehabilitation Medicine*, 40(2), 137-144. DOI: <http://dx.doi.org/10.2340/16501977-0134>

McCormack, H. M., David, J. D. L., & Sheather, S. (1988). Clinical applications of visual analogue scales: a critical review. *Psychological medicine*, 18(04), 1007-1019.

National Institute for Health and Care Excellence (2014). Osteoarthritis: care and management. NICE Guidelines CG177.

Peat, G., McCarney, R. & Croft, P. (2001). Knee pain and osteoarthritis in older adults: a review of community burden and current use of primary health care. *Annals of the Rheumatic Diseases*, 60(2), 91-97. DOI 10.1136/ard.60.2.91

NHS Choices (2015) <http://www.nhs.uk/Tools/Pages/Healthyweightcalculator.aspx> [19 June 2016]

Aquatic therapy for persistent knee pain

Petrick, M., Paulsen, T. & George, J. (2001). Comparison between quadriceps muscle strengthening on land and in water. *Physiotherapy*, 87(6), 310-317. DOI: [http://dx.doi.org/10.1016/S0031-9406\(05\)60765-](http://dx.doi.org/10.1016/S0031-9406(05)60765-1)

[1](#)

Pool Water Treatment Advisory Group (2013). *Swimming Pool Water*. PWTAG: London

Poyhonen, T., Keskinen, K., Kyrolainen, H., Hautala, A., Savolainen, J. & Malkia, E. (2001).

Neuromuscular function during therapeutic knee exercise under water and on dry land. *Archives of Physical Medicine and Rehabilitation*, 82(10), 1446-1452.

DOI: <http://dx.doi.org/10.1053/apmr.2001.25073>

Ratamess, N. A., Alvar, B. A., Evetoch, T. K., Housh, T. J., Kibler, W. B., & Kraemer, W. J. (2009).

Progression models in resistance training for healthy adults [ACSM position stand]. *Medicine and Science in Sports and Exercise*, 41(3), 687-708. DOI: 10.1249/MSS.0b013e3181915670

Silva, L. E., Valim, V., Pessanha, A. P., Oliveira, L. M., Myamoto, S., Jones, A. & Natour, J. (2008).

Hydrotherapy versus conventional land-based exercise for the management of patients with osteoarthritis of the knee: a randomized clinical trial. *Physical Therapy*, 88(1), 12-21. DOI:

10.2522/ptj.20060040

Wang, T. J., Belza, B., Elaine Thompson, F., Whitney, J. D., & Bennett, K. (2007). Effects of aquatic exercise on flexibility, strength and aerobic fitness in adults with osteoarthritis of the hip or knee. *Journal of advanced nursing*, 57(2), 141-152. DOI: 10.1111/j.1365-2648.2006.04102.x

Wang, T. J., Lee, S. C., Liang, S. Y., Tung, H. H., Wu, S. F. V., & Lin, Y. P. (2011). Comparing the efficacy of aquatic exercises and land-based exercises for patients with knee osteoarthritis. *Journal of Clinical Nursing*, 20(17-18), 2609-2622. DOI: 10.1111/j.1365-2702.2010.03675.x

Waller, B., Ogonowska-Slodownik, A., Vitor, M., Lambeck, J., Daly, D., Kujala, U. M. & Heinonen, A.

(2014). Effect of therapeutic aquatic exercise on symptoms and function associated with lower limb osteoarthritis: a systematic review with meta-analysis. *Physical Therapy*, 94, 1383-1395.

DOI: 10.2522/ptj.20130417

Aquatic therapy for persistent knee pain

Zweben, A., Fucito, L. M. & O'Malley, S. S. (2009) Effective strategies for maintaining research participation in clinical trials. *Drug information Journal*, 43(4), 459-67.

Acknowledgements:

The authors would like to thank Physiotherapy staff and patients at the University College London Hospital. Funding for an MSc in Musculoskeletal Physiotherapy was provided by the University College London Hospital's NHS trust and the Chartered Society of Physiotherapy Charitable Trust.

Disclosure statement:

No declarations of conflict.

Aquatic therapy for persistent knee pain

Table 1: Circuit of Aquatic Therapy exercises for people with Persistent Knee Pain

Warm up: Walking forwards, backwards, sideways

In standing:

- Hip and knee extension buoyancy resisted
- Knee flexion buoyancy assisted (hip in neutral)
- Knee extension buoyancy resisted (hip in neutral)
- Hip abduction
- Single leg stands +/- throw and catch ball
- Squats progressing to single leg squats
- Lunges
- Quadriceps, hamstring and gastrocnemius muscle stretch 2x30seconds

In sitting:

- Knee buoyancy assisted knee extension, buoyancy resisted knee flexion. Progressing to use flippers for resistance

Step:

- Step up
- Step down
- Step up and over

Supine supported by woggle/floats:

- Cycling, progression by incorporating flippers for resistance
- Hip extension buoyancy resisted

Dosage and Progression:

- To be performed at self-perceived 'somewhat hard' intensity on BORG scale (Dawes et al.

Aquatic therapy for persistent knee pain

2005)

- Progressed when rate of perceived exertion dropped below 'somewhat hard' by:
 - Increased repetition per set (8-12) or number of sets per exercises (1-3) (Ratamess et al. 2009)
 - Increasing speed to increase resistance (for example 1 repetition over 2 seconds progressed to 1 repetition over 1 second) in order to maintain rate of perceived exertion of at least 'somewhat hard'
 - Increased buoyancy aids to increase resistance
 - Move to shallower water to reduce effects of buoyancy

Aquatic therapy for persistent knee pain

Table 2. Sociodemographic and clinical outcome measures following 6 weeks of Aquatic Therapy or usual care

	Aquatic therapy			Comparison Group			
Outcomes	Baseline	6 weeks	Change	Baseline	6 weeks	Change	Effect size
							Hedges' <i>g</i>
Age (years)	64.3 (8.7)			62.3 (6.6)			
Weight (Kg)	78.1 (17.9)			88.1 (26.5)			
Height (m)	1.54 (0.04)			1.59 (0.06)			
BMI	32.4 (6.2)			34.6 (10.3)			
WOMAC – total	82.0 (19.1)	64.3 (18.2)	-17.7 (14.9)	65.0 (20.4)	67.6 (22.5)	2.6 (9.6)	0.16
WOMAC- Pain	16.7 (4.3)	11.6 (2.4)	- 5.1 (4.2)	14.3 (4.2)	14.6 (5.5)	0.3 (1.9)	0.73
WOMAC- stiffness	7.1 (1.3)	5.3 (2.3)	-1.9 (2.7)	6.6 (1.9)	6.9 (2.12)	0.3 (1.1)	0.72
WOMAC- function	58.1 (14.7)	47.4 (14.4)	-10.7 (8.9)	44.1 (16.7)	46.1 (16.3)	2.0 (9.6)	0.09
SF12 – mental	42.9 (16.8)	51.0 (12.0)	8.2 (9.2)	43.4 (13.1)	44.6(9.9)	1.2 (5.2)	0.58
SF 12 physical	33.4 (8.7)	37.7 (13.1)	4.3 (12.8)	34.6 (10.5)	34.6 (10.8)	0.01 (4.1)	0.26
VAS pain (cm)	7.9 (2.1)	4.5 (2.8)	-3.4 (3.9)	5.63 (2.6)	6.1 (2.5)	0.5 (1.1)	0.60
6MWD (m)	238.6 (85.5)	287.1 (63.2)	48.6 (58.1)	247.1 (98.4)	270.0 (74.8)	22.9 (42.7)	0.25

BMI – Body Mass Index, WOMAC - Western Ontario and McMaster Universities Osteoarthritis Index, SF12 - Short Form-12, 6MWD - Six minute walk distance and VAS - Visual Analogue Scale. All data presented as Mean (Standard Deviation)

Effect size: standardized mean difference between outcome scores of AT and usual care group at 6

weeks

Figure 1. Flow of participants

Aquatic therapy for persistent knee pain