King's Research Portal

DOI:

[10.1515/crelle-2017-0015](https://doi.org/10.1515/crelle-2017-0015)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Lekili, Y., & Polishchuk, A. (2019). A modular compactification of $M_{1,n}$ from A_∞ structures. *Journal für die Reine und Angewandte Mathematik*, 2019(755), 151–189. <https://doi.org/10.1515/crelle-2017-0015>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

A MODULAR COMPACTIFICATION OF $\mathcal{M}_{1,n}$ FROM A_∞ -STRUCTURES

YANKI LEKILI
ALEXANDER POLISHCHUK

ABSTRACT. We show that a certain moduli space of minimal A_∞ -structures coincides with the modular compactification $\overline{\mathcal{M}}_{1,n}(n-1)$ of $\mathcal{M}_{1,n}$ constructed by Smyth in [26]. In addition, we describe these moduli spaces and the universal curves over them by explicit equations, prove that they are normal and Gorenstein, show that their Picard groups have no torsion and that they have rational singularities if and only if $n \leq 11$.

INTRODUCTION

One of the motivations of the present work is to show that a study of the derived categories of coherent sheaves on such basic varieties as algebraic curves can uncover interesting geometry, including some aspects of the moduli spaces. The idea to study algebraic varieties via their derived categories, which has been around for a while (see [5]), recently got more focus and motivation coming from the homological mirror symmetry. In particular, it became clear that it is important to take into account the dg-enhancement, or the corresponding A_∞ -structure obtained by homological perturbation (in other words, one has to keep track of the higher Massey operations). Namely, if one takes a generator G of the derived category, then the corresponding Ext-algebra $\text{Ext}^*(G, G)$ has a structure of an A_∞ -algebra, from which the derived category can be recovered. This raises a natural question: what kind of A_∞ -algebras are obtained in this way, possibly for some specially chosen generators G . For example, for a smooth projective curve C we can take as a generator of the derived category the object

$$(0.0.1) \quad G(C, p_1, \dots, p_n) = \mathcal{O}_C \oplus \mathcal{O}_{p_1} \oplus \dots \oplus \mathcal{O}_{p_n},$$

where $p_1, \dots, p_n$ are distinct points of C . In the case $n = g$, the genus of C , the resulting A_∞ -algebras were studied in [16], [17] (for $g = 1$) and [22] (in general). The case of genus 0 curves was also studied in [22]. In this paper we consider the case when C is of genus 1 and n is arbitrary.

Note that to recover the derived category $D^b(C)$ from the A_∞ -algebra $E_{C, p_1, \dots, p_n}$ associated with the generator (0.0.1) one only needs the category of A_∞ -modules over $E_{C, p_1, \dots, p_n}$. On the other hand, this A_∞ -algebra itself, viewed up to a gauge equivalence, carries more information: in fact, one can recover the pointed curve $(C, p_1, \dots, p_n)$ from it. Moreover,

Y.L. is partially supported by a Royal Society Fellowship and a Marie Curie grant. A.P. is supported in part by the NSF grant DMS-1400390.

as was shown in [16] and [22], for $n = g$ one gets an equivalence between the appropriate moduli spaces of curves and moduli spaces of A_∞ -algebras.

In order to get such an equivalence, we allow the curves to be singular but also impose some restrictions on $(C, p_1, \dots, p_g)$ guaranteeing that the associative algebra structure on the $\text{Ext}^*(G, G)$ for $G = G(C, p_1, \dots, p_g)$ is independent of the pointed curve (whereas higher products do depend on it), and that G is indeed a generator of the perfect derived category of C . The latter property is equivalent to the ampleness of $\mathcal{O}_C(p_1 + \dots + p_g)$.

In this paper we study similar equivalence between moduli of curves and A_∞ -structures in the case of curves of arithmetic genus one with $n > 1$ (smooth) marked points. In this case the algebra $\text{Ext}^*(G, G)$ still does not depend on a curve provided one has $H^1(C, \mathcal{O}(p_i)) = 0$ for each of the marked points. In addition, we require $\mathcal{O}_C(p_1 + \dots + p_n)$ to be ample. We call the resulting moduli stack $\mathcal{U}_{1,n}^{\text{sns}}$ (“sns” stands for “strongly non-special”, since each p_i defines a non-special divisor).

Note that the relevance of our work to symplectic geometry is due to the fact that Fukaya category of n -marked (symplectic) torus gives rise to an A_∞ -structure in $\mathcal{U}_{1,n}^{\text{sns}}$. An attempt to directly compute this A_∞ -algebra runs into the well-known transversality problems: the constant maps contribute non-trivially to higher products (cf. [16], [17] for $n = 1$). To deal with this problem, one has to device a consistent set of perturbations which makes the computations hard (as one has to solve infinitely many PDEs in a consistent way). Our result tells that once we know that the cohomology algebra is isomorphic to $\text{Ext}^*(G, G)$ (which is easy to check), then we know that at the chain level the A_∞ -algebra arising from the Fukaya category corresponds to one of the curves in the moduli space $\mathcal{U}_{1,n}^{\text{sns}}$. From this characterization it follows that whatever one wants to compute for a given A_∞ -structure in our moduli space, in particular the one coming from the Fukaya category, we can do so using the *commutative* model given by the derived category of the corresponding curve. In the work in preparation [18], extending the ideas of [16], we will use this to establish a very precise form of homological mirror symmetry for n -punctured tori which is valid over $\mathbb{Z}$.

The paper consists of two parts. The first is a purely algebro-geometric study of the moduli stacks $\mathcal{U}_{1,n}^{\text{sns}}$ (without any reference to A_∞ -structures). Here our main result identifies $\mathcal{U}_{1,n}^{\text{sns}} \setminus \{C_{1,n}\}$, where $C_{1,n}$ is the *elliptic n -fold curve* (a certain generalization of the cuspidal cubic curve, see 1.5), with one of the alternative compactifications of $\mathcal{M}_{1,n}$ constructed and studied by Smyth in [26, 27]. Recall that for each m , $1 \leq m < n$, Smyth constructs the moduli stack $\overline{\mathcal{M}}_{1,n}(m)$ of *m -stable n -pointed curves*. The definition of m -stability involves restricting the type of singularity a curve can have (see Def. 1.5.1).

Smyth proves that these are proper irreducible Deligne-Mumford stacks over $\text{Spec}(\mathbb{Z}[1/6])$, that the corresponding coarse moduli spaces are projective and for $m \leq 10$ coincide with certain log canonical models of the Deligne-Mumford compactification $\overline{\mathcal{M}}_{1,n}$ proposed by Hassett and Keel ([27], [13]). Our interest in the current study is the case $m = n - 1$, and we consider a slight modification of the Smyth’s moduli stack, denoted by $\overline{\mathcal{M}}_{1,n}^\infty$ (see 1.5.3

for precise definition), which is a proper algebraic stack over $\mathrm{Spec}(\mathbb{Z})$ (resp., $\mathrm{Spec}(\mathbb{Z}[1/2])$ for $n = 2$, resp., $\mathrm{Spec}(\mathbb{Z}[1/6])$ for $n = 1$), such that $\overline{\mathcal{M}}_{1,n}(n-1) = \overline{\mathcal{M}}_{1,n}^\infty \times \mathrm{Spec}(\mathbb{Z}[1/6])$.

Theorem A[=Thm. 1.4.2+Thm. 1.5.7]. *Let us work over $\mathrm{Spec}(\mathbb{Z})$ for $n \geq 3$, over $\mathrm{Spec}(\mathbb{Z}[1/2])$ for $n = 2$, and over $\mathrm{Spec}(\mathbb{Z}[1/6])$ for $n = 1$. Let $\tilde{\mathcal{U}}_{1,n}^{sns} \rightarrow \mathcal{U}_{1,n}^{sns}$ be the $\mathbb{G}_m$ -torsor associated with the standard line bundle λ . Then $\tilde{\mathcal{U}}_{1,n}^{sns}$ is an affine scheme of finite type, defined by explicit equations. We have a natural equivalence*

$$(\tilde{\mathcal{U}}_{1,n}^{sns} \setminus \{C_{1,n}\})/\mathbb{G}_m \simeq \overline{\mathcal{M}}_{1,n}^\infty.$$

As a consequence of this approach to $\overline{\mathcal{M}}_{1,n}^\infty$ we are able to establish some additional results about geometry of this moduli space. For example, we show that for $n \geq 5$ it is a projective scheme given by explicit quadratic equations. We prove that $\overline{\mathcal{M}}_{1,n}^\infty$ is smooth for $n \leq 6$ and has rational singularities if and only if $n \leq 11$. For small values of n , we identify the moduli space explicitly. For example, we have an amusing identification

$$\overline{\mathcal{M}}_{1,6}^\infty \cong Gr(2, 5),$$

where $Gr(2, 5)$ stands for the Grassmannian of 2-planes in the 5-space. This extends the well known presentation of elliptic normal curves of degree 5 as linear sections of $Gr(2, 5)$ (see [10]) to singular curves (see Corollary 1.7.5).

A key structural result that we repeatedly use is the identification of the universal curve over $\overline{\mathcal{M}}_{1,n-1}^\infty$ with a blow-up of $\overline{\mathcal{M}}_{1,n}^\infty$ at $n-1$ special points, in such a way that the exceptional divisors of the blow-up correspond to the universal marked points (see Proposition 1.6.1). This also allows us to deduce that $\overline{\mathcal{M}}_{1,n}^\infty$ is normal and Gorenstein. In particular, we deduce that Smyth's moduli stacks $\overline{\mathcal{M}}_{1,n}(m)$ are normal and Gorenstein, which simplifies some statements in [27] formulated using the normalizations. In addition, we prove that the Picard group of $\overline{\mathcal{M}}_{1,n}^\infty$ is freely generated by λ (this was known rationally; we show that the Picard group has no torsion).

In the second part of the paper we identify the moduli stacks $\mathcal{U}_{1,n}^{sns}$ with an appropriate moduli of A_∞ -structures. Namely, we consider minimal A_∞ -algebra structures on the graded algebra $E_{1,n} = \mathrm{Ext}^*(G, G)$ for $G = G(C, p_1, \dots, p_n)$ given by (0.0.1). As we mentioned above, up to an isomorphism this algebra does not depend on $(C, p_1, \dots, p_n)$. Passing to the $\mathbb{G}_m$ -torsor $\tilde{\mathcal{U}}_{1,n}^{sns}$ allows to fix such an isomorphism canonically. Working over a field k , we prove using [22, Cor. 4.2.5] that the functor of minimal A_∞ -structures on $E_{1,n}$, viewed up to a gauge equivalence, is representable by an affine scheme $\widetilde{\mathcal{M}}_\infty(E_{1,n})$ of finite type over k .

Theorem B[=Thm. 2.2.8]. *Assume that $\mathrm{char}(k) \neq 2$ if $n = 2$ (resp., $\mathrm{char}(k) \neq 2, 3$ if $n = 1$). The affine scheme $\tilde{\mathcal{U}}_{1,n}^{sns} \times \mathrm{Spec}(k)$ is naturally isomorphic to the moduli scheme $\widetilde{\mathcal{M}}_\infty(E_{1,n})$ of minimal A_∞ -structures on the algebra $E_{1,n}$ up to a gauge equivalence.*

The proof follows the same strategy as in [22]: we reduce this to the study of the deformation functors around the most singular point of $\mathcal{U}_{1,n}^{sns}$ which corresponds to the trivial A_∞ -structure. Thus, using the tools developed in [22] we reduce the proof to studying the deformation functor of the elliptic n -fold curve.

Our work poses several natural questions. First, one can consider other alternative compactifications $\overline{\mathcal{M}}_{1,n}(m)$ constructed by Smyth and try to construct an interpretation in terms of A_∞ -structures. Given our explicit description of $\overline{\mathcal{M}}_{1,n}(n-1)$ by quadratic equations in projective space, one can ask whether one can calculate higher syzygies and whether the homogeneous coordinate algebra of $\overline{\mathcal{M}}_{1,n}(n-1)$ is Koszul. Another natural direction for a generalization is to study compactifications of $\mathcal{M}_{g,n}$ for $n > g$, related to appropriate A_∞ -structures—such a study was started by one of us in [23]. For $n < g$ the naive approach of studying A_∞ -structures on the corresponding Ext-algebra probably wouldn't work due to the fact that these A_∞ -structures are not determined (up to gauge equivalence) by any finite number of the higher products μ^n (see [9, Rem. 1.3.2.2]). This indicates that there are additional restrictions on A_∞ -structures coming from curves in this case.

1. CURVES OF ARITHMETIC GENUS 1 WITH n MARKED POINTS

1.1. Normal forms of pointed curves of arithmetic genus 1.

Let k be an algebraically closed field and let C be a reduced, connected projective curve over k of arithmetic genus 1 with n distinct smooth marked points: $p_1, \dots, p_n$. When C is smooth, $(C, p_1, \dots, p_n)$ defines a point of the moduli stack $\mathcal{M}_{1,n}$. To compactify $\mathcal{M}_{1,n}$ one has to allow C to be singular. More precisely, we are interested in *modular* compactifications of $\mathcal{M}_{1,n}$ obtained by specifying a deformation open class of curves satisfying the unique limit property in families (see [25]). Often such compactifications are constructed by identifying a class of singularities that C is allowed to have and then adding further restrictions on the configuration of the marked points guaranteeing the unique limit property. For example, the Deligne-Mumford moduli space of stable curves is a well known modular compactification for which C is allowed to have nodal singularities and $\omega_C(p_1 + \dots + p_n)$ is required to be ample ([8]). Smyth's moduli spaces $\overline{\mathcal{M}}_{1,n}(m)$ parametrize certain curves that are allowed to have elliptic l -fold points with $l \leq m$, in addition to nodes (see Section 1.5 below).

In our approach we start by considering the following requirements on $(C, p_1, \dots, p_n)$:

- (1) $h^0(\mathcal{O}_C(p_i)) = 1$ for all i .
- (2) $\mathcal{O}_C(p_1 + \dots + p_n)$ is ample.

We will see that the type of singularities of C will be determined by these conditions *a posteriori*.

We will first follow a pedestrian approach in constructing the moduli space of arithmetic genus 1 curves $(C, p_1, \dots, p_n)$ which satisfy the conditions (1) and (2). Recall that a reduced curve is automatically Cohen-Macaulay, so we have a relative dualizing sheaf ω_C on C . The condition (1) has a simple interpretation that we will use repeatedly.

Lemma 1.1.1. (i) *Let C be a reduced connected projective curve of arithmetic genus 1, p a smooth point on C . Then $h^0(\mathcal{O}_C(p)) = 1$ if and only if $h^1(\mathcal{O}_C(p)) = 0$ if and only if the restriction map*

$$H^0(C, \omega_C) \rightarrow \omega_C|_p$$

is an isomorphism.

(ii) *Let $\pi : C \rightarrow S$ be a flat projective morphism of relative dimension 1, with reduced connected geometric fibers of arithmetic genus 1, and let $p : S \rightarrow C$ be a section such that π is smooth near p . Then the following conditions are equivalent:*

- (1) *the natural map $\mathcal{O}_S \rightarrow \pi_*(\mathcal{O}_C(p(S)))$ is an isomorphism;*
- (2) *$R^1\pi_*(\mathcal{O}_C(p(S))) = 0$;*
- (3) *the natural map $\pi_*(\omega_C) \rightarrow p^*\omega_C$ is an isomorphism.*

Proof. (i) One uses the fact that $\chi(\mathcal{O}_C(p)) = 1$ (by Riemann-Roch theorem) and $h^1(\mathcal{O}_C(p)) = h^0(\omega_C(-p))$ (by Grothendieck-Serre duality). Since $H^0(C, \omega_C)$ is 1-dimensional, the restriction map to $\omega_C|_p$ is an isomorphism if and only if $h^0(\omega_C(-p(S))) = 0$.

(ii) The exact sequence

$$0 \rightarrow \omega_C(-p(S)) \rightarrow \omega_C \rightarrow p_*p^*\omega_C \rightarrow 0$$

shows that (3) implies that the map

$$R^1\pi_*(\omega_C(-p(S))) \rightarrow R^1\pi_*(\omega_C)$$

is an isomorphism. By Grothendieck duality, this is equivalent to (1). Conversely, if (1) holds then we get that the morphism of line bundles $\pi_*\omega_C \rightarrow p^*\omega_C$ is surjective, hence, an isomorphism. Thus, (1) and (3) are equivalent. The same exact sequence shows that (3) implies the vanishing of $\pi_*(\omega_C(-p(S)))$, and hence, by duality, of $R^1\pi_*(\mathcal{O}_C(p(S)))$. Conversely, (2) implies the similar condition for every fiber, hence, by part (i), we get the pointwise version of (3), and the condition (3) itself follows. $\square$

We will also use the following version of the residue theorem for singular curves.

Lemma 1.1.2. *Let $\pi : C \rightarrow S$ be a flat projective morphism of relative dimension 1, with connected reduced geometric fibers, and let $p_1, \dots, p_n : S \rightarrow C$ be sections such that π is smooth near each p_i . Then for any $\eta \in H^0(C \setminus \{p_1, \dots, p_n\}, \omega_{C/S})$ one has*

$$\sum_i \text{Res}_{p_i}(\eta) = 0.$$

Proof. Let $D = \sum n_i p_i$ be an effective relative divisor such that $\eta \in H^0(C, \omega_{C/S}(D))$. Note that $\underline{\text{Ext}}^i(\mathcal{O}_D, \omega_{C/S}) = 0$ for $i \neq 1$ and $\underline{\text{Ext}}^1(\mathcal{O}_D, \omega_{C/S})$ is supported at $p_1(S) \cup \dots \cup p_n(S)$. Therefore, we have the relative Serre duality pairing

$$\tau : \pi_* \underline{\text{Ext}}^1(\mathcal{O}_D, \omega_{C/S}) \otimes \pi_*(\mathcal{O}_D) \rightarrow R^1 \pi_* \omega_{C/S} \simeq \mathcal{O}_S.$$

By definition, its restriction τ_1 to the section 1 of $\pi_*(\mathcal{O}_D)$ is obtained by applying the functor

$$R\pi_* R\text{Hom}(\cdot, \omega_{C/S})$$

to the projection $\mathcal{O}_C \rightarrow \mathcal{O}_D$. Therefore, the exact sequence

$$0 \rightarrow \mathcal{O}_C(-D) \rightarrow \mathcal{O}_C \rightarrow \mathcal{O}_D \rightarrow 0$$

shows that the composition

$$\pi_*(\omega_{C/S}(D)) = \pi_* \underline{\text{Hom}}(\mathcal{O}_C(-D), \omega_{C/S}) \xrightarrow{\delta} \pi_* \underline{\text{Ext}}^1(\mathcal{O}_D, \omega_{C/S}) \xrightarrow{\tau_1} \mathcal{O}_S$$

is zero. The required formula will follow once we compute local contributions to $\tau_1(\delta(\eta)) = 0$ at each point p_i , corresponding to the canonical decomposition

$$\underline{\text{Ext}}^1(\mathcal{O}_D, \omega_{C/S}) \simeq \bigoplus_{i=1}^n \underline{\text{Ext}}^1(\mathcal{O}_{n_i p_i}, \omega_{C/S}).$$

Now the exact sequence

$$0 \rightarrow \mathcal{O}_C(-n_i p_i) \rightarrow \mathcal{O}_C \rightarrow \mathcal{O}_{n_i p_i} \rightarrow 0$$

induces an identification

$$p_* \underline{\text{Ext}}^1(\mathcal{O}_{n_i p_i}, \omega_{C/S}) \simeq p_*(\omega_{C/S}(n_i p_i)/\omega_{C/S}),$$

compatible with δ , and we claim that the projection to $\mathcal{O}_S$ induced by τ_1 is given by the residue at p_i . Indeed, we can replace C by an open neighborhood of p_i , smooth over S , in which case this is standard. $\square$

Let us fix a nonzero generator $\omega \in H^0(C, \omega_C)$. Note that by Lemma 1.1.1, ω does not vanish at any of the marked points p_i . Below we will use this generator to define residues of rational functions at points p_i .

The cases $n = 1, 2$ require special attention depending on the characteristic of k , which we will come back to later (see Sections 1.2 and 1.3). For now, we assume that $n \geq 3$.

Note that we have $H^1(C, \mathcal{O}_C(p_i)) = 0$, hence for each $i \neq j$ we have $h^1(\mathcal{O}_C(p_i + p_j)) = 0$ and $h^0(\mathcal{O}_C(p_i + p_j)) = 2$. By the residue theorem (see Lemma 1.1.2), for each $i \neq j$ there exist $h_{ij} \in H^0(C, \mathcal{O}(p_i + p_j))$ such that

$$\text{Res}_{p_i}(h_{ij}\omega) = 1 \quad \text{and} \quad \text{Res}_{p_j}(h_{ij}\omega) = -1.$$

Let us fix a choice of h_{1i} for $i = 2, \dots, n$. For distinct $i, j \geq 2$ we set

$$c_{ij} = h_{1i}(p_j).$$

The functions $1, h_{12}, h_{13}, \dots, h_{1n}$ form a basis of $H^0(C, \mathcal{O}(p_1 + \dots + p_n))$.

Set $D = p_1 + \dots + p_n$. It is easy to see that for each $N \geq 2$ the functions

$$1, h_{12}^r h_{13}, h_{12}^m, \dots, h_{1n}^m, \text{ where } 1 \leq m \leq N, 1 \leq r \leq N-1,$$

form a basis of $H^0(C, \mathcal{O}_C(ND))$. Let us normalize h_{12} and h_{13} (adding a constant to them) by requiring that

$$h_{12}(p_3) = h_{13}(p_2) = 0,$$

i.e., $c_{23} = c_{32} = 0$. Then $h_{12}h_{13}$ is regular near p_2 and p_3 , hence, for distinct indices $i, j \geq 2$ the element $h_{1i}h_{1j} - h_{12}h_{13}$ belongs to $H^0(C, \mathcal{O}_C(p_1 + p_i + p_j))$. Looking at the residues at p_i and p_j we see that

$$(1.1.1) \quad h_{1i}h_{1j} - h_{12}h_{13} = c_{ij}h_{1j} + c_{ji}h_{1i} + d_{ij}$$

for some constants d_{ij} . Similarly,

$$h_{12}h_{13}^2 - h_{12}^2h_{13} \in H^0(C, \mathcal{O}_C(2p_1 + p_2 + p_3)).$$

Hence, we should have a relation of the form

$$(1.1.2) \quad h_{12}h_{13}^2 - h_{12}^2h_{13} = ah_{12}h_{13} + bh_{12} + ch_{13} + d.$$

Thus, setting $x_i = h_{1i}$ for $i = 2, \dots, n$, we get relations of the form

$$(1.1.3) \quad x_i x_j = x_2 x_3 + c_{ij} x_j + c_{ji} x_i + d_{ij}, \text{ for } i < j, (i, j) \neq (2, 3),$$

$$(1.1.4) \quad x_2 x_3^2 = x_2^2 x_3 + a x_2 x_3 + b x_2 + c x_3 + d.$$

Let us normalize the choices of h_{1i} by requiring that

$$(1.1.5) \quad c_{32} = c_{i3} = 0 \text{ for } i \geq 2, i \neq 3.$$

Lemma 1.1.3. *Consider the algebra A over a commutative ring R with generators $x_2, \dots, x_n$ (where $n \geq 3$) and the defining relations (1.1.3), (1.1.4), where $c_{ij}, d_{ij}, a, b, c, d \in R$ and (1.1.5) is satisfied. Then the elements*

$$1, x_2^m x_3, x_i^m, \text{ for } i \geq 2, m \geq 1$$

form an R -basis in A if and only if the following relations hold:

$$(1.1.6) \quad \begin{aligned} a &= c_{3i} - c_{2i} - c_{i2}, \\ d_{ij} &= d_{2i} = -c, \\ d_{3i} &= b - c_{3i}c_{i2}, \\ d &= -c_{3i}c - c_{2i}d_{3i}, \\ c_{ij} + c_{ji} &= c_{3i} + c_{2j} + c_{j2}, \\ c_{ji}c_{2i} &= c_{3i}c_{2i} + c_{j2}c_{2i} + c_{ji}c_{2j} + d_{2j}, \\ c_{ji}c_{i2} + c_{ij}c_{j2} &= c_{i2}c_{j2} + b + c, \\ c_{ik}c_{jk} &= c_{ij}c_{jk} + c_{ji}c_{ik} + c_{3k}c_{2k} + d_{ij}, \end{aligned}$$

where $i, j, k \geq 4$ are distinct.

Proof. This is a standard application of the Gröbner basis technique. Namely, we order the variables naturally ($x_2 < x_3 < \dots < x_n$) and use the degree-lexicographical order on monomials in (x_i) . So, we have

$$1 < x_2 < x_3 < \dots < x_n < x_2^2 < x_2x_3 < \dots < x_2x_n < x_3^2 < \dots$$

Then, by the diamond lemma [6], it is enough to apply the relations to write the following monomials in two ways: $x_ix_jx_k$ and $x_2x_3^2x_m$ with $m \geq 4$. From equations (1.1.3) and (1.1.4) (and also (1.1.5)), for $i \geq 4$, we can simplify $x_2x_3x_i$ in the following two ways.

First reduce x_3x_i to get

$$x_2x_3x_i = x_2^2x_3 + c_{3i}x_2x_3 + c_{2i}c_{3i}x_i + (c_{3i}c_{i2} + d_{3i})x_2 + c_{3i}d_{2i}.$$

Alternatively, first reduce x_2x_i to get

$$x_2x_3x_i = x_2^2x_3 + (a + c_{i2} + c_{2i})x_2x_3 + c_{2i}c_{3i}x_i + (c + d_{2i})x_3 + bx_2 + (d + c_{2i}d_{3i}).$$

By equating the coefficients, we obtain the equations

$$\begin{aligned} c_{3i} &= a + c_{2i} + c_{i2}, \\ c + d_{2i} &= 0, \\ c_{3i}c_{i2} + d_{3i} &= b, \\ c_{3i}d_{2i} &= d + c_{2i}d_{3i}. \end{aligned} \tag{1.1.7}$$

Next, we consider $x_2x_ix_j$ for $i, j \geq 4$ and $i \neq j$. First reduce x_2x_j to get

$$\begin{aligned} x_2x_ix_j &= x_2^2x_3 + (c_{3i} + c_{2j} + c_{j2})x_2x_3 \\ &\quad + (c_{3i}c_{2i} + c_{j2}c_{2i} + c_{ji}c_{2j} + d_{2j})x_i \\ &\quad + c_{2j}c_{ij}x_j + (c_{i2}c_{j2} + c_{3i}c_{i2} + d_{3i})x_2 \\ &\quad + c_{3i}d_{2i} + c_{2j}d_{ji} + c_{j2}d_{2i}. \end{aligned}$$

Alternatively, first reduce x_ix_j to get

$$\begin{aligned} x_2x_ix_j &= x_2^2x_3 + (c_{ij} + c_{ji})x_2x_3 + c_{ji}c_{2i}x_i + c_{ij}c_{2j}x_j \\ &\quad + (c_{ij}c_{j2} + c_{ji}c_{i2} + d_{ij})x_2 + c_{ij}d_{2j} + c_{ji}d_{2i}. \end{aligned}$$

Equating the coefficients, we get the relations

$$\begin{aligned} c_{3i} + c_{2j} + c_{j2} &= c_{ij} + c_{ji}, \\ c_{3i}c_{2i} + c_{j2}c_{2i} + c_{ji}c_{2j} + d_{2j} &= c_{ji}c_{2i}, \\ c_{i2}c_{j2} + c_{3i}c_{i2} + d_{3i} &= c_{ij}c_{j2} + c_{ji}c_{i2} + d_{ij}, \\ c_{3i}d_{2i} + c_{2j}d_{ji} + c_{j2}d_{2i} &= c_{ij}d_{2j} + c_{ji}d_{2i}. \end{aligned} \tag{1.1.8}$$

One can do a similar analysis using $x_3x_ix_j$ for $i, j \geq 4$ and $i \neq j$. This gives

$$\begin{aligned} x_3x_ix_j &= x_2^2x_3 + (c_{3i} + c_{3j})x_2x_3 + (c_{2i}c_{3i} + c_{3j}c_{ji} + d_{3j})x_i \\ &\quad + c_{3j}c_{ij}x_j + (c_{3i}c_{i2} + d_{3i})x_2 + c_{3i}d_{2i} + c_{3j}d_{ij}. \end{aligned}$$

and

$$\begin{aligned} x_3x_ix_j &= x_2^2x_3 + (a + c_{ij} + c_{ji})x_2x_3 + c_{ji}c_{3i}x_i \\ &\quad + c_{3j}c_{ij}x_j + (c + d_{ij})x_3 + bx_2 + d + c_{ij}d_{3j} + c_{ji}d_{3i}. \end{aligned}$$

The new relations we get from these are

$$\begin{aligned} (1.1.9) \quad & c_{2i}c_{3i} + c_{3j}c_{ji} + d_{3j} = c_{ji}c_{3i}, \\ & c_{3i}d_{2i} + c_{3j}d_{ij} = c_{ij}d_{3j} + c_{ji}d_{3i}, \\ & c + d_{ij} = 0. \end{aligned}$$

To get the final equation in the statement we have to consider $x_ix_jx_k$ for $i, j, k \geq 4$. We can simplify $x_ix_jx_k$ in the following two ways. First reduce x_ix_j to get

$$\begin{aligned} x_ix_jx_k &= x_2x_3x_k + c_{ij}x_jx_k + c_{ji}x_ix_k + d_{ij}x_k \\ &= x_2^2x_3 + (c_{3k} + c_{ij} + c_{ji})x_2x_3 \\ &\quad + (c_{3k}c_{2k} + d_{ij} + c_{ij}c_{jk} + c_{ji}c_{ik})x_k \\ &\quad + c_{ij}c_{kj}x_j + c_{ji}c_{ki}x_i + (c_{3k}c_{k2} + d_{3k})x_2 \\ &\quad + c_{3k}d_{2k} + c_{ij}d_{jk} + c_{ji}d_{ik}. \end{aligned}$$

Alternatively, first reduce x_jx_k to get

$$\begin{aligned} x_ix_jx_k &= x_2x_3x_i + c_{jk}x_ix_k + c_{kj}x_ix_j + d_{jk}x_i \\ &= x_2^2x_3 + (c_{3i} + c_{jk} + c_{kj})x_2x_3 \\ &\quad + c_{jk}c_{ik}x_k + c_{kj}c_{ij}x_j \\ &\quad + (c_{3i}c_{2i} + c_{kj}c_{ji} + c_{jk}c_{ki} + d_{jk})x_i \\ &\quad + (c_{3i}c_{i2} + d_{3i})x_2 \\ &\quad + c_{3i}d_{2i} + c_{jk}d_{ik} + c_{kj}d_{ij}. \end{aligned}$$

The only new relation that we get from this is

$$(1.1.10) \quad c_{3k}c_{2k} + c_{ij}c_{jk} + c_{ji}c_{ik} + d_{ij} = c_{jk}c_{ik},$$

for distinct $i, j, k \geq 4$.

It turns out the ambiguities in reducing the monomials $x_2x_3^2x_m$ for $m \geq 4$ are already resolved by the above relations (we leave this as an exercise; this also follows from the observation in part (ii) of Proposition 1.1.5 below). It is easy to see that the equations (1.1.7), (1.1.8), (1.1.9), (1.1.10) simplify to the equations given in the statement of the lemma (the last equation in (1.1.8) and the first two equations in (1.1.9) are discarded as they are consequences of the other relations). $\square$

In particular, from equations (1.1.6), we can see that for $n \geq 4$ all the coefficients can be expressed in terms of the coefficients a, b, c, c_{i2}, c_{2i} and c_{ij} , where $4 \leq i < j$, which satisfy further polynomial relations. We can rewrite these relations as follows.

Definition 1.1.4. For $n \geq 3$, let U_n , be the affine scheme over $\mathbb{Z}$ defined by the equations (1.1.5) and (1.1.6) on coordinates $c_{ij}, d_{ij}, a, b, c, d$.

Proposition 1.1.5. (i) For $n = 3$ we have $U_3 \simeq \mathbb{A}_{\mathbb{Z}}^4$ with coordinates a, b, c, d .

For $n \geq 4$ let us set for each $i \geq 4$

$$c_i = c_{2,i}, \quad \bar{c}_i = c_{i,2}.$$

Let us also set $\bar{c} = b + c$. Then for $n \geq 4$ the ring of functions on U_n is generated by

$$a, c, \bar{c}, (c_i, \bar{c}_i)_{4 \leq i \leq n}, (c_{ij})_{4 \leq i < j \leq n},$$

with the defining relations

$$(1.1.11) \quad \begin{aligned} (c_j - c_i)c_{ij} &= (a + \bar{c}_i + c_j + \bar{c}_j)c_j - c, \\ (\bar{c}_i - \bar{c}_j)c_{ij} &= (a + c_i + \bar{c}_i + c_j)\bar{c}_i - \bar{c} \quad \text{for } 4 \leq i < j, \\ c_{ij}c_{jk} + c_{ji}c_{ik} - c_{ik}c_{jk} + (a + c_k + \bar{c}_k)c_k &= c, \end{aligned}$$

for $4 \leq i < j < k$, where

$$c_{ji} = a + c_i + \bar{c}_i + c_j + \bar{c}_j - c_{ij}$$

In particular,

$$U_4 \simeq \mathbb{A}_{\mathbb{Z}}^5$$

with coordinates $a, c, \bar{c}, c_4, \bar{c}_4$, and

$$U_5 \simeq \mathbb{A}_{\mathbb{Z}}^6$$

with coordinates $a, c_4, \bar{c}_4, c_5, \bar{c}_5, c_{45}$.

(ii) Consider the morphism $U_{n+1} \rightarrow U_n$ forgetting the coordinates $c_{n+1}, \bar{c}_{n+1}$ and $c_{i,n+1}$. Let also $\mathcal{C}_n \rightarrow U_n$ be the affine family of curves defined by (1.1.3), (1.1.4). Then the map

$$(1.1.12) \quad x_2 \mapsto c_{n+1}, \quad x_3 \mapsto a + c_{n+1} + \bar{c}_{n+1}, \quad x_i \mapsto c_{i,n+1}$$

defines an isomorphism $U_{n+1} \xrightarrow{\sim} \mathcal{C}_n$ over U_n .

Proof. (i) Eliminating the variables b, d_{ij} , as well as c_{3i} , we can rewrite (1.1.6) as

$$\begin{aligned} c_{ij} + c_{ji} &= a + c_i + \bar{c}_i + c_j + \bar{c}_j, \\ d + ac &= (a + c_i + \bar{c}_i)c_i\bar{c}_i - \bar{c}c_i - c\bar{c}_i, \\ c &= c_{ji}c_j + c_{ij}c_i - c_jc_i, \\ \bar{c} &= c_{ji}\bar{c}_i + c_{ij}\bar{c}_j - \bar{c}_j\bar{c}_i, \\ c_{ij}c_{jk} + c_{ji}c_{ik} - c_{ik}c_{jk} + (a + c_k + \bar{c}_k)c_k &= c. \end{aligned}$$

The last equation for a triple (i, j, k) together with the other equations imply the similar equation for any permutation of (i, j, k) . Thus, eliminating in addition c_{ji} for $j < i$ using the first set of equations above, we get the relations in (1.1.11) from the last three sets of equations.

It is easy to see that rewriting $(\bar{c}_i - \bar{c}_j)(c_j - c_i)c_{ij}$ in two ways using (1.1.11) we get the equation

$$(a + c_i + \bar{c}_i)c_i\bar{c}_i - \bar{c}c_i - c\bar{c}_i = (a + c_j + \bar{c}_j)c_j\bar{c}_j - \bar{c}c_j - c\bar{c}_j$$

for $i \neq j$. Hence, we can also eliminate d using the second set of equations above.

(ii) We have to compare the equations (1.1.3), (1.1.4) defining $\mathcal{C}_n$ with the equations (1.1.11) defining U_{n+1} . The case $n = 3$ is easy so let us assume $n \geq 4$. Then, the equation (1.1.4) follows from (1.1.3) for the pairs $(i, j) = (2, 4)$ and $(3, 4)$, Namely, the latter equations have form

$$x_2x_4 = x_2x_3 + \dots, \quad x_3x_4 = x_2x_3 + \dots$$

Multiplying the first equation with x_3 , the second—with x_2 , and subtracting, we get exactly (1.1.4). Next, using the relations (1.1.6) we can rewrite the equations (1.1.3) as

$$\begin{aligned} (1.1.13) \quad & x_2x_i = x_2x_3 + c_ix_i + \bar{c}_ix_2 - c, \quad 4 \leq i \\ & x_3x_i = x_2x_3 + (a + c_i + \bar{c}_i)(x_i - \bar{c}_i) + \bar{c} - c, \quad 4 \leq i \\ & x_ix_j = x_2x_3 + c_{ij}x_j + c_{ji}x_i - c, \quad 4 \leq i < j. \end{aligned}$$

It remains to observe that the equations (1.1.13) after the substitution (1.1.12) match the equations (1.1.11) with $j = n + 1$ in the first two equations and $k = n + 1$ in the third equation. $\square$

Note that there is a natural $\mathbb{G}_m$ -action on U_n that comes from rescaling the non-vanishing section ω of the dualizing sheaf on C . Correspondingly, the degree of the variables x_i are 1, the constants c_{ij} have degree 1, d_{ij} have degree 2, a has degree 1, b and c has degree 2, and d has degree 3. This action will play an important role in comparing this picture to the moduli of A_∞ -structures in Section 2.

Corollary 1.1.6. *Assume $n \geq 5$. For any field k the dimension of the Zariski tangent space of $U_n \times \text{Spec}(k)$ at the origin is equal to $(n-1)(n-2)/2$. The functions (of weight 1 with respect to the $\mathbb{G}_m$ -action)*

$$a, (c_i, \bar{c}_i)_{4 \leq i}, (c_{ij})_{4 \leq i < j}$$

form a minimal set of generators of the algebra of functions on U_n .

Proof. We have that the maximal ideal $\mathfrak{m}_0$ of the local ring at the origin is generated by $a, c, \bar{c}, c_i, \bar{c}_i, c_{ij}$. We see from the defining relations that c and $\bar{c}$ can be expressed in terms of the other generators. We get no additional linear dependences between the remaining generators in $\mathfrak{m}_0/\mathfrak{m}_0^2$. An easy count gives the result. $\square$

Corollary 1.1.7. *For $n \geq 5$ the graded ring $\mathcal{O}(U_n)$ is free over $\mathbb{Z}$ with the Hilbert series*

$$h_{U_n}(t) = \frac{1}{(1-t)^{n+1}} \cdot \prod_{k=3}^{n-3} (1 + kt + t^2).$$

Proof. Set $h_n = h_{U_n}$. Recall that we have a basis of $\mathcal{O}(\mathcal{C}_n)$ as an $\mathcal{O}(U_n)$ -module given by certain monomials in x_i . Since $\mathcal{C}_n = U_{n+1}$, this implies that

$$h_{n+1} = (1 + (n-1)t + nt^2 + nt^3 + \dots)h_n = \frac{1 + (n-2)t + t^2}{1-t} \cdot h_n.$$

Since $\mathcal{O}(U_5)$ is the ring of polynomials in 6 variables of degree 1, we have $h_5 = (1-t)^{-6}$, and the assertion follows by induction on n . $\square$

Sometimes the normalization (1.1.5) is not convenient. An alternative is to consider for $n \geq 5$ the affine space V_n (over $\mathbb{Z}$) with coordinates (c_{ij}) where $2 \leq i, j \leq n$, $i \neq j$, subject to the linear relations

$$c_{ij} + c_{ji} - c_{ik} - c_{ki} - c_{jk} - c_{kj} + c_{kl} + c_{lk} = 0$$

for any distinct i, j, k, l . We have a free action of $\mathbb{G}_a^{n-1}$ on V_n such that $(a_i)_{i \geq 2}$ acts by

$$c_{ij} \mapsto c_{ij} + a_i,$$

and the quotient $\overline{V}_n$ is still an affine space. Choosing $h_{1i} \in H^0(C, p_1 + p_i)$ with $\text{Res}_{p_1}(h_{1i}\omega) = 1$, and setting $c_{ij} = h_{1i}(p_j)$ gives a well defined point of $\overline{V}_n$ (since the only ambiguity is to add a constant to each h_{1i}). Note that this construction is compatible with the action of S_{n-1} permuting the points $p_2, \dots, p_n$ and indices $2, \dots, n$. Now Proposition 1.1.5 gives the following result.

Corollary 1.1.8. *The above construction gives a closed embedding $U_n \hookrightarrow \overline{V}_n$ for each $n \geq 5$, so that the projection $U_{n+1} \rightarrow U_n$ is compatible with the projection $\overline{V}_{n+1} \rightarrow \overline{V}_n$ omitting c_{ij} with $i = n+1$ or $j = n+1$.*

Example 1.1.9. Let us consider the wheel of $n \geq 3$ projective lines $C_1 \cup \dots \cup C_n$, where $1 \in C_i$ is glued to $0 \in C_{i+1}$ and $p_i = \infty \in C_i$. Let u_i denote the natural parameter on $\mathbb{A}^1 \subset C_i$. We have a global section ω of the dualizing sheaf on C given by

$$\omega|_{C_i} = \frac{du_i}{u_i(1-u_i)} = -d \ln(1-u_i^{-1}), \quad i = 1, \dots, n.$$

Then we have:

$$x_2 = h_{12} = \begin{cases} u_1 & \text{on } C_1, \\ 1 - u_2 & \text{on } C_2, \\ 0 & \text{on } C_i, \quad i > 2. \end{cases}$$

and for $j \geq 3$,

$$x_j = h_{1j} = \begin{cases} u_1 - 1 & \text{on } C_1, \\ 0 & \text{on } C_i, \quad 1 < i < j, \\ -u_j & \text{on } C_j, \\ -1 & \text{on } C_i, \quad i > j. \end{cases}$$

Now one easily checks that the defining relations of $C \setminus \{p_1, \dots, p_n\}$ become

$$(1.1.14) \quad \begin{aligned} x_2 x_j &= x_2 x_3, \quad 3 \leq j, \\ x_i x_j &= x_2 x_3 - x_j, \quad 3 \leq i < j, \\ x_2 x_3^2 &= x_2^2 x_3 - x_2 x_3. \end{aligned}$$

1.2. Case $n = 2$, $\text{char}(k) \neq 2$. In the $n = 2$ case we always assume that $\text{char}(k) \neq 2$.

Let t_1 be a formal parameter at p_1 such that $\text{Res}_{p_1} \omega/t_1 = 1$. Then there exists a function $f_1 \in H^0(C, \mathcal{O}_C(2p_1))$, such that at p_1 ,

$$f_1 \equiv \frac{1}{t_1^2} \pmod{k[[t_1]]}$$

(we use the fact that $\text{Res}_{p_1} f_1 \omega = 0$). Note that f_1 is unique up to adding a constant. Then the functions

$$1, f_1 h_{12}^r, h_{12}^m, \quad \text{where } 1 \leq m \leq N, 0 \leq r \leq N-2$$

form a basis of $H^0(C, \mathcal{O}_C(ND))$. Let us also normalize h_{12} so that at p_1

$$h_{12} \equiv \frac{1}{t_1} \pmod{t_1 k[[t_1]]}.$$

Then $f_1 - h_{12}^2$ is regular near p_1 . Now let us normalize f_1 so that

$$(f_1 - h_{12}^2)(p_1) = 0.$$

Then $f_1(f_1 - h_{12}^2) \in H^0(C, \mathcal{O}_C(p_1 + 2p_2))$, so we have

$$(1.2.1) \quad f_1^2 - f_1 h_{12}^2 = \alpha(f_1 - h_{12}^2) + \beta h_{12} + \gamma$$

for some constants α, β, γ .

The analog of Lemma 1.1.3 in this case states that for any choice of α, β, γ in a commutative ring R the R -algebra A generated by x and y subject to the defining relation

$$(1.2.2) \quad y^2 - yx^2 = \alpha(y - x^2) + \beta x + \gamma$$

has $(x^m, x^m y)$ as an R -basis. In fact, (1.2.2) is simply the unfolding of the tacnode singularity

$$(1.2.3) \quad y^2 - yx^2 = 0.$$

We extend Definition 1.1.4 by letting $U_2 = \mathbb{A}_{\mathbb{Z}}^3$ be the affine space generated by α, β , and γ . Note that we have a natural $\mathbb{G}_m$ -action such that $\deg(x) = 1$, $\deg(y) = \deg(\alpha) = 2$, $\deg(\beta) = 3$, $\deg(\gamma) = 4$.

Example 1.2.1. In the case when C is the wheel of $n = 2$ projective lines and 2 is invertible (see Example 1.1.9) we can use

$$(1.2.4) \quad h_{12} = \begin{cases} u_1 - \frac{1}{2} & \text{on } C_1, \\ \frac{1}{2} - u_2 & \text{on } C_2, \end{cases}$$

$$f_1 = \begin{cases} (u_1 - \frac{1}{2})^2 & \text{on } C_1, \\ \frac{1}{4} & \text{on } C_2. \end{cases}$$

Thus, the equation (1.2.1) in this case takes form

$$f_1^2 - f_1 h_{12}^2 = \frac{1}{4}(f_1 - h_{12}^2).$$

1.3. Case $n = 1$, $\mathbf{char}(k) \neq 2, 3$. The case $n = 1$ corresponds to the classical family of Weierstrass curves. We assume that $\mathbf{char}(k) \neq 2, 3$. Then we can choose a formal parameter t at $p = p_1$ such that

$$\omega \equiv dt \bmod t^4 k[[t]] \cdot dt,$$

where ω is a global section of the dualizing sheaf of C . The condition $h^1(C, \mathcal{O}_C(p)) = 0$ implies $h^0(C, \mathcal{O}_C(mp)) = m$ for $m \geq 1$. Hence, there exists a non-constant function (unique up to adding a constant) $x \in H^0(C, \mathcal{O}_C(2p))$ such that at p ,

$$x \equiv \frac{1}{t^2} \bmod k[[t]].$$

We can normalize it by adding a constant so that

$$x \equiv \frac{1}{t^2} \bmod t^2 k[[t]]$$

(the coefficient of t is zero since $\text{Res}_p(x^2 \omega) = \text{Res}_p(x^2 dt) = 0$). Similarly, since $h^0(C, \mathcal{O}_C(3p)) = 3$, there exists a unique function $y \in H^0(C, \mathcal{O}_C(3p))$ such that

$$y \equiv \frac{1}{t^3} \bmod tk[[t]].$$

Then $y^2 - x^3 \in H^0(C, \mathcal{O}_C(2p))$, so we have

$$(1.3.1) \quad y^2 - x^3 = \delta x + \epsilon$$

for some constants δ and ϵ . This also works in families over an affine base (see e.g., [22, Lem. 1.2.1]). Again, (1.3.1) (with δ and ϵ viewed as independent variables) is simply the unfolding of the cusp singularity

$$y^2 - x^3 = 0.$$

We extend Definition 1.1.4 to this case by letting $U_1 = \mathbb{A}_{\mathbb{Z}}^2$ be the affine space generated by δ and ϵ . Note that we have a natural $\mathbb{G}_m$ -action such that $\deg(x) = 2$, $\deg(y) = 3$, and $\deg(\delta) = 4$, $\deg(\epsilon) = 6$.

1.4. Moduli spaces. Let S be any scheme. Let $\mathcal{U}_{1,n}$ be the (non-separated) stack of flat, proper, finitely presented morphisms $\mathcal{C} \rightarrow S$ from an algebraic space $\mathcal{C}$ together with n sections, whose geometric fibers are reduced, connected curves of arithmetic genus 1. It is shown by Jack Hall in [25, appendix B] that $\mathcal{U}_{1,n}$ is an algebraic stack, locally of finite type over $\text{Spec } \mathbb{Z}$.

Definition 1.4.1. For $n \geq 1$, we define the moduli stack $\mathcal{U}_{1,n}^{sns}$ to be the open substack of $\mathcal{U}_{1,n}$ consisting of curves C of arithmetic genus 1 with n distinct smooth marked points $p_1, \dots, p_n$ such that

- (1) $h^0(\mathcal{O}_C(p_i)) = 1$ for all i and,
- (2) $\mathcal{O}_C(p_1 + \dots + p_n)$ is ample.

We also denote by $\tilde{\mathcal{U}}_{1,n}^{sns} \rightarrow \mathcal{U}_{1,n}^{sns}$ the $\mathbb{G}_m$ -torsor associated with a choice of a generator of the one-dimensional space $H^0(C, \omega_C)$, where ω_C is a dualizing sheaf.

Note that the condition (1) requires that p_i are non-special divisors, which is an open condition.

Theorem 1.4.2. For $n \geq 3$ the moduli stack $\tilde{\mathcal{U}}_{1,n}^{sns}$ is isomorphic to the affine scheme U_n over $\text{Spec}(\mathbb{Z})$ (see Def. 1.1.4), so that the open affine part of the universal curve $C \setminus D$, where $D = p_1 + \dots + p_n$, gets identified with the curve $\mathcal{C}_n \rightarrow U_n$ given by equations (1.1.3) (or equivalently, (1.1.13) for $n \geq 4$).

In the case $n = 2$ the moduli stack $\tilde{\mathcal{U}}_{1,n}^{sns}$ over $\text{Spec}(\mathbb{Z}[1/2])$ is isomorphic to the affine space $\mathbb{A}^3$ with coordinates α, β, γ , so that the affine universal curve $C \setminus D$ is given by (1.2.2).

In the case $n = 1$ the moduli stack $\tilde{\mathcal{U}}_{1,n}^{sns}$ over $\text{Spec}(\mathbb{Z}[1/6])$ is isomorphic to the affine space $\mathbb{A}^2$ with coordinates δ, ϵ so that the affine universal curve $C \setminus D$ is given by (1.3.1).

These isomorphisms are compatible with the $\mathbb{G}_m$ -actions described above.

Proof. This is similar to [22, Thm. 1.2.3]. We follow the line of argument given there. For simplicity let us assume that $n \geq 3$ —the cases $n = 1, 2$ can be analyzed similarly. Using the relative version (over an affine base $\text{Spec}(R)$) of the constructions that led to Lemma 1.1.3 and Proposition 1.1.5, we can associate with a family $\pi : C \rightarrow \text{Spec } R$ in $\tilde{\mathcal{U}}_{1,n}^{sns}$ an R -point of U_n . Thus, we obtain a functor $\tilde{\mathcal{U}}_{1,n}^{sns} \rightarrow U_n$.

Conversely, let $(a, c, \bar{c}, c_i, \bar{c}_i, c_{ij})$, where $4 \leq i < j$, represent an R -point of U_n . We consider the corresponding algebra A over R with generators $x_2, \dots, x_n$ and defining relations (1.1.13), or equivalently, (1.1.3), (1.1.4), where b, d, d_{ij} and c_{ij} for $i > j$ are determined from (1.1.6). Let $(F_m A)_{m \geq 0}$ be the increasing filtration on A associated with the generators $x_2, \dots, x_n$, so that $F_0 A = R \cdot 1$, $F_1 A / F_0 A = Rx_2 \oplus Rx_3 \oplus \dots \oplus Rx_n$ and for $m \geq 2$, $F_m A = (F_1 A)^m$. Lemma 1.1.3 implies that for $m \geq 2$, $F_m A / F_{m-1} A$ is freely generated over R by $x_2^{m-1} x_3$ and x_i^m for $2 \leq i \leq n$. Let

$$\mathcal{R}A := \bigoplus_{m \geq 0} F_m A$$

be the associated Rees algebra, and consider the corresponding projective scheme over $\text{Spec}(R)$,

$$C = \text{Proj}(\mathcal{R}A).$$

Let $T \in F_1 A$ be the element corresponding to $1 \in F_0 A \subset F_1 A$, and let $D = (T = 0)$ be the corresponding divisor in C . Then one has an isomorphism $C \setminus D \cong \text{Spec } A$ and the complementary closed set is given by

$$D \cong \text{Proj}(\bigoplus_{m \geq 0} F_m A / F_{m-1} A).$$

The graded algebra $\mathcal{R}A$ is the quotient of the polynomial ring $R[T, X_2, \dots, X_n]$ by the homogenization of the equations (1.1.3), (1.1.4),

$$(1.4.1) \quad X_i X_j - X_2 X_3 = c_{ij} T X_j + c_{ji} T X_i + d_{ij} T^2, \text{ for } i < j, (i, j) \neq (2, 3),$$

$$(1.4.2) \quad X_2 X_3^2 - X_2^2 X_3 = a T X_2 X_3 + b T^2 X_2 + c T^2 X_3 + d T^3.$$

We have n sections $p_i : \text{Spec } R \rightarrow D$ cut out by:

$$p_1 : T = 0, X_2 = X_3 = \dots = X_n,$$

$$p_i : T = 0, X_i \neq 0, X_2 = X_3 = \dots = X_{i-1} = X_{i+1} = \dots = X_n = 0 \text{ for } 2 \leq i \leq n.$$

Equations (1.4.1), (1.4.2) easily imply that

$$D = (T = 0) = \sqcup_{i=1}^n \text{im}(p_i).$$

Over a point $s \in \text{Spec } R$, the fibre C_s is given by the same equations over the residue field $k(s)$ of s . As $F_m A$ is a free R -module, we can easily compute the Hilbert function of C_s to be:

$$h_{C_s}(m) = \dim(F_m A \otimes_R k(s)) = mn \text{ for } m \geq 1$$

Hence, C_s is a reduced, connected, degree n curve in $\mathbb{P}_{k(s)}^{n-1}$ of arithmetic genus 1. We conclude that $\pi : C \rightarrow \text{Spec } R$ has relative dimension 1. Furthermore, $\mathcal{O}_{\mathbb{P}_R^{n-1}}(1)|_C = \mathcal{O}_C(1)$ is ample, so the divisor D is ample.

Note that as A is a free R -module, the morphism $\pi : C \setminus D \rightarrow \text{Spec } R$ is flat. For each $j \geq 2$ consider the distinguished open subset $V_j = \text{Spec } A_j \subset C$, where A_j is the degree 0 part of the localization $(\mathcal{R}A)_{X_j}$. Since $D \subset V_2 \cup \dots \cup V_n$, it suffices to check that each A_j is flat over R . We know that $(A_j)_{T/X_j}$ is flat over R since $\pi : C \setminus D \rightarrow \text{Spec } R$ is flat. On the other hand, $A_j/(\frac{T}{X_j}) \cong R \oplus R$ since $V_j \cap (T = 0)$ is the disjoint union of the section p_1 and p_j , and so $A_j/(\frac{T}{X_j})$ is also flat over R . Applying [22, Lem. 1.2.4] we conclude that A_j is flat over R .

Next, let us show that the projection $\pi : C \rightarrow \text{Spec } R$ is smooth near $p_1, \dots, p_n$. We can work with a geometric fiber of π , i.e., assume that R is an algebraically closed field. Let us show first the smoothness at p_j , where $j > 1$. The maximal ideal $\mathfrak{m}_{p_j}$ of the local ring at p_j is generated by $T/X_j, X_i/X_j$ for $i \neq j$. Suppose first that $j \geq 4$. Over the open set V_j we can write using (1.4.1),

$$\frac{X_i}{X_j} = \frac{X_2}{X_j} \frac{X_3}{X_j} + c_{ij} \frac{T}{X_j} + c_{ji} \frac{T}{X_j} \frac{X_i}{X_j} + d_{ij} \left(\frac{T}{X_j} \right)^2,$$

hence we have

$$\frac{X_i}{X_j} \equiv c_{ij} \frac{T}{X_j} \bmod \mathfrak{m}_{p_j}^2.$$

which implies that $\mathfrak{m}_{p_j}/\mathfrak{m}_{p_j}^2$ is generated by a single element, the image of T/X_j . Hence, C is smooth at p_j . For $j = 2$, the same argument works, except for the fraction X_3/X_2 : here we need to use that over V_2 one has

$$\frac{X_3}{X_2} = \left(\frac{X_3}{X_2}\right)^2 - a \frac{T}{X_2} \frac{X_3}{X_2} - b \left(\frac{T}{X_2}\right)^2 - c \left(\frac{T}{X_2}\right)^2 \frac{X_3}{X_2} - d \left(\frac{T}{X_2}\right)^3$$

(see (1.4.2)), hence $X_3/X_2 = 0 \in \mathfrak{m}_{p_2}^2$. A similar argument works for $j = 3$.

Now let us prove smoothness at p_1 . Note that p_1 lies in all of the open sets V_i for $i \geq 2$, so we can work on $V_2 \cap \dots \cap V_n$. The maximal ideal $\mathfrak{m}_{p_1}$ of the local ring at p_1 is generated by T/X_2 and $X_i/X_2 - 1$ for $3 \leq i \leq n$. For each $j \geq 4$ we have from (1.4.1)

$$\frac{X_j}{X_2} - \frac{X_3}{X_2} \equiv c_{2j} \frac{T}{X_2} \frac{X_j}{X_2} + c_{j2} \frac{T}{X_2} \bmod \mathfrak{m}_{p_1}^2 \equiv (c_{2j} + c_{j2}) \frac{T}{X_2} \bmod \mathfrak{m}_{p_1}^2.$$

On the other hand, dividing (1.4.2) by $X_2^2 X_3$ we get

$$\frac{X_3}{X_2} - 1 \equiv a \frac{T}{X_2} \bmod \mathfrak{m}_{p_1}^2.$$

Again, we conclude that T/X_2 generates $\mathfrak{m}_{p_1}/\mathfrak{m}_{p_1}^2$.

Next, we have to specify a choice of a global section of ω_C . We know that our family is of arithmetic genus 1. Thus, we can determine a global 1-form ω using the requirement that

$$\text{Res}_{p_1}(x_2 \omega) = 1$$

(where $x_2 = X_2/T$ is a function on C with simple poles at p_1 and p_2). The same reasoning also works in a family. Therefore, we conclude that $\pi : C \rightarrow \text{Spec } R$ defines an object of the moduli stack $\tilde{\mathcal{U}}_{1,n}^{sns}$.

Finally, observe that $H^0(C, \mathcal{O}(mD))$ can be identified with $F_m A$ inside the algebra A of functions on $C \setminus D$, by analyzing the polar conditions at the marked points $p_1, \dots, p_n$. Furthermore, we have

$$\text{Res}_{p_1}(x_i \omega) = 1 \text{ and } \text{Res}_{p_i}(x_i \omega) = -1 \text{ for } 2 \leq i \leq n,$$

as $x_i - x_2$ is regular at p_1 and $x_i = X_i/T$ has (simple) poles only at p_1 and p_i . Hence, it is clear that the functors that we constructed from U_n to $\tilde{\mathcal{U}}_{1,n}^{sns}$ and from $\tilde{\mathcal{U}}_{1,n}^{sns}$ to U_n are inverses of each other. $\square$

Corollary 1.4.3. *For $n \geq 3$ the scheme $\overline{\mathcal{U}}_{1,n}^{sns}$ is flat over $\mathbb{Z}$.*

Proof. This follows from the identification $\overline{\mathcal{U}}_{1,n}^{sns} \simeq U_n$, Corollary 1.1.7 (for the case $n \geq 5$), and isomorphisms $U_3 \simeq \mathbb{A}_{\mathbb{Z}}^4$, $U_4 \simeq \mathbb{A}_{\mathbb{Z}}^5$. $\square$

1.5. Comparison with Smyth's moduli spaces. For each $1 \leq m < n$ Smyth defined in [26] the notion of m -stability for n -pointed curves of arithmetic genus 1 and showed that the corresponding moduli stack $\overline{\mathcal{M}}_{1,n}(m)$ is an irreducible projective Deligne-Mumford stack over $\text{Spec } \mathbb{Z}[1/6]$. Below we recall the definition.

Let us denote by $C_{1,n}$ the singular curve corresponding to the point in the moduli space $\widetilde{\mathcal{U}}_{1,n}^{sns} \simeq U_n$ where all coefficients are zero. Thus, the curve $C_{1,n}$ has n smooth points at infinity $p_1, \dots, p_n$ such that $C_{1,n} \setminus \{p_1, \dots, p_n\}$ is given by the equations

$$x_i x_j = x_2 x_3$$

for $n \geq 4$ (where the indices $i < j$ vary in $[2, n]$), by the equation

$$x_2 x_3^2 = x_2^2 x_3$$

for $n = 3$, by the tacnode equation (1.2.3):

$$y^2 - yx^2 = 0$$

for $n = 2$, and finally for $n = 1$ we get the ordinary cusp

$$y^2 - x^3 = 0.$$

Alternatively, we can describe $C_{1,n}$ as the union of n generic lines passing through one point in the projective space $\mathbb{P}^{n-1}$. The arising singularity at this point is called the *elliptic n -fold point*, which is a Gorenstein singularity (see [26, Prop. 2.5]). We also refer to $C_{1,n}$ as the *elliptic n -fold curve*.

Next, let us recall (see [26, Lem. 3.1]) that each Gorenstein curve C of arithmetic genus 1 has the so-called *fundamental decomposition*

$$(1.5.1) \quad C = E \cup R_1 \cup \dots \cup R_k$$

where E , called the *minimal elliptic subcurve* of C , is a connected subcurve of arithmetic genus 1 with no disconnecting nodes, R_i are nodal curves of arithmetic genus 0, $R_i \cap R_j = \emptyset$ for $i \neq j$, and $R_i \cap E$ is a single point which is a node of C .

The notion of m -stability for an n -pointed curve of arithmetic genus 1 consists of the following three conditions:

Definition 1.5.1. A curve $(C, p_1, \dots, p_n)$ of arithmetic genus 1 (where all marked points are smooth and distinct) is said to be *m -stable* if

- (1) The curve C has only nodes and elliptic l -fold points, $l \leq m$, as singularities;
- (2) If $E \subset C$ is the minimal elliptic subcurve then

$$|E \cap \overline{C \setminus E}| + |E \cap \{p_1, \dots, p_n\}| > m;$$

- (3) One has $H^0(C, \mathcal{T}_C(-p_1 - \dots - p_n)) = 0$.

Note that in the original definition the condition (2) is required to hold for any connected subcurve of arithmetic genus 1. The fact that it is enough to require this condition for the minimal elliptic subcurve follows from [26, Lem. 3.5].

In [26, Lem. 3.10], Smyth shows that m -stability is a deformation-open condition, hence one can define $\overline{\mathcal{M}}_{1,n}(m)$ as an open substack of the stack $\mathcal{U}_{1,n}$ of n -pointed curves of arithmetic genus 1. In [26, Th. 3.18], Smyth verifies that $\overline{\mathcal{M}}_{1,n}(m)$ is an irreducible proper Deligne-Mumford stack over $\text{Spec } \mathbb{Z}[1/6]$.

The requirement for 6 to be invertible is caused by the pathology that the cuspidal curve $C_{\text{cusp}} = C_{1,1}$ has extra vector fields in characteristic 2, 3, while the tacnode $C_{\text{tn}} = C_{1,2}$ has extra vector fields in characteristic 2, which can lead to non-trivial infinitesimal automorphisms of $(C, p_1, \dots, p_n)$. Let us explain this phenomenon in more detail. Recall that the affine parts of C_{cusp} and C_{tn} are the plane curves given by

$$\begin{aligned} C_{\text{cusp}} \setminus \{p\} : y^2 &= x^3 \\ C_{\text{tn}} \setminus \{p_1, p_2\} : y^2 &= xy^2 \end{aligned}$$

It is easy to see that in $\mathbf{char}(k) \neq 2$ we have $\text{rk} H^0(C_{\text{tn}}, \mathcal{T}_{C_{\text{tn}}}) = 3$, while in $\mathbf{char}(k) \neq 2, 3$ we have $\text{rk} H^0(C_{\text{cusp}}, \mathcal{T}_{C_{\text{cusp}}}) = 2$ (see [26, Prop. 2.3]). In Lemma 1.5.2 below we write out explicitly the extra vector fields in $\text{char}(k) = 2, 3$ that prevent condition (3) of m -stability to hold for these curves unless there are sufficiently many marked points on them.

Let C be either the cusp or the tacnode. Let $\nu : \tilde{C} \rightarrow C$ be the normalization map. One has $\tilde{C} = \mathbb{P}^1$ for the cusp curve, and it is the disjoint union of two $\mathbb{P}^1$'s for the tacnode. Any vector field on C can be restricted to $C \setminus \text{Sing}(C) \cong \tilde{C} \setminus \nu^{-1}(\text{Sing}(C))$ and then extended to a rational vector field on $\tilde{C}$. This leads to a natural inclusion map:

$$\mathcal{T}_C \hookrightarrow \nu_* \mathcal{T}_{\tilde{C}} \otimes K(\tilde{C})$$

Lemma 1.5.2. *Consider the normalization map $\nu : \tilde{C} \rightarrow C_{\text{cusp}}$ defined in affine coordinates by $k[x, y]/(y^2 - x^3) \rightarrow k[t^2, t^3] \subset k[t]$ sending $(x, y) \rightarrow (t^2, t^3)$.*

If $\text{char}(k) = 2$, the natural inclusion $\mathcal{T}_{C_{\text{cusp}}} \rightarrow \nu_ \mathcal{T}_{\tilde{C}} \otimes K(\tilde{C})$ is generated by the vector fields*

$$t^2 \partial_t, t \partial_t, \partial_t, \frac{1}{t^2} \partial_t$$

which are images of the sections $x^2 \partial_y, y \partial_y, x \partial_y, \partial_y$ of $\mathcal{T}_{C_{\text{cusp}}}$.

In particular, there is no non-zero vector field on C_{cusp} which vanishes on 5 distinct points.

If $\text{char}(k) = 3$, the natural inclusion $\mathcal{T}_{C_{\text{cusp}}} \rightarrow \nu_ \mathcal{T}_{\tilde{C}} \otimes K(\tilde{C})$ is generated by the vector fields*

$$t^2 \partial_t, t \partial_t, \partial_t, \frac{1}{t} \partial_t$$

which are images of the sections $x^2 \partial_x, x \partial_x, y \partial_x, \partial_x$ of $\mathcal{T}_{C_{\text{cusp}}}$.

In particular, there is no non-zero vector field on $\mathcal{T}_{C_{\text{cusp}}}$ which vanishes on 4 distinct points.

Consider the normalization map $\nu : \tilde{C} \rightarrow C_{tn}$ defined in affine coordinates by $k[x, y]/(y^2 - yx^2) \rightarrow k[t] \oplus k[s]$ sending $x \rightarrow (t, s)$ and $y \rightarrow (t^2, 0)$.

If $\text{char}(k) = 2$, the natural inclusion $\mathcal{T}_{C_{tn}} \rightarrow \nu_* \mathcal{T}_{\tilde{C}} \otimes K(\tilde{C})$ is generated by vector fields

$$t^2 \partial_t, s^2 \partial_s, t \partial_t + s \partial_s, \partial_t + \partial_s$$

which are images of the sections $y \partial_x, (x^2 - y) \partial_x, x \partial_x, \partial_x$ of $\mathcal{T}_{C_{tn}}$.

In particular, there is no non-zero vector field on $\mathcal{T}_{C_{tn}}$ which vanishes on 5 distinct points.

Proof. This is an extension of [26, Prop. 2.3] to the case of $\text{char}(k) = 2, 3$ and follows from a similar calculation as given there. $\square$

Therefore, to include characteristics 2 (resp., 3) one can either throw away curves which have cusps or tacnodes with fewer than 5 (resp., 4) marked points which in general may result in losing properness, or to relax the condition (3) which would lead to an algebraic stack which is not a Deligne-Mumford stack. In our current study, we will only be concerned with the moduli stacks $\overline{\mathcal{M}}_{1,n}(n-1)$, and we choose the option of relaxing the condition (3). Thus, we propose the following version of these stacks over $\mathbb{Z}$.

Definition 1.5.3. The stack $\overline{\mathcal{M}}_{1,n}^\infty$ is the moduli stack of (reduced, connected projective) pointed curves $(C, p_1, \dots, p_n)$ of arithmetic genus 1 (where all marked points are smooth and distinct) such that

- (1)' The curve C has only nodes and elliptic l -fold points, $l < n$, as singularities;
- (2)' C has no disconnecting nodes (i.e., it coincides with its minimal elliptic subcurve);
- (3)' Every irreducible component of C contains at least one marked point.

We will see in Theorem 1.5.7 below that $\overline{\mathcal{M}}_{1,n}^\infty$ for $n \geq 3$ (resp., $\overline{\mathcal{M}}_{1,2}^\infty$) is proper over $\text{Spec}(\mathbb{Z})$ (resp., $\text{Spec}(\mathbb{Z}[1/2])$) and that in fact it is a projective scheme for $n \geq 5$. The following result compares it to Smyth's moduli stack $\overline{\mathcal{M}}_{1,n}(n-1)$ over $\text{Spec}(\mathbb{Z}[1/6])$.

Proposition 1.5.4. *Let $(C, p_1, \dots, p_n)$ be a reduced, connected projective curve of arithmetic genus 1 with smooth distinct marked points, over an algebraically closed field k . If $(C, p_1, \dots, p_n)$ is $(n-1)$ -stable then conditions (1)'–(3)' are satisfied. The converse is true under one of the following additional assumptions:*

- (a) $\text{char}(k) \neq 2, 3$;
- (b) $\text{char}(k) \neq 2$ and $n \geq 4$;
- (c) $n \geq 5$.

Proof. It is easy to see that for conditions (2) with $m = n - 1$ and (3) to be satisfied C has to coincide with its minimal elliptic subcurve. By [26, Lem. 3.3], this implies that we have the following possibilities for C :

- (i) a smooth elliptic curve;
- (ii) an irreducible rational nodal curve;
- (iii) a wheel of $\mathbb{P}^1$'s;
- (iv) C has an elliptic m -fold point p and the normalization of C at p consists of m distinct smooth rational curves $C_1, \dots, C_m$.

In each of these cases one can easily see that condition (3) implies (3)'. Conversely, if (1)'–(3)' are satisfied then to check (3) we use the above classification of minimal subcurves, together with [26, Cor. 2.4] (note that the argument of [26, Cor. 2.4] works also in the case $n = 2$, $\text{char}(k) = 3$) and Lemma 1.5.2. $\square$

Corollary 1.5.5. *There is a natural isomorphism $\overline{\mathcal{M}}_{1,n}(n-1) \simeq \overline{\mathcal{M}}_{1,n}^\infty \times \text{Spec}(\mathbb{Z}[1/6])$.*

The following observation will also be useful later.

Lemma 1.5.6. *For every curve C corresponding to a point of $\overline{\mathcal{M}}_{1,n}^\infty$, the dualizing sheaf ω_C is isomorphic to $\mathcal{O}_C$.*

Proof. This follows from the fact that C coincides with its minimal elliptic subcurve and from [26, Lem. 3.3]. $\square$

On the other hand, we can consider the GIT stability for the action of $\mathbb{G}_m$ on $\tilde{\mathcal{U}}_{1,n}^{sns}$. Since the degrees of all the coordinates $c_{ij}, d_{ij}, a, b, c, d$ on $\tilde{\mathcal{U}}_{1,n}^{sns}$ are positive, the GIT-semistable points (with respect to the identity character of $\mathbb{G}_m$) are exactly the points where not all coordinates vanish. Hence, this is precisely the open subscheme

$$\tilde{\mathcal{U}}_{1,n}^{sns} \setminus \{C_{1,n}\} \subset \tilde{\mathcal{U}}_{1,n}^{sns}$$

Theorem 1.5.7. *Let us work over $\text{Spec}(\mathbb{Z})$ for $n \geq 3$, over $\text{Spec}(\mathbb{Z}[1/2])$ for $n = 2$, and over $\text{Spec}(\mathbb{Z}[1/6])$ for $n = 1$. One has an isomorphism of stacks*

$$\overline{\mathcal{M}}_{1,n}^\infty \simeq (\tilde{\mathcal{U}}_{1,n}^{sns} \setminus \{C_{1,n}\})/\mathbb{G}_m$$

compatible with the inclusion into the stack $\mathcal{U}_{1,n}$ of all n -pointed curves of arithmetic genus 1. Hence, we have isomorphisms

$$\begin{aligned} \overline{\mathcal{M}}_{1,1}^\infty &\simeq \mathbb{P}(4, 6) \text{ over } \text{Spec}(\mathbb{Z}[1/6]), \\ \overline{\mathcal{M}}_{1,2}^\infty &\simeq \mathbb{P}(2, 3, 4) \text{ over } \text{Spec}(\mathbb{Z}[1/2]), \\ \overline{\mathcal{M}}_{1,3}^\infty &\simeq \mathbb{P}(1, 2, 2, 3), \\ \overline{\mathcal{M}}_{1,4}^\infty &\simeq \mathbb{P}(1, 1, 1, 2, 2), \\ \overline{\mathcal{M}}_{1,5}^\infty &\simeq \mathbb{P}^5, \end{aligned}$$

where $\mathbb{P}(d_1, \dots, d_k)$ denotes the weighted projective stack.

For $n \geq 6$, the stack $\overline{\mathcal{M}}_{1,n}^\infty$ is isomorphic to the n -dimensional irreducible projective variety in $\mathbb{P}_{\mathbb{Z}}^{n(n-3)/2}$ given by the equations obtained from (1.1.11) by eliminating variables c and $\bar{c}$. The corresponding line bundle $\mathcal{O}(1)$ is isomorphic to the standard tautological bundle λ . For every field k , the variety $\overline{\mathcal{M}}_{1,n}^\infty \times \text{Spec}(k)$ is still irreducible.

Remark 1.5.8. The identifications of $\overline{\mathcal{M}}_{1,n}^\infty$ for $n \leq 4$ show that it is not a Deligne-Mumford stack over $\text{Spec}(\mathbb{Z})$ (non-étale automorphism groups occur in characteristics 2 and 3). However, it is a “tame stack” in the sense of [1], which seems to be a better notion in positive and mixed characteristics.

Proof. First, we are going to check that any point $[(C, p_1, \dots, p_n)] \in \tilde{\mathcal{U}}_{1,n}^{sns}$, different from $C_{1,n}$, satisfies conditions (1)’–(3)’. Note that (3)’ holds by definition.

To see that C is Gorenstein, we need to check that the dualizing sheaf ω_C is locally free. As was mentioned above, we know that $C_{1,n}$ is Gorenstein by [26, Prop. 2.5], i.e. $\omega_{C_{1,n}}$ is locally free. It follows that any curve C in a neighborhood of $C_{1,n}$ is Gorenstein. Finally, note that the $\mathbb{G}_m$ -action brings an arbitrary point in $\tilde{\mathcal{U}}_{1,n}^{sns}$ to an isomorphic curve in a neighborhood of $C_{1,n}$.

Next, using the fact that $h^0(C, \mathcal{O}(p_i)) = 1$, we check condition (2)’. Indeed, each subcurve R_i in the fundamental decomposition (1.5.1) should have at least one marked point p_j since $\mathcal{O}(p_1 + \dots + p_n)$ is ample. But then for such a point we necessarily have $h^0(C, \mathcal{O}(p_j)) \geq h^0(R_i, \mathcal{O}(p_j)) = 2$, since the arithmetic genus of R_i is zero. Hence, C coincides with its minimal elliptic subcurve.

It follows that C can be either a smooth elliptic curve, an irreducible rational nodal curve, a wheel of $\mathbb{P}^1$ ’s, or a curve with a unique elliptic m -fold point for $m < n$ (see [26, Lem. 3.3]). This immediately gives (1)’.

Conversely, if conditions (1)’–(3)’ are satisfied then we can easily check that $h^0(C, \mathcal{O}(p_i)) = 1$ for each marked point p_i . Thus, we see that $\overline{\mathcal{M}}_{1,n}^\infty$ is an open substack in $\mathcal{U}_{1,n}^{sns}$ obtained by throwing away the elliptic n -fold point $C_{1,n}$.

The explicit identification of $\overline{\mathcal{M}}_{1,n}^\infty$ for $n \leq 5$ follows from the identification of U_n given in Proposition 1.1.5 for $n = 3, 4, 5$ and $n = 1, 2$ follows from the identification given by (1.3.1) and (1.2.2) (see Theorem 1.4.2).

The irreducibility of $\overline{\mathcal{M}}_{1,n}^\infty$ (resp., $\overline{\mathcal{M}}_{1,n}^\infty \times \text{Spec}(k)$) for $n \geq 6$ can be proved by the same method as in [26] (using the fact that the relevant curves are smoothable).

The identification $\mathcal{O}(1) \simeq \lambda$ follows from the fact that the pull-backs of both line bundles to $\tilde{\mathcal{U}}_{1,n}^{sns} \setminus \{C_{1,n}\}$ have natural trivializations, and the corresponding $\mathbb{G}_m$ -actions on the trivial line bundle are both given by the identity character $\mathbb{G}_m \rightarrow \mathbb{G}_m$. $\square$

In Proposition 1.7.1 below we will also prove an isomorphism

$$\overline{\mathcal{M}}_{1,6}^\infty \simeq Gr(2, 5),$$

where $Gr(2, 5)$ denotes the Grassmannian of 2-dimensional subspaces of a vector space of dimension 5.

Corollary 1.5.9. *For $n \geq 5$ the natural action of S_n on $\overline{\mathcal{M}}_{1,n}^\infty$ is induced by some automorphisms of the projective space $\mathbb{P}^{n(n-3)/2}$.*

Proof. Indeed, this follows from the fact that the projective embedding $\overline{\mathcal{M}}_{1,n}^\infty \hookrightarrow \mathbb{P}^{n(n-3)/2}$ is given by the S_n -equivariant line bundle λ . $\square$

Remark 1.5.10. In terms of the isomorphism $\overline{\mathcal{M}}_{1,n}^\infty \simeq (U_n \setminus \{0\})/\mathbb{G}_m$ (which is obtained by combining Theorems 1.4.2 and 1.5.7), the action of generators of S_n can be easily described. Namely, the subgroup of permutations fixing 1, 2, 3 acts by natural permutations of the coordinates $c_i, \bar{c}_i, c_{ij}$ (see Proposition 1.1.5). The transpositions (13), (23) and (24) act by the following involutions:

$$\begin{aligned} (13) : a &\mapsto a, \quad c \leftrightarrow \bar{c}, \quad c_i \leftrightarrow \bar{c}_i, \quad c_{ij} \mapsto c_{ji} = a + c_i + \bar{c}_i + c_j + \bar{c}_j - c_{ij}, \\ (23) : a &\mapsto -a, \quad c \mapsto c - \bar{c}, \quad \bar{c} \mapsto -\bar{c}, \quad c_i \mapsto a + c_i + \bar{c}_i, \quad \bar{c}_i \mapsto -\bar{c}_i, \quad c_{ij} \mapsto c_{ij} - \bar{c}_i, \\ (24) : a &\mapsto -a - 2c_4 - 2\bar{c}_4, \quad c \mapsto \bar{c} - \bar{c}_4(a + c_4 + \bar{c}_4), \quad \bar{c} \mapsto c - c_4(a + c_4 + \bar{c}_4), \\ &c_i \leftrightarrow c_{4i}, \quad \bar{c}_i \leftrightarrow \bar{c}_{4i}, \quad c_{ij} \mapsto c_{ij}, \quad \text{where } i, j \geq 5. \end{aligned}$$

These formulas can be checked using the equations of the relative curve (1.1.13). A more transparent way to see the action of the subgroup S_{n-1} fixing 1, is via the identification of the affine space containing U_n with the quotient $\bar{V}_n$ considered in Corollary 1.1.8.

We can now reprove the result of Smyth that the moduli stack $\overline{\mathcal{M}}_{1,n}(m)$ is smooth if and only if $m \leq 5$, and get some additional information on the singularities of $\overline{\mathcal{M}}_{1,n}(m)$ for $m > 5$.

First, we need a bit of deformation theory. Let us fix a field k , and for each $m \geq 1$ let us consider the deformation functor Def_m of the m -fold elliptic singularity over k (defined on local Artin algebras with the residue field k). We denote by B_m the base of the formal miniversal deformation of the elliptic m -fold singularity. Thus, B_m is the formal spectrum of R_m , a complete local Noetherian ring with the residue field k , and we have a formally étale morphism of functors

$$h_{R_m} \rightarrow \text{Def}_m$$

where h_{R_m} is the representable functor corresponding to R_m .

Let $(C, p_\bullet, v_\bullet)$ be a point of $U_n = \tilde{\mathcal{U}}_{1,n}^{sns}$, and let $\hat{\mathcal{O}}$ be the completion of its local ring. Let also $\hat{U}_n$ be the formal completion of U_n at this point, i.e., the formal spectrum of $\hat{\mathcal{O}}$. Let $q_1, \dots, q_r$ be all the non-nodal singular points of C , where q_i is the elliptic m_i -fold singularity. By [27, Lem. 2.1], the natural projection

$$(1.5.2) \quad p : h_{\hat{\mathcal{O}}} \rightarrow \prod_{i=1}^r \text{Def}_{m_i}$$

is formally smooth (we use the fact that the deformations of nodal singularities are unobstructed).

Lemma 1.5.11. *The morphism (1.5.2) factors through a formally smooth morphism $\tilde{p} : h_{\hat{\mathcal{O}}} \rightarrow h_R$, where*

$$v : h_R \rightarrow \prod_{i=1}^r \text{Def}_{m_i},$$

is the miniversal deformation with $R = \hat{\otimes} R_{m_i}$. Thus, $\tilde{p}$ corresponds to a formally smooth morphism

$$\hat{U}_n \rightarrow \prod_{i=1}^r B_{m_i}.$$

Proof. By definition, the morphism v is formally étale. In particular, it is formally smooth, so we can lift (non-uniquely) the projection p to a local homomorphism $R \rightarrow \hat{\mathcal{O}}$, so that we have a commutative triangle

$$(1.5.3) \quad \begin{array}{ccc} h_{\hat{\mathcal{O}}} & & \\ \tilde{p} \downarrow & \searrow p & \\ h_R & \xrightarrow{v} & \prod_{i=1}^r \text{Def}_{m_i} \end{array}$$

It then follows that the morphism $\tilde{p}$ is formally smooth. Indeed, since v is étale, and p is smooth, the morphism $\tilde{p}$ induces a surjection on tangent spaces, hence, we can apply [22, Lem. 4.5.3]. $\square$

The analog of Lemma 1.5.11 also holds with U_n replaced by $\overline{\mathcal{M}}_{1,n}(m)$. This implies that $\overline{\mathcal{M}}_{1,n}(m)$ is smooth if and only if $B_{m'}$ is smooth for $m' \leq m$. Since we have a formally smooth morphism from the completion of U_m at zero to B_m , this is equivalent to smoothness of $U_{m'}$ at zero for $m' \leq m$. Now we recall that U_m is smooth for $m \leq 5$ (see Proposition 1.1.5). On the other hand, U_6 is not smooth at zero, since it is 7-dimensional but the dimension of the tangent space is 10 (see Corollary 1.1.6). Thus, we recover the fact that $\overline{\mathcal{M}}_{1,n}(m)$ is smooth if and only if $m \leq 5$ (see [27, Cor. 4.17]).

In a similar vein we have the following results. In the rest of this subsection we work over an algebraically closed field.

Proposition 1.5.12. *The moduli stacks $\overline{\mathcal{M}}_{1,n}^\infty$ and the schemes U_n are smooth in codimension ≤ 6 .*

Proof. Use the stratification by the singularity type (see [27, Cor. 2.4]) and the fact that B_m is smooth for $m \leq 5$. $\square$

Proposition 1.5.13. *For $n \geq 3$ the scheme $U_n \simeq \tilde{U}_{1,n}^{sns}$ is Gorenstein and regular in codimension 1, hence, normal.*

Proof. Regularity in codimension 1 follows from Proposition 1.5.12. To prove the Gorenstein property we use the induction on n . We have seen in Proposition 1.1.5 that U_n is smooth for $n \leq 5$. Assume the assertion is true for U_n . By Theorem 1.4.2 and by Proposition 1.1.5(ii), we know that U_{n+1} is open in the universal curve over $U_n = \tilde{U}_{1,n}^{sns}$. So the morphism $U_{n+1} \rightarrow U_n$ is flat, with Gorenstein fibers (see the proof of Theorem 1.5.7) and Gorenstein base. Hence, U_{n+1} is Gorenstein too (see [12, Ch. V, Prop. 9.6]). $\square$

Corollary 1.5.14. *For $n \geq 5$ the projective scheme $\overline{\mathcal{M}}_{1,n}^\infty$ is normal and arithmetically Gorenstein, hence Gorenstein.*

Proof. Indeed, the homogeneous coordinate ring of $\overline{\mathcal{M}}_{1,n}^\infty$ is exactly the ring of functions on U_n . $\square$

Corollary 1.5.15. *The stacks $\overline{\mathcal{M}}_{1,n}(m)$ are normal and Gorenstein for $1 \leq m < n$.*

Proof. Recall that the Gorenstein property of a local Noetherian ring can be checked after passing to its completion (see [19, Thm.18.3]). Also, if $R \rightarrow S$ is a local formally smooth homomorphism of complete Noetherian rings, then R is Gorenstein if and only if S is Gorenstein (this follows from the results of [3]). Thus, by Lemma 1.5.11, to check the Gorenstein property it suffices to check that the base B_m of formal miniversal deformation of the elliptic m -fold point is Gorenstein. But we have a formally smooth morphism from the completion of U_m at zero to B_m (by the same Lemma), and U_m is Gorenstein by Proposition 1.5.13. Hence, B_m is also Gorenstein. On the other hand, as in Proposition 1.5.12, we see that stacks $\overline{\mathcal{M}}_{1,n}(m)$ are regular in codimension 1. Hence, we conclude that they are normal. $\square$

Remark 1.5.16. Over an algebraically closed field of characteristic zero, one has $\text{Pic}(\overline{\mathcal{M}}_{1,n}^\infty)_{\mathbb{Q}} = \mathbb{Q}$ (see [27, Prop. 3.2]) and the class of the canonical bundle of $\overline{\mathcal{M}}_{1,n}^\infty$ in $\text{Pic}(\overline{\mathcal{M}}_{1,n}^\infty)_{\mathbb{Q}}$ is $(n-11)\lambda$ (for $n \geq 5$ this follows from the description of $\overline{\mathcal{M}}_{1,n}^\infty$ as an explicit birational contraction of $\overline{M}_{1,n}$ in [27]). Below we will prove that in fact, $\text{Pic}(\overline{\mathcal{M}}_{1,n}^\infty) \simeq \mathbb{Z}$ and is generated by $\mathcal{O}(1) = \lambda$. We will also give an independent derivation of the formula for the canonical class (see 1.6.3).

1.6. Rational map from $\overline{\mathcal{M}}_{1,n}^\infty$ to $\overline{\mathcal{M}}_{1,n-1}^\infty$ and its applications. For each $1 \leq i < j \leq n$ let us consider the point $P_{ij} = P_{ij}(n) \in \overline{\mathcal{M}}_{1,n}^\infty$ corresponding to the elliptic $(n-1)$ -fold curve with the marked points p_i and p_j on the same component and exactly one marked point on each other component.

Let

$$\pi_n : \overline{\mathcal{M}}_{1,n}^\infty \dashrightarrow \overline{\mathcal{M}}_{1,n-1}^\infty$$

be the rational map corresponding to omitting the marked point p_n . Note that it is compatible via the isomorphism of Theorem 1.5.7 with the linear projection $U_n \rightarrow U_{n-1}$

omitting the coordinates c_n , $\bar{c}_n$ and c_{in} , $i = 4, \dots, n-1$ (here and below we use the coordinates introduced in Proposition 1.1.5(i)). Therefore, in terms of the natural embeddings $\overline{\mathcal{M}}_{1,n}^\infty \subset \mathbb{P}^{n(n-3)/2}$, $\overline{\mathcal{M}}_{1,n-1}^\infty \subset \mathbb{P}^{(n-1)(n-4)/2}$, the rational map π_n is induced by the linear projection

$$(1.6.1) \quad \mathbb{P}^{n(n-3)/2} \dashrightarrow \mathbb{P}^{(n-1)(n-4)/2}$$

along the $n-3$ -dimensional projective subspace $\mathbb{P}(K_n) \subset \mathbb{P}^{n(n-3)/2}$ where the linear subspace K_n is given by the equations $c_{ij} = 0$ for $4 \leq i < j \leq n-1$, $c_i = \bar{c}_i = a = 0$ for $4 \leq i \leq n-1$.

Proposition 1.6.1. *Assume that $n \geq 6$.*

(i) *The intersection $\overline{\mathcal{M}}_{1,n}^\infty \cap \mathbb{P}(K_n)$ is transversal and consists of the points $P_{in} = P_{in}(n)$, $i = 1, \dots, n-1$. More precisely, the homogeneous coordinates c_n , $\bar{c}_n$, c_{in} ($4 \leq i < n$) at these points are:*

$$\begin{aligned} P_{1n} : & \bar{c}_n = 0, c_{in} = c_n \neq 0; \\ P_{2n} : & \bar{c}_n = -c_n \neq 0, c_{in} = 0; \\ P_{3n} : & \bar{c}_n \neq 0, c_n = c_{in} = 0; \\ P_{in}, i \geq 4 : & c_n = \bar{c}_n = 0, c_{in} \neq 0, c_{jn} = 0 \text{ for } j \neq i. \end{aligned}$$

(i') *In terms of coordinates (c_{ij}) , where $2 \leq i, j \leq n$, $i \neq j$ (see Corollary 1.1.8), the point P_{i_0} , where $2 \leq i_0 \leq n$ is determined by the equations*

$$(1.6.2) \quad \begin{aligned} c_{ij} &= c_{ik}, \\ c_{ii_0} - c_{ik} &= c_{ji_0} - c_{jk}, \end{aligned}$$

where the indices i, j, k are distinct and different from i_0 .

(ii) *The rational map π_n is resolved by a diagram of regular maps*

$$(1.6.3) \quad \begin{array}{ccc} & \bar{\mathcal{C}}_{n-1} & \\ r_n \swarrow & & \searrow q_{n-1} \\ \overline{\mathcal{M}}_{1,n}^\infty & & \overline{\mathcal{M}}_{1,n-1}^\infty \end{array}$$

where $q_{n-1} : \bar{\mathcal{C}}_{n-1} \rightarrow \overline{\mathcal{M}}_{1,n-1}^\infty$ is the universal curve. Via the projection r_n , $\bar{\mathcal{C}}_{n-1}$ gets identified with the blow-up of $\overline{\mathcal{M}}_{1,n}^\infty$ at the $n-1$ points (P_{in}) , $i = 1, \dots, n-1$. The image of the canonical section $\sigma_i : \overline{\mathcal{M}}_{1,n-1}^\infty \rightarrow \bar{\mathcal{C}}_{n-1}$ (where $i = 1, \dots, n-1$) coincides with the exceptional divisor $E_i = r_n^{-1}(P_{in})$.

(iii) *One has $\pi_n(P_{ij}(n)) = P_{ij}(n-1)$ for $j \leq n-1$. Furthermore, $P_{ij}(n)$ is the singular point of the elliptic $(n-2)$ -fold curve $\pi_n^{-1}(P_{ij}(n-1))$. Let $C_{ij} \subset \pi_n^{-1}(P_{ij}(n-1))$ be the component containing p_i and p_j , and let $C_k \subset \pi_n^{-1}(P_{ij}(n-1))$ be the component containing p_k , where $k \neq i, j$. Then $r_n(C_k)$ is the line connecting $P_{kn}(n)$ and $P_{ij}(n)$, while $r_n(C_{ij})$ is a conic containing $P_{ij}(n)$, $P_{in}(n)$ and $P_{jn}(n)$.*

Proof. (i) Proposition 1.1.5(ii) shows that with respect to the homogeneous coordinates $y_2, \dots, y_{n-1}$ on the projective space $\mathbb{P}(K_n)$ given by

$$y_2 = c_n, \quad y_3 = c_n + \bar{c}_n, \quad y_i = c_{in} \text{ for } 4 \leq i < n,$$

the scheme-theoretic intersection $\overline{\mathcal{M}}_{1,n}^\infty \cap \mathbb{P}(K_n) \subset \mathbb{P}(K_n)$ is defined by the equations $y_i y_j = y_2 y_3$, for any $2 \leq i < j < n$. This immediately implies that this intersection is transversal and consists of the following n points:

$$\begin{aligned} P'_{1n} : y_2 = \dots = y_{n-1} &\neq 0; \\ P'_{in}, 2 \leq i < n : y_i &\neq 0, y_j = 0 \text{ for } j \neq i. \end{aligned}$$

It remains to show that $P'_{in} = P_{in}$. For this we use the equations (1.1.13) of the affine curve associated with each of the points P'_{in} . It is easy to check that this curve is a union of $n - 2$ lines and a conic C_{in} , which has as two points at infinity the marked points p_i and p_n . More precisely, with respect to the coordinates $x_2, \dots, x_n$ on the affine part of the corresponding curve this conic component is given by

$$\begin{aligned} C_{1n} : x_2 = \dots = x_{n-1}, x_2 x_n &= x_2^2 + c_n x_n; \\ C_{2n} : x_3 = \dots = x_{n-1} = 0, x_2 x_n &= c_n(x_n - x_2); \\ C_{3n} : x_2 = x_4 = \dots = x_{n-1} = 0, x_3 x_n &= \bar{c}_n(x_n - \bar{c}_n); \\ C_{in}, 4 \leq i < n : x_i x_n = c_{in}(x_n - x_i), x_j &= 0 \text{ for } j \neq i, n. \end{aligned}$$

(i') Applying a transposition swapping i_0 and n we can assume $i_0 = n$. Now the result follows from (i).

(ii) Let $r : B \rightarrow \overline{\mathcal{M}}_{1,n}^\infty$ be the blow-up at $\overline{\mathcal{M}}_{1,n}^\infty \cap \mathbb{P}(K_n) = \{P_{in} \mid i = 1, \dots, n-1\}$. We can realize both B and $\overline{\mathcal{C}}_{n-1}$ as (reduced) subschemes of the projective bundle

$$\mathbb{P}(\mathcal{O}^{n-2} \oplus \mathcal{O}(-1)) \rightarrow \overline{\mathcal{M}}_{1,n-1}^\infty.$$

Indeed, by (i), B is a closed subscheme of the blow-up $\tilde{B}$ of $\mathbb{P}^{n(n-3)/2}$ along the projective subspace $\mathbb{P}(K_n)$. The linear projection (1.6.1) extends to a regular map $\tilde{B} \rightarrow \mathbb{P}^{(n-1)(n-4)/2}$, which can be identified with the projective bundle $\mathbb{P}(\mathcal{O}^{n-2} \oplus \mathcal{O}(-1))$ over $\mathbb{P}^{(n-1)(n-4)/2}$. On the other hand, the embedding of the relative curve $\overline{\mathcal{C}}_{n-1}$ into the same projective bundle corresponds to the surjection

$$(x_2, \dots, x_{n-1}, 1) : q_{n-1}^*(\mathcal{O}^{n-2} \oplus \lambda) \rightarrow q_{n-1}^* \lambda(p_1 + \dots + p_{n-1}),$$

where $x_i = h_{1i}$ are rational functions defined in Sec. 1.1. Here, instead of normalizing h_{1i} by their residue at p_i , we view them as canonical morphisms

$$h_{1i} : \lambda^{-1} \simeq \mathcal{O}(p_i)|_{p_i} \rightarrow (q_{n-1})_* \mathcal{O}(p_1 + p_i - p_3), \text{ for } i \neq 3,$$

$$h_{13} : \lambda^{-1} \simeq \mathcal{O}(p_3)|_{p_3} \rightarrow (q_{n-1})_* \mathcal{O}(p_1 + p_3 - p_2)$$

(the isomorphisms with λ^{-1} follow from Lemma 1.1.1).

Let $H \subset \mathbb{P}(\mathcal{O}^{n-2} \oplus \mathcal{O}(-1))$ be the relative hyperplane at infinity, i.e., the image of the embedding $\mathbb{P}(K_n) \times \overline{\mathcal{M}}_{1,n-1}^\infty \hookrightarrow \mathbb{P}(\mathcal{O}^{n-2} \oplus \mathcal{O}(-1))$ corresponding to the embedding of bundles $\mathcal{O}^{n-2} \subset \mathcal{O}^{n-2} \oplus \mathcal{O}(-1)$. The intersection $H \cap \overline{\mathcal{C}}_{n-1}$ is exactly the union of the

canonical sections $\sigma_1, \dots, \sigma_{n-1}$. Note that $\bar{\mathcal{C}}_{n-1} \setminus H$ can be identified with the quotient of the affine family of curves

$$\mathcal{C}_{n-1} \setminus K_n \rightarrow U_{n-1} \setminus 0$$

by the action of $\mathbb{G}_m$. By Proposition 1.1.5(ii), we deduce the equality

$$\bar{\mathcal{C}}_{n-1} \setminus H = r^{-1}(\overline{\mathcal{M}}_{1,n}^\infty \setminus \mathbb{P}(K_n)).$$

Passing to closures we get the equality of the subschemes in $\mathbb{P}(\mathcal{O}^{n-2} \oplus \mathcal{O}(-1))$,

$$\bar{\mathcal{C}}_{n-1} = B.$$

The images of the canonical sections σ_i in $H \cap \bar{\mathcal{C}}_{n-1} \subset H = \mathbb{P}(K_n) \times \overline{\mathcal{M}}_{1,n-1}^\infty$ are given by the equations

$$\begin{aligned} \sigma_1 : y_2 = \dots = y_{n-1}; \\ \sigma_i, 2 \leq i < n : y_j = 0 \text{ for } j \neq i, \end{aligned}$$

where (y_i) are the homogeneous coordinates on $\mathbb{P}(K_n)$ defined in part (i). Since the restriction of the projection $r_n : \bar{\mathcal{C}}_{n-1} \rightarrow \overline{\mathcal{M}}_{1,n}^\infty \subset \mathbb{P}^{n(n-3)/2}$ to $H \cap \bar{\mathcal{C}}_{n-1}$ is given by the natural projection $\mathbb{P}(K_n) \times \overline{\mathcal{M}}_{1,n-1}^\infty \rightarrow \mathbb{P}(K_n) \subset \mathbb{P}^{n(n-3)/2}$, we deduce that $r_n(\sigma_i) = P'_{in} = P_{in}$ as claimed.

(iii) Without loss of generality we can assume that $i = 1$ and $j = n - 1$ (see Corollary 1.5.9). Then the fact that $\pi_n(P_{1,n-1}(n)) = P_{1,n-1}(n - 1)$ follows immediately from (i') and from Corollary 1.1.8. Next, as in (i) we see that the elliptic $(n - 2)$ -fold curve corresponding to $P_{1,n-1}(n - 1)$ is given by the equations

$$\begin{aligned} y_i y_j = y_2 y_3 \text{ for } 2 \leq i < j < n - 1, \\ y_2 y_3 + c_{n-1} y_{n-1} = y_i y_{n-1} \text{ for } 2 \leq i < n - 1. \end{aligned}$$

Using this one can easily check the remaining assertions. $\square$

Corollary 1.6.2. *For $n \geq 6$, the tangent cone to $\overline{\mathcal{M}}_{1,n}^\infty$ at the point P_{ij} is isomorphic to the affine cone over $\overline{\mathcal{M}}_{1,n-1}^\infty \subset \mathbb{P}^{(n-1)(n-4)/2}$.*

Proof. For $j = n$ this follows from Proposition 1.6.1(ii). The general case follows using the action of the symmetric group S_n on $\overline{\mathcal{M}}_{1,n}^\infty$. $\square$

We now show how the diagram (1.6.3) can be used in studying the geometric properties of our moduli spaces.

Proposition 1.6.3. *Assume $n \geq 5$.*

(i) *The canonical line bundle K on $\overline{\mathcal{M}}_{1,n}^\infty$ is $K \simeq \mathcal{O}(n - 11)$.*

(ii) *Over an algebraically closed field of characteristic zero or over $\text{Spec}(\mathbb{Z})$ one has*

$$\text{Pic } \overline{\mathcal{M}}_{1,n}^\infty = \mathbb{Z},$$

and this group is generated by the class $\mathcal{O}(1) = \lambda$.

(iii) Let us work over an algebraically closed field of characteristic zero. Then the variety $\overline{\mathcal{M}}_{1,n}^\infty$ has rational singularities if and only if $n \leq 11$.

Proof. For brevity we denote the maps in diagram (1.6.3) as $q = q_{n-1}$, $r = r_n$.

(i) Let $E \subset \overline{\mathcal{C}}_{n-1}$ be the exceptional divisor of the blow-up r (consisting of $n-1$ components). Since q is induced by the linear projection, we have

$$(1.6.4) \quad q^*\mathcal{O}(1) \simeq r^*\mathcal{O}(1)(-E).$$

Next, we note that by Lemma 1.5.6, the relative dualizing sheaf on the universal curve satisfies

$$\omega_q \simeq q^*\lambda = q^*\mathcal{O}(1).$$

Hence, we can prove our assertion by induction in n . For $n = 5$ this is true since $\overline{\mathcal{M}}_{1,5}^\infty \simeq \mathbb{P}^5$. Assuming that the canonical bundle on $\overline{\mathcal{M}}_{1,n-1}^\infty$ is $\mathcal{O}(n-12)$ we get

$$K_{\overline{\mathcal{C}}_{n-1}} \simeq \omega_q \otimes q^*\mathcal{O}(n-12) \simeq q^*\mathcal{O}(n-11).$$

Therefore, by (1.6.4), the canonical bundle of

$$\overline{\mathcal{M}}_{1,n}^\infty \setminus \{P_{1n}, \dots, P_{n-1,n}\} \simeq \overline{\mathcal{C}}_{n-1} \setminus E$$

is still $\mathcal{O}(n-11)$, and the induction step follows (recall that $\overline{\mathcal{M}}_{1,n}^\infty$ is normal by Corollary 1.5.14).

(ii) First, let us work over $\mathbb{C}$. We use the fact that the rational Picard group of $\overline{\mathcal{M}}_{1,n}^\infty$ is $\mathbb{Q}$ ([27, Prop. 3.2]). Let us show now that the group $\text{Pic}(\overline{\mathcal{M}}_{1,n}^\infty)$ has no torsion. Suppose ξ is a torsion line bundle on $\overline{\mathcal{M}}_{1,n}^\infty$. Then $L = r^*\xi$ is a torsion line bundle on $\overline{\mathcal{C}}_{n-1}$ with the property $L|_{E_i} \simeq \mathcal{O}$ for $i = 1, \dots, n-1$, where E_i are the components of the exceptional divisor. Recall that by Proposition 1.6.1(ii), E_i is the image of the i th canonical section of q . Next, we claim that $M := q_*(L(E_1))$ is a line bundle on $\overline{\mathcal{M}}_{1,n-1}^\infty$. Indeed, for every fiber C of q the line bundle $L|_C$ is torsion, hence, it has degree zero on every irreducible component of C . Let $p_1 = E_1 \cap C$. It is enough to show that $H^1(C, L|_C(p_1)) = 0$ (since then $H^0(C, L|_C(p_1))$ will be one-dimensional by Riemann-Roch). By Lemma 1.5.6, we have $\omega_C \simeq \mathcal{O}_C$, so by Serre duality we need to show the vanishing of $H^0(C, L^{-1}|_C(-p_1))$. But the line bundle $L^{-1}|_C(-p_1)$ has degree -1 on one of the irreducible components of C and degree 0 on the remaining components, so it has no global sections. Thus, $M = q_*(L(E_1))$ is a line bundle. Let D be the effective divisor given as the vanishing locus of the natural map $q^*M \rightarrow L(E_1)$, so that

$$(1.6.5) \quad L(E_1) \simeq q^*M(D).$$

Then on each fiber C of q we have $L|_C(p_1) \simeq \mathcal{O}(D)|_C$. We claim that the unique global section of $L|_C(p_1)$ vanishes at exactly one smooth point. This is easy to see when C is either irreducible or a wheel of projective lines. Suppose now that C is an elliptic m -fold curve, and let $C_1 \subset C$ be a component containing p_1 . It suffices to show that we cannot have a global section of $L|_C(p_1)$ vanishing at all the other components of C . Indeed, restricting such a global section to a neighborhood of the singular point q , we will get a

germ $f \in \mathcal{O}_{C,q}$ which restricts to zero on all branches but one and will have a nonzero derivative at q on the remaining branch. This contradicts the explicit description of $\mathcal{O}_{C,q}$ (see [26, Sec. 2] and Lemma 2.2.1(i) below), so no such global section exists. Hence, D defines a section of q , so it is isomorphic to $\overline{\mathcal{M}}_{1,n-1}^\infty$. Assume first that D is different from all the divisors E_i . Then the conditions $L|_{E_i} \simeq \mathcal{O}$ for $i = 1, \dots, n-1$ give isomorphisms

$$\begin{aligned}\mathcal{O}(-1) &\simeq M(q(D \cap E_1)), \\ \mathcal{O} &\simeq M(q(D \cap E_i)) \text{ for } i \geq 2\end{aligned}$$

on $\overline{\mathcal{M}}_{1,n-1}^\infty$. In particular, the Cartier divisors $q(D \cap E_2)$ and $q(D \cap E_3)$ are linearly equivalent. Since $\text{Pic}(\overline{\mathcal{M}}_{1,n-1}^\infty)_\mathbb{Q} = \mathbb{Q}$, every nonzero effective Cartier divisor on $\overline{\mathcal{M}}_{1,n-1}^\infty$ is ample. Since the divisors $q(D \cap E_2)$ and $q(D \cap E_3)$ do not intersect, we deduce that $D \cap E_2 = 0$, so $M \simeq \mathcal{O}$. But then we should have $\mathcal{O}(-1) \simeq \mathcal{O}(q(D \cap E_1))$ which is a contradiction. Suppose next that $D = E_i$ with $i \geq 2$. Then restricting (1.6.5) to E_1 and to E_i we get $\mathcal{O}(-1) \simeq M$ and $\mathcal{O} \simeq M \otimes \mathcal{O}(-1)$, which is a contradiction. It follows that $D = E_1$, i.e., $L = q^*M$. Restricting to E_1 we get $M \simeq \mathcal{O}$, hence, L is trivial.

Finally, to check that $\mathcal{O}(1)$ is a generator of $\text{Pic}(\overline{\mathcal{M}}_{1,n}^\infty)$, we observe that $\overline{\mathcal{M}}_{1,n}^\infty$ contains a projective line (with respect to the embedding given by $\mathcal{O}(1)$). Indeed, we can use of the lines contained in $\pi_n^{-1}(P_{ij}(n-1))$ (see Proposition 1.6.1(iii)). Hence, $\mathcal{O}(1)$ is not divisible in the Picard group of $\overline{\mathcal{M}}_{1,n}^\infty$.

The same result over $\text{Spec}(\mathbb{Z})$ follows by the standard method (see [21, p. 103]), using the irreducibility of the fibers of $\overline{\mathcal{M}}_{1,n}^\infty \rightarrow \text{Spec}(\mathbb{Z})$.

(iii) First, let us check that $\overline{\mathcal{M}}_{1,n}^\infty$ has rational singularities for $n \leq 11$, by induction on n . For $n \leq 6$ this is true since our moduli space is smooth. Assume $\overline{\mathcal{M}}_{1,n-1}^\infty$ has rational singularities. Let P be a point of $\overline{\mathcal{M}}_{1,n}^\infty$ which is different from all (P_{ij}) , so that the corresponding curve $(C, p_1, \dots, p_n)$ has at most elliptic $(n-2)$ -fold points as singularities. Applying the S_n -action to P we can assume that there is more than one marked point on the component of C containing p_n . Then viewing P as a point on $\overline{\mathcal{C}}_{n-1}$ we see that the projection $\overline{\mathcal{C}}_{n-1} \rightarrow \overline{\mathcal{M}}_{1,n-1}^\infty$ is smooth near P . Thus, P has an open neighborhood U which is smooth over $\overline{\mathcal{M}}_{1,n-1}^\infty$, hence U has rational singularities. It remains to check that $\overline{\mathcal{M}}_{1,n}^\infty$ has rational singularities near each of the points P_{ij} . Using the S_n -action, it is enough to consider the points P_{in} . Note that the projection $q : \overline{\mathcal{C}}_{n-1} \rightarrow \overline{\mathcal{M}}_{1,n-1}^\infty$ is smooth near E , hence $\overline{\mathcal{C}}_{n-1}$ has rational singularities near E . Now by Lemma 1.6.4 below, applied to the blow-up morphism r , $\overline{\mathcal{M}}_{1,n}^\infty$ has rational singularities near P_{in} if and only if

$$r_*K_{\overline{\mathcal{C}}_{n-1}} \simeq K_n,$$

where K_n is the canonical bundle of $\overline{\mathcal{M}}_{1,n-1}^\infty$. As we have seen in part (i),

$$K_{\overline{\mathcal{C}}_{n-1}} \simeq q^*\mathcal{O}(n-11) \simeq r^*K_n((11-n)E),$$

where we used (1.6.4). Thus,

$$r_*K_{\bar{C}_{n-1}} \simeq K_n \otimes r_*(\mathcal{O}((11-n)E)).$$

Since $H^0(E, \mathcal{O}_E(-i)) = 0$ for $i > 0$, we see that $r_*(\mathcal{O}(iE)) = \mathcal{O}$ for $i > 0$, which finishes the induction step. The same argument for $n = 12$ shows that $\bar{\mathcal{M}}_{1,12}^\infty$ does not have rational singularities at the special points P_{ij} (since $r_*(\mathcal{O}(-E)) \neq \mathcal{O}$). For any $n > 12$ we can find a point $P \in \bar{\mathcal{M}}_{1,n}^\infty$ projecting to a special point in $\bar{\mathcal{M}}_{1,12}^\infty$, such that an open neighborhood of P projects smoothly to $\bar{\mathcal{M}}_{1,12}^\infty$. Hence, $\bar{\mathcal{M}}_{1,n}^\infty$ will not have a rational singularity near P . $\square$

We have used the following result, well-known to the experts (see e.g., [15, Lem. 1, Thm. 3] for similar statements).

Lemma 1.6.4. *Let $\phi : Y \rightarrow X$ be a proper birational morphism of quasi-projective varieties over an algebraically closed field of characteristic zero. Assume that Y has rational singularities. Then X has rational singularities if and only if X is Cohen-Macaulay and $\phi_*\omega_Y \simeq \omega_X$.*

Proof. Let $\pi : Z \rightarrow Y$ be the resolution of singularities, $p = \phi \circ \pi$. Let us also set $n = \dim X = \dim Y = \dim Z$. Note that we have $R\pi_*\mathcal{O}_Z \simeq \mathcal{O}_Y$ since Y has rational singularities. By Grothendieck duality, this implies that $R\pi_*\omega_Z[n]$ is the dualizing complex of Y , which means, due to Grauert-Riemenschneider vanishing, that Y is Cohen-Macaulay and $\pi_*\omega_Z \simeq \omega_Y$.

Assume first that X has rational singularities. Then as above, we deduce that X is Cohen-Macaulay and $p_*\omega_Z \simeq \omega_X$. Hence,

$$\phi_*\omega_Y \simeq \phi_*\pi_*\omega_Z \simeq p_*\omega_Z \simeq \omega_X$$

as required.

Conversely, if X is Cohen-Macaulay and $\phi_*\omega_Y \simeq \omega_X$ then we get that $p_*\omega_Z \simeq \omega_X$, so by Grauert-Riemenschneider vanishing, we deduce that $Rp_*\omega_Z[n] = \omega_X[n]$ is the dualizing complex of X . Applying Grothendieck duality we obtain

$$Rp_*\mathcal{O}_Z \simeq R\text{Hom}(\omega_X[n], \omega_X[n]) \simeq \mathcal{O}_X,$$

so X has rational singularities. $\square$

1.7. Curves which are linear sections of $Gr(2, 5)$. In the case $n = 6$ we can identify the blow-up picture of Proposition 1.6.1 with a generic linear projection of the Grassmannian $Gr(2, 5)$ to $\mathbb{P}^5$.

Proposition 1.7.1. *There is an isomorphism $\bar{\mathcal{M}}_{1,6}^\infty \cong Gr(2, 5) \subset \mathbb{P}^9$, so that the map q_5 from the blow up of $\bar{\mathcal{M}}_{1,6}^\infty$ at 5 points to $\bar{\mathcal{M}}_{1,5}^\infty \simeq \mathbb{P}^5$ (see (1.6.3)) gets identified with a generic linear projection of $Gr(2, 5)$ (that can be defined over $\mathbb{Z}$).*

First, we consider the well known family of (degenerating) elliptic curves obtained by linear sections of $Gr(2, 5)$ (see e.g., [7, Sec. 2]). Namely, we consider a fixed linear subspace $L = \mathbb{P}^3 \subset \mathbb{P}^9$, intersecting $Gr(2, 5)$ at 5 distinct points $p_1, \dots, p_5$. Then for every 4-dimensional subspace $P \subset \mathbb{P}^9$, containing L , the intersection $Gr(2, 5) \cap P$ is a curve of arithmetic genus 1, containing the points $p_1, \dots, p_5$.

Lemma 1.7.2. *Let k be an algebraically closed field, and consider the above picture over k . Then for every 4-dimensional subspace $P \subset \mathbb{P}^9$ containing L , the curve $C = Gr(2, 5) \cap P$ with 5 marked points $p_1, \dots, p_5$ is 4-stable, i.e., defines a point of $\overline{\mathcal{M}}_{1,5}^\infty$.*

Proof. The curve $C = Gr(2, 5) \cap P$ is Gorenstein and is of arithmetic genus 1. Also, the line bundle $\mathcal{O}_C(p_1 + \dots + p_5) = \mathcal{O}_C(1)$ is ample. By [28, Prop. 2.3], the points $p_1, \dots, p_5$ are in general linear position. Hence,

$$h^0(\mathcal{O}_C(p_i)) = h^0(\mathcal{O}_C(1)(-p_1 - \dots - \hat{p}_i - \dots - p_5)) = 1$$

for $i = 1, \dots, p$. Therefore, $(C, p_1, \dots, p_5)$ corresponds to a point of $\mathcal{U}_{1,5}^{sns}$. By Theorem 1.5.7, it remains to check that we cannot have $C \simeq C_{1,5}$. Since C has degree 5, if this were the case C would have to be a union of 5 lines ℓ_i , connecting a point $q \in Gr(2, 5)$ with each of the five points $p_i = [U_i]$. Let $T \subset \mathbb{P}^9$ be the tangent space to $Gr(2, 5)$ at q . Then we have $\ell_i \in T$ for $i = 1, \dots, 5$, hence, $P \subset T$. It is well known that the intersection $Gr(2, 5) \cap T$ is 4-dimensional (in fact, it is the cone over the Segre embedding of $\mathbb{P}^1 \times \mathbb{P}^2$, see [2, Sec. 6.1.3]). Since P is a linear subspace of codimension 2 in T , this implies that $Gr(2, 5) \cap P$ has dimension ≥ 2 , which is a contradiction. $\square$

Proof of Proposition 1.7.1. It is enough to choose the embedding $L = \mathbb{P}^3 \subset \mathbb{P}^9$ defined over $\mathbb{Z}$ such that $Gr(2, 5) \cap L$ consists of 5 distinct points (over any algebraically closed field). Indeed, let $X \rightarrow Gr(2, 5)$ be the blow up along $p_1, \dots, p_5$. Then the fibers of the regular map $\pi : X \rightarrow \mathbb{P}^5$ (defined as a linear projection with the center L) are exactly linear sections of $Gr(2, 5)$ by 4-dimensional subspaces containing L . The exceptional divisors $E_1, \dots, E_5 \subset X$ give five non-intersecting sections of π . Thus, by Lemma 1.7.2, we get a family of 4-stable curves, which is the pull-back of the universal family with respect to some regular map $f : \mathbb{P}^5 \rightarrow \overline{\mathcal{M}}_{1,5}^\infty$. Recall that $\overline{\mathcal{M}}_{1,5}^\infty \simeq \mathbb{P}^5$ so that $\lambda = \mathcal{O}(1)$. We claim that $f^*\mathcal{O}(1) \simeq \mathcal{O}(1)$. Indeed, we have to calculate the line bundle λ associated with the family $\pi : X \rightarrow \mathbb{P}^5$. By Lemma 1.1.1(ii), we have an isomorphism

$$\lambda \simeq \omega_\pi|_{E_1} \simeq \mathcal{O}_X(-E_1)|_{E_i}$$

for any $i = 1, \dots, 5$. But $\mathcal{O}_X(-E_i)|_{E_i} \simeq \mathcal{O}(1)$ since E_i is the exceptional divisor of a blow-up. Hence, f is an isomorphism and the map $\pi : X \rightarrow \mathbb{P}^5$ is a universal curve over $\overline{\mathcal{M}}_{1,5}^\infty$. Now the identification of $\overline{\mathcal{M}}_{1,6}^\infty$ follows from Proposition 1.6.1.

It remains to give a subspace $\mathbb{P}^3 \subset \mathbb{P}^9$ with the required properties. We use homogeneous coordinates

$$[z_{12} : z_{13} : z_{14} : z_{15} : z_{23} : z_{24} : z_{25} : z_{34} : z_{35} : z_{45}]$$

on $\mathbb{P}^9$, so that the image of the Plücker embedding of $Gr(2, 5)$ is cut out by the equations

$$\begin{aligned} z_{12}z_{34} + z_{14}z_{23} &= z_{13}z_{24}, \\ z_{13}z_{45} + z_{15}z_{34} &= z_{14}z_{35}, \\ z_{12}z_{45} + z_{15}z_{24} &= z_{14}z_{25}, \\ z_{12}z_{35} + z_{15}z_{23} &= z_{13}z_{25}, \\ z_{23}z_{45} + z_{25}z_{34} &= z_{24}z_{35}. \end{aligned}$$

We choose the following four points on $Gr(2, 5)$:

$$\begin{aligned} p_1 &= [1 : 0 : -1 : 0 : 0 : 1 : 0 : 0 : 0 : 0], \\ p_2 &= [0 : 1 : 0 : -1 : 0 : 0 : 0 : 0 : 1 : 0], \\ p_3 &= [0 : 0 : 0 : 0 : 1 : 0 : 0 : 0 : 0 : 0], \\ p_4 &= [0 : 0 : 0 : 0 : 0 : 0 : 0 : 0 : 0 : 1], \end{aligned}$$

and let $L = \mathbb{P}^3$ be the linear subspace spanned by these four points in $\mathbb{P}^9$. Note that L is defined by the equations

$$z_{12} = z_{24}, \quad z_{13} = z_{35}, \quad z_{14} + z_{24} = 0, \quad z_{15} + z_{35} = 0, \quad x_{25} = 0, \quad x_{34} = 0.$$

In particular, L is defined over $\mathbb{Z}$. Since $Gr(2, 5) \subset \mathbb{P}^9$ has degree 5, one expects that L intersects $Gr(2, 5)$ in yet another point. Using the Plücker relations, it is easy to verify that indeed L intersects $Gr(2, 5)$ in exactly one other point, namely:

$$p_5 = [1 : -1 : -1 : 1 : 1 : 1 : 0 : 0 : -1 : -1].$$

□

Remark 1.7.3. Over an algebraically closed field of characteristic 0 the isomorphism of $\overline{\mathcal{M}}_{1,6}^\infty$ with $Gr(2, 5)$ follows immediately from the fact that $\overline{\mathcal{M}}_{1,6}^\infty$ is a smooth projective variety of dimension 6 and degree 5 in $\mathbb{P}^9$ (not contained in any hyperplane). Indeed, this is a part of the classification result of Fujita in [11].

Definition 1.7.4. Let C be a projective curve of arithmetic genus 1. We say that C is *4-prestable* if either C is irreducible with at most one nodal singularity, or, C is a wheel of ≤ 5 projective lines, or C is the elliptic m -fold curve with $m \leq 4$.

Note that over an algebraically closed field, a curve C is 4-prestable if and only if there exists 5 smooth points $p_1, \dots, p_5$ such that $(C, p_1, \dots, p_5)$ is 4-stable.

Corollary 1.7.5. *Let k be an algebraically closed field. A curve $C \subset \mathbb{P}^4$ can be obtained as a linear section of $Gr(2, 5)$ (in its Plücker embedding) if and only if C is 4-prestable.*

Remark 1.7.6. The fact that every 4-stable curve $(C, p_1, \dots, p_5)$ can be realized as a linear section of $Gr(2, 5)$ can also be proved directly using the fact that such curves are arithmetically Gorenstein in the projective embedding associated with $\mathcal{O}(p_1 + \dots + p_5)$. Namely, one can mimic the construction from [10, Sec. 4] of the Pfaffian presentation of

elliptic normal curves of degree 5 based on Buchsbaum-Eisenbud theorem on Gorenstein ideals of codimension 3 (see [4]). Let $I \subset S = k[x_0, \dots, x_4]$ be the homogeneous ideal corresponding to C . Then S/I has a minimal free resolution of the form

$$0 \rightarrow S(-5) \rightarrow S(-3)^5 \xrightarrow{f} S(-2)^5 \rightarrow S \rightarrow S/I \rightarrow 0,$$

where f is a skew-symmetric matrix of linear forms and I is generated by the principal 4×4 -Pfaffians of f . Interpreting f as a map $\tilde{f} : k^5 \rightarrow \wedge^2(k^5)$ we get a required linear section of the Grassmannian. In the case when C is the elliptic 5-fold curve, i.e., the union of 5 generic lines in $\mathbb{P}^4$, the minimal resolution still has the same form. However, in this case $\tilde{f}$ has a one-dimensional kernel, so it does not give an embedding of $\mathbb{P}^4$ into $\mathbb{P}^9$. Instead, in this case the Pfaffian presentation realizes C as a cone over 5 points in $\mathbb{P}^3$.

2. MODULI OF A_∞ -STRUCTURES

2.1. A_∞ -structures associated with curves.

For each curve $(C, p_1, \dots, p_n)$ corresponding to a point of $\mathcal{U}_{1,n}^{sns}$ we can consider the associative algebra $\text{Ext}^*(G, G)$, where

$$G = \mathcal{O}_C \oplus \mathcal{O}_{p_1} \oplus \dots \oplus \mathcal{O}_{p_n}.$$

It turns out that the fact that $h^0(\mathcal{O}_C(p_i)) = 1$ for all i implies that the algebra $\text{Ext}^*(G, G)$ is independent of the curve $(C, p_1, \dots, p_n)$, up to an isomorphism (the requirement that $\mathcal{O}_C(p_1 + \dots + p_n)$ is ample is not important at the moment).

More precisely, let us denote by $A_i \in \text{Hom}(\mathcal{O}_C, \mathcal{O}_{p_i}) \subset \text{Ext}^*(G, G)$ the natural generators. Also, a choice of nonzero tangent vectors at the marked points gives canonical generators B_i of the one-dimensional spaces $\text{Ext}^1(\mathcal{O}_{p_i}, \mathcal{O}_C) \subset \text{Ext}^*(G, G)$, so that the algebra $\text{Ext}^*(G, G)$ is generated by A_i and B_i over the subalgebra k^{n+1} (generated by the projectors to the summands of G). The algebra structure on $\text{Ext}^*(G, G)$ is given by the following easy computation, similar to the one in [9, Sec. 1.1].

Lemma 2.1.1. *Let $Q = Q_n$ be the quiver as in Figure 1. We identify A_i with the arrow from the central vertex to the vertex i and B_i with the arrow in the opposite direction. A choice of nonzero tangent vectors at all p_i 's gives rise to a canonical isomorphism of k -algebras*

$$\text{Ext}^*(G, G) = E_{1,n} := k[Q]/J_1,$$

where J_1 is the ideal generated by the relations

$$B_i A_i = B_j A_j, \quad A_i B_j = 0 \quad \text{for } i \neq j.$$

□

We will consider this as an isomorphism of graded algebras by declaring the gradings as $|A_i| = 0$ and $|B_i| = 1$. Note that this is different from the path-length grading.

We use the convention that the paths are composed from the right.¹ For example, $B_i A_i$ is the image under the product:

$$\text{Ext}^*(\mathcal{O}_{p_i}, \mathcal{O}_C) \otimes \text{Ext}^*(\mathcal{O}_C, \mathcal{O}_{p_i}) \rightarrow \text{Ext}^*(\mathcal{O}_C, \mathcal{O}_C)$$

of the generators $B_i \in \text{Ext}^*(\mathcal{O}_{p_i}, \mathcal{O}_C)$ and $A_i \in \text{Ext}^*(\mathcal{O}_C, \mathcal{O}_{p_i})$.

FIGURE 1. The quiver Q_n

Let R be any commutative ring. Recall that an A_∞ -algebra $\mathcal{A}$ over R is a graded R -module with a collection of R -linear maps

$$\mu^d : \mathcal{A}^{\otimes d} \rightarrow \mathcal{A}[2-d], \quad \text{for } d \geq 1,$$

i.e., μ^d lowers the degree by $d-2$. These maps are required to satisfy the A_∞ -associativity equations:

$$\sum_{m,n} (-1)^{|a_1|+\dots+|a_n|-n} \mu^{d-m+1}(a_d, \dots, a_{n+m+1}, \mu^m(a_{n+m}, \dots, a_{n+1}), a_n, \dots, a_1) = 0$$

In particular, $\mu^1 : \mathcal{A} \rightarrow \mathcal{A}[1]$ is a differential, i.e. $\mu^1 \circ \mu^1 = 0$, and the double product

$$(2.1.1) \quad a_2 \cdot a_1 = (-1)^{|a_1|} \mu^2(a_2, a_1)$$

on $\mathcal{A}$ is associative up to homotopy.

A *minimal A_∞ -algebra* $(\mathcal{A}, \mu^*)$ over a graded associative R -algebra $(A, \cdot)$ is an A_∞ -algebra structure on A with $\mu^1 = 0$ and such that (2.1.1) is the given double product on A .

When considering A_∞ -algebra structures on $E_{1,n}$ we in addition assume that they are *unital* with respect to the natural idempotents in $E_{1,n}$ corresponding to the vertices of the quiver Q_n , i.e., any operation μ^l , $l > 2$, that has one of these idempotents as one of the arguments, is required to vanish.

¹This is opposite to the convention adopted in [9] and [22], where we reverse the direction of the arrows in the quiver Q_n .

Here we do not delve into the theory of A_∞ -algebras, as they are well studied in the literature. A good reference is [24, Ch. 1] but the more relevant aspect of the theory for us is discussed in detail in [22, Sec. 4], which we refer to in this text.

Working over a field k , it turns out that if the first Hochschild cohomology group $HH^1(A)_{<0}$ vanishes then the set of minimal A_∞ -structures on the associative algebra A up to a gauge equivalence can be represented by an affine scheme (see [22, Cor. 4.2.5]). We denote this affine scheme by $\widetilde{\mathcal{M}}_\infty(A)$. Note that, in general, this scheme is constructed as the inverse limit

$$\widetilde{\mathcal{M}}_\infty(A) = \varprojlim_d \widetilde{\mathcal{M}}_d(A)$$

of affine schemes $\widetilde{\mathcal{M}}_d(A)$ of finite type which represents the set of minimal A_d -structures over A , hence is not necessarily of finite type.

As in [22, Section 3.1] (see also, [16, Section 5.1] for the special case $n = 1$), we can use a natural dg-resolution of $\text{Ext}^*(G, G)$ to construct an A_∞ -structure on $E_{1,n} \otimes R$ associated with a family of curves $(C, p_1, \dots, p_n)$ over $\text{Spec } R$ in $\widetilde{\mathcal{U}}_{1,n}^{sns}(R)$ (where R is a commutative ring). This is constructed by considering endomorphisms of G in a dg-enhancement of $D^b(C)$ and then applying the homological perturbation lemma (see [20], [14]) to get a minimal A_∞ -algebra, defined canonically up to a gauge equivalence.

Thus, we have a morphism of functors

$$(2.1.2) \quad \widetilde{\mathcal{U}}_{1,n}^{sns} \rightarrow \widetilde{\mathcal{M}}_\infty(E_{1,n}),$$

where $\widetilde{\mathcal{M}}_\infty(E_{1,n})$ is the functor associating with R the set of gauge equivalence classes of minimal A_∞ -structures on $E_{1,n} \otimes R$. We will show later using the results of [22, Sec. 4] that if we work over a field k then $\widetilde{\mathcal{M}}_\infty(E_{1,n})$ is representable by an affine k -scheme of finite type. Our main theorem in this section is that the map (2.1.2) is an isomorphism of affine schemes.

Lemma 2.1.2. *(i) The map (2.1.2) is compatible with the natural $\mathbb{G}_m$ -actions, where the action on $\widetilde{\mathcal{M}}_\infty(E_{1,n})$ is given by*

$$(2.1.3) \quad (\mu^n) \mapsto (\lambda^{n-2} \mu^n),$$

which is also the transformation induced by the rescaling $B_i \mapsto \lambda B_i$.

(ii) The A_∞ -structure on $E_{1,n}$ associated with the elliptic n -fold curve $C_{1,n}$ is trivial (up to a gauge equivalence).

Proof. (i) As in [22, Prop. 3.3.2] (see also [16] Lemma 5.2 for $n = 1$), one can check that it is possible to choose the homotopy operator needed to run the homological perturbation for the dg-algebra associated with the universal curve over the affine scheme $\widetilde{\mathcal{U}}_{1,n}^{sns} \simeq U_n$, in a $\mathbb{G}_m$ -equivariant way (where the $\mathbb{G}_m$ -action is extended naturally to the universal curve). Then, as in [22, Prop. 3.3.2], one checks that the resulting higher products μ^n

will have weight $n-2$ with respect to the $\mathbb{G}_m$ -action. It is easy to check that the rescaling $B_i \mapsto \lambda B_i$ produces the same transformation (2.1.3).

(ii) (See [22, Prop. 4.4.1] for a similar argument.) First, we observe that there is a natural $\mathbb{G}_m$ -action on $C_{1,n}$ (induced by the $\mathbb{G}_m$ -action on the universal curve over U_n), therefore, we get an induced $\mathbb{G}_m$ -action on the Ext-algebra $E_{1,n}$. It is easy to see that this action is given by λ^{deg} where deg is the cohomological grading on $E_{1,n}$. Using the $\mathbb{G}_m$ -equivariant homological perturbation as in (i), we obtain a $\mathbb{G}_m$ -equivariant A_∞ -structure on $E_{1,n}$. But μ^n lowers the cohomological degree by $n-2$, so only μ^2 can be nonzero. $\square$

Similarly to [22, Prop. 4.4.1] we get the following result connecting the associative algebra $E_{1,n}$ to the elliptic n -fold curve $C_{1,n}$.

Proposition 2.1.3. *We have an equivalence of perfect derived categories*

$$\mathrm{Per}(C_{1,n}) \simeq \mathrm{Per}(E_{1,n})$$

inducing a $\mathbb{G}_m$ -equivariant isomorphism

$$(2.1.4) \quad HH^*(C_{1,n}) \simeq HH^*(E_{1,n}),$$

so that the second grading on these spaces is given by the weights of the $\mathbb{G}_m$ -action.

2.2. Comparison of the moduli spaces. We have the following analog of [22, Lem. 4.4.2] (See also [26, Prop 2.3].)

Lemma 2.2.1. *Let $C = C_{1,n}$, where $n \geq 2$. We work over an arbitrary field k , except in the case $n = 2$, assume that $\mathrm{char}(k) \neq 2$. Recall that $D = p_1 + \dots + p_n$.*

(i) The algebra of functions on $C \setminus D$ can be identified with the subalgebra in $\prod_{i=1}^n k[x_i]$ consisting of $(f_i(x_i))$ such that $f_i(0) = f_j(0)$ for $i \neq j$ and

$$f'_1(0) = f'_2(0) + \dots + f'_n(0).$$

(ii) The one-dimensional space $H^1(C, \mathcal{O})$ has weight 1 with respect to the $\mathbb{G}_m$ -action.

(iii) The space $H^0(C, \mathcal{T})$, where $\mathcal{T}$ is the tangent sheaf, decomposes as a direct sum

$$H^0(C, \mathcal{T}) = H^0(C, \mathcal{T}(-D)) \oplus V,$$

where V is an n -dimensional subspace of weight 1 with respect to the $\mathbb{G}_m$ -action, such that the composition

$$V \rightarrow H^0(C, \mathcal{T}) \rightarrow H^0(D, \mathcal{T}|_D)$$

is an isomorphism. Furthermore,

$$H^0(C, \mathcal{T}(-D)) = H^0(C, \mathcal{T})^{\mathbb{G}_m}$$

and this space is spanned by the derivation coming from the $\mathbb{G}_m$ -action on C . The natural map $H^0(C, \mathcal{T}(nD)) \rightarrow H^0(C, \mathcal{T}(nD)|_D)$ is surjective for $n \geq 0$. Also, one has

$$H^0(C, \mathcal{T}(-2D)) = 0.$$

(iv) One has $H^1(C, \mathcal{T}) = 0$.

Proof. Let $C_i \simeq \mathbb{P}^1$, $i = 1, \dots, n$, be the irreducible components of C , and let $q \in C$ be the singular point. For $n \geq 3$ we number components in such a way that x_i is a coordinate on $C_i \setminus \{p_i\}$ for $i \geq 2$ and $x_j = 0$ on C_i for $j \geq 2$, $j \neq i$. The remaining component C_1 has the affine part $C_1 \setminus \{p_1\}$ given by $x_i = x_j$ for $i \neq j$, and we denote by x_1 the restriction of any of x_i 's to $C_1 \setminus \{p_1\}$. For $n = 2$, we let x_i be the natural coordinates on $C_i \setminus \{p_i\}$, $i = 1, 2$, obtained by restricting x (so that $y = 0$ on C_1 and $y = x^2$ on C_2). Let us also set $U = C \setminus D$, $U_i = C_i \setminus \{p_i\}$ and $V = C \setminus \{q\}$.

(i) Assume first that $n \geq 3$. Then the algebra $\mathcal{O}(C \setminus D)$ has the basis $1, x_i^m, x_2^m x_3$, where $i \geq 2$, $m \geq 1$. The projections $\mathcal{O}(C \setminus D) \rightarrow k[x_i]$ for $i \geq 2$ are given by $x_i \mapsto x_i$, $x_j \mapsto 0$ for $j \neq i$. The projection $\mathcal{O}(C \setminus D) \rightarrow k[x_1]$ sends all x_i to x_1 . Now the assertion follows immediately by considering the images of the basis vectors. In the case $n = 2$ we have the basis x^m, yx^m (where $m \geq 0$) on $\mathcal{O}(C \setminus D)$. The map

$$\mathcal{O}(C \setminus D) \rightarrow k[x_1] \oplus k[x_2]$$

sends x^m to (x_1^m, x_2^m) and yx^m to $(0, x_2^{m+2})$, and the assertion follows.

(ii) We can compute $H^1(C, \mathcal{O})$ using the covering of C by two affine open sets: $C = U \cup V$. Thus, this group is identified with the cokernel of the map

$$H^0(U, \mathcal{O}) \oplus H^0(V, \mathcal{O}) \rightarrow H^0(U \cap V, \mathcal{O}) = \prod_{i=1}^n k[x_i, x_i^{-1}].$$

Functions on $V = \sqcup_i C_i \setminus \{q\}$ map to collections $(P_i(x_i^{-1}))$, where P_i are arbitrary polynomials. A collection $(x_i Q_i(x_i))$, where Q_i are polynomials, comes from an element of $H^0(U, \mathcal{O})$ if and only if $Q_1(0) = 0$. Hence, the classes in $H^1(C, \mathcal{O})$ are represented by elements of the form (ax_1) with $a \in k$. It remains to observe that $(\lambda^{-1})^* x_1 = \lambda x_1$.

(iii) Let us first study derivations of the algebra $\mathcal{O}(U)$. Every such derivation restricts to a derivation of $\mathcal{O}(U \setminus q) = \prod_{i=1}^n \mathcal{O}(U_i)$. Assume first that $n \geq 3$. Then we get a collection of vector fields $v_i \in k[x_i, x_i^{-1}] \partial_{x_i}$. It is easy to see that such a collection extends to a derivation of $\mathcal{O}(U)$ if and only if there exists a constant $a \in k$ such that $v_i \in (ax_i + x_i^2 k[x_i]) \partial_{x_i}$. On the other hand, v_i is regular at infinity (i.e., extends to $C_i \setminus \{q\}$) if and only if $v_i \in x_i^2 k[x_i^{-1}] \partial_{x_i}$. Thus, an element of $H^0(C, \mathcal{T})$ corresponds to a collection of the form $(v_i = (ax_i + b_i x_i^2) \partial_{x_i})$; the subspace $H^0(C, \mathcal{T}(-D))$ consists of $(v_i = ax_i \partial_{x_i})$. On the other hand, for $n \geq 1$ the space $H^0(C, \mathcal{T}(nD))$ consists of $v_i = P_i(x_i) \partial_{x_i}$, where P_i are polynomials of degree $n + 2$ with $P_i(0) = 0$, $P_i'(0) = a$ (independent of i). This easily implies all our assertions.

(iv) As in (ii), we can use the covering $C = U \cup V$ to compute $H^1(C, \mathcal{T})$. Thus, we just need to see that every derivation of $\mathcal{O}(U \setminus q)$ is a sum of a derivation that is regular at infinity and a derivation that extends to U . But this follows easily from the explicit form of such derivations in (iii). $\square$

As in [22, Lem. 4.4.3], we deduce the following results about the Hochschild cohomology of $C_{1,n}$.

Corollary 2.2.2. *For $n \geq 3$, one has*

$$HH^1(C_{1,n})_{<0} = 0,$$

and the natural map

$$HH^2(C_{1,n}) \rightarrow HH^2(U)$$

is an isomorphism, where $U = C_{1,n} \setminus D$.

The same conclusions hold for $n = 2$ and $\mathbf{char}(k) \neq 2$ (resp., for $n = 1$ and $\mathbf{char}(k) \neq 2, 3$).

Proof. As before, we set $C = C_{1,n}$. We use the natural exact sequences

$$\begin{aligned} 0 \rightarrow H^1(C, \mathcal{O}) \rightarrow HH^1(C) \rightarrow H^0(C, \mathcal{T}) \rightarrow 0, \\ 0 \rightarrow H^1(C, \mathcal{T}) \rightarrow HH^2(C) \rightarrow HH^2(U) \rightarrow 0 \end{aligned}$$

(see [16, Sec. 4.1.3]). Now the assertions follows from Lemma 2.2.1(ii),(iii),(iv). For the case $n = 1$, see [16, Sec. 4.1, 4.2]. $\square$

Using the isomorphism (2.1.4) we deduce the vanishing

$$(2.2.1) \quad HH^1(E_{1,n})_{<0} = 0,$$

which implies the following result.

Lemma 2.2.3. *For $n \geq 3$ the functor $\widetilde{\mathcal{M}}_\infty(E_{1,n})$ of minimal A_∞ -structures on $E_{1,n}$ is represented by an affine k -scheme. The same conclusion holds for $n = 2$ and $\mathbf{char}(k) \neq 2$ (resp., $n = 1$ and $\mathbf{char}(k) \neq 2, 3$).*

Proof. This follows from [22, Cor. 4.2.5]) using the vanishing (2.2.1). $\square$

For every k -scheme X we denote by $\mathbf{L}_X$ the cotangent complex of X over k . We have the following analog of Lemma 4.4.5 of [22].

Lemma 2.2.4. *For $C = C_{1,n}$ the natural maps*

$$\begin{aligned} \mathrm{Ext}_C^1(\mathbf{L}_C, \mathcal{O}(-D - p_i)) \rightarrow \mathrm{Ext}_C^1(\mathbf{L}_C, \mathcal{O}(-D)) \rightarrow \mathrm{Ext}_U^1(\mathbf{L}_U, \mathcal{O}_U) \quad \text{and} \\ \mathrm{Ext}_C^2(\mathbf{L}_C, \mathcal{O}(-D - p_i)) \rightarrow \mathrm{Ext}_U^2(\mathbf{L}_U, \mathcal{O}_U) \end{aligned}$$

are isomorphisms, where p_i is any of the standard marked points on $C = C_{1,n}$.

Proof. The proof is similar to that of [22, Lem. 4.4.5]. Since $\mathbf{L}_C$ is a perfect complex, it is enough to show that the maps

$$\begin{aligned} \mathrm{Ext}_C^i(\mathbf{L}_C, \mathcal{O}(-D - p_i)) \rightarrow \mathrm{Ext}_C^i(\mathbf{L}_C, \mathcal{O}(-D)) \quad \text{and} \\ \mathrm{Ext}_C^i(\mathbf{L}_C, \mathcal{O}(nD)) \rightarrow \mathrm{Ext}_C^i(\mathbf{L}_C, \mathcal{O}((n+1)D)), \quad n \geq -1, \end{aligned}$$

are isomorphisms for $i = 1, 2$. Using the exact sequences

$$\begin{aligned} 0 \rightarrow \mathcal{O}(-D - p_i) \rightarrow \mathcal{O}(-D) \rightarrow \mathcal{O}_{p_i} \rightarrow 0, \\ 0 \rightarrow \mathcal{O}(nD) \rightarrow \mathcal{O}((n+1)D) \rightarrow \mathcal{O}_D \rightarrow 0, \end{aligned}$$

this reduces to the surjectivity statement in Lemma 2.2.1(iii) together with the surjectivity of the map

$$H^0(C, \mathcal{T}(-D)) \rightarrow H^0(C, \mathcal{T}(-D)|_{p_i})$$

which is checked similarly. $\square$

Next, we are going to compare the deformation theories of $\widetilde{\mathcal{U}}_{1,n}^{sns}$ and $\widetilde{\mathcal{M}}_{\infty}(E_{1,n})$. Our treatment here is parallel to Section 4.5 of [22] so we will be brief. A slight difference of our case from the one considered in [22] is in the identification of the tangent space to $\widetilde{\mathcal{U}}_{1,n}^{sns}$ at $C = C_{1,n}$. By Lemma 1.1.1, a choice of a nonzero global 1-form is equivalent to a choice of a nonzero tangent vector to one of the marked points. Thus, we can identify the tangent space to $\widetilde{\mathcal{U}}_{1,n}^{sns}$ with $\text{Ext}_C^1(\mathbf{L}_C, \mathcal{O}(-D - p_i))$ for any i —these spaces are canonically isomorphic. In fact, by Lemma 2.2.4, these spaces are naturally isomorphic to $\text{Ext}_U^1(\mathbf{L}_U, \mathcal{O}_U)$.

Let Art_k denote the category of local Artinian (commutative) k -algebras with the residue field k . We are going to compare two deformation functors,

$$F_{1,n}, F_{\infty} : \text{Art}_k \rightarrow \text{Sets}.$$

Here $F_{1,n}(R)$ is the set of isomorphism classes of families $C \rightarrow \text{Spec } R$ with sections $p_1, \dots, p_n$, which reduce to $C_{1,n}$ upon the specialization $R \rightarrow k$. Note that this is nothing but the fibre of $\widetilde{\mathcal{U}}_{1,n}^{sns}(R) \rightarrow \widetilde{\mathcal{U}}_{1,n}^{sns}(k)$ over the point corresponding to $C_{1,n}$.

Similarly, we define $F_{\infty}(R)$ as the fibre of $\widetilde{\mathcal{M}}_{\infty}(E_{1,n})(R) \rightarrow \widetilde{\mathcal{M}}_{\infty}(E_{1,n})(k)$ over the class of the trivial A_{∞} -structure. By Lemma 4.5.1 (i) of [22], these correspond to equivalence classes of minimal R -linear A_{∞} -structures on $E_{1,n}$, such that upon specialization $R \rightarrow k$ we get a formal A_{∞} -algebra, i.e. the A_{∞} -structure that is gauge equivalent to the trivial one.

Recall that the map

$$\widetilde{\mathcal{U}}_{1,n}^{sns}(k) \rightarrow \widetilde{\mathcal{M}}_{\infty}(E_{1,n})(k)$$

(see (2.1.2)) sends the point corresponding to $C_{1,n}$ to the class of the trivial A_{∞} -structure (see Lemma 2.1.2(ii)). Thus, we can consider the induced map

$$(2.2.2) \quad F_{1,n} \rightarrow F_{\infty}$$

of deformation functors on Art_k .

Proposition 2.2.5. *Assume that either $n \geq 3$, or $n = 2$ and $\text{char}(k) \neq 2$ (resp., $n = 1$ and $\text{char}(k) \neq 2, 3$). Then the morphism of deformation functors (2.2.2) is an isomorphism, and the tangent space to F_{∞} is naturally isomorphic to*

$$HH^2(E_{1,n})_{<0} = HH^2(E_{1,n})$$

Proof. The proof follows exactly the same line of argument as given in Prop. 4.5.4 of [22], using Corollary 2.2.2, Lemma 2.2.4 and Lemma 2.2.3 (the latter is needed to deduce that the functor F_{∞} is prorepresentable, hence, homogeneous). Note that, as we

observed above, the tangent space and obstruction space to $F_{1,n}$ can be identified with $\text{Ext}_C^1(\mathbf{L}_C, \mathcal{O}(-D - p_i))$ and $\text{Ext}_C^2(\mathbf{L}_C, \mathcal{O}(-D - p_i))$, respectively. $\square$

In particular, this leads to a computation of $HH^2(E_{1,n})$. Similarly to [22, Prop. 4.7.2] we can identify $HH_{<0}^2(E_{1,n})$ with the tangent space of the moduli scheme $\widetilde{\mathcal{U}}_{1,n}^{sns}$ at zero. Furthermore, since (2.1.2) is compatible with the $\mathbb{G}_m$ -action, this is a graded identification, hence, using Corollary 1.1.6 (resp. Theorem 1.4.2 for $n \leq 4$) we get the ranks of $HH^2(E_{1,n})$.

Corollary 2.2.6. *Over an arbitrary field k , we have for $n \geq 5$*

$$HH^2(E_{1,n}) = k^{(n-1)(n-2)/2}[1]$$

For $n = 3, 4$ we have:

$$HH^2(E_{1,4}) = k^3[1] \oplus k^2[2],$$

$$HH^2(E_{1,3}) = k[1] \oplus k^2[2] \oplus k[3].$$

For a field k with $\text{char}(k) \neq 2$,

$$HH^2(E_{1,2}) = k[2] \oplus k[3] \oplus k[4].$$

For a field k with $\text{char}(k) \neq 2$ or 3,

$$HH^2(E_{1,1}) = k[4] \oplus k[6].$$

$\square$

Remark 2.2.7. It was shown in [17] that for a field k with $\text{char}(k) = 2$ one has

$$HH^2(E_{1,1}) = k[1] \oplus k[3] \oplus k[4] \oplus k[6],$$

while for a field k with $\text{char}(k) = 3$ one has

$$HH^2(E_{1,1}) = k[2] \oplus k[4] \oplus k[6].$$

Using the methods of this paper (or computing using an explicit resolution as in [17]) one can show that for a field k with $\text{char}(k) = 2$ one has

$$HH^2(E_{1,2}) = k[1] \oplus k[2] \oplus k[3] \oplus k[4].$$

Finally, using the $\mathbb{G}_m$ -action as in the proof of Theorem A of [22], we deduce our second main result.

Theorem 2.2.8. *For $n \geq 3$, the map (2.1.2) induces an isomorphism of the moduli scheme $\widetilde{\mathcal{U}}_{1,n}^{sns}$ with the moduli scheme of minimal A_∞ -structures on $E_{1,n}$, up to a gauge equivalence. This isomorphism is compatible with the natural $\mathbb{G}_m$ -actions. The same conclusion holds for $n = 2$ and $\text{char}(k) \neq 2$ (resp., $n = 1$ and $\text{char}(k) \neq 2, 3$). $\square$*

As a consequence, we get an interpretation of the moduli space $\overline{\mathcal{M}}_{1,n}^\infty$ in terms of A_∞ -structures.

Corollary 2.2.9. *Under the assumptions of Theorem 2.2.8, the quotient stack*

$$(\widetilde{\mathcal{U}}_{1,n}^{sns} \setminus \{C_{1,n}\})/\mathbb{G}_m = \overline{\mathcal{M}}_{1,n}^\infty$$

is isomorphic to the moduli stack of non-formal minimal A_∞ -structures on $E_{1,n}$, up to gauge equivalence and rescaling, or equivalently up to an A_∞ -equivalence of A_∞ -structures over k^{n+1} .

Proof. We observe that all automorphisms of $E_{1,n}$ as an associative algebra over k^{n+1} have form

$$A_i \mapsto \lambda_i A_i, \quad B_i \mapsto \lambda \cdot \lambda_i^{-1} B_i,$$

for some invertible constants λ, λ_i . It is easy to check that the effect of such transformations on μ^n is exactly the rescaling $\mu^n \mapsto \lambda^{n-2} \mu^n$. $\square$

Remark 2.2.10. Note that for every subset $S \subset \{1, \dots, n\}$ we have a natural subquiver in Q_n such that the corresponding subalgebra is isomorphic to $E_{1,|S|}$. In particular, we have n subquivers $Q_{n-1}(i) \subset Q_n$ (where $i = 1, \dots, n$) that give embeddings of $E_{1,n-1}$ into $E_{1,n}$. Now given a minimal A_∞ -structure $\mu^\bullet$ on $E_{1,n}$, for each i we have a well defined restriction $\mu^\bullet|_{Q_{n-1}(i)}$, which is a minimal A_∞ -structure on $E_{1,n-1}$ (recall that we consider A_∞ -structures that are unital with respect to the idempotents in $E_{1,n}$). Therefore, we get maps

$$\tilde{\pi}_i : \widetilde{\mathcal{M}}_\infty(E_{1,n}) \rightarrow \widetilde{\mathcal{M}}_\infty(E_{1,n-1})$$

for $i = 1, \dots, n$. Under the isomorphism (2.1.2), the map $\tilde{\pi}_i$ corresponds to forgetting the marked point p_i (over some open locus including the smooth curves). Thus, for $i = n$ this morphism can be identified with the projection $U_n \rightarrow U_{n-1}$ of Proposition 1.1.5 (ii).

Next, we observe that by Proposition 2.1.3, we have that $HH^3(E_{1,n}) \simeq HH^3(C_{1,n})$ is finite-dimensional. Therefore, by [22, Cor. 4.2.6], there is a natural isomorphism of functors

$$\widetilde{\mathcal{M}}_\infty(E_{1,n}) \rightarrow \widetilde{\mathcal{M}}_d(E_{1,n})$$

for all $d \geq N(n)$ for some sufficiently large $N(n)$, where the forgetful map is given by

$$(\mu^i)_{3 \leq i} \rightarrow (\mu^i)_{3 \leq i \leq d}$$

In other words, the inverse limit $\varprojlim_d \widetilde{\mathcal{M}}_d(E_{1,n})$ stabilizes. We next determine the exact value of $N(n)$ for all n .

Theorem 2.2.11. *Over an arbitrary field k , we have for $n \geq 4$,*

$$\widetilde{\mathcal{M}}_\infty(E_{1,n}) \simeq \widetilde{\mathcal{M}}_d(E_{1,n}) \quad \text{for all } d \geq 4.$$

For $n = 3$, we have

$$\widetilde{\mathcal{M}}_\infty(E_{1,3}) \simeq \widetilde{\mathcal{M}}_d(E_{1,3}) \quad \text{for all } d \geq 5.$$

For $n = 2$, over a field k with $\mathbf{char}(k) \neq 2$, we have

$$\widetilde{\mathcal{M}}_\infty(E_{1,2}) \simeq \widetilde{\mathcal{M}}_d(E_{1,2}) \text{ for all } d \geq 6.$$

For $n = 1$, over a field k with $\mathbf{char}(k) \neq 2$ or 3 , we have

$$\widetilde{\mathcal{M}}_\infty(E_{1,1}) \simeq \widetilde{\mathcal{M}}_d(E_{1,1}) \text{ for all } d \geq 8.$$

Proof. Let $n \geq 5$. Abusing the notation we denote $U_n \times \mathrm{Spec}(k)$ simply as U_n . By Corollary 1.1.6, we know that

$$\Theta := \{a, (c_i, \bar{c}_i)_{4 \leq i \leq n}, (c_{ij}, \bar{c}_{ij})_{4 \leq i < j \leq n}\}$$

is a set of minimal generators of the algebra of functions on the affine scheme U_n , where $a, c_i, \bar{c}_i$ and $c_{ij}, \bar{c}_{ij}$ have degree 1 with respect to the natural $\mathbb{G}_m$ action. Furthermore, we have $k[U_n] = k[\Theta]/I$ where the ideal I is generated by quadratic relations. Now, the argument given in the proof of [22, Prop. 4.7.2] shows that the algebra of functions on $\widetilde{\mathcal{M}}_d(E_{1,n})$ is isomorphic

$$k[\Theta_{\leq d-2}]/I_{\leq d-2},$$

where $\Theta_{\leq d-2}$ (resp., $I_{\leq d-2}$) is the set of elements in Θ (resp., I) of degree $\leq d-2$. It follows that $\widetilde{\mathcal{M}}_\infty(E_{1,n}) \simeq \widetilde{\mathcal{M}}_d(E_{1,n})$ for $n \geq 5$ and $d \geq 4$.

The proof in the remaining cases is similar. Again we need to determine the algebra of functions on U_n in these cases. These have been worked out in Proposition 1.1.5 in the cases $n = 3, 4$. The assertion follows as above, since $k[U_4]$ (resp., $k[U_3]$) is free with generators of degrees ≤ 2 (resp., ≤ 3).

The case of $n = 2$ is worked out in Section 1.2 for $\mathbf{char}(k) \neq 2$: we get that $k[U_2] = \mathbb{A}^3$ with generators α, β and γ of degrees 2, 3 and 4, respectively. Hence, we have $\widetilde{\mathcal{M}}_\infty(E_{1,2}) \simeq \widetilde{\mathcal{M}}_d(E_{1,2})$ for all $d \geq 6$.

Finally, the case $n = 1$ was worked out in [17, Prop. 9] for $\mathbf{char}(k) \neq 2$ or 3 . (The general case is also studied in [16]). $\square$

Remark 2.2.12. We note that over a field k , the functor $\widetilde{\mathcal{M}}_3(E_{1,n})$ associating with R the set of gauge equivalence classes of minimal A_3 -structures on $E_{1,n} \otimes R$ is represented by the affine space $HH^2(E_{1,n})_{-1}$ (see [22, Thm. 4.2.4]). Assume that $n \geq 5$. By Corollary 2.2.6, it follows that

$$\widetilde{\mathcal{M}}_3(E_{1,n}) \simeq \mathbb{A}^{(n-1)(n-2)/2}$$

On the other hand, Theorem 1.4.2 together with Theorem 2.2.8 identifies the moduli scheme of A_∞ -structures on $E_{1,n}$ up to a gauge equivalence with the affine scheme

$$(2.2.3) \quad U_n \hookrightarrow \mathbb{A}^{(n-1)(n-2)/2}$$

In fact, this embedding can be identified with the natural map

$$\widetilde{\mathcal{M}}_\infty(E_{1,n}) \hookrightarrow \widetilde{\mathcal{M}}_3(E_{1,n})$$

which sends $(\mu^i)_{i \geq 3}$ to μ^3 by simply forgetting the higher products. Finally, note that if we interpret $U_n \simeq \widetilde{\mathcal{U}}_{1,n}^{sns}$ as moduli of curves then we can view (2.2.3) as an analog of the rational map from a $\mathbb{G}_m^g$ -torsor over $\mathcal{M}_{g,g}$ to $\mathbb{A}^{g^2-g}$, defined in [9] in terms of triple products.

REFERENCES

- [1] D. Abramovich, M. Olsson, A. Vistoli, *Tame stacks in positive characteristic* Ann. Inst. Fourier 58 (2008), 1057–1091.
- [2] M. A. Akivis, V. V. Goldberg, *Conformal Differential Geometry and Its Generalizations*, John Wiley & Sons, Inc., New York, 1996.
- [3] L. L. Avramov, H.-B. Foxby, *Locally Gorenstein Homomorphisms*, Amer. J. Math. 114 (1992), no. 5, 1007–1047.
- [4] D. A. Buchsbaum, D. Eisenbud, *Gorenstein ideals of height 3*, in *Seminar D. Eisenbud/B. Singh/W. Vogel*, Vol. 2, pp. 30–48, Teubner, Leipzig, 1982.
- [5] A. Bondal, D. Orlov, *Semiorthogonal decompositions for algebraic varieties*, preprint alg-geom/9506012.
- [6] G. M. Bergman, *The diamond lemma for ring theory*. Adv. in Math. 29 (1978), no. 2, 178–218.
- [7] I. Coskun, A. Prendergast-Smith, *Fano manifolds of index $n - 1$ and the cone conjecture*, Internat. Math. Res. Notices 2014, no. 9, 2401–2439.
- [8] P. Deligne, D. Mumford, *The irreducibility of the space of curves of given genus*, Inst. Hautes Études Sci. Publ. Math. No. 36 1969 75–109.
- [9] R. Fisette, A. Polishchuk, *A_∞ -algebras associated with curves and rational functions on $M_{g,g}$. I*. Compos. Math. 150 (2014), no. 4, 621–667.
- [10] T. Fisher, *Pfaffian presentations of elliptic normal curves*, Trans. AMS 362 (2010), 2525–2540.
- [11] T. Fujita, *On the structure of polarized manifolds with total deficiency one, II*, J. Math. Soc. Japan 33 (1981), no. 3, 415–434.
- [12] R. Hartshorne, *Residues and duality*, LNM 20, Springer-Verlag, 1966.
- [13] B. Hassett, *Classical and minimal models of the moduli space of curves of genus two*, in *Geometric methods in algebra and number theory*, Progr. Math. 235, 169–192. Birkhäuser, Boston, MA, 2005.
- [14] M. Kontsevich, Y. Soibelman, *Homological mirror symmetry and torus fibration*, in *Symplectic geometry and mirror symmetry (Seoul, 2000)*, 203–263, World Sci. Publishing, River Edge, NJ, 2001.
- [15] S. J. Kovacs, *A characterization of rational singularities*, Duke Math. J. 102 (2000), 187–191.
- [16] Y. Lekili, T. Perutz, *Arithmetic mirror symmetry for the 2-torus*, preprint arXiv:1211.4632.
- [17] Y. Lekili, T. Perutz, *Fukaya categories of the torus and Dehn surgery*, Proc. Natl. Acad. Sci. USA 108 (2011), no. 20, 8106–8113.
- [18] Y. Lekili, A. Polishchuk, *Arithmetic mirror symmetry for genus 1 curves with n marked points*, in preparation.
- [19] H. Matsumura, *Commutative ring theory*, Cambridge Univ. Press, 1986.
- [20] S. Merkulov, *Strong homotopy algebras of a Kähler manifold*, Internat. Math. Res. Notices 1999, no. 3, 153–164.
- [21] D. Mumford, *Stability of projective varieties*, L’Ens. Math. 23 (1977), no. 1-2, 39–110.
- [22] A. Polishchuk, *Moduli of curves as moduli of A_∞ -structures*, preprint arXiv:1312.4636.
- [23] A. Polishchuk, *Moduli of curves with nonspecial divisors and relative moduli of A_∞ -structures*, arXiv:1511.03797.
- [24] P. Seidel, *Fukaya categories and Picard-Lefschetz theory*, European Mathematical Society, Zürich, 2008.
- [25] D. I. Smyth, *Towards a classification of modular compactifications of $\mathcal{M}_{g,n}$* , Invent. Math. 192 (2013), no. 2, 459–503.

- [26] D. I. Smyth, *Modular compactifications of the space of pointed elliptic curves I*, Compos. Math. 147 (2011), no. 3, 877–913.
- [27] D. I. Smyth, *Modular compactifications of the space of pointed elliptic curves II*, Compos. Math. 147 (2011), no. 6, 1843–1884.
- [28] D. J. Wright, A. Yúkie, *Prehomogeneous vector spaces and field extensions*, Invent. Math. 110 (1992), no. 2, 283–314.

KING’S COLLEGE LONDON

UNIVERSITY OF OREGON