King's Research Portal

DOI:

[10.1016/j.cct.2017.05.011](https://doi.org/10.1016/j.cct.2017.05.011)

Document Version

Peer reviewed version

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Landoni, G., Lomivorotov, V., Pisano, A., Neto, C. N., Benedetto, U., Zoccai, G. B., Gemma, M., Frassoni, S., Agrò, F. E., Baiocchi, M., Barbosa Gomes Galas, F. R., Bautin, A., Bradic, N., Carollo, C., Crescenzi, G., Elnakera, A., El-Tahan, M. R., Fominskiy, E., Farag, A. G., ... Zangrillo, A. (2017). Mortality in caRdIAC surgery (MYRIAD): A randomized controlled trial of volatile anesthetics. Rationale and design. *CONTEMPORARY CLINICAL TRIALS*. <https://doi.org/10.1016/j.cct.2017.05.011>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Mortality in cardiac surgery (MYRIAD): A randomized controlled trial of volatile anesthetics. Rationale and design

Giovanni Landoni, Vladimir Lomivorotov, Antonio Pisano, Caetano Nigro Neto, Umberto Benedetto, Giuseppe Biondi Zoccai, Marco Gemma, Samuele Frassoni, Felice Eugenio Agrò, Massimo Baiocchi, Filomena R. Barbosa Gomes Galas, Andrei Bautin, Nikola Bradic, Cristiana Carollo, Giuseppe Crescenzi, Abeer Elnakera, Mohamed R. El-Tahan, Evgeny Fominskiy, Ahmed G. Farag, Gordana Gazivoda, Stefano Gianni, Evgeny Grigoryev, Fabio Guarracino, Sidik Hanafi, Weiqing Huang, Gudrun Kunst, Jan Kunstyr, Chong Lei, Rosalba Lembo, Zhi-jian Li, Valery Likhvantsev, Andrey Lozovskiy, Jun Ma, Fabrizio Monaco, Paolo Navalesi, Bukamal Nazar, Vadim Pasyuga, Elena Porteri, Colin Royse, Laura Ruggeri, Hynek Ryha, Fernanda Santos Silva, Luca Severi, Vladimir Shmyrev, Nikolay Uvaliev, Cheng Bin Wang, Chew-Yin Wang, Dario Winterton, Chow-Yen Yong, Jingui Yu, Rinaldo Bellomo, Alberto Zangrillo

PII: S1551-7144(16)30437-2
DOI: doi: [10.1016/j.cct.2017.05.011](https://doi.org/10.1016/j.cct.2017.05.011)
Reference: CONCLI 1565

To appear in: *Contemporary Clinical Trials*

Received date: 5 December 2016
Revised date: 14 May 2017
Accepted date: 17 May 2017

Please cite this article as: Giovanni Landoni, Vladimir Lomivorotov, Antonio Pisano, Caetano Nigro Neto, Umberto Benedetto, Giuseppe Biondi Zoccai, Marco Gemma, Samuele Frassoni, Felice Eugenio Agrò, Massimo Baiocchi, Filomena R. Barbosa Gomes Galas, Andrei Bautin, Nikola Bradic, Cristiana Carollo, Giuseppe Crescenzi, Abeer Elnakera, Mohamed R. El-Tahan, Evgeny Fominskiy, Ahmed G. Farag, Gordana Gazivoda, Stefano Gianni, Evgeny Grigoryev, Fabio Guarracino, Sidik Hanafi, Weiqing Huang, Gudrun Kunst, Jan Kunstyr, Chong Lei, Rosalba Lembo, Zhi-jian Li, Valery Likhvantsev, Andrey Lozovskiy, Jun Ma, Fabrizio Monaco, Paolo Navalesi, Bukamal Nazar, Vadim Pasyuga, Elena Porteri, Colin Royse, Laura Ruggeri, Hynek Ryha, Fernanda Santos Silva, Luca Severi, Vladimir Shmyrev, Nikolay Uvaliev, Cheng Bin Wang, Chew-Yin Wang, Dario Winterton, Chow-Yen Yong, Jingui Yu, Rinaldo Bellomo, Alberto Zangrillo, Mortality in cardiac surgery (MYRIAD): A randomized controlled trial

of volatile anesthetics. Rationale and design. The address for the corresponding author was captured as affiliation for all authors. Please check if appropriate. Concli(2017), doi: [10.1016/j.cct.2017.05.011](https://doi.org/10.1016/j.cct.2017.05.011)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Mortality in cardiac surgery (MYRIAD): A randomized controlled trial of volatile anesthetics. Rationale and design

Running head

Volatile anesthetics in cardiac surgery

Authors

Giovanni Landoni, MD, Prof¹; Vladimir Lomivorotov², MD, PhD; Antonio Pisano, MD³; Caetano Nigro Neto, MD⁴, PhD; Umberto Benedetto, MD, PhD⁵; Giuseppe Biondi Zoccai, MD, MStat⁶; Marco Gemma, MD¹; Samuele Frassoni, MSc¹; Felice Eugenio Agrò, Full Professor⁷; Massimo Baiocchi, MD⁸; Filomena R Barbosa Gomes Galas, MD, PhD⁹; Andrei Bautin, MD, PhD¹⁰; Nikola Bradic, MD¹¹; Cristiana Carollo, MD¹²; Giuseppe Crescenzi, MD¹³; Abeer Elnakera, Assistant Professor¹⁴; Mohamed R El-Tahan, Associate Professor¹⁵; Evgeny Fominskiy, MD, PhD¹⁶; Ahmed G. Farag, MD¹⁷; Gordana Gazivoda, MD¹⁸; Stefano Gianni, MD¹; Evgeny Grigoryev, MD, PhD¹⁹; Fabio Guarracino, MD²⁰; Sidik Hanafi, MD²¹; Weiqing Huang, Full Professor²²; Gudrun Kunst, MD²³; Jan Kunstyr, Associate Professor²⁴; Chong Lei, MD²⁵; Rosalba Lembo, MSc¹; Zhi-jian Li, Associate Professor²⁶; Valery Likhvantsev, MD²⁷, PhD, Professor; Andrey Lozovski, MD²⁸; Jun Ma, Full Professor²⁹; Fabrizio Monaco, MD¹; Paolo Navalesi, Full Professor³⁰; Bukamal Nazar, MD, Bch, FRCA³¹; Vadim Pasyuga, MD³²; Elena Porter, MD³³; Colin Royse, Full Professor³⁴; Laura Ruggeri, MD¹; Hynek Ryha, Assistant Professor³⁵; Fernanda Santos Silva, MD³⁶; Luca Severi, MD³⁷; Vladimir Shmyrev, MD, PhD²; Nikolay Uvaliev, MD³⁸; Cheng Bin Wang, Associate professor²⁹; Chew-Yin Wang, Full Professor³⁹; Dario Winterton, MD¹; Chow-Yen Yong, MD⁴⁰; Jingui Yu, Full Professor⁴¹; Rinaldo Bellomo, Full Professor⁴² Alberto Zangrillo, Full Professor¹

Affiliations

1 Department of Anesthesia and Intensive Care, IRCCS San Raffaele Scientific Hospital, via Olgettina 60 Milan, 20132 Italy

2 Siberian Biomedical Research Center, Department of Anesthesiology and Intensive Care, Novosibirsk, Russia

3 Cardiac Anesthesia and Intensive Care Unit, Monaldi Hospital A.O.R.N. "Dei Colli", via L. Bianchi Naples, 80131 Italy

4 Anesthesia Section of Cardiothoracic Surgery, Dante Pazzanese Institute of Cardiology, Sao Paulo, Brazil

5 University of Bristol, School of Clinical Sciences, Bristol Heart Institute, Bristol, United Kingdom

6 Department of Medico-Surgical Sciences and Biotechnologies, Sapienza University of Rome, Latina, and Department of AngioCardioNeurology, IRCCS Neuromed, Pozzilli, both in Italy

7 Anaesthesia, Intensive Care and Pain Management Department, University School of Medicine Campus Bio-Medico, Rome, Italy

8 Anesthesia and Intensive Care, S.Orsola-Malpighi university hospital, Bologna, Italy

9 Anesthesia and Intensive care, Heart Institute, University of Sao Paulo, Sao Paulo, Brazil

10 Head of the Research Department of Anesthesiology and Intensive Care, Federal Almazov North-West Medical Research Centre, Saint-Petersburg, Russia

11 Anesthesia and Intensive Care, University Hospital, Dubrava Croazia

12 Department of Medicine, Anesthesia and Intensive Care Unit, Padova University Hospital, Via Cesare Battisti 256, 35128 Padova

13 Anestesia e Terapia Intensiva Cardiochirurgica, Humanitas Research Hospital, Rozzano, Italy

14 Anesthesia and intensive care, Faculty of Medicine - Zagazig university, Zagazig, Egypt

15 Cardiothoracic Anaesthesia & Surgical Intensive Care, King Fahd Hospital of the University of Dammam, Al Khobar; Saudi Arabia

16 Department of Anaesthesia and Intensive Care, Academician EN Meshalkin Novosibirsk State Budget Research Institute of Circulation Pathology, Novosibirsk, Russia

17 Department of Anaesthesia and Intensive Care, King Abdullah Medical City "KAMC", Makkah, Saudi Arabia

18 Department of Anesthesiology and Intensive Care, Cardiovascular Institute 'Dedinje', Belgrade, Serbia

19 Deputy director for clinical and scientific affairs, Scientific Research Institute of Complex Issues of Cardiovascular Disease, Kemerovo, Russia

20 Department of Anaesthesia and Critical Care Medicine, Azienda Ospedaliero-Universitaria Pisana, Via Paradisa 2, 56123 Pisa, Italy

21 Department of Anaesthesiology and Intensive Care, Sarawak General Hospital, Kuching, Sarawak, Malaysia

22 Department of Anesthesiology, Wuhan Asia Heart Hospital, Wuhan, China

23 Department of Anaesthetics, King's College Hospital NHS Foundation Trust, London, United Kingdom

24 Department of Anesthesiology and Intensive Care, First Faculty of Medicine, Charles University in Prague and General University Hospital in Prague, Prague, Czech Republic

25 Department of Anesthesiology, 1st Affiliated Hospital of Fourth Military University, Xi'an, China

26 Department of Anesthesiology, The Second Xiangya Hospital of Central South University, Changsha, China

27 Anesthesiology e Intensive Care Department, Moscow Clinical Regional Research Institute, Shepkina st., 61/2, Moscow, Russia.

28 Department of Anesthesiology and Intensive Care, Ural Institute of Cardiology, Ekaterinburg, Russia

29 Center for Anesthesiology, Beijing Anzhen Hospital, Capital Medical University, Beijing, China

30 Anestesia e Rianimazione, Dipartimento Scienze Mediche e Chirurgiche, Università della Magna Graecia, Catanzaro, Italy

31 Cardiothoracic ICU and Anaesthesia, MKCC, West Riffa, Bahrain

32 Department of Cardiac Aneasthesia and Intensive Care, Astrakhan Federal Centre for Cardiac Surgery, Astrakhan, Russia

33 Anesthesia and Intensive Care, Azienda Ospedaliera Spedali Civili, Brescia, Italy

34 Department of Surgery Level 6 Clinical Medical Research Building, The University of Melbourne, Melbourne, Australia

35 Cardiothoracic Anesthesiology and Intensive Care, Institute for Clinical and Experimental Medicine, Prague, Czech Republic

36 Anesthetic Department, Hospital de Santa Maria, Lisboa, Portugal

37 Anesthesia and Intensive Care, Azienda ospedaliera San Camillo Forlanini, Roma, Italy

38 Department-anesthesia and intensive care, City Clinic Burgas, Burgas-Bulgaria

39 Anaesthesia and Intensive Care, University of Malaya, Kuala Lumpur, Malaysia

40 Anaesthesia and Intensive Care, Hospital Pulau Pinang, Georgetown, Malaysia

41 Department of Anesthesiology, Qilu Hospital of Shandong University, Shandong, China

42 The University of Melbourne, Melbourne, Victoria, Australia

Landoni Giovanni, MD

Department of Anesthesia and Intensive Care

San Raffaele Hospital

Via Olgettina 60 Milan, 20132 Italy

Phone: ++390226434524

Fax ++390226437178

Email: landoni.giovanni@hsr.it

ABSTRACT

Objective: There is initial evidence that the use of volatile anesthetics can reduce the postoperative release of cardiac troponin I, the need for inotropic support, and the number of patients requiring prolonged hospitalization following coronary artery bypass graft (CABG) surgery. Nevertheless, small Randomized Controlled Trials have failed to demonstrate a survival advantage. Thus, whether volatile anesthetics improve the postoperative outcome of cardiac surgical patients remains uncertain. An adequately powered randomized controlled trial appears desirable.

Design: Single blinded, international, multicenter randomized controlled trial with 1:1 allocation ratio.

Setting: Tertiary and University hospitals.

Interventions: Patients (n= 10,600) undergoing coronary artery bypass graft will be randomized to receive either volatile anesthetic as part of the anesthetic plan, or total intravenous anesthesia.

Measurements and main results: The primary end point of the study will be one-year mortality (any cause). Secondary endpoints will be 30-day mortality; 30-day death or non-fatal myocardial infarction (composite endpoint); cardiac mortality at 30 day and at one year; incidence of hospital re-admission during the one year follow-up period and duration of intensive care unit, and hospital stay. The sample size is based on the hypothesis that volatile anesthetics will reduce 1-year unadjusted mortality from 3% to 2%, using a two-sided alpha error of 0.05, and a power of 0.9.

Conclusions: The trial will determine whether the simple intervention of adding a volatile anesthetic, an intervention that can be implemented by all anesthesiologists, can improve one-year survival in patients undergoing coronary artery bypass graft surgery.

KEY WORDS:

Volatile anesthetics, Total intravenous anesthesia, Cardiac anesthesia, Cardiac surgery, Randomized trial, Intensive Care

INTRODUCTION

Coronary artery disease (CAD) remains one of the most common causes of death and significantly affects the use of health care resources. In the United States alone, it results in more than 397,000 Coronary Artery Bypass Grafting (CABG) surgeries per year.¹ There is initial evidence that the choice of anesthesia can influence survival in the specific setting of CABG. An international consensus conference considered volatile anesthetics among the few drugs, techniques or strategies that might reduce perioperative mortality in cardiac surgery and should be further studied.²

All volatile anesthetics have cardiac depressant effects but also a beneficial effect on the myocardial oxygen balance during ischemia.^{3,4} Animal trials have shown that volatile anesthetics can provide protection against the ischemia-reperfusion injury that occurs during cardiac surgery via preservation of mitochondrial function and improved cell survival.²² Furthermore volatile anesthetics reduce the inflammatory response seen in acute lung injury,^{5,6} and after brain,⁷ liver⁸ and kidney⁹ ischemia with associated clinical benefits. The most recent and comprehensive meta-analysis¹⁰ of 68 randomized controlled trials (RCTs) included 7,104 patients and compared volatile anesthetics with total intra-venous anesthesia (TIVA). It showed that in cardiac surgery volatile anesthetics are associated with reduced overall mortality (OR 0.55, 95% CI 0.35–0.85, $p=0.007$), reduced pulmonary complications (OR 0.71, 95% CI 0.52–0.98, $p=0.038$) and a reduction in other complications (OR 0.74, 95% CI 0.58–0.95, $p=0.020$). The largest multicenter study published so far on this topic by De Hert et al.¹¹ randomized 414 participants undergoing on-pump CABG. In this study one-year mortality was a secondary outcome and was different among groups (12.3% in the TIVA group, 6.9% in the Desflurane group, and 3.3% in the Sevoflurane group; $p=0.034$). A further recently published RCT suggested a mortality reduction at 1 year in the sevoflurane group when compared to the TIVA group¹². A previous large observational study¹³ and a long-term follow-up of a RCT¹⁴ found that the occurrence of complications in the postoperative period was an independent predictors of long-term survival, supporting the notion that perioperative organ

protection and reduction in postoperative complications with volatile anesthetics could decrease long-term mortality.

Therefore, we aim to carry out the MYRIAD trial (Mortality in caRdIac surgery. A randomized controlled trial of volatile anesthetics in cardiac surgery) a large multicenter RCT to identify whether a clinically important reduction in one-year mortality from 3% to 2% after CABG surgery can be achieved by including a volatile anesthetic as part of the overall anesthetic in patients receiving TIVA or volatile agents.

MATERIALS AND METHODS

Study design, approval, and registration

The planned study is a parallel group, randomized controlled, single-blinded multicenter trial with 1:1 allocation ratio. The study has been approved by the Human Research Ethics Committee of all the participating centers and is registered on clinicaltrials.gov as NCT02105610.

Study aim

The aim of our study is to test the hypothesis that volatile anesthetics can reduce one-year mortality from 3% to 2% in participants undergoing CABG, either with or without cardiopulmonary bypass (CPB).

Participants

We plan to enroll 10,600 participants with CAD undergoing elective CABG. Participants will be >18 years and undergoing scheduled isolated CABG (including multiple coronary artery bypass). The exclusion criteria are presented in Table 1.

Inclusion Criteria	Exclusion Criteria
Age > 18 years	Planned: <ul style="list-style-type: none"> - Valve surgery - Surgery of the aorta
Scheduled procedure	Unstable/ongoing angina
Isolated coronary artery bypass graft	Acute myocardial infarction (<1 month)
	Use of: <ul style="list-style-type: none"> - Sulfonylurea - Theophylline - Allopurinol
	Previous unusual response to an anesthetic agent
	Inclusion in other randomized controlled studies in the previous 30 days
	Any general anesthesia performed in the previous 30 days
	Emergency operation

	Kidney or liver transplant in medical history
	Liver cirrhosis (Child B or C)

Table 1: MYRIAD inclusion and exclusion criteria

Randomization, allocation and concealment

Subjects will be allocated according to a web-based centralized randomization service or by opening centrally provided sealed opaque envelopes with the use of a permuted-block design stratified according to center. We will use randomization blocks of 20. Patients will be unaware of group assignments. Anesthesiologists will provide the trial treatment intervention and, as a consequence, will know patients' group allocation but will not be involved in postoperative treatment, data collection, data entry or data analysis. Investigators and clinical personnel caring for patients, including intensive care physicians, will be blinded to the study drug for the duration of the trial. Data will be collected by trained observers who will not participate in patient care and will be blinded to patient allocation.

Interventions

Participants will be randomized to receive either anesthesia which includes a volatile agent or TIVA alone (Fig.1).

The volatile group will receive desflurane, isoflurane or sevoflurane to provide general anesthesia in addition to any intravenous agent (according to local protocols and expertise). The volatile agent will be administered for as long as possible (ideally from anesthesia induction to ICU sedation) and at the highest concentration permitted by local protocols and patient hemodynamics during at least one of the following time periods: anesthesia induction, pre-CPB, during CPB, after CPB in the surgical theatre, and in the ICU.

Within the volatile group the following strategies are strongly suggested but not mandatory: a) at least 1 minimal alveolar concentration (MAC) for at least 30 minutes (this is the minimum

concentration demonstrated to be cardioprotective in experimental studies); b) discontinuation of the volatile agent for at least 15 minutes before CPB (a wash-out period before ischemia seems to be a prerequisite for the preconditioning phenomenon); c) wash-in/wash-out periods, defined as: volatile administration (at least 0.5 MAC) for at least three periods of 10 minutes, interspersed by wash-out periods of 10 minutes or more.

The aim of these strategies is to enhance the cardio-protective properties of volatile agents^{23,24} without significantly modifying the local protocols and without affecting patient safety.

The TIVA group will receive any intravenous agent and no volatile agent. Agents for TIVA will be administered as both target-controlled infusions or manually controlled infusions according to local protocols and expertise.

In case of repeated operation during the first hospitalization the patients will follow the study allocated anesthesia.

This is a pragmatic study. Accordingly, we have chosen not to require or define a strict anesthetic protocol. This allows all patients to be treated according to the best available treatment available in each center. We consider that this approach adds external validity to future findings.

CABG: Coronary artery bypass graft; TIVA: Total intravenous anesthesia.

Figure 1: MYRIAD Flow chart.

All participants will receive perioperative intensive treatment according to their institutional practice, including general anesthesia, pacing, inotropic drugs, mechanical ventilation, postoperative sedation/analgesia, diuretics, intravenous fluids, antibiotics and invasive monitoring. Such treatment will include but not be limited to, invasive arterial pressure, electrocardiogram, central venous pressure, cardiac output, pulse oximetry, temperature, urine output, arterial blood

gases and frequent routine laboratory examinations. No additional intervention or laboratory examination will be performed on participants outside of institutional practice.

Data will be collected at the end of surgical intervention, at ICU discharge, and at hospital discharge. We will record data about dosage, timing and mode of administration of all drugs used for the anesthesia. Surgical characteristics including CPB and aortic cross-clamping duration will also be collected. With regards to volatile anaesthetic use, we will collect data on what agent was administered, what dose, for how long and at what time points (induction, before CPB or the start of the anastomosis, during CPB or anastomosis, after CPB or anastomosis, and in ICU). If available, we will be collecting the baseline creatinine value. In case of myocardial infarction we will be collecting the cardiac biomarker value. Follow-up at 30 days and at one year will focus on the adverse cardiac events, hospital readmissions and survival.

Outcomes

We hypothesize that volatile anesthetics will reduce one-year mortality from any cause in participants undergoing CABG surgery.

Secondary endpoints will include: 30-day all-cause mortality; 30-day non-fatal myocardial infarction and 30-day death (composite endpoint); cardiac mortality at 30 days and at one year; hospital re-admission during the follow-up period; ICU and hospital stay. We also collect the number of adverse events: stroke, delirium, postoperative cognitive impairment, acute renal failure, surgical revision for bleeding, high dose inotropic drugs and the use of intra-aortic balloon pump or other mechanical circulatory support. Definitions of the outcomes are presented in the Supplementary material. To perform 30-days and one-year follow-up telephone contact (patient and relatives) will be used. In case loss to follow-up by telephone, the following methods will be used to establish vital status at one year: contacting the patient's general practitioner, contacting the city municipality, and sending a letter to the home address of the patient.

Statistical analysis and sample size estimates

An epidemiologist with extensive experience in designing, conducting and analyzing clinical trials, not involved in patient management, and blinded to the assigned intervention will be responsible for the statistical analysis.

Data will be stored electronically via a web based CRF and analyzed using STATA (Stata Statistical Software: version 14, College Station, TX, USA). We will not apply any imputation for missing data. All data will be analyzed according to the intention-to-treat principle, beginning immediately after randomization.

Demographic and baseline disease characteristics will be summarized with the use of descriptive statistics. Categorical variables will be reported as absolute numbers and percentages. Unadjusted univariate analyses, to compare the two treatment groups, will be based on Chi-square or Fisher's exact test. Relative risks and 95% confidence intervals will be calculated by means of the two-by-two table method with the use of log-normal approximation. Continuous variables will be reported as mean $\pm$ standard deviation (SD) or median and interquartile range (IQR). Normality will be evaluated using visual histogram evaluation and a Q-Q plot. Between-group differences will be evaluated using the T test or Wilcoxon signed rank test, in accordance with normality of the distribution. A logistic regression model using a stepwise selection will be used to estimate the treatment effect and predictors of mortality. The pre-randomization clinical data and center will be entered into the model if their univariate p value is less than 0.1 and there is no correlation between them. Collinearity and overfitting will be assessed using a stepwise regression model and Pearson correlation test. The treatment group (volatile anaesthetics or TIVA) will be forced into the multivariate model. If the outcome event proves to be rare, a Poisson regression model will be used. A classic logistic regression will be performed with a consistent number of events and the number of covariates in the model will be decided based on the number of outcome events.

On the basis of the latest large RCTs comparing CABG versus percutaneous interventions^{15,16,17,18} or comparing different CABG techniques,¹⁹ we hypothesize a one-year mortality of 3% in the control (TIVA) group. Following the results of a recent meta-analysis²⁰ and one large retrospective observational study²¹ demonstrating reduced mortality with the use of volatile anesthetics, we hypothesize a reduction in mortality from 3 to 2% in the volatile anesthetic group. Sample-size calculation is based on Pearson's Chi-square test with a two-sided alpha error of 0.05 and 90% power. We calculated that we would need a sample size of 5,300 participants per group using the continuity correction resulting in the total study population of 10,600 patients. Despite recent literature on sample size calculation in multicenter RCTs²⁵, we were unable to use this novel method in our competitive-enrolment trial as we did not know a priori the number of participating centers and the number of patients per center. We therefore used the conservative approach described above, which results in a greater sample size.

An independent safety committee will perform three interim analyses after recruitment of 25% (n=2,650), 50% (n=5,300) and 75% (n=7,950) of patients. Data evaluation at each interim analysis will be based on the alpha spending function concept, according to Lan and De Mets²⁶, and will employ O'Brien-Fleming Z-test boundaries²⁷, which are very conservative early in the trial. For the first interim analysis the efficacy stopping rule would require an extremely low P value ($P < 0.000015$). For the second interim analysis $P < 0.003$ will be taken as efficacy stopping rule. For the third interim analysis $P < 0.02$ will be taken as efficacy stopping rule. Investigators will be kept blind to the interim analysis results.

The independent safety committee will also perform conditional power analyses in order to evaluate potential interruption for futility issues in the trial. Conditional power will be calculated by assuming that the proportion of outcomes will follow the observed trend.

Moreover, if during the first interim analysis the independent safety committee will observe a similar direction and magnitude of the study technique effect on 30-day and 1-year mortality, they

will be in the position to use 30-day mortality data to suggest study continuation or interruption in the following interim analyses.

All data analyses will be carried out according to a pre-established analysis plan. Because organ protection elicited by volatile anesthetics can be modulated by several clinical factors we pre-specified subgroups analyses that are summarized in Table 2.

The dosage and mode of anesthetic administration could be confounders in the trial. Therefore, we will also separately analyze: centers using volatile agent also during CPB; centers that routinely do not use propofol as induction agent in the volatile group; centers with a high overall mortality; centers that use volatile anesthetic throughout all the procedure (before CPB, during CPB and after CPB); centers using TIVA as main anesthetic technique agent before study initiation; centers using volatile agent as main anesthetic technique before study initiation.

Planned subgroup analyses	
Ejection Fraction <40%	
Patients with diabetes mellitus	
Age <60 years	
Previous:	
- Myocardial infarction	
- Vascular Surgery	
- Stroke or transient ischemic attack	
Chronic kidney disease	
Perioperative beta-blocker use	
On-pump CABG vs Off-pump CABG	
Bypass grafts ≥ 4	
Drug used as volatile agent:	
- Desflurane	
- Isoflurane	
- Sevoflurane	
Drug used as hypnotic agent:	
- Propofol	

- Midazolam
Drug used as opioid agent:
- Remifentanyl
- Fentanyl
- Sufentanyl
Volatile administration strategy:
- At least 1 MAC of volatile agent for 30 minutes
- 15 minutes wash-out period of volatile agent
- Planned wash-in/wash-out period of the volatile agent

CABG: Coronary artery bypass graft; MAC: Minimum alveolar concentration.

Table 2: Planned subgroups analyses. Ejection Fraction will be measured by echocardiography.

Monitoring of the study

Auditors will verify adherence to required clinical trial procedures and will confirm accurate data collection according to the Good Clinical Practice (GCP) guidelines. Study monitoring and follow-up protocols, from the initial set-up to final reporting, will be fulfilled according to current National and International requirements.

Ethical aspects

This is a randomized trial of different anesthesiological strategies that have been used for decades on hundreds of millions of participants. The incidence of adverse events such as malignant hyperthermia, allergy and propofol syndrome is negligible, and unavoidable if the patient has to undergo general anesthesia. Subjects will not experience any tangible additional risk because of the trial.

Trial data will be stored in an electronic database with no patient identifiers (a numeric code will be used).

Study initiation, timing, participating centers and source of funding

The study started after Ethical Committee approval from each contributing recruiting center. Consecutive participants who sign the written informed consent, aged 18 years or older will be enrolled. The study progress will be updated monthly. The first 4500 participants were randomized by April 2017 in 32 hospitals and 13 countries. The number of participating centers is continuously increasing, as no *a priori* limit to the number of participating centers has been established.

The authors are solely responsible for the design and conduct of this study, all study analyses and drafting and editing of the paper.

This trial is funded by the Italian Ministry of Health (RF-2010-2318290).

DISCUSSION

The important innovation in this large multicenter RCT is that it potentially provides anesthesiologists with evidence for choosing anesthetics that will lead to the best clinical outcome for their patients. To the best of our knowledge this is the largest RCT ever performed on anesthetic drugs.

The inclusion of a volatile anesthetic for CABG is a simple technique that can be applied to all patients, as anesthesiologists worldwide are trained in both types of anesthesia delivery, and the equipment for both anesthesia techniques are readily available. If the hypothesis is proven correct and mortality is reduced, this simple intervention can save over 2,500 lives each year worldwide and contribute to reduced health care cost.

The design of the study is deliberately pragmatic rather than strictly controlled. By allowing a range of anesthetic drugs and techniques used by participating institutions, the feasibility and external validity is maximized. The inclusion of off-pump CABG is justified by preliminary randomized evidence on the efficacy of volatile agents in this group of patients²⁸⁻³¹, and differences between techniques will be explored in a subgroup analysis.

The tradeoff for a pragmatic design is to have a conservative estimate of mortality and power the study appropriately. Although we are studying an anesthetic intervention, the outcome is highly relevant to cardiologists and cardiac surgeons, as survival after coronary intervention is the primary reason for performing the operation.

LIMITATIONS

A possible limitation, that, in our opinion, is also potentially a strength of our study, is that we decided not to mandate a strict anesthetic protocol, including different opioids, induction agents, cardioplegia fluids, on- or off-pump procedures. This allows all patients to be treated according to the best available practice in each center. It also adds external validity to our findings.

As the cardioprotective effect of volatile agents may also be diminished by the lack of a strict protocol to use volatile agents, we plan to collect data on the dose, length, and timing of administration of various volatile agents, in order to better understand how these variables might influence the potential benefits of volatile drugs.

CONCLUSIONS

The MYRIAD trial will be the first adequately powered RCT comparing the effects of volatile and total intravenous anesthetics on survival after CABG. If the predicted effect is proven, approximately 2,500 lives could be saved each year worldwide.

ACKNOWLEDGEMENT

This trial is funded by the Italian Ministry of Health (Grant: RF 2010-28).

We want to thank Paola Zuppelli and Lara Sussani for their precious collaboration in relation to this RCT.

REFERENCES

- [1] Mozaffarian D., Benjamin E.J., Go AS, et al. Heart Disease and Stroke Statistics—2015 Update A Report From the American Heart Association. *Circulation*. 2015;131:00-00.
- [2] Landoni G., Augoustides J.G., Guarracino F., et al. Mortality reduction in cardiac anesthesia and intensive care: results of the first International Consensus Conference. *HSR Proc Intensive Care Cardiovasc Anesth*. 2011;3(1):9-19.
- [3] Ross S. et al. Protective effects of anaesthetics in reversible and irreversible ischaemia-reperfusion injury. *Br J Anaesth* 1999; 82:622–32.
- [4] Kunst G., Klein A.A.. Peri-operative anaesthetic myocardial preconditioning and protection - cellular mechanisms and clinical relevance in cardiac anaesthesia. *Anaesthesia*. 2015 Apr;70(4):467-82.
- [5] Fortis S, Spieth PM, Lu WY, et al. Effects of anesthetic regimes on inflammatory responses in a rat model of acute lung injury. *Intensive Care Med*. 2012 Sep;38(9):1548-55
- [6] Schilling T, Kozian A, Senturk M, et al. Effects of volatile and intravenous anesthesia on the alveolar and systemic inflammatory response in thoracic surgical patients. *Anesthesiology*. 2011 Jul;115(1):65-74
- [7] Yang Q, Dong H, Deng J, et al. Sevoflurane preconditioning induces neuroprotection through reactive oxygen species-mediated up-regulation of antioxidant enzymes in rats. *Anesth Analg*. 2011 Apr;112(4):931-7
- [8] Beck-Schimmer B, Breitenstein S, Bonvini JM, et al. Protection of pharmacological postconditioning in liver surgery: results of a prospective randomized controlled trial. *Ann Surg*. 2012 Nov;256(5):837-44

- [9] Lee HT, Kim M, Kim J, Kim N, Emala CW. TGF-beta1 release by volatile anesthetics mediates protection against renal proximal tubule cell necrosis. *Am J Nephrol*. 2007;27(4):416-24
- [10] Uhlig C, Bluth T, Schwarz K, et al. Effects of Volatile Anesthetics on Mortality and Postoperative Pulmonary and Other Complications in Patients Undergoing Surgery: A Systematic Review and Meta-analysis. *Anesthesiology*. 2016 Jun;124(6):1230-45
- [11] De Hert S. et al. A comparison of volatile and non-volatile agents for cardioprotection during on-pump coronary surgery. *Anaesthesia*. 2009,(64):953-60.
- [12] Likhvantsev VV, Landoni G, Levikov DI, Grebenchikov OA et al. Sevoflurane versus total intravenous anesthesia for isolated coronary artery bypass surgery with cardiopulmonary bypass: a randomized trial. *J Cardiothorac Vasc Anesth*. 2016 Oct;30(5):1221-7.
- [13] Khuri SF, Henderson WG, DePalma RG, et al. Determinants of long-term survival after major surgery and the adverse effect of postoperative complications. *Ann Surg*. 2005 Sep;242(3):326-41
- [14] Rhodes A, Cecconi M, Hamilton M, et al. Goal-directed therapy in high-risk surgical patients: a 15-year follow-up study. *Intensive Care Med*. 2010 Aug;36(8):1327-32
- [15] Serruys PW, Morice MC, Kappetein AP, et al; SYNTAX Investigators. Percutaneous coronary intervention versus coronary-artery bypass grafting for severe coronary artery disease. *N Engl J Med*. 2009;360(10):961-972
- [16] Park SJ, Ahn JM, Kim YH, et al. Trial of everolimus-eluting stents or bypass surgery for coronary disease. *N Engl J Med*. 2015 Mar 26;372(13):1204-12
- [17] Kapur A, Hall RJ, Malik IS, et al. Randomized comparison of percutaneous coronary intervention with coronary artery bypass grafting in diabetic patients. 1-year results of the CARDia (Coronary Artery Revascularization in Diabetes) trial. *J Am Coll Cardiol*. 2010 Feb 2;55(5):432-40

- [18] Farkouh ME, Domanski M, Sleeper LA, et al. Strategies for multivessel revascularization in patients with diabetes. *N Engl J Med*. 2012 Dec 20;367(25):2375-84
- [19] Shroyer AL, Grover FL, Hattler B, et al. On-pump versus off-pump coronary-artery bypass surgery. *N Engl J Med*. 2009 Nov 5;361(19):1827-37
- [20] Landoni G, Greco T, Biondi-Zoccai G et al. Anaesthetic drugs and survival: a Bayesian network meta-analysis of randomized trials in cardiac surgery. *Br J Anaesth* 111 (6): 886–96 (2013).
- [21] Jakobsen C.J., Berg H., Hindsholm K.B., Faddy N., Sloth E.. The influence of propofol versus Sevoflurane anesthesia on outcome in 10,535 cardiac surgical procedures. *J Cardiothorac Vasc Anesth*. 2007 Oct; 21(5):664-71.
- [22] Andrews D.T., Royse C., Royse A.G.. The mitochondrial permeability transition pore and its role in anaesthesia-triggered cellular protection during ischaemia-reperfusion injury. *Anaesth Intensive Care*. 2012 Jan; 40(1):46-70.
- [23] Muntean D.M. et al. Volatile anaesthetics and cardioprotection - lessons from animal studies. *Fundam Clin Pharmacol*. 2013;27:21-34.
- [24] Lange M. et al. Desflurane-induced preconditioning has a threshold that is lowered by repetitive application and is mediated by beta 2-adrenergic receptors. *J Cardiothorac Vasc Anesth*. 2009; 23:607-13.
- [25] Vierron E. and Giraudeau B. Sample size calculation for multicenter randomized trial: taking the center effect into account. *Contemp Clin Trials* 2007; 28(4):451-8.
- [26] De Mets D.L., Pocock S.J., Julian D.G.. The agonising negative trend in monitoring of clinical trials. *Lancet* 1999; 354:1983-8.

- [27] O'Brien P.C. and Fleming T.R. "A multiple testing procedure for clinical trials." *Biometrics*, 35, pages 549-556.
- [28] Guarracino F et al. Myocardial damage prevented by volatile anesthetics: a multicenter randomized controlled study. *J Cardiothorac Vasc Anesth* 2006; 20(4):477-83.
- [29] Wang J et al. Sevoflurane at 1 MAC provides optimal myocardial protection during off-pump CABG. *Scand Cardiovasc J* 2013; 47(3):175-84.
- [30] Tempe DK et al. Myocardial protection with isoflurane during off-pump coronary artery bypass grafting: a randomized trial. *J Cardiothorac Vasc Anesth* 2011; 25(1):59-65.
- [31] Guerrero Orriach JL et al. Prolonged sevoflurane administration in the off-pump coronary artery bypass graft surgery: beneficial effects. *J Crit Care* 2013; 28(5):879.e13-8.