King's Research Portal

DOI:

[10.1108/IJHCQA-03-2016-0025](https://doi.org/10.1108/IJHCQA-03-2016-0025)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Knight, A., Littlejohns, P., Poole, T-L., Leng, G., & Drummond, C. (2017). The NICE alcohol misuse standard – evaluating its impact. *INTERNATIONAL JOURNAL OF HEALTH CARE QUALITY ASSURANCE*, 30(3), 260 - 273. <https://doi.org/10.1108/IJHCQA-03-2016-0025>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

International Journal of Health Care Quality Assurance

The NICE alcohol misuse standard – evaluating its impact

Alec Knight, Peter Littlejohns, Tara-Lynn Poole, Gillian Leng, Colin Drummond,

Article information:

To cite this document:

Alec Knight, Peter Littlejohns, Tara-Lynn Poole, Gillian Leng, Colin Drummond, (2017) "The NICE alcohol misuse standard – evaluating its impact", International Journal of Health Care Quality Assurance, Vol. 30 Issue: 3, pp.260-273, <https://doi.org/10.1108/IJHCQA-03-2016-0025>

Permanent link to this document:

<https://doi.org/10.1108/IJHCQA-03-2016-0025>

Downloaded on: 10 July 2017, At: 05:28 (PT)

References: this document contains references to 10 other documents.

The fulltext of this document has been downloaded 214 times since 2017*

Users who downloaded this article also downloaded:

(2017),"Editorial", International Journal of Health Care Quality Assurance, Vol. 30 Iss 3 pp. 206-207 https://doi.org/10.1108/IJHCQA-01-2017-0011

(2017),"Embedding learning from adverse incidents: a UK case study", International Journal of Health Care Quality Assurance, Vol. 30 Iss 3 pp. 216-223 https://doi.org/10.1108/IJHCQA-05-2016-0060

Access to this document was granted through an Emerald subscription provided by All users group

For Authors

If you would like to write for this, or any other Emerald publication, then please use our Emerald for Authors service information about how to choose which publication to write for and submission guidelines are available for all. Please visit www.emeraldinsight.com/authors for more information.

About Emerald www.emeraldinsight.com

Emerald is a global publisher linking research and practice to the benefit of society. The company manages a portfolio of more than 290 journals and over 2,350 books and book series volumes, as well as providing an extensive range of online products and additional customer resources and services.

Emerald is both COUNTER 4 and TRANSFER compliant. The organization is a partner of the Committee on Publication Ethics (COPE) and also works with Portico and the LOCKSS initiative for digital archive preservation.

*Related content and download information correct at time of download.

The NICE alcohol misuse standard – evaluating its impact

Alec Knight

Health Services and Population Research Department, King's College London, London, UK

Peter Littlejohns

Department of Primary Care and Public Health Sciences, King's College London, London, UK

Tara-Lynn Poole

University College London Medical School, University College London, London, UK

Gillian Leng

National Institute for Health and Care Excellence, London, UK, and

Colin Drummond

Addictions Department, King's College London, London, UK

Abstract

Purpose – The purpose of this paper is to explore factors affecting implementing the National Institute for Health and Care Excellence (NICE) quality standard on alcohol misuse (QS11) and barriers and facilitators to its implementation.

Design/methodology/approach – Qualitative interview study analysed using directed and conventional content analyses. Participants were 38 individuals with experience of commissioning, delivering or using alcohol healthcare services in Southwark, Lambeth and Lewisham.

Findings – QS11 implementation ranged from no implementation to full implementation across the 13 statements. Implementation quality was also reported to vary widely across different settings. The analyses also uncovered numerous barriers and facilitators to implementing each statement. Overarching barriers to implementation included: inherent differences between specialist vs generalist settings; poor communication between healthcare settings; generic barriers to implementation; and poor governance structures and leadership.

Research limitations/implications – QS11 was created to summarise alcohol-related NICE guidance. The aim was to simplify guidance and enhance local implementation. However, in practice the standard requires complex actions by professionals. There was considerable variation in local alcohol commissioning models, which was associated with variation in implementation. These models warrant further evaluation to identify best practice.

Originality/value – Little evidence exists on the implementing quality standards, as distinct from clinical practice guidelines. The authors present direct evidence on quality standard implementation, identify implementation shortcomings and make recommendations for future research and practice.

Keywords Evidence-based practice, Service quality, Service delivery, Standards, National Health Service, Quality standards

Paper type Research paper

Introduction

Alcohol-related ill-health and harm is high and is increasing in England (National Audit Office, 2008). Alcohol-related harm's total annual cost to English society is estimated at £21 bn, with alcohol misuse costing the NHS £3.5 bn a year (Public Health England, 2014).

In London, alcohol misuse has been identified as a public health priority (London Health Board, 2013). An average borough in South London is likely to face £14 m costs each year in alcohol-related health, £47 m a year in alcohol-related crime and £30 m a year in alcohol-related reduced productivity. Furthermore, research shows 40 per cent of attendances at emergency departments in two South London boroughs were alcohol-related (Drummond *et al.*, 2014) and admissions to acute care account for 14 per cent within King's Health Partners and 50 per cent of adult mental health admissions in South West London (Barnaby *et al.*, 2003). While measures to address alcohol misuse in the NHS and more broadly are well evidenced (Pilling *et al.*, 2011; University of Stirling, 2013), how to implement these measures is poorly applied and understood (Drummond *et al.*, 2011).

In England and Wales, the National Institute for Health and Care Excellence (NICE) produces evidence-based guidance documents and associated quality standards – prioritised statements derived from guidance, designed to drive measurable quality improvements within a health or care area. They are relevant to numerous audiences, including health and social care commissioners, care providers and service users. In August 2011, National Institute for Health and Care Excellence published a quality standard (QS11) focussing on alcohol dependence and harmful alcohol use. The QS11 statements are in five broad categories: identification and assessment in all settings; assessment in specialist services; medically assisted alcohol withdrawal; conditions comorbid with alcohol use; and interventions for alcohol misuse. By ensuring that health and social care services are commissioned, delivered and used in accordance with these five categories, the guidance aims to “contribute to improving the effectiveness, safety and experience of care for harmful drinkers and people with alcohol dependence” (p. 7) and “contribute to reducing alcohol-related hospital admissions and readmissions to hospital” (p. 8). Close implementation and adherence to QS11 should, therefore, contribute to alcohol healthcare service improvements in South East London. The degree to which South East London services accord with the QS11 statements is unknown. To address this knowledge gap, we designed a research study on QS11 adherence in Southwark, Lambeth and Lewisham boroughs. Within this geographical area, our study addressed two primary research questions:

- RQ1. To what extent and to what quality level are QS11 quality statements implemented?
- RQ2. What are the barriers and facilitators to better or fuller QS11 quality statement implementation?

Methods

To investigate QS11 implementation, we designed a qualitative, structured-interview study of individuals with direct experience commissioning, delivering or using alcohol healthcare services. The methodological orientation underpinning the study was content analysis (Hsieh and Shannon, 2005) and our geographical focus was Southwark, Lambeth and Lewisham London boroughs.

Developing the interview guide

The QS11 guidance contains 13 quality statements in five domains (Table I). We developed an interview guide based on these statements, comprising 52 questions (each statement had four probes). We tested the interview guide with eight interviewees (who did not participate in the main study). The interviewer (AK) read each quality statement in turn to the pilot study interviewees and then asked four questions. Two probes elicited information about current implementation, two probes about implementation barriers and facilitators.

Table I.
NICE quality
standard 11 – quality
statements

Identification and assessment in all settings

- 1 Health and social care staff receive alcohol awareness training that promotes respectful, non-judgmental care of people who misuse alcohol
- 2 Health and social care staff opportunistically carry out screening and brief interventions for hazardous and harmful drinking as an integral part of practice
- 3 People who may benefit from specialist assessment or treatment for alcohol misuse are offered referral to specialist alcohol services and can access specialist alcohol treatment

Assessment in specialist services

- 4 People accessing specialist alcohol services receive assessments and interventions delivered by appropriately trained and competent specialist staff
- 5 Adults accessing specialist alcohol services for alcohol misuse receive a comprehensive assessment that includes the use of validated measures
- 6 Children and young people accessing specialist services for alcohol use receive a comprehensive assessment that includes the use of validated measures
- 7 Families and carers of people who misuse alcohol have their own needs identified, including those associated with risk of harm, and are offered information and support

Medically assisted alcohol withdrawal

- 8 People needing medically assisted alcohol withdrawal are offered treatment within the setting most appropriate to their age, the severity of alcohol dependence, their social support and the presence of any physical or psychiatric comorbidities
- 9 People needing medically assisted alcohol withdrawal receive medication using drug regimens appropriate to the setting in which the withdrawal is managed in accordance with NICE guidance

Conditions comorbid with alcohol misuse

- 10 People with suspected, or at high risk of developing, Wernicke's encephalopathy are offered thiamine in accordance with NICE guidance

Interventions for alcohol misuse

- 11 Adults who misuse alcohol are offered evidence-based psychological interventions, and those with alcohol dependence that is moderate or severe can in addition access relapse prevention medication in accordance with NICE guidance
- 12 Children and young people accessing specialist services for alcohol use are offered individual cognitive behavioural therapy, or if they have significant comorbidities or limited social support, a multicomponent programme of care including family or systems therapy
- 13 People receiving specialist treatment for alcohol misuse have regular treatment outcome reviews, which are used to plan subsequent care

Our pilot interviewees provided feedback on probe wording, which we used to update the interview guide before conducting interviews. The pilot interviewees also commented that most participants would not have direct experience in relation to all 13 statements. Consequently, we decided to ask each interviewee to specify which statements they felt able to discuss from direct personal experience. We then focussed our interview questions specifically on those statements. Therefore, the quality statements addressed in each interview differed depending on interviewee experience.

Selecting and recruiting the sample

Our sampling frame was organisations in Southwark, Lambeth and Lewisham employing staff involved in commissioning or delivering healthcare services for hazardous or harmful drinkers. This included local authorities, NHS-funded and third sector organisations involved in primary, secondary or specialty care. Within this sampling frame, we used a snowball sampling method to identify potential participants and a maximum-variation purposive sampling method to select participants according to numerous criteria such as geographical location, professional background, experience and seniority. We used the snowball method to

identify participants, as the population we were sampling was not well delineated. We also had limited personal connections to people in the population. We used the purposive sampling method to select participants as we wanted to control who participated in the research so that we could represent broad perspectives. After each pilot and data collection interview, we asked our interviewees to provide names and contact details for individuals with significant experience relevant to our study (i.e. participant identification). Following discussion among the research team, individuals were purposively selected based on their personal characteristics in comparison with interviewee characteristics (i.e. participant selection). Where possible, we asked the recommending interviewee to provide a personal e-mail introduction to selected participants. In all other cases, we approached selected participants directly by e-mail, explaining that a previous interviewee had recommended him or her. Geographical, healthcare sector and the interviewees' professional characteristics are presented in Table II. Our sample comprised 12 men and 26 women.

Provisional data analysis started after completing the first interview. We stopped recruiting participants when we reached data saturation (the point at which no new data emerged) both in interview themes and names generated through participant identification. The sampling approach led to a high response rate, with 45 identified and 38 participating in the interview study (84 per cent response rate). Reasons for non-participation were non-specific declines (3), non-response (2) and sickness-absence during data collection (2). Where selected participants were not able to participate, we recruited substitutes with similar characteristics.

Setting and data collection

Interviews were conducted between March and July 2014, either in a private room in the interviewee's workplace if available, or in the interviewer's university office. All interviews were conducted individually and face-to-face. Participation was voluntary and informed

	Interviewees
<i>Borough</i>	
Southwark	11
Southwark and Lambeth	11
Lewisham	9
Lambeth	7
<i>Sector</i>	
NHS/tertiary healthcare	12
Local authorities and commissioning	11
Third sector healthcare	5
NHS/primary healthcare	5
NHS/secondary healthcare	3
Service user groups	2
<i>Professional group</i>	
Medical doctors	8
Nurses	8
Public health professionals	7
Tertiary care professionals	7
Commissioners	5
Service user representatives	2
Clinical psychologist	1

Notes: Some interviewees held positions in multiple sectors, particularly NHS primary, secondary and tertiary healthcare. The figures recorded here refer to an interviewee's primary role, or his/her role at interview

Table II.
Sample – geographical,
healthcare sector and
professional
characteristics

consent was obtained from interviewees. Interviews lasted 22-82 minutes (mean 52 minutes), were recorded and transcribed verbatim by a professional transcriber. The interviewer was an academic psychologist with no clinical or managerial role, who used the interview guide to structure the interview.

Data analysis

Two data judges (AK and T-LP) qualitatively analysed the interviews using conventional and directed content analysis methods. The framework for the directed content analysis was a 13×4 matrix, reflecting QS11's 13 quality statements (rows) and the four probe questions about implementation quality and quantity, barriers and facilitators (columns). There was also space for general comments about all 13 statements that did not fit into any column. This analysis provided information that allowed us to answer *RQ1* and *RQ2*. Judges independently read through each interview transcript and extracted data from the interview onto a blank 13×4 data extraction table. Judges completed data extraction tables and met regularly to compare analyses. Discrepancies were discussed and resolved. Judges met periodically with the data auditor (PL) to discuss the analysis. The data auditor's role was to check the analysis process for rigor and to make technical inputs (e.g. to suggest alternative labels or themes). Conventional content analysis was also conducted on interview data when data extraction tables were completed. This analysis provided supplementary information to allow us to provide a rich answer to *RQ2*. Judges independently noted the overarching themes (across quality statements and interviewees). During their meetings, judges compared emerging overarching themes. Discrepancies were discussed and resolved by agreeing on a composite themes list endorsed by both judges. Amending the list was repeated until no new themes emerged from the data and until no further changes needed to be made to accommodate both judges' suggestions (i.e. saturation). The data auditor checked the composite themes lists and made suggestions for alternative categories and labels.

Results

Directed content analysis

Table III synthesises the most common responses about implementation (quality and extent), barriers and facilitators for each quality statement. There were large differences in implementation across the statements. Implementation ranged from nil (statement 6) through partial or patchy (statement 1), to full (statement 4). For some statements, there was also disagreement between interviewees about how widely those practices were implemented (e.g. statement 8). Differences within and between statements were also present in the implementation quality reports; e.g., wide variations in average practice quality were reported in relation to some statements (e.g. statement 2), while average practice quality in other areas was reported to be high (statement 5).

Conventional content analysis

In addition to the statement-level analysis, our thematic-level analysis uncovered seven themes: implementation variability across and within borough, sector and care setting; specialist vs generalist settings; communication between healthcare professionals in different settings; generic barriers to implementation; children and young people (CYP); governance structures and leadership; and service users' views. These themes are described below.

Implementation variability across and within borough, sector, care setting. A striking theme arising from our interviews was the variability between three contiguous boroughs in relation to the quality and extent that the QS11 quality statements were implemented. For example, interviewees reported particularly high opportunistic screening levels for hazardous and harmful drinking in one borough owing to financial incentives unique to that locality (quality statement 2).

Statement 1 – health and social care staff receive alcohol awareness training that promotes respectful, non-judgmental care of people who misuse alcohol

Quality of implementation

Differences in perceived quality of training between different types of organisations (e.g. primary care vs third sector)

Training sometimes focussed on the effects of alcohol and where to find information and services rather than promoting respectful and non-judgemental care

Online training viewed as inadequate compared to face-to-face training

Services do not always report on how training is used, meaning information may not be available on quality of implementation

Training gap surrounding binge drinking in young people

Extent of implementation

Training coverage reported as very patchy within and across some boroughs, professional groups, and organisations

Some groups reported to have low coverage of training (e.g. nurses, health visitors, drug workers), while others had high coverage (e.g. medical doctors, adult services, local authority employees)

Barriers to implementation

Absence of training providers

Resources and prioritising

Time and workloads

Inability to access training provision

Emphasis on other substances for young people

Poor governance of third sector organisations

Facilitators to implementation

Professional development plans

Financial incentive plan (e.g. CQUIN)

Prioritising training (making alcohol a top priority)

Making training a strategic priority for CCGs

Statement 2 – health and social care staff opportunistically carry out screening and brief interventions for hazardous and harmful drinking as an integral part of practice

Quality of implementation

Financial incentive may encourage widespread screening but may also lead to a tendency to screen patients without providing necessary advice or referral. Payments also only received for new registrants, and there was no payment for intervention after screening

Mixed evidence that services use appropriate screening tools

Good integration of screening into existing systems and audits reported in some localities

Extent of implementation

Reports of outreach programmes being conducted outside of healthcare settings

Routine practice in specialist services; good implementation in secondary care and in other settings with a specialist healthcare professional

Poor implementation in young people's services and when GPs see young people

Barriers to implementation

Capacity and workload

Lack of accountability and consistency or recording information across settings

Services for CYP focus on other substances. Interventions in schools may miss those in need. CYP may also not be honest and there are difficulties involving parents

Facilitators to implementation

Use of the same screening tool across services

Easy to use apps to facilitate interventions

Writing screening/brief interventions into trust policy

Financial incentives

Passionate individual members of staff

Public health awareness campaigns to raise staff awareness

Experience and history of delivering interventions

IJHCQA
30,3

266

Statement 3 – people who may benefit from specialist assessment or treatment for alcohol misuse are offered referral to specialist alcohol services and can access specialist alcohol treatment

Quality of implementation

- Reports of inappropriate referrals to A and E from GPs
- Appropriate referrals are better when a specialist is available
- Absence of a clear pathway for referrals for CYP

Extent of implementation

- Patchy implementation, better in areas with specialists in primary care and areas with financial incentives
- Poor/zero implementation for CYP. CAMHS is the default service for CYP

Barriers to implementation

- Poor information about and communication between services, lack of confidence referral is being made to the correct place
- Services being housed in unattractive buildings and areas
- Service users reluctant to access services in mixed drug and alcohol service facilities due to stigma
- Lack of a named person in services for clients to contact
- Lack of understanding and confidence in how to tackle alcohol, or not knowing where to refer to
- Demand outstripping supply and waiting lists being long
- Regular re-commissioning of services
- Inappropriate referral due to staff wanting to offload work they consider outside of their remit

Facilitators to implementation

- Financial incentives (e.g. CQUIN)
- Specialist staff (e.g. mental health nurse) to make better assessments and improve referrals

Statement 4 – people accessing specialist alcohol services receive assessments and interventions delivered by appropriately trained and competent specialist staff

Quality of implementation

- Very high compliance in NHS settings
- Third sector provision reported to be compliant after being audited by commissioners
- Parity in quality between NHS and third sector provision is unknown

Extent of implementation

- Full implementation in NHS settings
- Key staff reported to be appropriately trained in third sector services

Barriers to implementation

- None reported

Facilitators to implementation

- Professional development structures
- Writing training into contracts
- Commissioners asking for evidence of appropriate training

Statement 5 – adults accessing specialist alcohol services for alcohol misuse receive a comprehensive assessment that includes the use of validated measures

Quality of implementation

- Common measures used across NHS settings ensures measures are valid
- Validity of measures in third service provision is achieved through random checks of care plans by commissioners

Extent of implementation

- High level of compliance as staff are trained in drug and alcohol treatment

Barriers to implementation

- Some providers may be resistant to showing commissioners their case files during audits

Facilitators to implementation

- Several different screening tools are available
- Formal processes are established to use the tools
- Measuring outcomes facilitated a change in service quality
- 360-degree appraisals
- Strong management can create an atmosphere of professionalism

Table III.

(continued)

Statement 6 – children and young people accessing specialist services for alcohol use receive a comprehensive assessment that includes the use of validated measures

Quality of implementation

Zero (no specialist services exist)

Extent of implementation

Zero (no specialist services exist)

Barriers to implementation

CYP services are highly structured and rigid, particularly in relation to using validated measures

Forms often incomplete as CYP can be unreliable to return to services

Integration and communication between services (e.g. social workers, youth offending service)

Services have lost their impetus (e.g. National Treatment Agency for Substance Misuse)

Referrals threshold into CAMHS is so high that they always decline referrals from GPs

CAMHS not seeing itself as part of the mainstream healthcare system

Different measures used in different services

Facilitators to implementation

Compliance is high in Youth Offending Services as attendance is not optional

Putting the focus on the key worker to fill out assessments

Statement 7 – families and carers of people who misuse alcohol have their own needs identified, including those associated with risk of harm, and are offered information and support

Quality of implementation

Needs are often acknowledged but may not be directed towards formal use of the service

Where services for families are used, there are reports that they are very successful

Although pathways for families exist they are rarely used

Reactive approach is taken to families; children are supported through safeguarding

Extent of implementation

Services range from very difficult to access to fairly easy to access. However, actual use of the services may be limited due to limited capacity

A&E provide information about support for family and carers

Safeguarding covers support for CYP, but this relies on clinicians contacting other services

Barriers to implementation

People who misuse alcohol often have normal lives and jobs, making it difficult to recognise when they are developing a problem

Carers may be appearing to cope and not see the need for support

The method used to identify carers' needs is laborious and intrusive

Confidentiality surrounding the alcohol user. This makes it difficult to record data and difficult to audit implementation

Training for supporting families and carers takes a long time

Lack of resources

Family members may have problems being treated in the same place as the service user

Facilitators to implementation

Writing into organisational policy the provision of information to carers

Statement 8 – people needing medically assisted alcohol withdrawal are offered treatment within the setting most appropriate to their age, the severity of alcohol dependence, their social support and the presence of any physical or psychiatric comorbidities

Quality of implementation

Referrals to appropriate settings are improved by a community nurse facilitating the process

Short in-patient detoxes may not be long enough and may cause patients to relapse as a result

Psychiatric in-patients receive good treatment as they are treated as mental health in-patients for detox

Extent of implementation

Gaps in services in some areas; moderate-to-good coverage in other areas

The appropriateness of the setting may depend on capacity

Differential success between in-patient and community detoxing

Where a specialist nurse makes the assessment implementation coverage is good

(continued)

Table III.

IJHCQA
30,3

268

Barriers to implementation

- Poor communication and lack of integration between services
- Division of services causes confusion as to where to send patients
- Disagreement over who pays for services
- Poor visibility of community services in some localities
- Lack of clarity over who is responsible to deliver services
- Stigma may prevent patients from accessing certain services

Facilitators to implementation

- Presence of a community nurse
- Having a long-term relationship with clients

Statement 9 – people needing medically assisted alcohol withdrawal receive medication using drug regimens appropriate to the setting in which the withdrawal is managed in accordance with NICE guidance

Quality of implementation

- Only one drug available in some localities
- Reports that GPs prescribe detox drugs poorly
- Regimens are appropriate when prescribed from a specialist service

Extent of implementation

- Generally, very well run in localities where services are run by NHS
- Variable implementation in locality run by third sector providers as access to appropriate settings can be problematic
- Good implementation in in-patient setting

Barriers to implementation

- GPs not offering detox because they do not feel confident to do so
- Lack of training for GPs
- Cost of drugs on GPs drugs budgets may mean they prefer for specialist services to offer the treatments

Facilitators to implementation

- Change in triage standards gave managers the opportunity to make improvements in NHS services

Statement 10 – people with suspected, or at high risk of developing, Wernicke's encephalopathy are offered thiamine in accordance with NICE guidance

Quality of implementation

- Generally thought to be good, but worse among some professional groups such as social work

Extent of implementation

- In NHS specialist services, it was reported that there was full implementation and it was very unlikely that patients would be missed
- Reports of good implementation in treatment services and poorer implementation in the community

Barriers to implementation

- Lack of awareness of the availability of thiamine
- Mixed views about the effectiveness of thiamine
- Confusion about whose responsibility it is to offer thiamine

Facilitators to implementation

- None reported

Statement 11 – adults who misuse alcohol are offered evidence-based psychological interventions, and those with alcohol dependence that is moderate or severe can in addition access relapse prevention medication in accordance with NICE guidance

Quality of implementation

- Reports of different practice from NICE recommendations
- IAPT input may help but it is not specific to alcohol
- Waiting lists reported in some localities

Extent of implementation

- Large gaps exist in provision, with some reports of zero provision in some localities
- Some services offered in some localities; other services may not be offered at all
- Relapse prevention medication is offered very sparingly
- Reports of good implementation where dual-diagnosis staff are present
- Psychological services are rare within addictions services

Table III.

(continued)

Barriers to implementation

- Language barriers mean some communities do not understand what services do or what is available
- Lack of trust among immigrant communities
- Waiting lists for psychological services
- Training skills are poor in addiction services and training is out-dated
- Shift of emphasis from secondary to primary care resulted in lack of specialist knowledge
- GP appointment time slots too short to explore mental health needs properly

Facilitators to implementation

- Multilingual healthcare staff
- Training all staff in relapse prevention
- Signposting in A and E

Statement 12 – children and young people accessing specialist services for alcohol use are offered individual cognitive behavioural therapy, or if they have significant comorbidities or limited social support, a multicomponent care programme including family or systems therapy

Quality of implementation

- Zero (no specialist services exist)
- Children may be referred to the third sector, where training can be poor

Extent of implementation

- Zero (no specialist services exist)
- Low confidence among interviewees that this is happening locally

Barriers to implementation

- Family and systems therapy viewed as only useful for a small number of young people
- CYP's confidentiality
- Difficult of outreach
- CYP do not want to talk about alcohol and drug use in schools due to zero tolerance policies
- Parents want to feel they are doing a good job and ignore signs of alcohol misuse

Facilitators to implementation

- None reported

Statement 13 – people receiving specialist treatment for alcohol misuse have regular treatment outcome reviews, which are used to plan subsequent care

Quality of implementation

- Care plans may be recorded as paper or electronic notes, which can make it hard to use Treatment Outcomes Profile (TOPs) forms
- High level of compliance may mask the fact that TOPs forms may not be used properly

Extent of implementation

- Good-to-full implementation in NHS settings
- High level of compliance in specialist services, but a lack of follow-up when patients leave A and E

Barriers to implementation

- TOPs is time-consuming to complete
- Lack of acknowledgement of the need for treatment outcome reviews in non-specialist settings
- Looking at a patient's combined mental and physical health needs is not commonplace in A and E unless a staff member there is particularly passionate about this view

Facilitators to implementation

- Performance management
- Financial incentives

Table III.

Specialist healthcare services provided for alcohol misuse are determined by commissioners at borough level, which resulted in dramatically different service delivery models between boroughs. Alcohol services are delivered by a single third sector organisation in Lewisham, while in Southwark and Lambeth, they are delivered by a collaborative between numerous NHS and third sector organisations. Accordingly, our interviewees highlighted significant differences between the boroughs in adherence to quality statements about specialist treatment. At sector level (i.e. NHS-funded vs third sector organisations), it is difficult to draw

conclusions about implementation differences, as few interviewees had direct experience providing or using services in both. However, at the setting level (i.e. primary, secondary and specialty care) our interviewees often described significant differences (e.g. across different GP surgeries or accident and emergency (A&E) departments). Often these differences in implementation were attributed to specialist staff (e.g. a dual-diagnosis nurse or a consultant psychiatrist with an interest in alcohol misuse).

Specialist vs generalist settings. Another key feature in the responses was a perceived wide gap in quality and extent that QS11 was implemented in specialist settings (e.g. community drug and alcohol teams) vs generalist settings (e.g. GP surgeries and A&E departments). Almost universally, specialist settings were described as having closer and better adherence to QS11, while implementing the quality statements was more variable, patchy or lower quality in generalist settings.

Communication between healthcare professionals in different settings. Poor integration and communication between staff different healthcare settings stood out as significant barriers to implementing QS11. Interviewees described this as resulting from regular service re-commissioning. For example, interviewees described inappropriate referrals to A&E departments from GPs, while GPs complained about a lack of confidence in referring patients (quality statement 3). Lewisham's new integrated service made a concerted effort to ensure a clear alcohol pathway, with training including what services are available and how to signpost patients to services. Nevertheless, our GP interviewees gave mixed feedback on how useful these are in assisting with referrals (quality statement 3). Where interviewees reported good compliance with NHS treatment outcome reviews, they also reported poor follow-up from A&E department staff (quality statement 13). Weak integration and communication between service staff was also reported to have affected the extent to which people received medically assisted alcohol withdrawal in an appropriate setting (quality statement 8). Service users and their carers reported being confused and unclear about treatment pathways. Community or dual-diagnosis nurses were said to improve service communication and integration.

Generic barriers to implementation. Several generic barriers and facilitators to implementation were described by many interviewees. Typical implementation barriers across boroughs, sectors and settings included insufficient time and problems with accessing specialist services or support at certain times, work pressures and stress, and inadequate non-financial resources such as staffing levels. Across many QS11 quality statements, facilitators to better or fuller implementation included financial incentives, performance managing staff for a specific behaviour or outcomes, and obliging staff to undertake an activity as standard operating procedure (e.g. required paperwork).

CYP. The picture described to us by our interviewees was complex regarding CYP alcohol misuse services. Most interviewees had no experience dealing with CYP who misuse alcohol, their families or carers. The experienced few reported that key quality statements had not been implemented. One interviewee suggested that might be due to few CYP presenting with alcohol misuse problems. Nonetheless, there are no clear pathways for referrals within the healthcare system and professionals were unsure what screening tools are appropriate for CYP. Interviewees expressed concern that owing to the focus in youth services on illegal substances, alcohol was socially acceptable and is normalised by youth workers. Interviewees reported that alcohol misuse problems might most often be recognised once a young person was referred to Youth Offending Services (YOS).

The mechanisms for dealing with CYP substance misuse do not fall under the NHS umbrella, except for CAMHS. Although CAMHS is the CYP default service, interviewees reported that CAMHS staff do not see themselves as part of the mainstream healthcare system, which affects integration and communication between alcohol misuse service staff

(quality statement 6). Interviewees described the threshold for referral to CAMHS from GPs as so high that referrals are often declined. Interviewees also noted that most children's services were in the third sector, where worker training was less formalised. Consequently, interviewees expressed concern about a training gap around CYP binge drinking (quality statement 1), in stark contrast to YOS, where CYP receive comprehensive assessment and care as they are obligated to attend and undertake rehabilitation (quality statement 6).

Interviewees expressed concern that CYP needs as carers' concerns were not recognised, which acts as a barrier to them being offered appropriate support. Schools were where most CYP needs were identified. However, stigma and secrecy attached to alcohol misuse, plus loyalty towards family members, may make it unlikely that problems will be identified. The harm risk identified through safeguarding young carers' emotional needs may not be identified (quality statement 7). Furthermore, referral to safeguarding services relies on clinicians going out of their way to contact external service staff.

There was recognition among interviewees, who spoke about CYP issues regarding supporting family members and carers. Most services incorporate carer need assessments and services to support families, including having expert family therapists, family and carer groups, and behavioural therapy for couples. Where services were used, they were often reported as being useful or successful. However, interviewees told us how health and social care culture is still focussed on the service user. Thus, some interviewees reported difficulty for families and carers accessing services because triage for family members and carers is mostly carried out via having an existing service user as a patient. Consequently, interventions with family members and carers were described as akin to using a sledgehammer to crack a nut, e.g., in South London and Maudsley NHS Foundation Trust (SLAM), an electronic patient-journey record is opened for families and carers, where information about the intervention may not require the same detail as for a service user. Confidentiality was also considered to be a barrier to families and carers accessing services. Interviewees described how service users do not always want families to know about their situation and family members concerned for loved ones find that health and social care workers cannot discuss patients. Finally, some interviewees were concerned that family members and carers may not want to receive treatment or support in the same physical location as service users (quality statement 7).

Governance structures and leadership. Commissioners reported a correlation between service improvement and proactive governance. For example, identification and brief advice training is a strategic intervention being rolled out across all boroughs. Interviewees reported that there was good follow-up about implementing training through auditing and monitoring trainers' work. Consequently, interviewees reported that one facilitator to effective training is through Clinical Commissioning Group (CCG) staff making it a strategic priority (quality statement 1). Another initiative introduced in one borough was a financial incentive for GPs to conduct opportunistic screening. Implementation was audited through client satisfaction surveys and staff were asked to evaluate the brief intervention process through feedback questionnaires. Our interviewees reported that the scheme resulted in increased use and improved opportunistic screenings, resulting ultimately in quality statement 2 being implemented better.

Service users' views. Service users and carers reported difficulty accessing services because an individual service user may not want to associate himself or herself with other service users. This is particularly the case where drug and alcohol services are mixed, housed in unattractive buildings and areas (quality statement 3). Some interviewees reported concerns around whether family members and carers were happy to receive support in the same location as the service user (quality statement 7). Stigma around services was reported to prevent patients from accessing certain services for medically assisted withdrawal (quality statement 8).

Discussion

Our directed content analysis provided information about how widely and how well 13 quality QS11 statements were implemented. Implementation quality and extent varied widely between statements, and for some there were mixed reports about how widely and how well recommended practices were adhered to. For each statement, we asked about implementation barriers and facilitators, which allowed us to explore possible mechanisms or explanations for these findings. For example, interviewees believed that some differences in implementation quality and extent were attributable to inherent differences between healthcare settings (i.e. primary, secondary and specialist care), different delivery systems for specialist care in different geographical locations (i.e. NHS vs third sector provision) and the key leaders and staff presence or absence promoting healthcare services for people who misuse alcohol. Significantly, the QS11 sections relating to CYP and their families or carers (statements 6 and 12) were noted as being particularly poorly implemented – attributed to a discrepancy between how QS11 prescribes services for CYP to be delivered, and how services were commissioned and delivered locally. Relatively few interviewees spoke at length about CYP services and those who did, mentioned poor focus on alcohol misuse among CYP compared to older adults or other substance abuse (e.g. cannabis and other illegal substances).

Directed content analysis findings were supplemented by conventional content analysis, which delineated overarching barriers and facilitators to QS11 implementation. Generic barriers to implementation such as insufficient time and resources were often cited by our interviewees. Variability in service quality provided in different locations, sectors and settings were highlighted, with closest adherence to the guidelines being reported in specialist settings for treating people who misuse alcohol, and less adherence in more generalist settings. Poor or patchy QS11 implementation was often attributed to weak communication between healthcare professionals in different sectors or settings. Conversely, good and widespread QS11 implementation, and relatively good communication between healthcare professionals in different sectors and settings was often attributed to good governance, and strong and focussed leadership at various levels within the care pathway.

Our study has two major strengths. First, it provides a multi-professional perspective on QS11 implementation, considering insights from various stakeholders including commissioners, healthcare professionals and service users. Second, independent researchers conducted the interviews. Responses were likely to be more frank than if they had been conducted by professionals working in the area. Similarly, disinterested researchers conducted the analysis. They were not motivated to report positive or negative findings in any direction. Two limitations, however, may restrict how far our findings are generalised. First, our study was based on interviews with 38 key informants in three South East London boroughs. Therefore, our work requires replication. Second, our sampling strategy was designed to gain insights into 13 quality statements. Nevertheless, we obtained less data in relation to certain quality statements (i.e. CYP) than we did in relation to others (e.g. identification and assessment in all settings). Therefore, some analyses were based on less data from fewer sources than were other analyses. We did not report interviewee suppositions about differences between boroughs, sectors or settings that were not supported by direct experience or that were objectively verifiable (e.g. differences between boroughs in commissioning structure or service provision).

The NICE Standards summarise numerous individual NICE guidance. The aim was to simplify the guidance and enhance local implementation. However, in practice, each standard (containing several statements) requires complex actions by numerous professionals and institutions. It could be argued that standards are aimed at those responsible for commissioning the whole service (or care pathway), while the individual guidance documents are geared towards providers. We identified considerable variation in local alcohol commissioning models associated with varying implementation. The differing models

warrant further evaluation to identify which support best practice. It is likely that to improve NICE standards implementation, CCGs, health and well-being boards may need to refer to standards explicitly in their alcohol strategies. Further research is needed to assess whether standards provide added value to the individual guidance documents. It appears that services for young people are an area for further work to understand the most effective service models.

Acknowledgement

This study was supported by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care, South London, at King's College Hospital NHS Foundation Trust. Peter Littlejohns was the founding Clinical and Public Health Director of the National Institute of Health and Care Excellence (NICE) from 1999 to 2011. In this role he designed the process and methods for the development of NICE guidelines and was the executive director responsible for the Citizens Council and the R&D programme. Gillian Leng is an Executive Director at NICE with responsibility for implementing NICE guidance and quality standards. Colin Drummond was Chair of the NICE Standards Committee. The views expressed are the authors' and not necessarily the NHS, NIHR or Department of Health.

References

- Barnaby, B., Drummond, C., McCloud, A., Burns, T. and Omu, N. (2003), "Substance misuse in psychiatric inpatients: comparison of a screening questionnaire survey with case notes", *British Medical Journal*, Vol. 327, pp. 783-784.
- Drummond, C., Deluca, P., Coulton, S., Bland, M., Cassidy, P., Crawford, M., Dale, V., Gilvarry, E., Godfrey, C. and Heather, N. (2014), "The effectiveness of alcohol screening and brief intervention in emergency departments: a multicentre pragmatic cluster randomized controlled trial", *PLoS One*, Vol. 9 No. 6.
- Drummond, C., Gual, A., Goos, C., Godfrey, C., Deluca, P., Von Der Goltz, C., Gmel, G., Scafato, E., Wolstenholme, A. and Mann, K. (2011), "Identifying the gap between need and intervention for alcohol use disorders in Europe", *Addiction*, Vol. 106 No. S1, pp. 31-36.
- Hsieh, H.-F. and Shannon, S.E. (2005), "Three approaches to qualitative content analysis", *Qualitative Health Research*, Vol. 15 No. 9, pp. 1277-1288.
- London Health Board (2013), "Health in London – evidence paper", London Health Board, London.
- National Audit Office (2008), "Reducing alcohol harm: health services in England for alcohol misuse", The Stationery Office, London.
- National Institute for Health and Care Excellence (2011), "Alcohol-use disorders: diagnosis and management", available at: guidance.nice.org.uk/qs11 (accessed 13 July 2016).
- Pilling, S., Whittington, C., Taylor, C. and Kendrick, T. (2011), "Identification and care pathways for common mental health disorders: summary of NICE guidance", *British Medical Journal*, Vol. 342, pp. 1203-1209.
- Public Health England (2014), "Alcohol treatment in England 2013-14", available at: nta.nhs.uk/uploads/adult-alcohol-statistics-2013-14-commentary.pdf (accessed 13 July 2016).
- University of Stirling (2013), "Health first: an evidence-based alcohol strategy for the UK", available at: www.stir.ac.uk/media/schools/management/documents/Alcoholstrategy-updated.pdf (accessed 13 July 2016).

Corresponding author

Alec Knight can be contacted at: alec.knight@kcl.ac.uk

For instructions on how to order reprints of this article, please visit our website:

www.emeraldgrouppublishing.com/licensing/reprints.htm

Or contact us for further details: permissions@emeraldinsight.com